

# **Association between Serum Alfa Fetoprotein and Placenta Previa -Benghazi Medical Center**

## **Abstract :**

The independent risk factor for placenta Previa is a previous caesarean section. The risk increases with the number of caesarean sections performed. Placenta Previa (P.P) occurs when the placenta is wholly or partially implanted in the lower uterine segment. The exact cause of P.P is not known, but its association with various risk factors such as advanced maternal age, multiparity, previous miscarriages, previous caesarean section, and cigarette smoking is well documented. It is concluded that placenta Previa is associated with advanced maternal age, history of previous abortion, multiparity, history of previous antepartum & postpartum hemorrhage, preterm deliveries and previous history of cesarean section.

**Keywords:** Placenta Previa, maternal age, abortion, cesarean section

## **INTRODUCTION:**

Placenta Previa (P.P) occurs when the placenta is wholly or partially implanted in the lower uterine segment. The exact cause of P.P is not known, but its association with various risk factors such as advanced maternal age, multiparity, previous miscarriages, previous caesarean section, and cigarette smoking is well documented. Traditionally, PP is classified as "complete " When the placenta completely covers

the internal cervical os, "partial " when the placenta partially covers the os, "marginal " when the lower edge of the placenta just reaches the os , and "low – lying" when the placenta is in the lower segment but does not reach the internal os. Complete and partial PP are considered "major placenta Previa", while marginal PP and low lying placenta are considered "minor placenta Previa ". In recent years, ultrasound scanning has led to more accurate localization of the placenta. Nonetheless, the relationship between the different types of PP, associated risk factors, and pregnancy outcome is poorly characterized. Although the clinical course of placental Previa is highly suggestive, the etiology of this condition still retracts obscure. The strongest connection was found between previous history of cesarean section, high parity, and advanced maternal age, but the strength of the connection varies from study to study.<sup>1</sup> Antepartum hemorrhage forms one of the most dangerous and devastating group of disorders in obstetrics. Placenta Previa occurs in approximately 1 in 300 deliveries. At least 5% of such pregnancies have associated placental invasion (placenta accreta), which can necessitate hysterectomy. The independent risk factor for placenta Previa is a previous caesarean section. The risk increases with the number of caesarean sections performed. The incidence of placenta accreta is 3% in women with placenta Previa and one previous cesarean section. The risk continues to increase with each additional cesarean section up to 67% in women with a placenta Previa and five or more cesarean sections.<sup>2</sup>

**Associations' Fetal abnormality:** The Fate of fetal abnormality is approximately doubled in women with placenta Previa. Intra uterine growth restriction: Is common in

women with multiple bleeds from a placenta Previa. The overall rate is 15%. Maternal serum alpha - fetoprotein are of great association with P.P specifically in Women With unexplained elevated its screening level. Ten percent of women with a bleeding P.P will have a co-existent abruption.<sup>1</sup>

**Diagnosis:** The possibility of placenta Previa should not be dismissed until sonographic evaluation has clearly proved its absence. The simplest, safest, and most accurate method of placental localization is provided by trans abdominal sonography.

Alpha-fetoprotein (AFP) is a serum glycoprotein that was discovered early in human fetal serum by Bergstrand and Czar in 1956n. It is the chief mammalian tumor associated fetal protein found in adult's blood (by a small amount of 10-20 ng/ml).

Maternal serum alpha fetoprotein is elevated in some pregnancy complications such as spontaneous abortion, pre-eclampsia, gestational hypertension, preterm delivery and premature rupture of membranes (PROM). Also, poor maternal and fetal outcome is strongly related to the elevation of maternal AFP (probably as a result of placental injury). In addition, a significant association was found between increasing maternal AFP and the greater likelihood of persistent placenta Previa. Furthermore, the risk for abnormal placental adherence was increased in women with an elevated maternal serum AFP level, especially in the presence of a placenta previa.<sup>3</sup>

## **AIMS OF THE STUDY:**

To evaluate the relationship between maternal serum alpha-fetoprotein and the risk of persistent placenta Previa and outcome

## REVIEW OF LITERATURE:

Study in India by Pooja Verma et al, they found that the level of Maternal Serum Alfa Fetoprotein (MSAFP ) was higher in 14 out of 15 cases (93.3%) of placenta Previa with placental adherence. There was significant surgical intervention (80%) and increased maternal morbidity (68.8%) in the study group with placental adherence and raised MSAFP respectively. They concluded that the MSAFP is an important biomarker for prognostication of placental adherence in low lying placenta.<sup>2</sup>

Study by [E L Koster](#) et al included 275 women with Previa at 15-20 weeks' gestation, 33 (12%) had Previa at delivery. Trend analysis revealed a greater likelihood of persistent Previa with increasing MSAFP values ( $p=0.01$ ). Mid-trimester MSAFP  $<1$  multiple of the median was associated with a decreased incidence of persistence of 4%, significantly less than the risk at  $> \text{ or } =1$  multiple of the median (16%;  $p=0.01$ ). They concluded that there is an association between

increasing MSAFP values and greater likelihood of persistent placenta Previa. An MSAFP value  $<1$  multiple of the median is associated with a reduction in the risk of persistence of Previa to delivery.<sup>4</sup>

Study by Hümeysra Öztürk et al was found a significant relationship between adverse pregnancy outcomes and abnormal elevation of Human Chorionic Gonadotropin (HCG) and AFP levels in the second trimester. In cases of isolated elevation of HCG, preeclampsia and preterm labor/spontaneous preterm birth rate were slightly higher than in the control group ( $p=0.043$ ,  $p=0.015$ ), while Intra Uterine Growth Retardation (IUGR), Premature rupture of Membrane (PPROM), placental abruption, and intrauterine fetal death rates were all similar ( $p=0.063$ ,  $p=0.318$ ,  $p=1.00$ ,  $p=0.556$ ). In case having an elevation in both markers, increased rate of obstetric complications has been observed. A significant relationship was found between the high levels of maternal serum AFP and HCG multiple of the median and poor pregnancy outcomes like preeclampsia, IUGR, PPRM, intrauterine fetal death ( $p=0.003$ ,  $p=0.001$ ,  $p=0.040$ ,  $p=0.006$ ). They concluded that no definitive follow-up and treatment protocols have been established for patients at increased risk. In light of these findings, it is recommended to inform and educate patients about the most likely signs and symptoms of complications, to make more often antenatal visits, to perform more frequent ultrasound examination (fetal growth, AFI, etc.), NST, arterial/venous doppler, biophysical profile, and cervical length measurements in high-risk group.<sup>5</sup>

Study by [E L Butler](#) included 107 women with placenta Previa delivered during the study. Fourteen (13%, 95% CI 7%, 21%) had MSAFP at least 2.0 multiple of the median. They were significantly more likely than those with lower MSAFP levels to have one or more of the following outcomes: hospitalization for antepartum bleeding before 30 weeks' gestation (50% versus 15%), delivery before 30 weeks' gestation (29% versus 5%), or preterm delivery for pregnancy-associated hypertension before 34 weeks' gestation (14% versus none). The MSAFP cut-off of 2.0 multiple of the median provided the best combination of sensitivity and specificity for those outcomes, using receiver operating characteristic curves. They concluded that women with placenta Previa who also have high MSAFP levels are at increased risk of bleeding in the early third trimester and preterm birth. <sup>6</sup>

Risk factors of placenta Previa: mostly described are; Prior cesarean delivery, prior pregnancy terminations, history of intrauterine surgery, tobacco use, multiple gestation, multiparity, increased maternal age. <sup>7</sup>

The classic clinical presentation of placenta Previa is painless bleeding in the late second trimester or early third trimester. However, some patients with placenta

Previas will experience painful bleeding, possibly the consequence of uterine contractions or placental separation, some other cases could experience no bleeding at all before labour. There is a possibility that placenta Previa may lead to an unstable lie or mal presentation in late pregnancy. <sup>8</sup>

## **SUBJECTS AND METHODS:**

### **STUDY DESIGN:**

**Type of study:** A prospective study will included women with singleton pregnancies diagnosed by sonographic have evidence of placenta previa ,admitted to Maternity department ( Benghazi Medical Center (BMC) during 1<sup>st</sup> Sep 2021 to 28<sup>th</sup> Feb. 2021 ( six months) and will be followed till delivery .

### **Participants:**

All mothers admitted to Maternity department with singleton pregnancies beyond 24 weeks gestation will be selected in this study and proofed with ultra sound had placenta previa (the most recent ultrasound result will be dependable if the patients had more than one ultra Sound) .

**Exclusion criteria: Multiple pregnancy**

**Procedure:**

Full medical and obstetric history will be taken from all patients, general and obstetric examination will be done, MSAFP will be measured. Mothers will be follow till delivery.

**Data collection:** All needed data will be collected in proforma in appendix I.

**Data Analysis:**

Data will be analysed using (SPSS) statistical package of social science program version 23. The statistical analysis included:

I. Descriptive Statistics: Including (Mean value, Standard deviation, Number and Percentage.

II. Inferential Statistics: will be used when needed as t- test and Chi-square, P-value will be considered significant when  $\leq 0.05$ .

Data will be presented in form of tables and figures, where the figures will be done by Microsoft Excel 2010.

**Time table:** This study will be done during period from August 2020 to January 2021 explained in following time

**Approval:**

Approval of the study obtained from the manager of the hospital & informed consent obtained from the mothers.

## RESULTS

The study included 103 placenta Previa cases and 96 control cases.

**Table1: Placenta Previa cases & their control group**

Group	No	%
Placenta Previa cases	103	51.8
Control	96	48.2
Total	199	100.0

The mean age  $\pm$  standard deviation of cases was  $34.89 \pm 5.817$  years as compared to their control  $31.98 \pm 6.476$  years. This difference was statistically significant,  $P= 0.001$  as shown in table 2.

**Table 2: Age in years of Placenta Previa cases & their control group**

<b>Placenta Previa cases</b>		<b>Control</b>	
<b>N= 103</b>		<b>N=96</b>	
Mean	Standard deviation	Mean	Standard deviation
34.89	5.817	31.98	6.476

$P= 0.001$

The mean number of gravidity  $\pm$  standard deviation of cases was nearly similar to their controls ( $5.17 \pm 2.43$  &  $5.17 \pm 2.74$  years respectively). This difference was not statistically significant,  $P= 0.996$ . Regarding parity, the mean number of parity  $\pm$  standard deviation of cases was  $3.39 \pm 1.94$  as compared to  $3.60 \pm 2.33$  of controls. The difference was not statistically significant,  $P= 0.478$ . The mean number of abortions  $\pm$  standard deviation of cases was  $0.91 \pm 1.23$  as compared to  $0.57 \pm 0.98$ , this difference was statistically significant.

**Table 3: Gravidity, parity and number of abortions of Placenta Previa cases & their control group**

Variable	Groups	Mean	Standard deviation	P value
<b>Gravida</b>	Placenta Previa cases N= 103	5.17	2.43	P= .996
	Control N=96	5.17	2.74	
Parity	Placenta Previa cases N= 103	3.39	1.94	P= .478
	Control N=96	3.60	2.33	
Number of abortions	Placenta Previa cases N= 103	.91	1.23	P=0.034
	Control N=96	.57	.98	

The most common blood group among both cases & controls was group O positive, 59.0 % were controls as compared to 41.0% were cases, followed by blood group A positive, 51.9% were controls as compared to 48.1% were cases and the least common blood group was AB negative, only one case had this blood group. These differences were not statistically significant, P= 0.06. Table 4

**Table 4: Blood groups of Placenta Previa cases & their control group**

Group	Blood group							
	B positive	B negative	A positive	A negative	AB positive	AB negative	O positive	O negative
Case No	20	0	28	7	15	1	25	7
Case %	(54.1%)	0.0%	51.9%	87.5%	57.7%	100.0%	41.0%	70.0%
Control No	17	2	26	1	11	0	36	3
Case %	45.9%	100.0%	48.1%	12.5%	42.3%	0.0%	59.0%	30.0%
Total No	37	2	54	8	26	1	61	10

Total %	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
---------	--------	--------	--------	--------	--------	--------	--------	--------

P= 0.06

Regarding history of preterm labor of both cases & controls, all preterm labor cases were recorded among placenta Previa cases and none of the control cases had history of preterm labor. This difference was statistically significant, P= 0.002, as shown by table 5.

**Table 5: History of preterm labor of Placenta Previa cases & their control group**

Group	History of preterm labor	
	Yes	No
Case No	10	93
Case %	100.0%	49.2%
Control No	0	96
Case %	0.0%	50.8%
Total No	10	189
Total %	100.0%	100.0%

P= 0.002

Regarding history of preterm labor of both cases & controls, all preterm labor cases were recorded among placenta Previa cases and none of the control cases had history of preterm labor. This difference was statistically significant,  $P= 0.002$ , as shown by table 6.

**Table 6: History of preterm labor of Placenta Previa cases & their control group**

Group	History of preterm labor	
	Yes	No
Case No	10	93
Case %	100.0%	49.2%
Control No	0	96
Case %	0.0%	50.8%
Total No	10	189
Total %	100.0%	100.0%

$P= 0.002$

Regarding history of preterm labor of both cases & controls, all preterm labor cases were recorded among placenta Previa cases and none of the control cases had history of preterm labor. This difference was statistically significant,  $P= 0.002$ , as shown by table 7.

**Table 7: History of preterm labor of Placenta Previa cases & their control group**

Group	History of preterm labor	
	Yes	No
Case No	10	93
Case %	100.0%	49.2%
Control No	0	96
Case %	0.0%	50.8%
Total No	10	189
Total %	100.0%	100.0%

Regarding history of complications in the past deliveries of both cases & controls, all complications were recorded among placenta Previa cases and none of the control cases had history of past pregnancy complications. The complications included; postpartum hemorrhage, blood transfusion, urinary tract infection, adhesions, pre-eclamptic toxemia, perineal tear, abruptio placenta & adhesions, and wound infection. These differences were statistically highly significant,  $P= 0.001$ , as shown by table 8.

**Table 8: History of complications in the past deliveries of Placenta Previa cases & their control group**

Group	History of any complication in the past delivery							
	Post-partum hemorrhage	Urinary tract infection	Blood transfusion	Adhesions	Pre-eclamptic toxemia	Perineal tear	Abruptio placenta & adhesions	Wound infection
Case No	2	1	2	2	1	1	1	1
Case %	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%

Control No	0	0	0	0	0	0	0	0
Case %	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
Total No	2	1	2	2	1	1	1	1
Total %	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%

P=0.001

Regarding history of bleeding in early pregnancy of both cases & controls, all cases with a history of bleeding in the early present pregnancy cases were recorded among placenta Previa cases and none of the control cases had history of preterm labor. This difference was statistically highly significant, P= 0.000, as shown by table 9.

**Table 9: History of Bleeding in early pregnancy among Placenta Previa cases & their control group**

Group	History of Bleeding in early pregnancy	
	Yes	No
Case No	24	79
Case %	100.0%	45.1%
Control No	0	96
Case %	0.0%	54.9%
Total No	24	175
Total %	100.0%	100.0%

P= 0.000

Regarding history of bleeding in the second & third trimester of both cases & controls, all cases with a history of bleeding in the second & third trimester of the present pregnancy were recorded among placenta Previa cases and none of the control cases had history of preterm labor. This difference was statistically highly significant,  $P= 0.000$ , as shown by table 10.

**Table 10: History of Bleeding in in the second & third trimester among Placenta Previa cases & their control group**

Group	History of Bleeding in the second & third trimester	
	Yes	No
Case No	99	4
Case %	100.0%	4.0%
Control No	0	96
Case %	0.0%	96.0%
Total No	99	100
Total %	100.0%	100.0%

$P= 0.000$

Regarding history of passive smoking of both cases & controls, 54.1% of cases with a history of passive smoking were recorded among placenta Previa cases and 45.9 % of the control cases had history of passive smoking. A higher proportion of cases who had no history of passive were control group (78.6%) compared to 21.4% who were placenta Previa cases. These differences were statistically significant,  $P= 0.025$ , as shown by table 11.

**Table 11: History of passive smoking among Placenta Previa cases & their control group**

Group	History of passive smoking	
	Yes	No
Case No	100	3
Case %	54.1%	21.4%
Control No	85	11
Case %	45.9%	78.6%
Total No	185	14
Total %	100.0%	100.0%

$P= 0.025$

Regarding history of antepartum hemorrhage & LAP of both cases & controls, all cases with a history of antepartum hemorrhage & LAP of the present pregnancy were recorded among placenta Previa cases and none of the control cases had history of antepartum hemorrhage & LAP of the present pregnancy. These differences were statistically highly significant,  $P=0.000$ , as shown by table 12.

**Table 12: History of Bleeding in in the second & third trimester among Placenta Previa cases & their control group**

Group	History of Antepartum hemorrhage & LAP	
	Yes	No
Case No	98	5
Case %	100.0%	5.0%
Control No	0	96
Case %	0.0%	95.0%
Total No	98	101
Total %	100.0%	100.0%

$P=0.000$

Regarding type of placenta of both cases & controls, all cases with low lying placenta, all marginal placenta cases, partial placenta and central placenta were recorded among placenta Previa cases. All cases with anterior & posterior fundal placenta were recorded among control group. These differences were statistically highly significant,  $P= 0.000$ , as shown by table 13.

**Table 13: Type of placenta among Placenta Previa cases & their control group**

Group	Type of placenta					
	Low lying placenta	Marginal placenta	Partial placenta	Central placenta	Anterior fundal placenta	Posterior fundal placenta
Case No	58	2	2	41	0	0
Case %	100.0%	100.0%	100.0%	100.0%	0.0%	0.0%
Control No	0	0	0	0	95	1
Case %	0.0%	0.0%	0.0%	0.0%	100.0%	100.0%
Total No	58	2	2	41	95	1
Total %	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%

$P=0.000$

Regarding history of postpartum hemorrhage of both cases & controls, all cases with a history of postpartum hemorrhage of the present pregnancy were recorded among placenta Previa cases and none of the control cases had history of postpartum hemorrhage of the present pregnancy. These differences were statistically highly significant,  $P= 0.002$ , as shown by table 14.

**Table 14: History of Present pregnancy postpartum hemorrhage among Placenta Previa cases & their control group**

<b>Group</b>	<b>History of postpartum hemorrhage</b>	
	Yes	No
Case No	10	93
Case %	100.0%	49.2%
Control No	0	96
Case %	0.0%	50.8%
Total No	10	189
Total %	100.0%	100.0%

$P= 0.002$

Regarding preterm delivery of the present pregnancy of both cases & controls, all preterm deliveries were recorded among placenta Previa cases and none of the control cases had preterm delivery. These differences were statistically significant,  $P= 0.000$ , as shown by table 15.

**Table 15: History of preterm delivery of the present pregnancy of Placenta Previa cases & their control group**

Group	History of preterm delivery of the present pregnancy	
	Yes	No
pCase No	27	76
Case %	100.0%	44.2%
Control No	0	96
Case %	0.0%	55.8%
Total No	27	172
Total %	100.0%	100.0%

P= 0.000

Regarding history of placenta accrete of the present pregnancy of both cases & controls, all of placenta accrete cases were recorded among placenta Previa cases and none of the control cases had preterm delivery. These differences were statistically significant, P= 0.000, as shown by table 16.

**Table 16: History of placenta accrete of the present pregnancy of Placenta Previa cases & their control group**

Group	History of Present pregnancy placenta accrete of the present pregnancy	
	Yes	No
Case No	14	89
Case %	100.0%	48.1%
Control No	0	96
Case %	0.0%	51.9%
Total No	14	185
Total %	100.0%	100.0%

P= 0.000

Regarding Mode of delivery of the present pregnancy of both cases & controls, all of the placenta Previa cases had cesarean section compared to none of the control cases. These differences were statistically significant,  $P= 0.000$ , as shown by table 17.

**Table 17: Mode of delivery of the present pregnancy of Placenta Previa cases & their control group**

Group	Mode of delivery of the present pregnancy	
	Normal delivery	Cesarean Section
Case No	9	94
Case %	8.6%	100.0%
Control No	96	0
Case %	91.4%	0.0%
Total No	105	94
Total %	100.0%	100.0%

$P= 0.000$

Regarding history of cesarean hysterectomy of the present pregnancy of both cases & controls, all of the placenta Previa cases had cesarean hysterectomy compared to none of the control cases. These differences were statistically significant,  $P= 0.000$ , as shown by table 18.

**Table 18: History of cesarean hysterectomy of the present pregnancy of Placenta Previa cases & their control group**

Group	Cesarean hysterectomy of the present pregnancy	
	Yes	No
Case No	12	91
Case %	100.0%	48.7%
Control No	0	96
Case %	0.0%	51.3%
Total No	12	187
Total %	100.0%	100.0%

$P= 0.000$

Regarding history of cesarean hysterectomy of the present pregnancy of both cases & controls, all of the placenta Previa cases had cesarean hysterectomy compared to none of the control cases. These differences were statistically significant,  $P= 0.000$ , as shown by table 19.

**Table 19: History of cesarean hysterectomy of the present pregnancy of Placenta Previa cases & their control group**

Group	Cesarean hysterectomy of the present pregnancy	
	Yes	No
Case No	12	91
Case %	100.0%	48.7%
Control No	0	96
Case %	0.0%	51.3%
Total No	12	187
Total %	100.0%	100.0%

$P= 0.000$

Regarding history of blood transfusion of the present pregnancy of both cases & controls, 90.3% of the placenta Previa cases had history of blood transfusion compared to 9.7% of the control cases. These differences were statistically significant,  $P= 0.000$ , as shown by table 20.

**Table 20: History of blood transfusion of the present pregnancy of Placenta Previa cases & their control group**

Group	Blood transfusion	
	Yes	No
Case No	28	75
Case %	90.3%	44.6%
Control No	3	93
Case %	9.7%	55.4%
Total No	31	168
Total %	100.0%	100.0%

$P= 0.000$

## DISCUSSION

Placenta Previa can result in life-threatening maternal complications such as hemorrhage and shock and similarly adverse infant outcomes such as prematurity, stillbirth and neonatal death

Regarding maternal age the present study showed that placenta cases had higher mean age as compared to control cases similarly, ZAHRA et al in her Master degree thesis had advanced maternal age among the studied cases associated with Placenta.<sup>10</sup>

The present study showed that cases with placenta Previa had a history of prior cesarean section, advanced age, multiparity, and history of passive smoking. Many studies showed that placenta Previa include a history of prior cesarean delivery, uterine injuries, increasing age and multiparity.  
11-13

The current study illustrated that advanced maternal age was associated with increased risk of placenta Previa, similarly, Rosenberg et al., showed that advanced maternal age has been associated with a slight increase in the risk of placenta Previa, and also could be due to effect of multiparity. Similarly, Surraya concluded that there's an association between incidence of placenta Previa and increasing parity.<sup>14</sup>

A systematic review and meta-analysis of 22 studies including over 2 million deliveries indicated that the incidence of placenta Previa increases from 10 in 1000 deliveries with one previous caesarean delivery to 28 in 1000 with three or more caesarean deliveries.<sup>15</sup>

Wright et al., reported that the mean blood loss for PAS disorders cases undergoing cesarean hysterectomy was 3000ml, whereas the mean required packed red blood cell (PRBC) units for transfusion was 5 units. An estimated blood loss of  $\geq 5000$  mL was found in about 41.7% of women with a known diagnosis of PAS disorders.<sup>16, 17</sup>

In the current study there was all of women with a history of abortion in the placenta Previa group than in the control group. Bulk studies concluded that a history of abortion has a contributing role in placenta Previa in the succeeding pregnancy.<sup>18-22</sup>

## **CONCLUSION**

It is concluded that placenta Previa is associated with advanced maternal age, history of previous abortion, multiparity, history of previous antepartum & postpartum hemorrhage, preterm deliveries and previous history of cesarean section. Similarly, history of present bladder injury , blood transfusion, premature delivery and history of antepartum hemorrhage in the present pregnancy

## **RECOMMENDATIONS**

From the present study, it is recommended that there is a need for comprehensive obstetrics care to appropriately treat Placenta Previa and its complications. This calls for proper examination and education during the antenatal care.

## References:

- 1-Wasan Hameed Salman, Faiz Al –Waeely. (2019). Predisposing factors and pregnancy outcome in different types of Placenta Previa. *Int. J. Adv. Res. Biol. Sci.* 6(2): 13-23.
- 2- Pooja Verma, Kavita N. Singh, Vineeta Ghanghoriya .To study analyze maternal serum alpha-fetoprotein as a biomarker of placental adherence in low lying placenta. *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*  
*Verma P et al. Int J Reprod Contracept Obstet Gynecol. 2016 Jun;5(6):1959-1963*
- 3 -Mohammed Hany Mosbeh, Mohammed Abdallah Mohammed, Mo'men Mohamed Hassan, Ahmed Rabie Abd El-Raheim, Heba Reda Mohammed. Alpha-Fetoprotein for Prediction of Placenta Accreta in Women with Complete Placenta Previa Centralis: A Prospective Study. *Indian Journal of Public Health Research & Development*, March 2020, Vol. 11, No.03
- 4-E L Koster , J S Dashe, D D McIntire, R M Ramus. Association of maternal serum alpha-fetoprotein with persistent placenta previa. *J Matern Fetal Neonatal Med* 2004 Jul; 16(1):3-7.
- 5-Hümeyra Öztürk, Salim Erkaya, Sibel Altınbaş Burak Karadağ, Nazan Vanlı Tonyalı, Demet Özkan. The role of unexplained high serum alpha-fetoprotein (AFP)

and human chorionic gonadotropin (hCG) levels in the second trimester to determine poor obstetric outcomes. J Turk Soc Obstet Gynecol 2014; 3:142-7

6- E L Butler<sup>1</sup>, J S Dashe, R M Ramus. Association between maternal serum alpha-fetoprotein and adverse outcomes in pregnancies with placenta previa. Obstet Gynecol 2001 Jan; 97(1):35-8.

7- Placenta Praevia and Placenta Accreta: Diagnosis and Management (Green-top Guideline No. 27a). Available from:

<https://obgyn.onlinelibrary.wiley.com/doi/pdf/10.1111/1471-0528.15306>. Accessed

in 20 / 12/ 2020

8- Oyelese Y, Smulian JC. Placenta Previa, Placenta Accreta, and Vasa Previa. J Obstetrics & Gynecology. 2006; 107(4):927- 41.

9- Dulay A T. Placenta Previa. 2022 Merck Sharp & Dohme Corp 2022 Merck Sharp & Dohme Corp., a subsidiary of Merck & Co., Inc., Kenilworth, NJ, USA

**10- ZAHRA R.:** Advanced maternal age and placenta previa for women giving birth in Finland; a register-based cohort study. UNIVERSITY OF EASTERN FINLAND, Faculty of Health Sciences Public health. Master's thesis, 2017.

11- Ananth CV, Smulian JC, Vintzileos AM. The association of placenta previa with history of cesarean delivery and abortion: a metaanalysis. Am J Obstet Gynecol 1997; 177:1071–8.

12- Ananth CV, Savitz DA, Luther ER. Maternal cigarette smoking as a risk factor for placental abruption, placenta previa, and uterine bleeding in pregnancy. Am J Epidemiol 1996; 144: 881–9.

13- Ananth CV, Wilcox AJ, Savitz DA, Bowes WA Jr, Luther ER. Effect of maternal age and parity on the risk of uteroplacental bleeding disorders in pregnancy. Obstet

Gynecol 1996;88:

511–6.

14-Marshall N, Fu R, Guise J. (2011): Impact of multiple cesarean deliveries on maternal morbidity: a systematic review. Am J Obstet Gynecol; 205:262.e1–8.

UNDER PEER REVIEW

- 15-Wright J, Pri-Paz S, Herzog T, Shah M, Bonanno C, Lewin S. (2011): Predictors of massive blood loss in women with placenta accreta. *Am J Obstet Gynecol*; 205:38.e1-6.
- 16-Judson E, Polyakov A, Lawrence A.(2008): Intra-abdominal haemorrhage at 17 weeks' gestation caused by placenta percreta: a case report. *Royal Austr N Zeal Col Obstet Gynecol*; 48:218–224.
- 17- Lal A, Hibbard J. (2015): Placenta previa: an outcome-based cohort study in a contemporary obstetric population. *Arch Gynecol Obstet*; 292:299–305.
- 18-Tuzovic L, Djelmis J, Ilijic M. Obstetric risk factors associated with placenta previa development: case-control study. *Croat Med J*. 2003; 44: 728-733.
- 19-Oya A, Nakai A, Miyake H, Kawabata I, Takeshita T. Risk factors for peripartum blood transfusion in women with placenta previa: a retrospective analysis. *J Nippon Med Sch*. 2008; 75: 146-151.
- 20-Shaikh S. Frequency of placenta previa in multigravida at tertiary care hospital. *Int J Cur Res Rev*. 2014; 6: 39-43.
- 21-Karami M, Jenabi E. Placenta previa after prior abortion: a meta-analysis. *Biomed Res Ther*. 2017; 4: 1441-1450.
- 22-Hung TH, Hsieh CC, Hsu JJ, Chiu TH, Lo LM, Hsieh TT. Risk factors for placenta previa in an Asian population. *International Journal of Gynecology and Obstetrics*. 2007; 97: 26-30.

UNDER PEER REVIEW