

Original Research Article

EVALUATION OF THE CHEMICAL QUALITY OF MILK POWDER SOLD IN OPEN MARKETS IN ABA, ABIA ATATE, NIGERIA

ABSTRACT

Objective: This study investigated some chemical parameters of milk powder sold in Aba market.

Materials and Methods: A total of six (6) brands of milk powder, namely: Peak, Three Crown, Loyal, Cow Bell, Jago, and Dano milk, packaged in affordable sachets, were purchased from the new market (Abia Ohuru) in Aba town, Abia State. The samples were emptied from the sachet into sterile plastic containers, coded, sealed and transported to the laboratory immediately for analysis. Proximate, mineral, and vitamin compositions were determined using standard methods. Anova was used to separate the means and a significant difference was judged at $p < 0.05$

Result The obtained values for almost all parameters were not significantly different (p) among the samples examined. However, the obtained values were slightly lower than the values written on the labels. The obtained values and the values on the labels were not significantly different from the CODEX Alimentarius and NFDAC set standards for milk powder, except for vitamin A values for almost all the samples.

Conclusion: The quality of milk powder sold in the Aba market can be considered to be good enough for human consumption, although slight quality degradation was found in protein, carbohydrates, fat, and especially in the vitamin A and B content of the samples. Exposure to harsh conditions during storage and retailing processes could be a possible reason.

Keywords: Evaluation, Nutrient composition, Milk powder, Open Market.

INTRODUCTION

Milk and dairy products are consumed by Nigerians of all ages and ethnic groups. Nigerians drink milk and dairy products that are either produced locally or imported from other parts of the world. Northerners who raise cattle have traditionally enjoyed fresh milk and locally produced

milk products for many decades. (Akinyosoye, 2004). Globalization and international trade have led to the widespread consumption of milk and dairy products across the country in recent decades.

Many Nigerians now drink milk from animals. According to a survey, consumption of various types of imported milk and milk products is highest in the south eastern geopolitical zone.

(Akinyosoye, 2004). The bulk of milk imported into the country is powdered milk (milk powder). Milk powder may be whole or skim milk powders (SMPs). Skim milk powder has a maximum shelf life of 3 years, while whole milk powder has a maximum shelf life of about three months.

The bulk of milk imported into the country is powdered milk (milk powder). Milk powder may be whole or skim milk powders (SMPs). Skim milk powders has a maximum shelf life of 3 years, while whole milk powder has a maximum shelf life of about three months (Flegam and Oluwaniyi, 2015).

Dehydration or drying is used to reduce bulk for easier transportation and to minimize water activity to limit the growth of microorganisms, extending the shelf life of the product. (Augustin and Carven, 2003). Thus, milk powder is the product of dehydration of fluid milk to about 97% solids (FAO, 2003).

Although milk has a low microbial load and low water activity, milk powder is not sterile. Under certain conditions, the quality of milk changes, and this changes its health effects.

In developing countries, subjecting milk powders to extreme weather conditions is prevalent. Prolonged storage at high temperatures and exposure to ultraviolet rays cause quality deterioration. Milk sugar and milk proteins under prolonged storage at high temperatures interact, resulting in a complex sequence of chemical reactions known as the non-enzymatic browning or Millard reaction. This can result in the loss of lysine and a change in the nutritional content of milk proteins (adirusmadi.blogspot.com.2013). Milk fat rancidity has been linked to prolonged storage and exposure to light. Furthermore, storage of milk powders at room

temperature causes a noticeable alteration in color and flavor. High temperatures and high water activity accelerate microbiological spoilage of milk powder (adirusmadi.blogspot.com, 2013).

Besides, adulterants like soap, acid, starch, table sugar, and chemicals are often added to milk either to increase its shelf-life or to increase the profit of the producer. (Azad and Ahmed, 2016). Poor storage conditions and adulterants lower the quality of milk and milk products. Milk quality degradation is very common globally. Lack of routine quality control, the perishability nature of milk and efforts to bridge the gap between demand and supply are some of the possible reasons behind milk quality degradation (Kamthania *et al.*, 2014).

To ensure that consumers are always offered good quality milk in developed countries, strict monitoring is done routinely until good quality milk gets to the consumers. Such strict monitoring is lacking in developing countries.

The aim of the study was to evaluate the chemical quality of milk obtained from the open market and to compare the quality with standard quality and the values written on the label of each sample.

MATERIALS AND METHODS

The Study Area

Aba Metropolis

The study was carried out in Aba, a strategically selected site in the south-eastern geopolitical zone of Nigeria.

Businessmen and women who are engaged in importation, production, processing, and marketing businesses dwell in Aba. Milk powder is one of the commodities imported into Aba. Processing of milk powder into dairy products also goes on in Aba at micro or macro levels. Hawkers, retailers in the various markets in Aba, supermarkets, fast foods, and restaurants all market milk and milk products in Aba.

The majority of Aba residents consume milk and dairy products marketed in Aba, particularly children and young adults. High-calorie, salty, fatty, and sugary foods, as well as a sedentary lifestyle, are popular among Aba residents, as they are with many other urban dwellers in Sub-Saharan Africa (Chukwuonye *et al.*, 2013). And as a result, diet and lifestyle-related diseases are major health concerns for Aba residents (Chukwuonye *et al.*, 2013; Ogah, 2013).

Procurement of Dried milk Samples.

Six (6) of the most commonly consumed dried milk powder were purchased from the new market (Ahia Ohuru), a purposefully selected market in Aba. The samples of milk were labeled, put inside a sterile plastic container and transported immediately to the laboratory for analysis. Peak, Three Crowns, Jago, Loyal, Cowbell milk, and Dano milk were the dried milk powders purposefully chosen.

Chemical Quality

Proximate Analysis

The AOAC (2010) methods were used to determine the: moisture, crude protein, fat, crude fiber and ash content of the dried milk powder. The oven method was used for moisture determination, The Kjeldahl method was used to calculate crude protein, and The AOAC (2010) soxhlet extraction method was used to determine the fat content. Oven drying, digestion and extraction methods were used to determine the crude fiber, ash content was determined by weighing, incinerating, cooling and re- weighing. According to Oyenuga (1968), the

carbohydrate content was determined by difference: % Carbohydrates = (% moisture + % fat + % ash + % protein + % crude fiber) – 100. The calorific content of the sample was calculated using the values obtained for protein, fat, and carbohydrate. Determination of Minerals: The phenanthroline method of Lee and Stumm (1960) was used to evaluate iron. The method described by Pearson (1976) was used to calculate calcium levels. A flame photometer was used to determine potassium, as defined by Osborne and Voogt (1978). Zinc was determined using an atomic absorption spectrophotometer (Hitachi model A-1800) and flame atomization. The sodium content of dried milk powder was determined using the AOAC flame photometry process. The colorimeter was used in accordance with the AOAC (2010) procedure to determine the vitamin A. The AOAC (2010) scalar analyzer method was used to assess thiamine, riboflavin amount and vitamin C levels. The solid non-fat content was calculated using the difference method (SNF). The reduction method was used to calculate lactose.

Data Analysis

Data was entered, organized and analyzed using Statistical Package for Social Scientists SPSS Inc., Chicago, Illinois, USA) Version 15.0. Numerical data obtained from the various analyses was expressed as means and standard deviation. Descriptive statistics of mean, standard deviation and student's -T student were calculated the absolute differences.

RESULTS

Proximate Composition of Dried Milk

Table 1 shows the proximate composition of dried milk commonly consumed by the students. There were significant differences ($p < 0.05$) in the proximate composition of dried milk commonly consumed by the students. Moisture content was highest (4.05^a) in Three crown milk and lowest (3.08^b) in Dano milk. The protein content of the samples ranged (20.41%) in Dano milk to (23.97%) in Three Crown milk. The obtained protein values were lower than the values written on the label of each sample and the standard value of 24%. Fat was highest (28.39%) in Three crown milk and slowest (25.79%) in Peak milk. The obtained values were lower than the values on the labels of each sample, but higher than the stipulated 24% fat. Also, ash content was highest (5.78%) in Dano milk and lowest (4.97%) in Three crown milk. The values obtained were lower than the standard protein value of 6.5%. Not all the samples had ash content written

on their labels. Similarly, carbohydrate was highest (40.71%) in Dano milk and lowest (36.45%). Furthermore, protein content of the entire samples except in cowbell milk (24.07%) was lower than the recommended minimum value of not less than 24% (NAFDAC, 2019) and of minimum of 34% of CODEX (2014) and also was lower than what is written in the label (26g, 32g, 23.5g, 25g, and 23.5g 23.6 g in loyal, Dano, Jago, Peak, Three crown and cowbell milk respectively.

Tables: 1: Proximate Composition of Dried Milk Samples

Brand (%)	MC (%)	CP (%)	FAT (%)	CF (%)	ASH (%)	CHO (%)	EV (%)
Three crown	4.05 ^{ab}	23.97 ^{ab}	28.39 ^{ab}	0.165 ^{ab}	4.99 ^{ab}	38.55 ^{ab}	516.65 ^{ab}
Dano milk	3.98 ^b	20.41 ^{bc}	27.55 ^{bc}	0.340 ^{bc}	5.78 ^{bc}	40.71 ^{bc}	487.21 ^b
Jago	4.12 ^{cb}	23.66 ^{cb}	26.87 ^c	0.165 ^a	5.13 ^c	37.58 ^a	492.34 ^c
Cowbell milk	4.03 ^{ab}	23.07 ^{ab}	27.54 ^{bc}	0.110 ^c	5.62 ^a	36.45 ^c	502.20 ^a
Peak milk	4.11 ^{ca}	23.81 ^a	25.79 ^a	0.105 ^a	5.31 ^c	39.62 ^b	482.67 ^c
Loyal	3.52 ^b	23.96 ^c	27.52 ^b	0.080 ^c	5.52 ^b	38.54 ^a	499.67 ^{ab}
S value	4.00	24.00	24.00	0	6.5	38	496
LSD:	0.08	0.21	0.085	0.065	0.095	0.20	0.07

Values are given as the mean and L.S. D.; Mean scores having different superscript letters on the same column differ significantly at 5% significant level.

Footnote: moisture content (MC), crude protein (CP), crude fiber (CF), carbohydrate (CHO), energy value (EV)

Mineral Composition of Milk samples

Table 2 shows Mineral Composition of Milk samples. Calcium ranged from (572mg) in cowbell milk to (867mg) in loyal milk. The values obtained are similar to the values written on the labels, but slightly lower than the standard value of 950mg/100g. The values obtained for sodium ranged from (37.03mg) in Jago milk to (44.02 mg) in Cowbell Milk. Not all the samples had sodium value written on its label. The obtained values differed from the standard value of

37mg/100g. The potassium content obtained for the samples was highest (1320mg) in loyal milk and lowest (1257 mg) in three crown milk. The values obtained were lower than standard value for potassium (133mg/100g). Not all the samples have the potassium content written on the label. Zinc value ranged from (2.57 mg) in three crown to (2.90 mg) in cowbell milk. The values were lower than the standard value for zinc of 3.1mg/100g for. Not all the samples had zinc value written on the label.

Table 2: Mineral Composition of Dried Milk Samples

	Ca	Na	K	Fe	Zn
Brand	mg/100g	mg/100g	mg/100g	mg/100g	mg/100g
Three cr	805.22 ^{ab}	38.45 ^{ba}	1257 ^{ac}	0.515 ^d	2.57 ^a
Jago	833.37 ^{bc}	37.03 ^{bc}	1281 ^{cb}	0.530 ^d	2.71 ^b
Dano	806.02 ^c	39.04 ^b	1321 ^c	0.360 ^d	2.66 ^c
Peak	834.55 ^a	37.92 ^c	1284 ^{cb}	0.420 ^d	2.73 ^b
Cowbell	843.85 ^b	44.02 ^a	1281 ^{cb}	0.585 ^d	2.90 ^e
Loyal	867.61 ^c	41.62 ^b	1320 ^c	0.490 ^d	2.85 ^e
Std					
value	950mg/100g	37mg/100g	1330mg/100g	0.50mg/100g	3.1mg/100g
LSD	0.685	0.485	0.495	0.5	0.125

Values are given as the mean and L.S. D.; Mean scores having different superscript letters on the same column differ significantly at 5% significant level Footnote: calcium (Ca), sodium (Na), potassium (K), Iron (Fe), zinc (Zn)

Vitamin Composition of Milk Samples

Table 3 shows vitamin composition of Milk samples. The vitamin A content obtained ranged from (579i μ) in Dano milk to (2475^b i μ) in cowbell milk. The values obtained were lower than the values written on the label of each sample. The obtained values, the values written on the labels and the standard value for vitamin A all differed significantly (P<0.05). Vitamin B₁, was

highest (0.89mg) in peak milk, lowest (0.09 mg) in cowbell milk. The values differed significantly from the standard value for vitamin B₁, of (0.26mg). Not all the milk samples had the vitamin B₁ content written on the label. Vitamin B₂ was highest (1.44 mg) in loyal milk and lowest (0.72 mg) cowbell milk. The values obtained were lower than the standard value of 1.45mg. Not all the sample had to vitamin B₂ content writhen on the label. The vitamin B₃ content obtained ranged from (0.20 mg) in cowbell milk to (0.27 mg) in loyal milk. The values obtained were lower than the standard value of 0.68mg/100. Not all the sample had the value of vitamin B₃ written on the label. Vitamin C content ranged from (0.61 mg) in three crown milk to (0.87 mg) in peak milk. The values obtained were lower than standard value of 2.2mg/100g for vitamin C. Not all the samples had vitamin C content written on the labels.

Table 3: Vitamin Composition of Dried Milk

Brand	Vitamin A	Vitamin B ₁	Vitamin B ₂	Vitamin B ₃	Vitamin C
	(1U) (mg)/100g)	(1U) (mg)/100g)	(1U) (mg)/100g)	(1U) (mg)/100g)	(1U) (mg)/100g)
Three Crown	741 ^a	0.85 ^a	0.79 ^a	0.210 ^a	0.61 ^a
Cowbell	2475 ^b (iμ)	0.09 ^b	0.72 ^b	0.20 ^b	0.64 ^a
Jago	662 ^c	0.18 ^c	0.75 ^c	0.23 ^b	0.69 ^c
Loyal	2480 ^d	0.10 ^d	1.44 ^d	0.27 ^c	0.81 ^d
Peak	735 ^e	0.89 ^e	0.85 ^e	0.25 ^c	0.87 ^e
Dano	579 ^f	0.76 _f	1.42 ^c	0.22 ^b	0.81 ^f
Std value	11091/100g	0.26g	1.45mg	0.68mg/100g	2.2mg100g
LSD:	0.08	0.215	0.085	0.055	0.065

Values are given as the mean and L.S. D.; Mean scores having different superscript letters on the same column differ significantly at 5% significant level.

Table 4: NUTRITIONAL COMPOSITION OF SAMPLED MILK POWDER FROM THE LABELS

	Loyal	Peak	Dano	Three crown	Cow bell	Jago
Energy	507kcal	490kcal	500kcal	520kcal	516	520
Fat	28g	26g	28	28	30	30
Protein%	24g	24g	22g	23.5	23.6g	32
Carb	39g	40g	41g	39.9g	37.9g	40
Lactose	NS		36g	37.8g	NS	NS
Vita A	2500iu/RE	753 μ g	600/2000 iu	753mg/2500 iu	2500 iu	735
D ₃	350	9mg	10/400i μ	8.9mg/350m	200mg	10mg
E	NS	0.53mg	NS	12.6mg	4.5 μ g	NS
K	3mg	NS	NS	NS	12 μ g	NS
Folic	6mg	88mg	NS	20mg	200mg	8mg
Calcium	870	.84g	800mg	800mg	580	NS
Dietary fiber	NS	0	NS	NS	NS	NS
Vita C	7mg	88mg	NS	NS	18	15mg
B ₁	0.13mg	0.99mg	NS	.93mg	0.1mg	NS
B ₂	1.5mg	0.91mg	NS	NS	1mg	NS
B ₁₂	2.5mg	3.9mg	NS	2.5mg	1.6	NS
Phosphorus	620	NS	NS	620mg	750mg	NS
Zinc	2.5mg	NS	2.5mg	2.5mg	1.8mg	NS
Moisture	3g	NS	NS	NS	< 3%	4g
B ₃	0.3mg	NS	NS	NS	200	NS
Sodium	240	NS	NS	NS	180	NS

Footnote: NS; not stated

DISCUSSION

Proximate analysis of dried milk samples

The moisture content within normal ranges found in this study is not consistent with the findings of Khalid *et al.*, (2009) who reported lower moisture content in dried milk sampled in Sudan. Low moisture content implies low water activity levels, which does not permit microbial growth but might cause changes to occur in milk powders after production. It is the lipids and the non-lipid constituents of milk which undergo such changes.

Normally, the rates of such reactions are accelerated with the elevation of storage temperature and oxygen.

Furthermore, the milk fat and milk protein of most of the samples were found to be slightly lower than the fat content written on the label of each sample. This agrees with the findings of Ahmed *et al.*, 2005 and Ahmed and Shakoori, 2002), who discovered quality depreciation in dried whole milk, but disagrees with the findings of Kajal *et al.*, (2012), who discovered protein and fat content within normal ranges in dried milk samples from Mymensingh, Bangladesh.

Protein quality degradation was also found. Protein quality deterioration manifests as the Millard reaction. This reaction leads to nutritional quality reduction (Ahmed *et al.*, 2005). Loss of nutritive value after production primarily concerns loss of available lysine due to Millard reactions. Storage at 20°C temperature, at normal water content does not cause an appreciable loss. Extensive Millard reactions can cause a decrease in protein digestibility and the formation of weak mutagens (Ahmed *et al.*, 2005; DT-6 lesson 42 (2012).

The digestibility of protein and the accessible lysine content of milk powders have been reported to decrease after manufacture and while in storage. The moisture content, storage period, and storage temperature all influence these reductions. Higher moisture levels and longer storage times at high temperatures have been found to hasten degradation. Because storage temperatures in tropical markets are high and unmanaged, the observed decline in the quality of dried whole milk samples necessitates increased supervision and education of the marketers. The mineral and vitamin content of dried milk samples declined as well, according to the results of the vitamin and mineral study. The nutritional quality of milk powder is affected when it is stored in adverse conditions during transportation, storage and distribution. All of the samples showed a considerable loss in vitamin A content, as well as a modest fall in vitamin B1, B2, B3, and C

content, which agrees with the finding of Akpanyung (2006). The decrease in mineral content is also similar with the findings of Akpanyung (2006), who observed an average calcium concentration of 11,108 mg/kg in powdered milk samples purchased from Nigerian local markets. The mean Fe content in this study, ranging from (0.36mg/100g to 0.58mg/g), is higher than that reported by Semaghiul *et al.*, (2008) (21.73 ppm), Salah *et al.*, (2013) (20.41 ppm) and Perween *et al.*, 2011 (4.33 mg/l). Akpanyung (2006) reported a mean Zn content of 126.1mg/kg, which is similar to that obtained in this study. But it is significantly higher than that reported by Semaghiul *et al.*, (2008) of 3.24 ppm.

The higher values written on the label of all the milk samples, compared to the values obtained for vitamin A and water-soluble vitamins in this study, are an indication of quality deterioration. High temperature storage affects water-soluble vitamins (Wikipedia, the free encyclopedia, 2019), causing a reduction in thiamin, vitamin B₁₂, and vitamin B₁₂. According to DT-6 lesson 42 (2012), exposure of milk powders to high temperatures and light causes vitamin and mineral losses. Also, exposure of milk powders to sunlight causes vitamin A destruction by the ultraviolet rays of sunlight.

Protecting milk powders against high temperatures and light helps to minimize loss of light-sensitive vitamins, in particular riboflavin. Tropical market places are characterized by high temperatures and the exposure of milk powder to light. Milk powders stored under adequate conditions (no light, minimal oxygen, low moisture) will likely have minimal loss of vitamins. It is suggested that to protect light-sensitive vitamins, exposure of powders to such conditions should be avoided (DT-6 lesson 42, 2012). Besides, periodic monitoring of milk at the various distribution chains is paramount.

Conclusion

The quality of milk powder sold in the Aba market can be considered to be fairly good for human consumption, although slight quality degradation was found in protein, carbohydrates, fat, and especially in the vitamin _A and _B content of the samples. Exposure to harsh conditions (high temperatures and exposure to ultra violet light) during storage and retailing processes could be the possible reason. Milk powder in the market places in tropical zones like Nigeria should be

marketed in a more conducive condition. Milk powder marketed in a cool, dry environment will not have an appreciable nutrient loss.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

- Akinyosoye V.O (2013) Demand for Dairy Products in Nigeria Evidence from the Nigerian Living Standard Survey. Journal of Economics in Rural Development. Vol. 16. No.1
- Azad T and Ahmed (2016). Common milk adulteration and their detections techniques. Food contamination 3, 26. Available @ <https://doi.org/10.1186/54055-016-0045-3> retrieved 2 oct, 2019.
- Food and agriculture organisation (FAO, 2013) milk and milk products. Available from <http://www.fao.org/dairy-production.products/> on adirumadi.blogspot.com. (2013).
- Kamthania M, Saxena J, Saxena K, Sharma DK (2014). Methods of Detection and Remedial Measure. Internal Journal of Engineering Technology. Res. 1: 15-20.
- www.usdec.org(2001). An overview of change in the characteristics, functionality and Nutritional value of skim milk powder (SMP) during storage.