

Effect of processing methods on the nutritional composition of ripe pumpkin fruit

ABSTRACT

This research was carried out to assess processing methods on the nutritional composition of ripe pumpkin (*C. moschata*). Thermal processing (steam blanching and drying) was used to prepare pulp and powder. A significant difference was observed in moisture content between fresh, pulp and powder i.e. 88.14 %, 88.93 % and 6.12 % respectively. The heat-sensitive nutrients (ascorbic acid and β -carotene) were significantly decreased after processing to a pulp (13.88 mg/ 100 g) and powder (10.77 mg/ 100 g) in comparison to fresh fruit (15.81 mg/ 100 g). The ash, crude fibre, fat, protein, total soluble solids, titratable acidity, total carbohydrate, energy, reducing sugars and total sugars significantly increased in powder as compared to fresh and pulp due to moisture content difference. The colour was recorded to be (L^* 36.83, a^* 4.44, b^* 40.13), (L^* 0.19, a^* 8.12, b^* 9.60) and (L^* 7.41, a^* 1.23, b^* 12.83) for fresh, pulp and powder, respectively. The chroma (C^*) analyzed was 40.37, 12.57 and 12.89, respectively while hue angle (h°) was 83.69, 49.77 and 84.52, respectively while the browning index (BI) was 0.75, - 0.43 and 0.35 for fruit, pulp and powder respectively. Several value-added products such as pumpkin concentrate, jam, juice, syrup, chutney, confectioneries, bakery products, ready to cook instant food premixes as well as reconstituted products and weaning foods with improved vitamin A content and minerals.

Keywords: Ripe pumpkin, *Cucurbita moschata*, pumpkin pulp, pumpkin powder, β -carotene

1. INTRODUCTION


Pumpkin, also known as auyama, squash, or sambo, belongs to the family of cucurbits and is classified into the species *Cucurbita maxima*, *Cucurbita pepo*, *Cucurbita moschata*, and *Cucurbita mixta*. Worldwide production is approximately 27,643,932 tonnes which produced a

total of 2,042,955 tonnes/ ha with an average of 13.53 t/ha [1]. Ripe fruit is characterized by the hard outer cover, with yellow to orange-fleshed, firm texture and flavour depending on a different location. Based on the nutritional aspect, it contains low carbohydrates 8.8 % in comparison to other vegetables. Other nutrients like protein, fat, and fibre are less than two per cent. According to [2], it is the richest source of minerals such as potassium (439 mg), calcium (26 mg) and phosphorus (17 mg). Furthermore, it contains the highest amount of β -carotene which is converted to vitamin A. Apart from this, vitamin C, vitamin E, lycopene and dietary fibre were found in higher amounts by [3]. Many nutritional studies have shown that traditional vegetables, which are often overlooked by urban populations are an essential source of nutrients and vitamins for the rural population in low-income countries [4]. Since the fruit pulp has a low lipid concentration, lipids are mobilized and retained in the seeds, making the fruits a healthy food for people who are overweight. Excess fat intake has been linked to atherosclerosis, cancer and ageing in the cardiovascular system [5]. As a result, pumpkin diets should be recommended to minimize the risk of the aforementioned ailments in humans. Moisture content reduction in fruits and vegetables extends their shelf life, reduces enzymatic browning and prevents microbial spoilage. Blanching and KMS solution deeping enhanced the structural changes in the food that facilitate the dehydration process as well as enzymatic inactivation [6]. The aim of this study was to determine the effect of processing methods on the nutritional quality of ripe pumpkin (*C. moschata*).

2. MATERIALS AND METHODS

2.1 Preparation of ripe pumpkin powder

Ripe pumpkin pulp was prepared according to the method described by [7].



Cooking in the pressure cooker after adding water (@ (5 %))



Allow cooling



Grinding in a mixer cum grinder



Sieving



Ripe pumpkin pulp



Labelling



Storage in refrigerator

Flow chart 1: Unit operations for the preparation of pumpkin pulp

2.2 Preparation of ripe pumpkin pulp


Ripe pumpkin powder was prepared as per the method described by [8].

Ripe pumpkin fruit



Washing





Flow chart 2: Unit operations for the preparation of pumpkin powder

2.3 Chemical analysis

The moisture content (%), ash (%), protein (%) and minerals (iron mg/100 g) was determined as per the method suggested by AOAC [9]. Crude fibre (%) was analyzed as per [10], Crude fat (%) was determined using [11] method. Ranganna (2009) procedure was employed in scrutinizing β -carotene (mg/100 g), total carbohydrates (%) whereas total energy (Kcal/100 g) was calculated by the differential method as per [12] method. Ascorbic acid, titratable acidity and Total Soluble Solids (TSS) were determined as per the procedure given by [13], reducing sugars and total sugars were analyzed according to the method suggested by [14]. The colour of fresh, pulp and powder was measured in a Lovibond Colour Tintometer Model PFX-I series spectrocolourimeter in which RYBN colour units were obtained along with CIE readings i.e. L^* , a^* and b^* values. Each sample was measured three times for colour [15]. Changes in colour (ΔE), chroma (C^*), and hue angle (h^0) were calculated as per the formula proposed by Goswami *et al.* (2015) [16].

2.4 Data analysis

The chemical parameters were analyzed by Complete Randomized Design (CRD) and sensory evaluation was analyzed using Randomized Block Design (RBD) as described by [17,

18] respectively. The means were separated for comparison by Tukey's honest significant difference (HSD) and the statistical significance was defined as $p \leq 0.05$.

3. RESULTS AND DISCUSSION

3.1 Moisture (%)

The results obtained for chemical characteristics of ripe pumpkin, pulp and powder are depicted in Table 1. The findings revealed moisture content of 88.14 ± 1.34 , 88.93 ± 0.08 and 6.12 ± 0.05 per cent in pumpkin fruit, pulp and powder, respectively. The values analyzed are in the range given by [19] for pumpkin fruit, lower than the result of [8], higher than the observations of [20] and almost similar to the data of [21] for fruit. [7] noticed the value closer to the current data in the pulp while [22] recorded a range inclusive to present findings in the pulp. [23] gave a higher value in comparison to the current results in powder. while [20] reported a higher value in powder while [24] and [25] examined lower values as compared to the present study in powder.

3.2 Total soluble solids (°B)

The data presented in Table 1 observed total soluble solids to be 7.90 ± 0.03 , 5.40 ± 0.05 and 52.80 ± 0.32 °B in ripe pumpkin, pulp and powder, respectively. The findings are near to the results of [20] in fruit but lower than the values given by [7] and Dhiman *et al.* (2018a) [21] in pulp and powder, respectively.

3.3 Total sugars

The pumpkin fruit, pulp and powder contain 4.81 ± 0.11 , 3.42 ± 0.19 and 40.32 ± 0.75 per cent total sugars as per the current study. These findings are higher than the value reported [27] and near to the results of [8] in fruit. [7] has detected almost near results to current data in the pulp while [20] documented higher value in the pulp. Furthermore [25] studied a lower value than the present study in powder.

3.4 Reducing sugars (%)

Reducing sugars of 1.98 ± 0.05 , 2.15 ± 0.03 and 16.40 ± 0.05 per cent were noticed in pumpkin fruit, pulp and powder, respectively. These results are higher than the value noted by Rao and Sharma (2013) but lower than the observations of [8] in fruit while almost in confirmation with the results of [7] in the pulp. [20] found a higher value in contrast to the

present findings whereas, [25] revealed a lower value in comparison to current findings in powder.

3.5 Titratable acidity (%)

The value for titratable acidity recorded was 0.06 ± 0.01 , 0.05 ± 0.001 and 0.83 ± 0.01 per cent in pumpkin fruit, pulp and powder, respectively which is lower than the value noted by [26] in ripe pumpkin. The data is almost in line with the results of Rana (2020) for pulp while in conformity with the results of [20] for powder.

3.6 Crude protein (%)

The crude protein content perceived in pumpkin fruit, pulp and powder are 1.99 ± 0.27 , 1.72 ± 0.02 and 5.04 ± 0.06 per cent, respectively. The values analyzed in pumpkin fruit are higher than the findings of [21] while in the range of [19] but lower than the observations of [8]. In pulp, results are higher than the value of [7] but lower than the range detected by [22]. Usha *et al.* (2010) [23] and [20] have noticed higher values as compared to the present study in powder.

3.7 Crude fat (%)

The crude fat content was found to be 0.80 ± 0.32 , 0.51 ± 0.01 and 2.72 ± 0.27 per cent in pumpkin fruit, pulp and powder, respectively. The results are in the range given by [19] in pumpkin fruit while higher than the value of [8]. The fruit pulp exhibited a value in the range of [22] but higher than the data revealed by [7]. The fat content of powder is reflecting a higher value as compared to the findings of [23] and [20] but is lower than the values noted by [24] and [25].

3.8 Crude fibre (%)

The analysis of pumpkin fruit, pulp and powder showed the crude fibre content of 0.85 ± 0.05 , 0.49 ± 0.03 and 4.91 ± 0.01 per cent, respectively. The values for fruit have been found lower in comparison to the range of [19] while higher than the observations of Dhiman *et al.* (2018a and 2020) in fruit. Further, [7] noticed almost similar results to the present data in the pulp while [22] gave a higher range of results in contrast to current findings in the pulp. [23], [20], , (2019), [24] and [25] have evaluated lower values as compared to the current data in powder.

3.9 Total carbohydrate (%)

The data (Table1) represented the total carbohydrate content of 8.21 ± 0.26 , 8.06 ± 0.03 and 81.54 ± 0.94 per cent in pumpkin fruit, pulp and powder, respectively. The values

determined in the present study are higher than the range given by [27], but in the range of [19] while lower than the results of [21] in fruit. [23] and [25] have exhibited lower data as compared to present findings in powder.

3.10 Ascorbic acid (mg/100 g)

The ascorbic acid content in pumpkin fruit, pulp and powder was examined to be 14.22 ± 1.02 , 12.01 ± 0.07 and 10.26 ± 0.70 mg/100 g, respectively. The results are lower than the values reported in the studies of [26], [28] and [21] while almost near observations were found by [20] in pumpkin fruit. [7] observed a lower value in pulp as compared to present observations while [20] have given almost near data for powder.

3.11 β -carotene (mg/100 g)

The mean value revealed for β -carotene values of pumpkin fruit, pulp and powder was 15.81 ± 0.34 , 13.88 ± 0.06 and 10.77 ± 0.31 mg/100 g, respectively. The results for pumpkin fruit reflected a higher value as compared to as recorded by [26] but in the range of [19]. An almost similar value was analyzed by [29] while a somewhat lower amount was recorded by [21] in pumpkin fruit. The results of [7] reported a comparatively lower value in the pulp. [23] and [20] have obtained lower values in comparison to current observations in powder. The present findings exceed the results detected by [24] and [25] in powder.

3.12 Ash (%)

The results for ash content showed a value of 0.86 ± 0.06 , 0.78 ± 0.02 and 4.58 ± 0.26 per cent, respectively in pumpkin fruit, pulp and powder. These results are in the range of findings by [19] but higher than those recorded in the study of [29] and Dhiman *et al.* (2020) on pumpkin fruit. The current observations for pulp are in compliance with [7] and [22]. [23] and [24] evaluated higher values in powder as compared to the present study. The findings are also almost near to the data noted by [20] but higher than the finding reported by [25] in powder.

3.13 Total energy value (Kcal/100 g)

The total energy value calculated in pumpkin fruit, pulp and powder weres 44.60 ± 1.70 , 41.75 ± 0.12 and 351.16 ± 1.24 Kcal/100 g, respectively. The findings of [21] exhibited a higher value as compared to the present observations in pumpkin fruit.

Table 1: Nutritional characteristics of fresh and processed pumpkin fruit

| Parameters | Ripe pumpkin fruits | | |
|--------------|---------------------|--------------------|-------------------|
| | Fresh (WB) | Pulp (WB) | Powder (DWB) |
| Moisture (%) | 88.14 ± 1.34^b | 88.93 ± 0.08^a | 6.12 ± 0.05^c |

| | | | |
|--------------------------------------|---------------------------|---------------------------|----------------------------|
| Total Soluble Solids (TSS °B) | 7.90 ± 0.03 ^b | 5.40 ± 0.05 ^c | 52.80 ± 0.32 ^a |
| Total sugars (%) | 4.81 ± 0.11 ^b | 3.42 ± 0.19 ^c | 40.32 ± 0.75 ^a |
| Reducing sugars (%) | 1.98 ± 0.05 ^c | 2.15 ± 0.03 ^b | 16.40 ± 0.05 ^a |
| Titrateable acidity (%) | 0.06 ± 0.01 ^c | 0.05 ± 0.001 ^b | 0.83 ± 0.01 ^b |
| Crude protein (%) | 1.99 ± 0.27 ^b | 1.72 ± 0.02 ^c | 5.04 ± 0.06 ^a |
| Crude fat (%) | 0.80 ± 0.32 ^b | 0.51 ± 0.01 ^c | 2.72 ± 0.27 ^a |
| Crude fibre (%) | 0.85 ± 0.05 ^b | 0.49 ± 0.03 ^c | 4.91 ± 0.01 ^a |
| Total carbohydrates (%) | 8.21 ± 0.26 ^b | 8.06 ± 0.03 ^c | 81.54 ± 0.94 ^a |
| Ascorbic acid (mg/ 100 g) | 14.22 ± 1.02 ^a | 12.01 ± 0.07 ^b | 10.26 ± 0.70 ^c |
| β-carotene (mg/ 100 g) | 15.81 ± 0.34 ^a | 13.88 ± 0.06 ^b | 10.77 ± 0.31 ^c |
| Ash (%) | 0.86 ± 0.06 ^b | 0.78 ± 0.02 ^c | 4.58 ± 0.26 ^a |
| Total energy (Kcal/ 100 g) | 44.60 ± 1.70 ^b | 41.75 ± 0.12 ^c | 351.16 ± 1.24 ^a |

WB=wet basis and DWB= dry weight basis

The means sharing the same superscript letter in rows are not significantly different from each other(Tukey's HSD test, $p \leq .05$).

3.14 Colour

Colour is often used as an indication of the quality and freshness of food products. It is a critical parameter in defining the uses and acceptability of a product. The colour was recorded to be (L* 36.83, a* 4.44, b* 40.13), (L* 0.19, a* 8.12, b* 9.60) and (L* 71.41, a* 1.23, b* 12.83) for fresh, pulp and powder respectively (Table 2) and pictorial representation of the same is depicted in Fig 1, 2 and 3. Ripe pumpkin fruit has clear yellow to orange colour. Minimal changes to this attribute after processing is desirable [29]. a* and b* represent the redness and yellowness of the product, respectively while L* indicates the lightness. The increment of L* and a* values in powder and pulp means a more pure and intense yellow colour. In this case, positive values indicate red. The higher b* values indicated yellowness (Table 1). Therefore, yellowness was highly observed in fresh and powder than pulp due to enzymatic browning which occurs in pulp samples during preparation. The changes in redness and yellowness of pumpkin powder can be evaluated by chroma. The higher value of chroma was obtained in fresh (40.37) followed by powder (12.89) which indicated a more pure and intense colour [30]. Hue angle (h°) which is the dimension of the colour perceived was observed higher in powder (84.52), fresh (83.69) and pulp (49.77). The higher the hue angle the pure the colour perceived and vice versa. In pulp samples, enzymatic browning may be due to the effect of low hue value compared to fresh and powder. The browning index (BI) was 0.75, -0.43 and 0.35, respectively for fresh, pulp and powder. The positive value in fresh and powder indicate there was no colour defect but the negative value signifies a defect of colour due to enzymatic browning during processing.

Table 2: Colour of ripe pumpkin fruits

| Food material | Colour | | | | | |
|---------------|--------|------|-------|-------------|----------------|---------------------|
| | L* | a* | b* | Chroma (c*) | Hue angle (h°) | Browning Index (BI) |
| Fresh | 36.83 | 4.44 | 40.13 | 40.37 | 83.69 | 0.75 |
| Pulp | 0.19 | 8.12 | 9.60 | 12.57 | 49.77 | -0.43 |
| Powder | 71.41 | 1.23 | 12.83 | 12.89 | 84.52 | 0.35 |




Fig. 1. CIE readings of ripe pumpkin fruit




Fig. 2. CIE readings of ripe pumpkin pulp




Fig. 3. CIE readings of ripe pumpkin powder

4. CONCLUSION

The results have shown that steam blanching of pumpkin slices followed by deeping in Potassium Metabisulfite (KMS) solution before drying produces high quality and stable colour

of the final products. Processing pumpkin into pulp and powder will extend the keeping quality and enhance utilization into different value-added products after harvest hence food and nutritional security. This could lead to efficient and profitable utilization of ripe pumpkin fruit thereby ensuring reduction of postharvest losses. Pumpkin flour can act as an important food ingredient as a source of β -carotene for improving the body's immune system, vision and preventing Vitamin A deficiency (VAD), especially in young children. If pumpkin fruits will be processed immediately after harvest, post-harvest losses will be minimized, food and nutrition security will be improved as well as people's income through value-added products.

REFERENCES

1. Food and Agriculture Organization, FAO, 2018. FAOSTAT-statistics. Retrieved from. <http://www.fao.org/faostat/en/#data/QC>.
2. Rodríguez RR, Valdes R, Ortiz GS. Características agronomicas y calidad nutricional de los frutos y semillas de calabaza *Cucurbitasp*. Revista Colombiana de Ciencia Animal – RECIA. 2018;10(1): 86–97.
3. Ward E. Nutrients Your Diet May Be Missing. Available online <http://www.medicinenet.com/script/main/art.asp?article key=7671>. 2007.
4. Food and Agriculture Organization of the United Nations, Wild food plants: agrobiodiversity strategies to combat food insecurity and HIV/AIDS impact in rural Africa, Food and Agriculture Organization, 2003.
5. Chuwa C, Dhiman AK, Kathuria D, Mwita MA., Gautam S. Food Fibres: A solution to combat non-communicable diseases. *Nutrition and Metabolism: An Open Access*; 2020;DOI: 10.29011/NMOA-105.100005
6. Morais DR, Rotta EM, Sargi SC, Bonafe EG, Suzuki RM, Souza NE, Matsushita M, Visentainer JV. Proximate composition, mineral contents and fatty acid composition of the different parts and dried peels of tropical fruits cultivated in Brazil. *Journal of the Brazilian Chemical Society*. 2017;28:308-18.
7. Rana S. Development of muffins from ripe pumpkin (*Cucurbita moschata*). M Sc. Thesis. Department of Food Science and Technology. Dr YS Parmar University of Horticulture and Forestry, Nauni, Solan. 2020; pp.67-74.
8. Dhiman AK, Vidiya N, Attri S, Ramachandran P. Studies on development and storage stability of dehydrated pumpkin-based instant soup mix. *Journal of Applied and Natural Science* 2017;9:1815-20.
9. AOAC, 2012. *Official Methods of Analysis of the Association of Official Analytical Chemists*. 19th ed, Association of Official Analytical Chemists, Washington DC

10. AOAC, 2010. *Official Methods of Analysis of the Association of Official Analytical Chemists*. 18th ed, Association of Official Analytical Chemists, Washington DC
11. AOAC. 2009. *Official Methods of Analysis of the Association of Official Analytical Chemists* 6th edn International.. Inc, USA IL
12. AOAC, 2006. *Official Methods of Analysis of the Association of Official Analytical Chemists*. 18th ed, Association of Official Analytical Chemists, Washington DC
13. AOAC, 2004. *Official Methods of Analysis of the Association of Official Analytical Chemists*. 11th ed, Association of Official Analytical Chemists, Washington DC
14. Dubois M, Gilles KA, Hamilton JK, Rebers PT, Smith F. Colorimetric method for determination of sugars and related substances. *Analytical chemistry* 1956;28:350-56.
15. Ranganna S. *Handbook of analysis and quality control of fruit and vegetable products*. 2nd edn. Tata McGraw Hill Publishing Company Limited, New Delhi, India. 2009.
16. Goswami D, Gupta RK, Mridula D, Sharma M, Tyagi SK. Barnyard millet-based muffins: Physical, textural and sensory properties. *LWT-Food Science and Technology*. 2015;64(1):374-380.
17. Cochran WG and Cox CM. *Experimental Designs*. John Wiley and Sons, New York. 1967;pp. 171-217.
18. Mahony MO. Sensory evaluation of food. In: *Statistical Methods and Procedures*. Marcel Dekker Inc, New York. 1985; pp. 168-69.
19. Karanja JK, Mugendi BJ, Khamis FM, Muchugi AN. Nutritional Evaluation of Some Kenyan Pumpkins (*Cucurbita* spp.). *International Journal of Agriculture and Forestry*. 2014;4:195-200.
20. Dhiman AK, Babita K, Attri S, Ramachandran P. Preparation of pumpkin powder and seed kernel powder for supplementation in weaning mix and cookies. *International Journal of Chemical Studies* 2018a;6:167-75.
21. Dhiman AK, Chauhan P, Attri S, Kathuria D, Ramachandran P, Sharma A. Effect of pretreatment and drying methods on nutritional composition of ripe pumpkin (*Cucurbita moschata*). *International Journal of Current Microbiology and Applied Science* 2020;9:2536-52.
22. Nakazibwe I, Olet EA, Kagoro-Rugunda, G. Nutritional physico-chemical composition of pumpkin pulp for value addition: Case of selected cultivars grown in Uganda. *African Journal of Food Science* 2020;14:233-43.
23. Usha R, Lakshmi M, Ranjani M. Nutritional, sensory and physical analysis of pumpkin flour incorporated into weaning mix. *Malaysian Journal of Nutrition* 2010;16:379-87.

24. Khatib SE, Muhieddine M. Nutritional profile and medicinal properties of pumpkin fruit pulp. In: *The Health Benefits of Foods-Current Knowledge and Further Development*. IntechOpen. 2019;pp. 1-20.
25. Khan MA, Mahesh C, Vineeta P, Sharm, GK, Semwal AD. Effect of pumpkin flour on the rheological characteristics of wheat flour and biscuit quality flours. *Journal of Food Processing and Technology*. 2019;10:45-50.
26. Rao TVR, Sharma S. Nutritional quality characteristics of pumpkin fruit as revealed by its biochemical analysis. *International Food Research Journal*. 2013;20:2309-16.
27. Blessing AC, Ifeanyi UM, Chijioke OB. Nutritional evaluation of some Nigerian pumpkins (*Cucurbita spp.*). *Fruit, Vegetable and Cereal Science and Biotechnology* **5**:64-71.
28. Dhiman, A.K., Babu, N.G., Attri, S., Ramachandran, P., 2018b. Preparation of pumpkin pulp and effect of different preservation methods on chemical and sensory properties during storage. *Journal of Pharmacognosy and Phytochemistry*. 2011;7:943-49.
29. Thakur BR, Singh RK, Nelson PE. Quality attributes of processed tomato products. A Review. *Food Research International* 1996;12:375–401.
30. Pomeranz Y, Meloan C. “Food Analysis. Theory and Practice.” The Avi Pub. Co., Inc., London. 1971.