

Bioactive constituents and biochemical assessment of *Argemone mexicanana* linn and *Utrica dioica* leaf extracts using male albino rat

Abstract

Utrica dioica (stinging nettle) and *Argemone mexicanana* linn are medicinal plants used in most part of Nigeria in treatment of diseases. The present work evaluated bioactive constituents and biochemical effect of *Argemone mexicanana* linn and *Utrica dioica* extracts using male albino rat model. A total of eighteen male albino rats were used for the study. Group I served as control, Group II received 300mg/kg of *Utrica dioica* while Group III received 300mg/kg body weight of *Argemone mexicanana* linn. Preliminary phytochemical study showed the strong presence of flavoniod, tannin, saponin, alkaloid, cyanogenic glycoside and oxalate while the quantitative phytochemical and vitamin results showed high amount of these active compounds. Biochemical results from this study showed very low hepatotoxic and renal toxicity profile while the lipid profile indicates hypocholestrolemia potentials of the plants. This study suggest that even though there was low observed toxic response from intake of *Utrica dioica* (stinging nettle) and *Argemone mexicanana* linn, proper care should be taken regarding the dose used for medicinal purposes as continuous usage at high quantity may predispose the human health.

Key words: Bioactive, *Utrica dioica* , *Argemone mexicanana* ,biochemical

Introduction

In most developing countries of the world, there is an increasing demand for plant based remedies for human health problem. This upsurge in affinity for herbal remedy is probably due to easy accessibility, non-involvement of expert consultation, inexpensive as well as inadequacy of the primary health care (Udoamaka and Jose, 2014). Many medicinal plants are traditionally used in treatment of various human diseases. *Urtica dioica* (stinging nettle), is a perennial plant belonging to the genus *Urtica* and of the family *Urticaceae*. The stem is erect and the leaves are dark-green (Ahmed and Parasuraman, 2014). *Argemone mexicana* Linn a prickly annual herb belonging to the family papaveraceae and is widely distributed throughout the subtropical and tropical regions of the world. This work is aimed at profiling the bioactive constituents of these plants as well as their biochemical effect using male albino rats.



Fig 1. *Argemone mexicanna linn*



Fig 2 *Urtica dioica*

Method of analysis

Sample collection

Fresh leaf of *Urtica dioica* and *Argemone mexicanna linn* were harvested from Umuoda Amuzu in Aboh Mbaise L.G.A of Imo State. The samples were carefully washed with clean tap water before analysis.

Preparation of ethanol leaf extracts of *Utrica dioca* and *Argemone mexicanna linn*

Exactly 200g of the powdered plant were measured into a conical flask and 500ml of 70% ethanol were added and left at room temperature for 48 hours. The extract was filtered. The filtrate was evaporated to dryness on a water bath (50°C) to give the crude extract, which the mass was determined

Experimental design

Eighteen healthy male albino rats of about six weeks old were used for the study. The animals were randomly divided into three groups of six rats each. They were housed in standard cages and allowed to acclimatize to laboratory conditions for one week prior to commencement of experiment. The animals were allowed free access to commercial rat feed and water *ad libitum*. Group I served as control, Group II received 300mg/kg body weight of ethanol leaf extract of *Utrica dioca* while group III received 300mg/kg body weight of ethanol leaf *Argemone mexicanna linn*. At the end of the experiment which lasted for twenty eight days, the animals were sacrificed in accordance with the National Institute of Health guidelines for the care and use of laboratory animals (NIH, 1985). Blood samples were collected by cardiac puncture into anticoagulant free tubes with cork for biochemical analysis.

Determination of phytochemical constituents

In determination of preliminary qualitative phytochemical constituents, Adetugi and Popoola(2001) was adopted in determination of saponin and tannin while Sofaworo, 1993 was employed to determine alkaloid. Trease and Evans, (1989) was used to evaluate flavanoid and oxalate while the method of Stephen (1970) was used to determine cyanogenic glycoside.

Quantitative phytochemical composition

Obadoni and Ochuko(2001) was adopted in evaluation of saponin and oxalate content while Boham and Kocipai (1974) was used to determine flavonoid composition. The method of Harbone (1973) was employed in alkaloid and cyanogenic glycoside determination while Van-Burden and Robinson, (1981) was used to determine tannin content

Determination of vitamin composition

The method of AOAC(1975) was employed to determine riboflavin, tocopherol and retinol content while Barakat (1973) was used to determine thiamine, niacin and ascorbic acid content.

Biochemical analysis

Serum alanine aminotransferase (ALT) and serum aspartate aminotransferase activities were determined by the method of Reitman and Frankel (1957) while Alkaline phosphatase activity was determined by Bassey *et al.*, (1946) using commercial diagnostic kit. Serum total protein was determined by the method of Henry *et al.*, (1974) while lipid profile- Total cholesterol, high density lipoprotein, Low density lipoprotein and serum triglyceride was determined using Randox reagent kit by the method of Fossati and Prencipe, (1982). The assay of total bilirubin was done using the method of Jendrasik and Groff, (1938). Urea was done using Urease Berthlor method while creatinine was determined by the method of Heinegard and Triderstorm (1973).

Expression of results and statistical analysis

Results are presented as mean of triplicate determinations $\pm$ standard deviation. Statistical analysis was done using students package for social sciences version 20 computer software. Means were separated using One way analysis of variance.

Results

Table 1. Preliminary qualitative constituents of *Utrica dioca* and *Argemone mexicanna* linn

Saponin	+	+
Flavoniod	++	++
Alkaloid	++	+
Tannin	++	++
Cyanogenic glycoside	+	+
Oxalate	+	++

Table 2. Quantitative phytochemical constituents of *Utrica dioca* and *Argemone mexicanna* linn(%)

Saponin	8.84±0.32	2.01±0.01
Flavoniod	23.68± 0.33	14.22±0.02
Alkaloid	7.26±0.14	4.36±0.09
Tannin	14.31±0.31	12.82±0.08
Cyanogenic glycoside	1.79±0.01	3.32±0.20
Oxalate	8.53±0.01	7.53±0.21

Values are mean of triplicate determination ± standard deviation

Table 3. Vitamin composition of *Utrica dioca* and *Mexicanna linn*

Riboflavin	2.68±0.02	1.80±0.03
Thiamine	4.41±0.01	3.90±0.21
Niacin	19.00±0.04	7.12±0.00
Ascorbic acid	1.90±0.00	1.50±0.05
Tocopherol	13.45±o.11	11.8±0.02
Retinol`	4.40±0.03	3.80±0.01

Values are mean of triplicate determination ± standard deviation

Table 4: Liver integrity of rats administered *Utrica dioca* and *Argemone mexicanna linn* extract.

Groups	AST (IU/L)	ALT (IU/L)	ALP (IU/L)	Total Protein	Albumin (mg/dl)	Total bilirubin
--------	------------	------------	------------	---------------	-----------------	-----------------

				(mg/dl)		
Group I	61.51±0.46	33.94±0.12	83.21±0.07	9.13±0.56	7.01±0.01	0.19±0.10
Group II	62.62±0.91*	34.50±0.14*	84.07±0.09*	9.31±0.27*	7.41±0.05*	0.20±0.01*
Group III	63.03±0.04*	36.63±1.09*	82.01±0.04*	9.47±0.32*	7.65±0.15*	0.21±0.04*

Values are mean of triplicate determination ± standard deviation.

* indicate no significant difference from the control.

Legend:

Group I = Control

Group II = 300mg/kg body weight of ethanol leaf extract of *Utrica dioca*

Group III= 300mg/kg body weight of ethanol leaf extract of *Argemone mexicanna linn*

Table 5: Renal function of rats administered *Utrica dioca* and *Argemone mexicanna linn* extract

Groups	Urea (mg/dl)	Creatinin (mg/dl)
Group I	32.12±0.09	0.45±0.21
Group II	34.03±0.16*	0.47±0.23*
Group III	34.16±0.14*	0.49±0.25*

Values are mean of triplicate determination ± standard deviation.

* indicate no significant difference from the control.

Legend:

Group I = Control

Group II = 300mg/kg body weight of ethanol leaf extract of *Utrica dioca*

Group III= 300mg/kg body weight of ethanol leaf extract of *Argemone mexicanna linn*

Table 6. Lipid profile of rats administered *Utrica dioca* and *Argemone mexicanna linn* extract

Groups	TC(mg/dl)	HDL(mg/dl)	LDL(mg/dl)	TG(mg/dl)
Group I	91.02±0.85	30.29±1.41	21.31±0.04	77.13±0.6
Group II	80.12±1.71*	33.07±1.02*	16.16±0.15*	72.38±1.01*
Group III	79.05±1.98*	35.21±0.767*	14.01±1.21*	790.55±0.55*

Values are mean of triplicate determination ± standard deviation.

* indicate significant difference from the control

Legend:

Group I = Control

Group II = 300mg/kg body weight of ethanol leaf extract of *Utrica dioca*

Group III= 300mg/kg body weight of ethanol leaf extract of *Argemone mexicanna linn*

Discussion

Phytochemicals or plant chemical are bioactive non-nutrient plant compound that provide a range of activities against oxidative damage (Liu,2013). Findings from this study showed both qualitative and quantitative phytochemical constituents of *Utrica dioca* and *Argemone*

mexicana linn. The increased presence of flavonoids shows that extracts of these plants may have anti-inflammatory potentials. These phytochemicals have been reported to play an important role in detoxification of reactive oxygen species (Chandra *et al.*, 2009). Vitamins are antioxidants that play an important role in cellular defense strategy against oxidative stress (Fikriye and Omer, 2005). Findings from the study showed appreciable amount of vitamin in the two plants studied. This shows that extracts of these plants can serve as free radical scavenger. The liver is prone to xenobiotic-induced injury because of its central role in xenobiotic metabolism and analysis of liver enzymes such as AST, ALT and ALP have been reported as a rough proportionality of the extent of liver damage (Iweala and Osundiya, 2010). Friday (2004) noted that cell-derived enzymes have high activity in cells and spill in plasma when the cell is damaged or the enzymes produced in excess. Result of the liver enzymes (AST, ALT and ALP) shows no significant increase in their activities relative to control ($P < 0.05$). The observed normalcy in enzyme activity may have resulted from phytochemicals substances present in *Utricularia dioica* and *Argemone mexicana* linn. This shows there was little or no cellular impairment that may lead to loss of functional integrity of the liver. Total proteins are rough measures of protein status but reflect major functional changes in liver integrity (Iweala and Osundiya, 2010). The measurement of the level of total protein and total bilirubin is used to assay for the synthetic and excretory function of the liver. Total protein obtained from the present study showed no significant difference ($P < 0.05$) compared to control. This may as well be an indication of no impairment of hepatocyte since total protein is synthesized by the liver cells. Iweala and Osundiya, (2010) opined that alteration in the synthesis of total protein could have consequences in overall physiological function of the animals. Albumin result obtained from this study could further portray the medicinal efficacy of *Utricularia dioica* and *Argemone mexicana* linn as results showed no significant levels of albumin. This shows that the extracts did not induce much hepatocellular injury in animals as decrease in albumin level is reported to signify liver injury (Saba *et al.*, 2010). An increase in total bilirubin may indicate that the excretory function of the liver has been affected by damage caused to the liver or it could also indicate that a toxic substance might have competed and displaced the binding of bilirubin on albumin or the uptake of bilirubin is inhibited. Total bilirubin level was not significantly increased. This however suggests that intake of the extract may not be associated with diseases like jaundice. The Kidney is highly susceptible to toxicants for two reasons. A high volume of blood flows through it and it filters large amounts of toxins which can concentrate in the kidney tubules causing: decreased ability to excrete body wastes, inability to maintain body fluid and electrolyte balance and

decreased synthesis of essential hormones. The renal function result obtained from this study (Table 5) shows that there were no significant increase in urea and creatinine levels compared to control ($P < 0.05$). It has been reported that high blood urea is associated with increased tissue protein catabolism, excess breakdown of blood protein and diminished excretion of urea (Friday, 2004). Hence justifying the medicinal usage of these plants. Non significant increase in creatinine level relative to control may be an indication of none haepatocellular damage by the extract. Lipid profile results presented in Table (6) shows significant increase in HDL, decrease in total cholesterol, LDL and TG. Increased lipids in the blood are considered to increase the risk of myocardial infarction (Anuradha *et al.*, 2001). Cholesterol is an essential substance involved in many cellular functions, including the maintenance of membrane fluidity, production of vitamin D on the surface of the skin, production of hormones and possibly helping cell connections in the brain (Ugwu *et al.*, 2011). Increased level of circulating cholesterol and its accumulation in heart tissues are well known for their role in cardiovascular damage (Grover *et al.*, 2001). The decrease in cholesterol level is an indication that extract from the plants could be have hypocholesterolemic effect. Triglyceride accumulation in serum has been reported to play a major role in coronary heart diseases (Mahale *et al.*, 2008). The result show a decrease in triglyceride in a dose dependent ratio. High Density Lipoprotein (HDL) level is a very well known measure of cardiac health due to their strong inverse relationship with coronary artery disease. The protective role of HDL-C against CVD has been suggested to occur in several ways such as serving as transport particles for excess cholesterol to the liver, where it is converted into bile acids and excreted (Ugwu *et al.*, 2011). There was a significant increase in the HDL levels of groups II and III relative to control ($P < 0.05$). This could be as a result of some bioactive compounds in *Utrica dioca* and *Argemone mexicanana* linn which may have increased the HDL level which is responsible for transport of cholesterol out of the cell. The results of lipid profile suggest that *Utrica dioca* and *Argemone mexicanana* linn extracts may have cholesterol reducing properties which is necessary in prevention of arteriosclerosis.

Conclusion

The result of this study has shown that *Utrica dioca* and *Argemone mexicanana* linn ethanol extract at 300mg per kg body does not show evidence of cumulative toxicity and may as well be useful in reduction of hypercholesterolemia. This however may suggest that increased and prolonged intake of these plant extracts may have adverse health effects.

References

- Adetugi, S and Popoola, C (2001). Food analysis and instrumentation (Theory and Practical) 1st edition, Napthal print, Surulere, Lagos Nigeria. Pp 140-160
- Anuradha, C.V and Ravikumar, P (2001). Restoration on tissue antioxidants by fenugreek seeds (*Trigonella Foenum Graecum*) in alloxan-diabetic rats. *Indian Journal of Physiology and Pharmacology*, **45**:408–20.
- AOAC(1975) Official method of analysis. 17th edition. Washington DC
- Barakat, M.Z., Shehab, S.K., Darnish, N and Zahermy, E.I (1993). Determination of ascorbic acid from plants. *Analyst Biochemistry*, **153**: 225-245
- Bassey, O.A., Lowery, O.H and Brook M (1946). One point colometric method of determining alkaline phosphatase in serum or plasma. *Journal of Biological Chemistry*, **164**:321.
- Boham, A.B and Kocipai, A.C (1974). Flavonoid and condensed tannin from leaves of Hawaiian vaccinium vaticulum and vaccinium calycinum. *Pacific Science*, **48**: 458-463
- Chandra R., Bharagava R.N., Yadav S and Mohan D (2009). Accumulation and distribution of toxic metals in wheat (*Triticum aestivum* L) and Indian mustard (*Brassica campestris* L) irrigated with distillery and tannery effluents. *Journal of Hazard Matter*. **162**: 1514-1521.
- Disbrey, D.D and Rack, J.H (1970) Histological laboratory methods. Livingstone Edinburgh. Pp 56-128.
- Ekpenyong, C. E, Akpan, E. E and Udoh, N. S (2012). Phytochemistry and Toxicity Studies of *Telfairia Occidentalis* Aqueous Leaves Extract on Liver Biochemical Indices in Wistar Rats. *American Journal of Medicine and Medical Sciences* , **2**(5): 103-110..
- Fossati, P and Prencipe, L (1982). Serum triglycerides determined colorimetrically. *Journal of Clinical Chemistry*, **28**: 2077-2080.
- Friday, E.U. (2004): Clinical uses of enzymes, In: Conceptual Enzymology, Innolox printers and publishers, Calabar. Pp. 190-212.
- Grover, J. K, Yadav, S and Vats, V (2002). Medicinal plants of India with anti-diabetic potential. *Journal of Ethnopharmacology*, **81**:81–100.
- Harbone, J.B (1973). Phytochemical methods. A guide to modern techniques of plant analysis. Chapman and Hall London. PP 19-34

- Heinegard, D and Tiderstorm, K (1973).Determination of serum creatinin by direct colorimetric method. *Journal of Clinical Chemistry Acta*, **43**:305-310
- Henry, R.F., Shrin N and Winkelman (1971).Clinical chemistry principles and techniques 2nd Eds, Herper and Row Hagerstown.
- Iweala, E.E and Osundiya A.O (2010). Biochemical, haematological and histological effects of diatary supplementation with leaves of *Gnetum africanum* Welw on paracetamol-induced hepatotoxicity. *International Journal of Pharmacology*, **6**(6): 872-879.
- Jendrassik, L and Groff, P (1938).Colorimetric method for measurement of birubin. *Biochemistry Journal*, **297**:61-89
- Liu, HR (2013). Health benefits of fruits and fruits and vegetables are from additive and synergistic combinations of phytochemicals. *Am J. Clin Nutr.* **78**:517-520
- Mohala, D.S.,Dewani, A.P., Saoji, A.N and Yeole, P.G (2008).Antihyperlipidemic activity isolated constituents from *Lageneria* species in albino rats. *International Journal of Green Pharmacology*, **2**: 807-817.
- NIH(1985).National Research Council Guide for the care and use of laboratory animals. Publication no: 85-123(rev). National Institute Health, Bethesda, M.D.
- Obadoni, B and Ochuko, P (2001). Phytochemical studies and comparison efficacy of the crude extract of some homeostatic plants in Edo and Delta State. *Global Journal of Pure and Applied Science*, **8**: 203-208
- Reitman, S and Franklin, S (1957). A colorimetric method for the determination of serum glutamic oxaloacetate and glutamic pyruvic transaminases. *American Journal of Clinical Pathology*, **28**: 56-58.
- Saba, A.B., Oyagbemi, A.A and Azeez, O.I (2010).Amelioration of carbon tetrachloride-induced hepatotoxicity and haemotoxicity by aqueous leaf extract of *Cnidioscolus aconitifolius* leaf in rats. *Nigerian Journal of Physiological Society*, **25**: 139-147.
- Sofowara, A (1993).Medicinal plants and traditional medicine in Africa. Spectrum books Ltd Ibadan, Nigeria. Pp 191-289.
- Stephen, A (1970). Phytochemical methods. Chapman and Hill London. P113
- Tease, G.E and Evans, W.C (1989). Textbook of Pharmacology. 12th edition Baliene Tindall: London

Udoamaka, F. E and Jose, M. P (2014). The use of plants in the traditional management of diabetes in Nigeria: Pharmacological and toxicological considerations. *Journal of Ethnopharmacology*, **155**:857-924

Ugwu, C. E., Olajide, J. E., Alumana, E. O and Ezeanyik, L. U. S(2011). Comparative effects of the leaves of *Vernonia amygdalina* and *Telfairia occidentalis* incorporated diets on the lipid profile of rats. *African Journal of Biochemistry Research* , **5**(1): 28-32.

Van-Burden, T.P and Robinson, W.C (1981).Formation of complexes between protein and tannin acid. *Journal of Agriculture and Food Chemistry*, **1**:77-82

UNDER PEER REVIEW