

Assessment of surface and groundwater quality for suitability of irrigation purposes collected from some selected sources of Eastern India

Abstract

The assessment of surface and groundwater for irrigation purpose was an important step towards sustainable water resource management for agricultural crops in the Eastern part of India. Therefore, the study was conducted during the pre-monsoon period in the month of April-May, 2019 to monitor the quality of different sources of water for assessing its suitability for irrigation purposes in Odisha state of eastern India. Twenty seven surface water and twenty one groundwater samples were collected and analyzed for chemical parameters. The suitability of the surface and groundwater for irrigation were evaluated based on pH, residual sodium bicarbonate (RSBC), electrical conductivity (EC), sodium adsorption ratio (SAR), Adjusted SAR (Adj. SAR), residual sodium carbonate (RSC), soluble sodium percentage (SSP), permeability index (PI), Kelly's ratio (KR), Chloride (Cl^-), boron (B), nitrate (NO_3^-), iron (Fe), Manganese (Mn), zinc (Zn), copper (Cu), chromium (Cr) and cadmium (Cd). Results showed that three rivers Mahanadi, Brahmani and Bansadhara were found medium salinity ($\text{EC } 0.25\text{--}0.75 \text{ dSm}^{-1}$) and all the rivers have no sodicity problem with RSC within the permissible limit of 2.5 meqL^{-1} , and free of chloride and boron toxicity. All the surface water samples were safe with respect to Zn (2 mgL^{-1}), Cu (0.2 mgL^{-1}), Fe (5 mgL^{-1}), Mn (0.2 mgL^{-1}) and Cr (0.1 mgL^{-1}). Four rivers were exceeded the maximum permissible limit of Cd. Groundwater samples collected from the three farms of Odisha University of Agriculture and Technology (OUAT) at Bhubaneswar were found pH below the permissible limit of 8.5. Four groundwater sources at Sambalpur, Rourkela, Gajapati and Ganjam were found medium salinity ($\text{EC } 0.25\text{--}0.75 \text{ dSm}^{-1}$) which could be safely used for irrigation with management. In other cases, the salinity level was safe ($\text{EC } <0.25 \text{ dSm}^{-1}$). Groundwater sample of Putibandha, Sambalpur, Rourkela and Ganjam where the salinity measured medium in terms of EC also have sodicity problem. The water of Rourkela and Ganjam has RSC more than safe limit but less than the maximum permissible limit (MPL). But RSC in Groundwater of Sambalpur exceeded the MPL with KR falling in the bad category and SSP in doubtful category. All the groundwater samples were found below the toxicity level and safe with respect to Cl^- , NO_3^- , B, Zn (2 mgL^{-1}), Cu (0.2 mgL^{-1}), Fe (5 mgL^{-1}), Mn (0.2 mgL^{-1}) and Cr (0.1 mgL^{-1}) where Cd content of three sources exceeded the MPL (OUAT Agronomy Farm 0.011 mgL^{-1} , Putibandh of Sambalpur 0.011 mgL^{-1} and Rourkela 0.010 mgL^{-1}). It is concluded that all the surface water samples collected from nine different rivers are found good quality and suitable for irrigation purpose except some specific limitations associated with certain sources which could be managed. Among all the groundwater sources, groundwaters of Putibandh area in Sambalpur district are found very poor in quality and unsuitable for irrigation.

Keywords : Surface water . groundwater . quality . irrigation.

Introduction

Water is a most important natural resource; it is an essential human need and is a valuable national asset. In addition to drinking, it is required for other human activities like, agriculture, industry, bathing, cooking, washing, recreation, navigation, fisheries etc [1, 2]. Groundwater plays an important role all over the world for the survival of both flora and fauna. India is one of the largest users of groundwater, particularly for drinking and agriculture purposes [3]. Groundwater has become the major source of water use in the agricultural sector in many countries where river and drainage systems are not sufficient. Therefore, poor groundwater quality for

irrigation purpose is a matter of worry in recent years [4]. The groundwater quality assessment based on different agriculture indices has been studied in different parts of world [5, 6]. There are a number of reports on the assessment of groundwater quality from Cameroon [7], Ghana [8], Bangladesh [9], Tamil Nadu [10, 11, 12, 2], Bhatina, South west Punjab. [13], Madhya Pradesh [14, 15]) and Sant Ravidas Nagar, Bhadohi, Uttar Pradesh [16]. In Uttarakhand, irrigation water quality assessment has been done in Doon Valley [17], Nainital [18], Almora [19, 20], Haridwar [21], Udham Singh Nagar [22]. In Odisha, physico-chemical analysis of surface and ground water of Bargarh district [23], Keonjhar city [24]. Surface water quality is a very sensitive and global environmental issue that is important for long-term economic development and environmental sustainability [25, 26, 27]. Awareness and attention to water irrigation quality have increased worldwide in recent years, and new approaches have been developed to achieve the sustainable management of water resources [28, 29]. In the same context, the shortage of water resources has become a big problem in many countries, particularly under continued population growth, accelerated industrialization, rapid urbanization, and global climate change [30, 31]). Therefore, water scarcity and sustainable irrigation water management have become global challenges for sustainable agriculture development in order to produce sufficient food to satisfy the population's food requirements [32, 33, 34]. In Odisha, there are mostly three sources of water rivers, surface storage and ground water. Odisha is neither abundant nor certain about its water resources and it is not a good sign for the future. According to the source of water, the quality and characteristics vary mainly due to varying geology and climate. On a local level also there may be differences due to weather and the source from surface water or groundwater with varying geology. The plant growth is affected due to the chemical parameters of irrigation water through toxicity and deficiency directly and by altering the availability of nutrients indirectly. Anthropogenic activities within river basins, erosion, and atmospheric depositions are also the main negative impacts on the water quality of most the reservoirs [35]. Anthropogenic influence as well as natural processes degrade surface waters and impair their use for drinking, industrial, agricultural and recreational purposes [36]. Urban runoffs and sewage disposal in the catchment area of rivers also contribute to poor water quality [37]. Industrial wastewater, runoff from the agricultural lands and disposal of municipal sewage are the most vulnerable for water pollution [38]. The soil salinity is also an important factor for the determination of water quality. A study on quality of major surface water and ground water sources is therefore very important for managing the irrigation for the present situation. To the best of our knowledge, no comprehensive study has been made on water quality of major rivers and ground water used for irrigation in Odisha. Therefore, in the present investigation, an attempt has been made to assess the quality of different water resources and classification of water based on their suitability for irrigation purpose collected from different places of Odisha.

Materials and methods

Study area

Odisha is located on the eastern coast of India, between 17°31" and 22°31" N latitude and 81°31" and 87°31" E longitude. It covers 155,707 km², which is about 4.74% of the area of India. The climate is tropical with 1450mm average rainfall. The state is spread over an area of 155,707 km² and 1030 km from north to south and 500 kilometers from East to West. Its coastline is 480 km long. The state has 30 districts. Locations of water sample collection from different river basins are decided based on the physiographic data and maps of rivers, and according to its place of tributaries where it flows and covers maximum area. Physiographically, Odisha consists of coastal plains, central plateaus, central hilly regions, flood plains, and uplands.

Sampling and analysis

Surface water samples were collected during April-May, 2019 from middle reach of nine major rivers *viz.* Mahanadi, Rushikulya, Brahmani, Subarnarekha, Kolab, Bansadhara, salandi, Bahuda, and Ib of Odisha. Thus, Twenty seven samples were collected from nine surface water sources. Similarly, groundwater samples were collected at two hours interval from seven borewells, three borewells of Odisha University of Agriculture and Technology Research Farm *viz.* Agronomy Farm, Central Farm and Orchard, one each at Sambalpur, Rourkela, Gajapati and Ganjam districts of Odisha. Twenty-one groundwater samples (borewell) were collected in pre-monsoon season in the year 2019 from different sites. The samples of surface and ground water were collected in pre-washed polyethylene narrow mouth bottles of 500 ml capacity and stored (three times rinsed with same water before collecting the samples). Location of sampling sites of surface water and groundwater were

measured by using global positioning system (GPS) for most of the samples (Fig. 1). The method involved in analysis of water samples are depicted in Table 1 and 2.

Statistical interpretation

Entire data obtained in the present study was analyzed using completely randomized design (CRD). To evaluate significant differences among the sites for all water quality variables, data of nine treatments of different rivers and three replication of each river and seven treatments of different groundwater samples and three replication of each were analyzed using one-way analysis of variance (ANOVA) at 0.05% and 0.01% level of significance [39]. Data's for different parameters of water samples were presented as mean values.

Results and discussion

The results obtained from laboratory analysis of various chemical parameters and calculated indices values collected from all the sources and locations are given in Table 3 to 11 and Supplementary data Table 1 to 2. To ascertain the suitability of collected surface and groundwater samples for irrigation purposes was discussed on comparison with recommended standard values given by different authors.

pH

In the present study area, the pH showed that all the surface water collected from canals and rivers at different points varied from 6.23 to 8.05 with a mean of 7.29 (Table 3), and 5.68-8.10 for groundwater with a mean of 7.04 (Table 4). Highest pH of 8.02 was measured in the water of Ib river and lowest 6.51 for the river near Janiguda farm, Kolab. Among the groundwater samples, the highest pH of 8.02 was found in the bore well water of Sambalpur, Putibandh area and lowest of 5.97 measured in the borewell, water of Agronomy farm, OUAT, Bhubaneswar. The pH of water is an important index of hydrogen ion activity and it is the resulting value of the acid-base interaction of several minerals and organic components in water. pH is an important ecological factor and universally to express the intensity of the acid and alkaline condition of the water samples. According to Food and Agriculture Organization (FAO) recommendation the permissible limit of pH of irrigation water is 6.5-8.5 [40, 41]. All the surface water samples collected in our study had pH between 6.23 and 8.05 and ground water, 5.68 to 8.1 (Fig. 2). The present investigation showed that all the surface water sources tested had pH value within the safe limit except the water of Kolab which had pH of 6.47 which was less than the permissible limit of 6.5 and ground water used for irrigation in Central Farm and Agronomy Farm of OUAT have pH less than the permissible limit of 6.5. But all other samples collected from industrial and urban areas had alkaline and significantly higher pH (7.48-8.02).

Electrical Conductivity (EC)

The EC values of water sample during monitoring periods ranged in between from 0.068-0.705 dSm⁻¹ and mean is 0.25 dSm⁻¹ of surface water and 0.045-0.723 dSm⁻¹ with mean 0.31 dSm⁻¹ of groundwater (Table 3 and 4). Highest EC of 0.57 dSm⁻¹ was measured in the water of Bramhani river at Rourkela-3 (sector-7) of Sundergarh district and lowest of 0.07 dSm⁻¹ in the water of Salandi river collected at Keonjhar and among all groundwater sources highest EC of 0.70 dSm⁻¹ was found in Putibandh area of Sambalpur and lowest of 0.06 dSm⁻¹ in the Orchard of OUAT at Bhubaneswar. The permissible range of EC is 0.25-0.75 dSm⁻¹ [42]. So, all the samples fall into C1 (low salinity) and C2 (medium salinity water some leaching sensitive crops) classes of salinity hazard classification by USDA [43] (Table 7).

Carbonate (CO₃²⁻) and Bicarbonate (HCO₃⁻)

The presence of CO₃²⁻ and HCO₃⁻ ions in irrigation water has a significant influence on the concentration of calcium and magnesium content of water and soil permeability after its application. Results showed that CO₃²⁻ was absent in almost all the water samples except few groundwater samples where it ranged from 0.99 to 5.40 meq L⁻¹ with a mean of 1.12 meq L⁻¹ (Table 3 and 4). Highest carbonate content of 4.59 meq L⁻¹ was

found in groundwater samples of Putibandh area of Sambalpur district. The HCO_3^- ranged from 0.99 to 2.97 meq L^{-1} with a mean of 2.09 meq L^{-1} for surface samples (Table 4). In case of ground water samples, HCO_3^- varied from 0.99 to 2.97 meq L^{-1} with a mean value of 1.46 meq L^{-1} (Table 6). Alkalinity in water is mainly due to CO_3 , HCO_3 and OH^- content. Carbonate content should be between 0-1 meqL^{-1} [6]. There was no carbonate in all the surface water and ground water except Samabalpur (Putibandh), Rourkela -3 (sector-7) and Gajapati (Paralakhemundi) had higher mean CO_3^{2-} concentration with highest of 4.59 meqL^{-1} measured at Sambalpur which is more than the safe limit 1.0 meqL^{-1} . These samples had also higher pH values.

Basic Cations (Na^+ , K^+ , Ca^{2+} and Mg^{2+})

In this study, four cations which have a direct and indirect influence on quality of irrigation water have been measured and the analysis results of all surface water and groundwater have been presented in Table 3 and 4. The concentration of four cations viz. Na^+ , K^+ , Ca^{2+} and Mg^{2+} ranged from 0.09 to 2.54, 0.018 to 0.65, 0.23 to 1.2, 0.09 to 1.2 meqL^{-1} with respective mean values of 0.59, 0.078, 0.66, 0.35 meqL^{-1} for surface water and ranges of 0.112 to 3.04, 0.005 to 0.078, 0.24 to 1.72, 0.13 to 0.54 meqL^{-1} with respective mean values 0.98, 0.037, 0.861, 0.28 meqL^{-1} in groundwater. In irrigation water, the permissible limit for Na^+ , K^+ , Ca^{2+} and Mg^{2+} are 200, 30, 80 and 35 mg L^{-1} [44, 45].

Specific Elements/Ions Chloride (Cl^-)

The chloride content ranges from 21.27-67.36 mgL^{-1} with a mean of 36.37 mgL^{-1} for surface water and 3.54 to 60.26 mgL^{-1} with a mean of 39.98 mgL^{-1} for groundwater samples with highest concentration found in Kolab river near Janiguda of Koraput, lowest in Subernarekha river of Mayurbhanj. Among the groundwater samples collected highest was found in Rourkela-3 (sector-7) and lowest in the groundwater of OUAT Horticulture orchard, Bhubaneswar (Table 3 and 4). The chloride concentration serves as an indicator of pollution by sewage, and industrial effluents. In this study all the samples are within safe limit based upon suitability classification on the basis of chloride concentration ($< 10 \text{ meqL}^{-1}$ or 350 meqL^{-1}). Chloride content in well water of Behampur (11.52 meqL^{-1}), Jaipatna (21.20 meqL^{-1}), Odagaon (10.37 meqL^{-1}), Puri (18.44 meqL^{-1}) canal water from Soro (10.37 meqL^{-1}) pond water of (21.2 meqL^{-1}) are found unsuitable for irrigation purposes[13].

Boron (B)

B content ranges from 0.0225-0.934 mgL^{-1} with a mean of 0.28 mgL^{-1} for surface water (Table 3) and 0.043-0.912 mgL^{-1} with a mean of 0.26 mgL^{-1} for groundwater samples (Table 4). The highest concentration was found in Salandi river (Keonjhar) and lowest in Rushikulya river among surface water samples, and groundwater samples highest was found in the water of OUAT Agronomy farm at Bhubaneswar and lowest in Putibandh of Sambalpur district. B is also present in irrigation water as unionized boric acid expressed as boron element in mgL^{-1} . Sensitivity to boron encompasses a wide variety of field and tree crops, although fruit, nut and berry crops are particularly sensitive. In the present investigation, there is no B toxicity as all the samples had boron less than 2 mgL^{-1} [46].

Nitrate (NO_3^-)

Nitrate content ranges from 0.98-3.23 mgL^{-1} with a mean of 2.10 mgL^{-1} for surface water (Table 3) and 2.96-8.2 mgL^{-1} with a mean of 4.75 mgL^{-1} (Table 4) for groundwater samples with the highest concentration found in Ib river, lowest in Salandi river, Keonjhar. Among ground water samples highest was measured in the water of the bore well in Horticulture Orchard of Bhubaneswar and lowest in the groundwater of Rourkela -3 (sector-7). In the present study, all the samples are safe with respect to NO_3^- ($< 30 \text{ mgL}^{-1}$) [46].

Micronutrients and Heavy metals

The content of trace and heavy metals (Zn, Cu, Fe, Mn, Cr and Cd) in water sources were analysed which is presented in Table 3 and 4. The results revealed that Cr concentration was non-detectable in AAS in

both surface and groundwater samples. The ranges of Zn, Cu, Fe, Mn and Cd were 0.002- 0.043, 0.032-0.052, 0.036-0.184, 0.01-0.056 and 0.001-0.035 mgL⁻¹ respectively in surface water and 0.007-0.044, 0.026-0.045, 0.033-1.09, 0.01-0.11 and 0.007- 0.012 mgL⁻¹ in groundwater samples respectively. All the water samples both surface and ground water were below the toxicity level and safe with respect to Zn, Cu, Fe, Mn, Cr except Cd which is more than permissible limit for the Kolab, Bahuda, Rushikulya and Mahanadi river and from ground water samples Agronomy field, OUAT, Rourkela-3 sector-7 and Putibandh of Sambalpur samples (Fig. 7).

Sodium Adsorption Ratio (SAR)

Results showed (Table 5 and 6) that SAR values varied from 0.16 to 2.73 with mean 0.79 for surface water and 0.18 to 3.26 with a mean of 1.33 for groundwater. The suitability of the well water samples was evaluated by determining the Sodium Adsorption Ratio value which is an expression of concentration of Na⁺ relative to Ca²⁺ and Mg²⁺ and they were categorized under different classes based on salinity and alkalinity hazards. U.S Salinity laboratory proposed that SAR as more reliable criteria for evaluating Na hazard in irrigation water. According to Richards [47], water samples having SAR 10-18 are of medium sodium hazard and these waters cannot be used for fine textured soils. Water with SAR less than 10 has no sodium hazard. According to Richards, SAR values measured in the sampled waters in surface water and ground water was found no sodicity. All the samples are within the permissible range of FAO, and all the samples are found to be in S1 class of sodicity hazard classification of USSL (1954) as all the values are within 0-10. Ground water of Sambalpur having EC 0.70 dSm⁻¹ and SAR 3.20 is classified under moderate sodicity category as per the guidelines [48, 46].

Adjusted SAR (SAR_{adj.})/ Adj.R_{Na}

Results showed that SAR_{adj.} (Table 5 and 6) ranged from 0.12 to 2.65, with a mean of 0.73 for surface water and 0.13 to 3.18 with a mean of 1.01 for groundwater. In the present study, all the surface and groundwater was found in S1 category according to the guidelines [49]. Adjusted SAR (SAR_{adj.}) is a better criterion for assessing the sodicity effect of irrigation water than SAR. Irrigation of Ca-rich or Mg-rich soil with water containing carbonate and bicarbonate ions will form insoluble calcium and magnesium carbonates resulting in a decrease in the concentration of Ca and Mg used in the SAR calculation. This leads to an increase in the value of measured SAR which does not show the real picture of sodicity under such situation. Irrigation water contains sufficient quantities of sulphate and bicarbonate ions to produce precipitation of calcium sulphate and calcium carbonate that remove calcium from solution and hence markedly increase sodium hazards. SAR under this situation may not give the correct sodium hazard since it is the SAR_{sw} value rather than the SAR_{iw} value that affect the ESP of the soil and an eventually the soil permeability. Hence, a new index called adjusted SAR [50] and is calculated from procedure [51] is recommended but both the older SAR procedure and the new adjR_{Na} are acceptable, with a preference expressed towards the adjR_{Na} because it and the Ca_x of Supplementary Table 3 offer a better insight into the change in calcium in the soil-water due to addition by dissolution of calcium from soil carbonates and silicates, or loss of calcium from soil-water by precipitation as carbonates [46].

Permeability Index (PI)

The PI values ranged from 65.71 to 305.38%, with mean 143.28% for surface water and 77.79 to 160.27% with mean, 112.19% for groundwater (Table 5 and 6). Highest and lowest PI was found in Subarnarekha river and Bahuda river respectively among surface water samples and among groundwater samples highest and lowest PI were measured in the groundwater collected the Central farm, OUAT, Bhubaneswar and Rourkela-3 (sector-7), respectively. The Permeability Index (PI) value is used to evaluate the sodium hazards of irrigation water. High sodium in the irrigation water can cause severe soil permeability problem. Permeability is affected not only by high sodium but also by CO₃²⁻ and HCO₃⁻ content in water. Long term irrigation affects the permeability of the soil due to presence of Na⁺, Ca²⁺, Mg²⁺ and HCO₃⁻ ions in water. Therefore, the PI values can be effectively used to determine the suitability of surface water and groundwater to be used for irrigation purpose. Water can be classified as class I, class II and class III. Class I is excellent when PI more than 75%,

class II is good with PI 25-75%, and Class III unsuitable when PI is less than 25% [52, 53]. All the samples are fall in class I (more than 75%) category of classification.

Residual sodium carbonate (RSC)

Results showed that the range of RSC was found to be 0.75 to 2.53 meqL⁻¹ with a mean of 1.075 meqL⁻¹ for surface water and -0.25 to 4.49 meqL⁻¹ with a mean of 1.44 meqL⁻¹ for groundwater (Table 5 and 6). It is used to predict the additional sodium hazard associated with CaCO₃ precipitation involve a calculation of the residual sodium carbonate. RSC is another alternative measure of the sodium content in relation to calcium and magnesium. The proportion of bicarbonate ion higher than calcium ions is considered undesirable because after evaporation of irrigation water bicarbonate ions tend to precipitate calcium ions. Therefore, the effect of bicarbonate together with carbonates evaluated through RSC is a better indicator of sodium hazard of irrigation water. Two samples collected from this river from sites at Mayurbhanj are found to be above the permissible limit out of all surface water samples. For ground water samples the highest RSC of 4.49 meqL⁻¹ was found in Putibandh area, Sambalpur district. Samples from Putibandh found above the permissible limit of 2.5 meqL⁻¹ [54]. Samples from river Mahanadi, Brahmani, Subarnarekha, Ib and groundwater of Rourkela and Gamjam found to be above safe limit (1.25 - 2.5 meqL⁻¹) but within maximum permissible limit, as shown in Table 7 and Fig. 4.

Residual Sodium Bicarbonate (RSBC)

RSBC ranges from 0.21-2.74 meqL⁻¹ with a mean of 1.43 meqL⁻¹ for surface water and -0.73–1.95 meqL⁻¹ with a mean of 0.68 meqL⁻¹ for groundwater samples presented in Table 5 and 6. Among surface water samples highest RSBC was measured in Subarnarekha river, lowest in Bahuda river, and among groundwater samples highest found in Gajapati and lowest at Putibandh, Sambalpur. Since carbonate ions do not occur very frequently in appreciable concentrations, and as bicarbonate ions do not precipitate magnesium ions, the residual sodium bicarbonate was calculated [55, 56] suggested that alkalinity hazard should be determined through the index called Residual Sodium Bicarbonate (RSBC). Water with RSBC <5, 5–10 and >10 meqL⁻¹ is considered safe, marginal, and unsatisfactory, respectively. In the present study area, all the samples are within safe limit and are classified as low alkaline water (< 2.5 meqL⁻¹).

Soluble Sodium Percentage (SSP)

Results presented in Table 5 and 6 reveal that SSP ranges from 13.33% to 65.98%, with a mean 36.73 % for surface water and 13.49% to 654 % with a mean of 44.16% for groundwater. Highest SSP from surface water samples found in Bramhani river of Banei and lowest in Bahuda river of Gajapati and from all the groundwater samples, highest SSP found in Putibandh area, Sambalpur and lowest in Horticulture orchard of OUAT at Bhubaneswar. Excess of sodium ions characterizes the water as saline or alkaline depending upon its occurrence in association with chloride/ sulphate or carbonate/ bicarbonate ions. The quality of irrigation water used to be evaluated with respect to sodium based on soluble sodium percentage (SSP) since a high value indicates soft water and low value hard water. Classification of irrigation water based on SSP (Table 7) [57]. In the present study, highest SSP from surface water samples found in Bramhani river of Banei and lowest in Bahuda river of Gajapati and from all the groundwater samples, highest SSP found in Putibandh area, Sambalpur and lowest in Horticulture orchard of Odisha University of Agriculture Technology at Bhubaneswar. All the surface water and ground water samples fall in excellent, good and permissible range of SSP except Putibandh of Sambalpur (Table 7; Fig. 5). High SSP reduces permeability of soil and eventually results in soil with poor conditions of drainage [58].

Kelley's Ratio (KR)

Results showed that KR content ranges from 0.11 to 1.46 with a mean of 0.55 for surface water and 0.13 to 1.78 (Table 5 and 6) with a mean of 0.78 for groundwater samples. The highest value of KR was found in Rushikulya river, lowest found in Bahuda river among all surface water samples, and among groundwater highest found in Putibandh, Sambalpur and lowest in Rourkela-3 (sector-7). The level of Na⁺ measured against

Ca^{2+} and Mg^{2+} is known as Kelley's ratio, based on which irrigation water can be rated Kelley's ground water having less than one is considered as good for irrigation (Kelley [59]). $\text{KR} > 1$ indicates an excess level of Na^+ in waters. Therefore, water with a $\text{KI} \leq 1$ has been recommended for irrigation, while water with $\text{KI} \geq 1$ is not recommended for irrigation due to alkali hazards [60, 61]. KR content ranged from 0.11 to 1.46 with a mean of 0.54 for surface water and 0.13 to 1.78 with a mean of 0.78 for groundwater samples. In the present study, all the sources of surface water have KR less than 1 and hence recommended for irrigation. Similarly, among the ground water sources, water collected from Sambalpur (1.66) and Gajapati (1.12) had KR more than 1 (Table 7; Fig. 6).

In the present study area, all the surface water samples are of good quality and suitable for irrigation purpose as based upon all the parameters the samples are within safe limit. Based upon SSP only Bahuda river is of excellent quality, mahanadi, subarnarekha, salandi is of good quality and others are within permissible range according to the classification [57] while studying the water quality of Puri district. Based upon EC all the samples fall in low salinity class except Mahanadi, Bramhani, Bansadhara river which falls in medium salinity class, according to class [47]. The SAR, Adj. SAR and PI falls in excellent category of suitability classification and based upon all other parameters the samples are safe for irrigation purposes (Table 8). The ground water samples are suitable for irrigation purpose based upon all parameters except Putibandh of Sambalpur basing upon the RSC falls in unsuitable category [54], based upon SSP under doubtful class, based on KR bad category and sample of Gajapati based upon KR which falls in bad category [62]. Based upon EC all the samples fall under low salinity class except Sambalpur, Rourkela-3 (sector-7), Ganjam, Gajapati which fall under medium salinity class [47]. Based upon SSP only sample of Orchard fall under excellent category, sample of Sambalpur under doubtful category and all other samples are under permissible category of suitability classification [57]. Based upon SAR, Adj.SAR, PI all the samples fall under excellent category of suitability classification (Table 9).

Conclusions

It can be concluded from the present study area that all the surface water samples collected from 27 points of nine different rivers are of good quality and suitable for irrigation purpose on the basis of quality assessment parameters except some specific limitations associated with certain sources which can be overcome with management. Based upon salinity measured in terms of EC all rivers except three rivers Mahanadi, Brahmani and Bansadhara have no salinity hazard. These three rivers are of medium salinity ($0.25\text{--}0.75 \text{ dSm}^{-1}$) but the water can be safely used for irrigation with management. In other cases, the salinity level is safe ($<0.25 \text{ meqL}^{-1}$). Water samples of all rivers except Subarnarekha, Brahmani, Ib and Mahanadi have no sodicity hazard. The water of these three rivers was above the safe limit but within the permissible limit of RSC, 2.5 meqL^{-1} but can be used with management. When measured on individual basis two samples collected from Subarnarekha river from two sites at Mayurbhanj are above the permissible limit of 2.5 meqL^{-1} and thus not suitable for irrigation. The water of all the nine rivers collected from different places of their mid-reach have no chloride and boron toxicity. Nitrate was much below the maximum permissible level. All the surface water samples were below the toxicity level and safe with respect to Zn (2 mgL^{-1}), Cu (0.2 mgL^{-1}), Fe (5 mgL^{-1}), Mn (0.2 mgL^{-1}) and Cr (0.1 mgL^{-1}). But water of four major rivers exceeded the maximum permissible limit of Cd. These are in the order: Rushikulya, $0.034 \text{ mgL}^{-1} > \text{Kolab}, 0.023 \text{ mgL}^{-1} > \text{Mahanadi}, 0.014 \text{ mgL}^{-1} > \text{Bahuda}, 0.011 \text{ mgL}^{-1}$. In the other hand, groundwater samples collected from the three Farms of OUAT at Bhubaneswar are acidic in reaction and are below the permissible limit of 6.5–8.5 with respect to other characteristics, they are safe to use.

Ethics approval Not applicable.

References

1. Suresh R, Prabakaran R, Babu K, Selvaraj B. Spatial analysis of groundwater quality for Chinnar river sub basin, Tamilnadu, India using GIS. *Int J Mod Res Rev*, 2015; 3:1034-1037.

2. Ramamoorthy P, Backiaraj S, Ajithkumar R. Evaluation of ground water quality for drinking and irrigation suitability: A case study in Marakkanam Block, Villupuram district, Tamilnadu, India. *J Ind Pollut Control*, 2018;34(2): 2159-2163.
3. Sarkar AA, Hasan A. Water quality assessment of a ground water basin in Bangladesh for irrigation use. *Pak J Biol Sci.* 2006; 9 (9): 1677-1684.
4. Rawat KS, Singh SK, Gautam SK. Assessment of groundwater quality for irrigation use: a peninsular case study. *Appl Water Sci.* 2018; 8: 233.
5. Raihan F, Alam JB. Assessment of groundwater quality in Sunamganj of Bangladesh. *Iran J Environ Health Sci Eng*, 2008; 5(3): 155–166.
6. Sarkar AA, Hasan A. Water quality assessment of a ground water basin in Bangladesh for irrigation use. *Pak J Biol Sci*, 2006; 9(9): 1677-1684.
7. Andrew AA, Shimada J, Hosono T. Evaluation of ground water quality and its suitability for drinking and agriculture uses in banana plain (Mbanga Njmba,) of the Cameroon volcanic line. *Environ Geochem Health*, 2013; 33 (6): 559–575.
8. Ackah M, Agymang O, Anim AK. Assessment of ground water quality for drinking and irrigation the case study of Tweimann Oyrifa community Ga east municipality Ghana. *Int J. Ecol Environ Sci*, 2011; 1 (3-4): 185–194.
9. Baher MM, Reja MS. Hydrochemical characteristics and quality assessment of shallow ground water in a coastal area of south west Bangladesh. *Environ Earth Sci*, 2010; 61(5):
10. Ahamad J, Longnathan K, Anathakrishnan S. A comparative evaluation of ground water suitability for drinking and irrigation purpose in Pugalur area Karur district Tamilnadu, India. *Appl Sci Res*, 2013; 5 (1): 213–223.
11. Nepolian M, Chidambaram S, Thivya et al. C. Assessment of hydrochemical and qualities studies in ground water of Villupuram district Tamilnadu, India. *International Research Journal of Earth Sciences*, 2016; 4(3): 1-10
12. Samaahayira S, Florence AC. Evaluation of ground water quality for irrigation purpose Rediyarchatram block of Dindigul district Tamilnadu, India. *Int Res J Earth*, 2017.
13. Sharma AD, Rishi SM, Kessari T. Evaluation of ground water quality and suitability for irrigation and drinking purpose in south west Punjab, India using hydrochemical approach. *Appl Water Sci*, 2017; 7: 3137-3150.
14. Tripathi AK, Mishra UK, Mishra A, Dubey P. Assessment of groundwater quality of Gurh Tehseel, Rewa district Madhya Pradesh, India. *Int J. Sci. Eng Res*, 2012; 3(9): 1-12.
15. Mishra SK, Mishra UK, Tripathy AK, Singh AK, Mishra AK. Hydrogeochemical studies in Ground water quality in and sajjanpur area Satna district Madhya Pradesh. *Int J. Sci.* 2013.
16. Mahadev S, Ahamad A, Kushwaha J, Singh P, Mishra PK. Geochemical assessment of ground water quality for its suitability for drinking and irrigation purpose in rural areas of Sant Ravidas Nagar (Bhadohi), Uttar Pradesh. *Geol Ecol Landsc*, 2018; 2(2):127-136.
17. Dudeja D, Bartarya SK, Biyani, AK. Hydrochemical and water quality assessment of groundwater in Doon valley of outer Himalaya, Uttarakhand, India. *Environ Monit Assess*, 2011; 181(1-4): 183.
18. Jain CK, Bandopadhyay A, Bhadra A. Assessment of ground water quality for irrigation purpose district in

- Nainital Uttarakhand, India. *J Indian Water Resources Soc.* 2012; 32: (3-4): 8-14.
19. Bhandari, NS, Nayal, K. Correlation study on physicochemical parameters and quality assessment of Kosi river water, Uttarakhand. *J Chem*, 2008; 5 (2): 342-346.
 20. Seth R, Sharma MM, Gupta VK, Singh P, Singh R, Gupta S. Application of technique in the assessment of ground water quality of Udham Singh Nagar Uttarakhand. *Water Qual Expos Hea*, 2014; 6(4): 199-216.
 21. Bahukhandi, KD, Bartarya, SK, Siddiqui, NA. Assessment of ground and surface water quality Haridwar district, Uttarakhand. *Int J Chemtech Res.* 2017; 10 (10): 95–118.
 22. Singh KK, Tewari G, Kumar S. Evaluation of Groundwater Quality for Suitability of Irrigation Purposes: A Case Study in the Udham Singh Nagar, Uttarakhand. *J Chem* Article ID 6924026, 2020;15 <https://doi.org/10.1155/2020/6924026>
 23. Mahananda MR, Mohanty BP, Behera NR. Physico-chemical analysis of surface and ground water of Bargarh district, Orissa, India. *Int J Recent Res Appl Stud*, 2010; 2(3): 285-295.
 24. Madhav S, Kumar A, Kushawaha J, Ahamad A, Singh P, Dwivedi SB. Geochemical assessment of groundwater quality in Keonjhar City, Odisha, India. *Sustain Water Resour Manag*, 2020;6(46): 1-11. <https://doi.org/10.1007/s40899-020-00395-7>
 25. Nagy-Kovács Z, Davidesz J, Czihát-Mártonné K, Grischek T. Water quality changes during riverbank filtration in Budapest, Hungary. *Water*, 2019; 11(2): 302. <https://doi.org/10.3390/w11020302>
 26. Sandhu C, Grischek T, Börnick H, Feller J, Sharma SK. A water quality appraisal of some existing and potential riverbank filtration sites in India. *Water*, 2019; 11(2): 215. <https://doi.org/10.3390/w11020215>
 27. El-Zeiny AM, Abd El-Hamid HT, El-Alfy, MA. Anthropogenic impacts on water quality of river Nile and marine environment, Rosetta branch using geospatial analyses. *Res J Environ Sci*, 2018;47(3-4): 89-101. DOI: 10.21608/joese.2018.158202
 28. Shirmohammadi B, Malekian A, Salajegheh A, Taheri B, Azarnivand H, Malek Z, Verburg PH. Scenario analysis for integrated water resources management under future land use change in the Urmia Lake region, Iran. *Land Use Policy*, 2020; 90, 104299. <https://doi.org/10.1016/j.landusepol.2019>
 29. El-Alfy MA, Hasballah AF, Abd El-Hamid HT, El-Zeiny AM. Toxicity assessment of heavy metals and organochlorine pesticides in freshwater and marine environments, Rosetta area, Egypt using multiple approaches. *Sustain Environ Res*, 2019; 29: 19. <https://doi.org/10.1186/s42834-019-0020-9>
 30. Osman R, Ferrari E, McDonald S. Water scarcity and irrigation efficiency in Egypt. *Water Economics and Policy*, 2016; 2(4) 1650009. <https://doi.org/10.1142/S2382624X16500090>
 31. Wang F, Wang X, Zhao Y, Yang ZF. Long-term water quality variations and chlorophyll a simulation with an emphasis on different hydrological periods in Lake Baiyangdian, northern China. *J Environ Inform*, 2012; 20(2): 90–102 doi:10.3808/jei.201200223
 32. Wallace JS. Increasing agricultural water use efficiency to meet future food production. *Agric Ecosyst Environ*, 2000; 82(1-3): 105–119. [https://doi.org/10.1016/S0167-8809\(00\)00220-6](https://doi.org/10.1016/S0167-8809(00)00220-6)
 33. Molden D. *Water for Food Water for Life: A Comprehensive Assessment of Water Management in Agriculture*; Routledge: Abingdon, UK. 2013.
 34. Movilla-Pateiro L, Mahou-Lago XM, Doval MI, Simal-Gandara J. Toward a sustainable metric and indicators for the goal of sustainability in agricultural and food production. *Critical Reviews in Food*

- Science and Nutrition, 2020; 61(7): 1108-1129. DOI: 10.1080/10408398.2020.1754161
35. Haydar S, Arshad M, Aziz JA. Evaluation of drinking water quality in urban areas of Pakistan: A Case Study of Southern Lahore. *Pakistan J Eng Appl Sc.* 2009; 5:16-23.
 36. Kumar A, Dua A. Water quality index for assessment of water quality of river Ravi at Madhopur (India). *Glob J Environ Sci.*, 2010; 8(1): 49-57. DOI: 10.4314/gjes.v8i1.50824
 37. Central Pollution Control Board of India Status of groundwater quality in India part-II, groundwater quality series: gwqs/10/2007-2008. New Delhi, 2008.
 38. Singh KK, Tewari G, Kumar S. Evaluation of Groundwater Quality for Suitability of Irrigation Purposes: A Case Study in the Udham Singh Nagar, Uttarakhand. *J. Chem.* 2012; Article ID 6924026, 2020:15 <https://doi.org/10.1155/2020/6924026>
 39. Zar JH. Biostatistical analysis (5th ed.). Upper Saddle River, New Jersey: Prentice Hall.
 40. Ayres RS, Westcott DW. Water quality for agriculture. F.A.O. Irrigation and Drainage Paper No. 29, Rev., 1985; 1. F.A.O., Rome, 2009:174.
 41. Kundu S. Assessment of surface water quality for drinking and irrigation purposes: A case study of Ghaggar river system surface water. *Bull Env Pharmacol Life Sci.* 2012; 1 (2): 1-1.
 42. Ayers RS, Westcot DW. Water quality for agriculture. FAO Irrigation and Drainage Paper 29, Rev. 1. Rome, Italy: Food and Agricultural Organization of the United Nations, 1994.
 43. USSL Staff. Diagnosis and improvement of saline and alkali soils. USDA Handbook No 60. Washington DC, USA, 1954: 160 pp.
 44. Duncan, RR, Carrow, RN, Huckle M. Under standing water quality and guideline to management (An overview of challenges of water uses on golf course 21st Century), USGA, Far Hills, NJ, USA, 2000.
 45. Sharifi Z, Sinegani AAS. Aresenic and other water quality indicators of groundwater in an agriculture area of Qorveh Plain, Kurdistan, Iran. *Am Eurasian J Agric Environ Sci.* 2012; 12(4): 548–555.
 46. Ayres RS, Westcott DW. Water quality for agriculture. F.A.O. Irrigation and Drainage Paper No. 29, Rev., 1985; 1. F.A.O., Rome, 174 p.
 47. Richards LA. Diagnosis on improvement of saline and alkali soils, USDA., Handbook No.60, Agri. Handbook .U.S. Dep. Agric., 1954; 160.
 48. Miller RW, Gardiner DT. Soils in our environment. 9th edition. Prentice Hall-Inc., Upper Sddle River, New Jersey 07458. ISBN 0-13-020036-0, 2007; p-452.
 49. Gupta IC. Use of saline water in agriculture in arid and semi-arid zones of India. 1979; ref 16, 210.
 50. Rhoades JD. Reclamation and management of salt-affected soils after drainage. Proc. First Annual Western Provincial Conference on Rationalization of Water and Soil Research and Management, Alberta, Canada. 1982; 123–197.
 51. Suarez DL. Relation between pHc and sodium adsorption ratio (SAR) and an alternative method of estimating SAR of soil or drainage waters. *Soil Sci Soc. Am. J.*, 1981; 45(3): 469-475.
 52. Doneen LD. Notes on water quality in agriculture. Water Science and Engineering. Paper, 4001. University of California., 1964:10.

53. Al-Amry AS. Hydro geochemistry and groundwater quality assessment in an arid region: A case study from Al Salameh area, Shabwah, Yemen. The 3rd International Conference on Water Research and Arid Environment and the 1st Arab Water Forum. 2008.
54. Eaton FM. Significance of carbonates in irrigation waters. Soil Sci. 1950;69:123-134. <https://doi.org/10.1097/00010694-195002000-00004>
55. Gupta SK., Gupta IC. Management of saline soils and water. Oxford and IBH publication Company. New Delhi, India, 1987:399.
56. Gupta SK. Variations of water table in Yamuna drainage basin of Haryana-implications and management strategies. Paper presented at the Seminar on Strategies for Irrigation Water Management, Patna, 1983.
57. Behera B, Das M, Rana, GS. Groundwater quality for drinking and irrigation purpose in Damojodi area, Koraput District, Odisha, India. World Appl Sci. J. 2016; 34 (6): 695-703.
58. Laze P, Rizani S, Ibraliu A. Assessment of irrigation water quality of Dukagjin basin in Kosovo. J Int Sci Publ, 2016; 4: 544-551.
59. Kelly WP. Permissible composition and concentration of irrigated waters. In: Proceedings of the A.S.C.F, 1940;607.
60. Ramesh K, Elango L. Groundwater quality and its suitability for domestic and agricultural use in Tondiar river basin, Tamil Nadu, India. Environ Monit Assess, 2012; 184: 3887-3899.
61. Karanth KR. Groundwater assessment development and management. Tata McGrawHill Publishing Company Ltd., New Delhi, 1987; p 725.
62. Shah S., Mistry N. Groundwater quality assessment for irrigation use in Vadodara district, Gujarat, India. Int J Agril Biosys Eng., 2013; 7(7): 719 – 724.

Table 1 Methodology for analysis of chemical parameters

Sl. No.	Parameters	Method used	Reference
1	pH	Glass Electrode pH meter	Jackson, 1973

2	EC	Conductivity meter	Jackson, 1973
3	Carbonate and Bicarbonate	Volumetric Rapid titration Method	Jackson, 1973
4	Chloride	Silver Nitrate Titration Method	Jackson, 1973
5	Nitrate	Colorimetric method using phenol disulphonic acid	Page et al.,1982
6	Calcium and Magnesium	Versenate titration method	Jackson, 1973
7	Potassium	Flame photometer method	Page et al.,1982
8	Boron	Colorimetrically using Azomethine-H	Bingham, 1982
9	Fe, Mn, Zn, Cu, Cd, Cr	Atomic absorption spectrophotometer	Lindsay and Norvell,1978

Table 2 Methodology for analysis of different indices

Indices	Formula	Parameters used in the calculation and their units	Reference
Sodium Adsorption Ratio (SAR)	$SAR = \frac{Na^+}{\frac{\sqrt{Ca^{2+} + Mg^{2+}}}{2}}$	Na ⁺ , Ca ²⁺ , Mg ²⁺ are in meq L ⁻¹	Richards (1954); Todd (1980)
Adjusted Sodium Adsorption Ratio (Adj ^R Na/ Adj. SAR)	$Adj^R Na = \frac{[Na^+]}{\frac{\sqrt{Ca_x^{2+} + Mg^{2+}}}{2}}$	Na ⁺ , Ca _x ²⁺ , Mg ²⁺ are in meq L ⁻¹	Suarez (1981), Ayers and Westcot (1994)
Soluble Sodium Percentage (SSP)	$SSP = \frac{Na^+ + K^+}{Ca^{2+} + Mg^{2+} + Na^+ + K^+} \times 100$	Na ⁺ , Ca ²⁺ , Mg ²⁺ are in meq L ⁻¹	Doneen (1964); Todd (1980)
Permeability Index (PI)	$PI = \frac{Na^+ + \sqrt{HCO_3^-}}{Ca^{2+} + Mg^{2+} + Na^+} \times 100$	HCO ₃ ⁻ , Ca ²⁺ , Mg ²⁺ , Na ⁺ are in meq L ⁻¹	Doneen (1964)
Residual Sodium Carbonate (RSC)	$RSC = (CO_3^{2-} + HCO_3^-) - (Ca^{2+} + Mg^{2+})$	CO ₃ ²⁻ , HCO ₃ ⁻ , Ca ²⁺ , Mg ²⁺ , are in meq L ⁻¹	Eaton (1950); Richards (1954)
Residual Sodium Bicarbonate (RSBC)	$RSBC = (HCO_3^- - Ca^{2+})$	HCO ₃ ⁻ , Ca ²⁺ are in meq L ⁻¹	Gupta and Gupta (1987)
Kelly's Ratio (KR)	$KR = \frac{Na^+}{Ca^{2+} + Mg^{2+}}$	Na ⁺ , Ca ²⁺ , Mg ²⁺ are in meq L ⁻¹	Kelly (1963)

Table 3 Chemical parameters of the collected surface water samples from different water sources measured during Pre Monsoon period, 2019

Sl. No.	Rivers	Mean values	pH	EC (dSm ⁻¹)	CO ₃ ⁻²	HCO ₃ ⁻¹	Na	K	Ca	Mg	Cl ⁻¹	B	NO ₃ ⁻	Zn	Cu	Fe	Mn	Cd	Cr	
			meq L ⁻¹							mg L ⁻¹			mg L ⁻¹							
1	Mahanadi		7.38	0.27	0	2.64	0.58	0.115	0.71	0.37	38.40	0.23	2.31	0.013	0.035	0.173	0.011	0.014	ND*	
2	Rushikulya		7.54	0.22	0	2.31	1.52	0.075	0.55	0.91	29.54	0.05	1.25	0.017	0.048	0.043	0.034	0.034	ND	
3	Bramhani		7.09	0.57	0	2.64	0.93	0.079	1.10	0.13	34.38	0.16	2.96	0.011	0.044	0.038	0.054	0.005	ND	
4	Subernarekha		7.30	0.21	0	2.64	0.16	0.069	0.29	0.29	23.63	0.12	2.19	0.008	0.042	0.170	0.014	0.008	ND	
5	Bahuda		7.69	0.09	0	1.32	0.30	0.071	1.03	0.49	36.63	0.15	2.07	0.009	0.043	0.045	0.015	0.011	ND	
6	Kolab		6.51	0.15	0	1.65	0.42	0.052	0.28	0.39	46.08	0.09	2.18	0.040	0.046	0.169	0.017	0.023	ND	
7	Bansadhara		7.37	0.45	0	1.98	0.81	0.101	1.08	0.22	37.81	0.73	1.79	0.013	0.050	0.047	0.016	0.007	ND	
8	Salandi		6.75	0.07	0	0.99	0.10	0.100	0.33	0.22	29.54	0.89	1.10	0.020	0.037	0.181	0.017	0.002	ND	
9	Ib		8.02	0.21	0	2.64	0.53	0.044	0.61	0.13	50.81	0.10	3.07	0.003	0.036	0.108	0.014	0.006	ND	
Grand mean			7.29	0.25	0	2.09	0.59	0.078	0.66	0.35	36.31	0.28	2.10	0.015	0.042	0.108	0.021	0.012		
SEm (±)			0.081	0.05	0	0.40	0.24	0.008	0.08	0.08	4.76	0.04	0.13	0.001	0.002	0.002	0.001	0.001		
CD (0.05)			0.242	0.14	0	1.19	0.72	0.023	0.240	0.25	1585	0.12	0.38	0.002	0.005	0.006	0.003	0.002		
Standard value			6.5-8.5	0.25-0.75	0-1	0-10	0 – 40	0-0.051	0-20	0-5	350	0-2	5-30	2.00	0.20	5.00	0.20	0.01		
Observed Range			6.23 - 8.05	0.068-0.705	0	0.99 - 2.97	0.09-2.54	0.018-0.65	0.23-1.2	0.09-1.2	21.27-67.36	0.0225-0.934	0.98-3.23	0.002-0.043	0.032-0.052	0.036-0.184	0.01-0.056	0.001-0.035		

*ND: Non detectable

Table 4 Chemical parameters of the collected groundwater samples measured during Pre Monsoon period, 2019

Sl. No.	Ground water source	Mean values	pH	EC (ds/m)	CO ₃ ⁻²	HCO ₃ ⁻²	Na	K	Ca	Mg	Cl ⁻¹	B	NO ₃ ⁻	Zn	Cu	Fe	Mn	Cd	Cr	
					meq L ⁻¹						mg L ⁻¹			mg kg ⁻¹						
1	Agronomy field, OUAT, Bhubaneswar		5.97	0.21	0.00	1.65	0.73	0.055	1.01	0.22	37.81	0.86	4.74	0.018	0.032	0.959	0.018	0.011	ND	
2	Sambalpur, Putibandh		8.02	0.70	4.59	1.32	3.01	0.018	1.61	0.22	55.53	0.05	5.33	0.042	0.038	0.035	0.109	0.011	ND	
3	Rourkela -3 (sector-7)		7.89	0.30	1.65	1.32	0.50	0.06	0.98	0.33	56.60	0.13	3.06	0.008	0.036	0.036	0.018	0.010	ND	
4	Orchard, OUAT, Bhubaneswar		6.46	0.06	0.00	1.32	0.13	0.016	0.41	0.43	22.42	0.13	7.92	0.033	0.043	0.130	0.037	0.008	ND	
5	Central Farm, OUAT, Bhubaneswar		5.97	0.11	0.00	0.99	0.40	0.018	0.33	0.18	24.34	0.21	3.99	0.023	0.035	1.033	0.016	0.009	ND	
6	Ganjam, Krishna Nagar		7.48	0.48	0.00	1.98	1.17	0.074	1.18	0.29	47.26	0.39	4.84	0.028	0.033	0.036	0.012	0.008	ND	
7	Gajapati ,Paralakhemundi		7.50	0.31	1.65	1.65	0.89	0.016	0.48	0.31	35.92	0.09	3.41	0.035	0.028	0.046	0.011	0.009	ND	
Grand mean			7.04	0.31	1.12	1.46	0.98	0.037	0.86	0.28	39.98	0.26	4.75	0.026	0.035	0.325	0.032	0.009		
SEm (±)			0.076	0.01	0.37	0.36	0.02	0.003	0.05	0.03	4.10	0.01	0.12	0.001	0.001	0.012	0.001	0.000		
CD (0.05)			0.228	0.03	1.12	1.06	0.08	0.010	0.15	0.10	12.30	0.05	0.38	0.003	0.004	0.036	0.002	0.001		
Standard value			6.5-8.5	0.25-0.75	0-1	0-10	0-40	0-0.051	0-20	0-5	140-700	0-2	5-30	2.00	0.20	5.00	0.20	0.01		
Observed Range			5.68-8.1	0.045-0.723	0.99-5.4	0.99 - 2.97	0.112-3.04	0.005-0.078	0.24-1.72	0.13-0.54	3.54-60.26	0.043-0.912	2.96-8.2	0.007 - 0.044	0.026 - 0.045	0.033-1.09	0.01-0.11	0.007-0.012		

*ND: Non detectable

Table 5 Calculated indices of the collected surface water samples from different sources during pre-monsoon period 2019

Sl. No.	Rivers	Mean values	SAR	Adj. SAR	RSC (meq L ⁻¹)	PI (%)	KR	RSBC (meq L ⁻¹)	SSP (%)
1	Mahanadi		0.80	0.76	1.56	133.63	0.55	1.93	39.49
2	Rushikulya		1.68	1.60	0.84	106.52	0.93	1.76	43.60
3	Bramhani		1.18	1.19	1.41	118.26	0.75	1.54	45.06
4	Subernarekha		0.32	0.28	2.05	254.97	0.32	2.35	29.90
5	Bahuda		0.35	0.28	-0.20	79.39	0.19	0.29	19.27
6	Kolab		0.74	0.57	0.98	153.36	0.65	1.37	41.47
7	Bansadhara		1.01	0.90	0.68	104.77	0.63	0.90	41.28
8	Salandi		0.19	0.13	0.44	169.25	0.18	0.66	26.80
9	Ib		0.87	0.82	1.90	169.38	0.72	2.03	43.74
Grand mean			0.79	0.73	1.07	143.28	0.55	1.43	36.73
SEm (±)			0.25	0.25	0.42	19.15	0.13	0.42	4.82
CD (0.05)			0.76	0.75	1.28	57.40	0.41	1.27	14.47
Standard value			10 -18	10-20	<1.25	>25	<1	<3	20-40
Observed range			0.16- 2.73	0.12-2.65	-0.75 - 2.53	65.71-305.38	0.11- 1.46	0.21-2.74	13.33- 60.36

Table 6 Calculated indices of the collected groundwater samples of different sources during pre-monsoon period 2019

Sl. No.	Ground water source	Mean values	SAR	Adj.SAR	RSC (meq L ⁻¹)	PI (%)	KR	RSBC (meq L ⁻¹)	SSP (%)
1	Agronomy field, OUAT, Bhubaneswar		0.93	0.81	0.41	99.87	0.59	0.64	56.92
2	Sambalpur, Putibandh		3.16	2.61	4.08	85.82	1.66	0.29	62.43
3	Rourkela -3(sector-7)		0.62	0.49	1.65	90.47	0.38	0.33	29.92
4	Orchard, OUAT, Bhubaneswar		0.20	0.16	0.47	130.63	0.16	0.90	15.02
5	Central Farm, OUAT, Bhubaneswar		0.78	0.54	0.47	152.78	0.76	0.66	45.90
6	Ganjam, Krishna nagar		1.78	1.27	2.15	97.37	0.79	0.80	45.64
7	Gajapati, Paralakhemundi		1.87	1.16	0.85	128.38	1.12	1.16	53.32
Grand mean			1.33	1.01	1.44	112.19	0.78	0.68	44.16
SEm (±)			0.13	0.12	0.44	8.97	0.03	0.38	1.97
CD (0.05)			0.40	0.37	1.33	26.90	0.09	1.16	5.90
Standard value			10 -18	10-20	<1.25	>25	<1	<3	20-40
Observed range			0.18-3.26	0.13-3.18	-0.25 - 4.49	77.79-160.27	0.1375 - 1.78	-0.73-1.95	13.49-654

Table 7 Classification of surface and ground water samples based on EC, SAR, Adj. SAR, RSC, SSP and KR

Parameter/Index	Quality	Surface water sources	% of total surface water samples	Ground water sources	% of total groundwater samples
EC range (dSm⁻¹) (Richards, 1954)					
< 0.25	Low salinity (C1)	6 samples (Rushikulya, Bahuda, Subarnarekha, Kolab, Salandi and Ib)	66.67	3 samples (OUAT Agronomy Field, OUAT Orchard and OUAT-Central farm)	42.86
0.25-0.75	Medium salinity water some leaching for sensitive crops (C2)	3 samples (Mahanadi, Bansadhara and Bramhani)	33.33	4 samples (Sambalpur, Rourkela, Ganjam and Gajapati)	57.14
0.75-2.25	High salinity (C3)	Nil	-	Nil	-
> 2.25	Very high salinity (C4)	Nil	-	Nil	-
SAR (Richards, 1954; Todd, 1980)					
< 10	Excellent (S1)	9 samples (Mahanadi, Rushikulya, Bramhani, Subarnarekha, Bahuda, Kolab, Bansadhara Salandi and Ib)	100	7 samples (OUAT Agronomy Field, Sambalpur, Rourkela, OUAT Orchard, OUAT Central farm, Ganjam and Gajapati)	100
10-18	Good (S2)	Nil		Nil	
19-26	Doubtful (S3)	Nil		Nil	
> 26	Unsuitable (S4)	Nil		Nil	
Adj. RNa/ Adj. SAR (Gupta, 1979)					
< 10	Normal (S1)	9 samples (Mahanadi, Rushikulya, Bramhani, Subarnarekha, Bahuda, Kolab, Bansadhara Salandi and Ib)	100	7 samples (OUAT Agronomy Field, Sambalpur, Rourkela, OUAT Orchard, OUAT Central farm, Ganjam and Gajapati)	100
10-20	Medium (S2)	Nil		Nil	
20-30	High (S3)	Nil		Nil	
30-40	Very High (S4)	Nil		Nil	
> 40	Unsuitable (S5)	Nil		Nil	
RSC (meq L⁻¹) (Eaton, 1950; Richards, 1954)					
< 1.25	Safe	5 samples (Rushikulya, Bahuda, Kolab,	55.56	4 samples (OUAT-Agronomy Field, OUAT	57.14

		Bansadhara and Salandi)		Orchard, OUAT-Central farm and Gajapati)	
1.25-2.5	Can be used with management	4 samples (Mahanadi, Brahmani, Subarnarekha, Ib)	44.44	2 samples (Rourkela, Ganjam)	28.57
> 2.5	Unsuitable	Nil		1 sample (Sambalpur)	14.29
SSP range (%) (Wilcox, 1954; Behera <i>et al.</i>, 2016)					
< 20	Excellent	1 sample (Bahuda)	11.11	1 sample (OUAT Orchard)	14.29
20-40	Good	3 samples (Subarnarekha, Salandi and Mahanadi)	33.33	1 sample (Rourkela)	14.29
40-60	Permissible	2 samples (Rushikulya and Bramhani) 3 samples (Kolab, Bansadhara and Ib)	55.56	4 samples (OUAT-Agronomy Field, OUAT-Central farm, Ganjam and Gajapati)	57.14
60-80	Doubtful	Nil	-	1 sample (Sambalpur)	14.29
KR range (Kelly, 1963)					
≤ 1	Good	9 samples (Mahanadi, Rushikulya, Bramhani, Subarnarekha, Bahuda, Kolab, Bansadhara, Salandi and Ib)	100	5 samples (OUAT Agronomy field Bhubaneswar, Rourkela - 3(sector-7), Orchard OUAT Bhubaneswar, Central Farm OUAT Bhubaneswar, and Ganjam)	71.43
> 1	Bad	Nil	-	2 samples (Sambalpur and Gajapati)	28.57

EC electrical conductivity, *SAR* sodium adsorption ratio, *Adj. SAR* adjusted sodium adsorption ratio, *RSC* residual sodium carbonate, *SSP* soluble sodium percentage, *KR* kelly's ratio

Table 8 Suitability classification of surface water samples basing upon different parameters

Rivers	SAR	Adj.SAR	RSC	PI	KR	RSBC	SSP	pH	EC	HCO ₃	Cl	B	NO ₃
Mahanadi	Ext	Ext	S	Ext	G	S	G	S	MS	S	S	S	S
Rushikulya	Ext	Ext	S	Ext	G	S	Pl	S	LS	S	S	S	S
Bramhani	Ext	Ext	S	Ext	G	S	Pl	S	MS	S	S	S	S
Subernarekha	Ext	Ext	S	Ext	G	S	G	S	LS	S	S	S	S
Bahuda	Ext	Ext	S	Ext	G	S	Ext	S	LS	S	S	S	S
Kolab,nagavali	Ext	Ext	S	Ext	G	S	Pl	S	LS	S	S	S	S
Bansadhara	Ext	Ext	S	Ext	G	S	Pl	S	MS	S	S	S	S
Salandi	Ext	Ext	S	Ext	G	S	G	S	LS	S	S	S	S
Ib	Ext	Ext	S	Ext	G	S	Pl	S	LS	S	S	S	S

Table 9 Suitability classification of ground water samples basing upon different parameters

Ground water source	SAR	Adj.SAR	RSC	PI	KR	RSBC	SSP	pH	EC	HCO ₃	Cl	B	NO ₃
Agronomy field, OUAT	Ext	Ext	S	Ext	G	S	Pl	A	LS	S	S	S	S
Sambalpur, Putibandh	Ext	Ext	US	Ext	B	S	Db	S	MS	S	S	S	S
Rourkela-3, sector-7	Ext	Ext	S	Ext	G	S	Pl	S	MS	S	S	S	S
Orchard, OUAT, Bhubaneswar	Ext	Ext	S	Ext	G	S	ext	A	LS	S	S	S	S
Central Farm, OUAT, Bhubaneswar	Ext	Ext	S	Ext	G	S	Pl	A	LS	S	S	S	S
Ganjam, Krishna nagar, Berhampur	Ext	Ext	S	Ext	G	S	Pl	S	MS	S	S	S	S
Gajapati Paralakhemundi	Ext	Ext	S	Ext	B	S	Pl	S	MS	S	S	S	S

A Acidic, Db Doubtful, Ext Excellent, Pl Permissible, LS low Salinity, MS Medium Salinity, G good, B Bad, S safe, US Unsuitable

EC electrical conductivity, SAR sodium adsorption ratio, Adj. SAR adjusted sodium adsorption ratio, RSC residual sodium carbonate, RSBC residual sodium bicarbonate, PI permeability index, SSP soluble sodium percentage, KR kelly's ratio,

Table 10 Suitability of water of nine major rivers of the state for irrigation

Rivers	Acidity/ alkalinity (pH, RSBC)	Salinity (EC)	Sodicity (RSC, SAR, adj. SAR PI, KR, SSP)	Specific ion Toxicity (Cl, B, NO ₃)	Trace element and Heavy Metal Toxicity (Fe, Mn, Zn, Cu, Cd, Cr)	Water Quality
Mahanadi	Safe	Medium	RSC > SL but < MPL	safe	Cd > MPL	Can be used with Management
Rushikulya	Safe	Safe	safe	safe	Cd > MPL	Can be used with Management
Bramhani	Safe	Medium	RSC > SL but < MPL	safe	Safe	Can be used with Management
Subernarekha	Safe	Safe	RSC > SL but < MPL	safe	Safe	Can be used with Management
Bahuda	Safe	Safe	safe	safe	Cd > MPL	Can be used with Management
Kolab	Safe	Safe	safe	safe	Cd > MPL	Can be used with Management
Bansadhara	Safe	Medium	safe	safe	Safe	Suitable
Salandi	Safe	Safe	safe	safe	Safe	Suitable
Ib	Safe	Safe	RSC > SL but < MPL	safe	Safe	Can be used with Management

EC electrical conductivity, *SAR* sodium adsorption ratio, *Adj. SAR* adjusted sodium adsorption ratio, *RSC* residual sodium carbonate, *RSBC* residual sodium bicarbonate, *PI* permeability index, *SSP* soluble sodium percentage, *KR* kelly's ratio

Table 11 Suitability of water of seven ground water sources of the state for irrigation

Ground water source	Acidity/alkalinity (pH, RSBC)	Salinity (EC)	Sodicity (RSC, SAR, SARadj, PI, KR, SSP)	Specific ion Toxicity (Cl, B, NO ₃)	Trace element and Heavy Metal Toxicity (Fe, Mn, Zn, Cu, Cd, Cr)	Water Quality
Agronomy field, OUAT, Bhubaneswar	Acidic	Safe	safe	safe	Cd > MPL	Can be used with Management
Putibandh, Sambalpur	safe	Medium	RSC > MPL KR - Bad SSP-Doubtful	Safe	Cd > MPL	Unsuitable
Rourkela-3, sector-7	safe	Medium	RSC > SL but < MPL	safe	Cd > MPL	Can be used with Management
Orchard, OUAT, Bhubaneswar	Acidic	Safe	safe	safe	Safe	Can be used with Management
Central Farm, OUAT, Bhubaneswar	Acidic	Safe	safe	safe	Safe	Can be used with Management
Ganjam, Krishna nagar, Berhampur	safe	Medium	RSC > SL but < MPL	safe	Safe	Can be used with Management
Gajapati Paralakhemundi	safe	Medium	KR- Bad	safe	Safe	Can be used with Management

EC electrical conductivity, SAR sodium adsorption ratio, Adj. SAR adjusted sodium adsorption ratio, RSC residual sodium carbonate, RSBC residual sodium bicarbonate, PI permeability index, SSP soluble sodium percentage, KR kelly's ratio




Fig. 1 Location of the study area (surface and ground water sampling sites)




Fig. 2 pH of different surface and ground water samples used for irrigation




Fig. 3 Electrical conductivity (EC) of different surface and ground water samples used for irrigation




Fig. 4 Residual sodium carbonate (RSC) of different surface and ground water samples used for irrigation




Fig. 5 Soluble sodium percentage (SSP) of different surface and ground water samples used for irrigation

UNDER PEER REVIEW




Fig.6 Kelly's ratio (KR) of different surface and ground water samples used for irrigation




Fig. 7 Cadmium (Cd) content of different surface and ground water samples used for irrigation