

## Original Research Article

### **“Studies on Genetic Variability and Divergence in Mustard (*Brassica Juncea* L.).”**

#### **ABSTRACT**

The phenotypic coefficient of variation was greater than genotypic coefficient of variation for all the traits. The high magnitude of both coefficients was recorded in case of number of primary number, secondary, number of siliquae per plant, seed yield per plant and harvesting index. High heritability with high genetic advance was recorded for number of primary branches, number of secondary branches, length of main raceme, number of siliquae in main raceme, number of siliquae per plant, number of seeds per siliquae, biological yield, 1000 seed weight, seed yield per plant and harvesting index. In the present findings the significant phenotypic correlation of seed yield per plant were found positive for the characters Days to 50% flowering, plant height, number of primary branches, number of secondary branches, length of main raceme, number of siliquae per plant, length of siliqua, number of seeds per siliquae, harvesting index and seed yield per plant showed positive significant correlation with seed yield per plant. The path coefficient analysis of different characters revealed that highest positive direct effect on seed yield per plant per plant was exerted by harvest index per plant followed by biological yield and negative direct effect on seed yield / plant was exhibited by number of siliquae per plant followed by length of siliquae, days to 50% flowering, number of secondary and number of primary branches. Genetic divergence analysis based on Mahalanobis  $D^2$  statistics, twenty genotypes were grouped into five clusters in mustard. The highest intra-cluster distance was recorded in cluster II followed by cluster I, while the lowest value was recorded in case of cluster III, IV and V. The maximum inter cluster distance was recorded between cluster III and cluster V followed by cluster IV and cluster V. The minimum inter cluster distance was observed between cluster I and cluster III followed by cluster II and cluster V.

*Keywords: Indian mustard, variability, genetic advance, heritability, correlation, path analysis and diversity.*

#### **INTRODUCTION**

Oilseed Brassicas, also known as rapeseed-mustard, are a significant category of oilseed crops worldwide, consisting of eight cultivated crops from the Brassiceae tribe of the Brassicaceae family

(Cruciferae). The term "mustard" comes from the European practise of preparing a heated paste by combining the sweet "must" of old wine with crushed seeds of black mustard (*Brassica nigra*) [1]. In India, Indian mustard (*Brassica juncea* (L.) $2n=36$ ) is a major oilseed crop. It is grown for oil, condiments, and medicinal purposes. However, the crop is primarily grown to produce edible vegetable oil. Despite sharing the same family and genus, rapeseed and mustard have distinct botanical characteristics. Rapeseed is an annual herb that grows to a height of 45 to 150 cm. In most situations, a waxy covering covers the stems. Plant leaves are easily distinguished from mustard (rai) plants. Rapeseed leaves are sessile, glabrous, and hairy. The stalk is partially or completely grasped by the lowest segment of the blade (lamina). The fruits are thicker than mustard (rai) and feature a beak that is one-third to half the fruit's length. The seeds are golden or brown in colour, with a smooth seed coat. Mustard plants are tall (90-200 cm), erect, and highly branched. Instead of being dilated at the base and clasping like rapeseed, the leaves are stalked, broad, and pinnatifid. The fruits are narrow, ranging in length from 2 to 6.5 cm, with strong ascending or erect stems and short, thick beaks. The seed has a rough seed coat and is brown to dark brown in colour. Mustard is the world's third most significant oilseed crop, behind soybeans and palm oil. Among the several oilseed crops produced globally, the estimated area, production, and yield of mustard were 34.88 mha, 69.22 mt, and 1.98 Mt/ha, respectively [2]. Rapeseed production, area, and yield in India are 8.30 million hectares, 1.3 million tonnes, and 1.1 million tonnes per acre, respectively. Where rapeseed yield is lower than that of other crops.

Genetic variety and diversity are crucial requirements for crop development because they allow for more selective breeding. Because seed yield is a polygenic characteristic that frequently leads to changes in other characters, the link between seed yield and other traits is useful for selecting the proper selection in breeding programmes. During the selection process, correlation analysis measures the degree, direction, and strength of the association between two or more variables. Path coefficient analysis quantifies the direct and indirect impacts of numerous independent variables on a dependent variable [3]. As a result, the current study is being carried out to estimate the heritability and diversity of genotypes for yield and yield contributing variables, as well as to assess the amount of direct and indirect reasons of linkage among various characters through path analysis in mustard.

## **MATERIALS AND METHODS**

The present investigation entitled "Genetic Variability and Genetic Divergence in mustard in Mustard" was carried out under, Department of Plant Breeding & Genetics at Agriculture Farm, School

of Agriculture, Lovely Professional University, Jalandhar (Punjab) during the *rabi* 2021-22. The experimental design comprised 20 different genotypes of Mustard (Table 1) were grown in a RBD with three replications during *rabi* 2021-22. Each plot consists of two rows of 1.8 m length. Five competitive plants were selected from each replication for 14 quantitative traits *viz*: days to 50% flowering, days to maturity, plant height, number of primary branches per plant, number of secondary branches per plant, number of siliqua per plant, number of seed per siliqua, siliqua length of siliqua, number of siliqua on main raceme, 1000- seed weight, biological yield per plant, harvest index, seed yield per plant. The mean values of each genotype were computed for statistical analysis. The standard method of analysis of variance was given [4], phenotypic and genotypic coefficient of variation, heritability (Broad Sense) and genetic advance as percent of mean were estimated by the formula as suggested by [5] and [6]. The formula of genotypic correlation coefficients was estimated by [7]. Path analysis along with genotypic correlation coefficient is applied to know the direct and indirect effects of the components on yield as suggested by [8] and illustrated by [3]. The replicated data were subjected to genetic divergence analysis using Mahalanobis's D<sup>2</sup> - statistic [9].

## RESULTS AND DISCUSSION

Analysis of variance revealed that the treatments were highly significant for all the traits except length of siliquae and 1000 seed weight under study *viz*. days to 50% flowering, days to maturity, plant height, length main raceme, number of siliquae on main raceme, number of siliquae per plant, number of seeds per siliqua, biological yield per plant, seed yield per plant and harvest index which signified that genotype is highly variable for the observed traits. Similar results were also obtained by [10],[11],[12] and [13] in which analysis of variance showed significant differences for all the characters under their study.

In the present study, the phenotypic coefficient of variation (PCV) was greater than genotypic coefficient of variation (GCV) for all the traits number of primary numbers, secondary, number of siliquae per plant, seed yield per plant and harvesting index. The high magnitude of both coefficients was recorded for traits; (Table 3). The estimates of phenotypic and genotypic coefficient of variability for of the characters under study are in accordance with earlier reports, [14],[15],[16],[17],[18],[19],[11] and [20].

High heritability with high genetic advance was recorded for number of primary branches, number of secondary branches, length of main raceme, number of siliquae in main raceme, number of siliquae per plant, number of seeds per siliquae, biological yield, 1000 seed weight, seed yield per plant and harvesting index (Table 3). This might be attributed to additive gene action responsible for their expression and hence, phenotypic selection for their amenability, can be brought about. Similar result found, [19] for 1000 seed weight, [21] for number of Siliqua per plant, [22] for 50% flowering, [23],[24] for 50% flowering, plant height, seeds per Siliqua, 1000 seed weight, [25] for seed yield per plant, number of secondary branches, 1000 seed weight, number of seeds per Siliqua, primary branches per plant and Siliqua length, [26],[27],[28] and [29].

The significant phenotypic correlation of seed yield per plant was found positive for the characters namely, number of primary branches per plant, number of secondary branches, number of siliquae on main raceme, number of siliquae per plant, length of siliqua, number of seeds per siliqua, 1000 seed weight, biological yield per plant and harvest index (Table 4). This suggests that these characters should be considered while selecting for improvement in seed yield per plant provided the character should show high variability, which is the basis for selection. However, days to maturity and plant height showed negative correlation with seed yield per plant. In the present findings, the magnitudes of genotypic correlation coefficient were greater than corresponding phenotypic correlation coefficient (in general also) similar findings by [11],[30],[20],[13] and[19].

The path coefficient analysis of different characters revealed that positive direct effect on seed yield per plant per plant was observed by harvest index per plant followed by biological yield and negative direct effect on seed yield / plant was exhibited by number of siliquae per plant followed by length of siliquae, days to 50% flowering, number of secondary and number of primary branches. (Table 5). We can conclude based on present finding that those all character which had positive direct effect on seed yield per plant that universal traits for overall improvement. Negative direct effect on seed yield per plant was exerted by days to 2 maturity, number of secondary branches per plant, number of siliqua on main raceme and length of siliqua. In the present findings, the similar results are also reported by Direct and positive effect by [30],[31],[13],[19],[20],[32] and [31] viz seed yield per plant.

Genetic divergence analysis based on Mahalanobis D2 statistics, twenty genotypes were grouped into five clusters in mustard (Table 6). The highest intra-cluster distance was recorded in cluster II followed by cluster I, while the lowest value was recorded in case of cluster III, IV and V.(Table 7). The genotypes showing maximum diversity from clusters I& II could be utilized directly for future hybridization programs. In conformity to the present investigation, similar findings were found by [33],[34] and [35].

It was observed that biological yield per plant was the highest contributor towards divergence followed by all except Length of main raceme (Table8) don't contribute significantly to the total divergence. Similarly, Devi *et al.*, (2017) was observed, biological yield, Harvest index 1000 seed weight and seed yield per plant were the major contributors for genetic diversity among the genotypes. The cluster mean values for 14 characters of 20 genotypes have been represented in (Table 8).

**Table 1: Details of the Genotypes is given below.**

S.NO	ENTRIES	PEDIGREE / SOURCE (DRMR)
1	Laxmi	DRMR

2	Geeta	DRMR
3	Gujarat Mustard-1	DRMR
4	Aravali	DRMR
5	Maya	DRMR
6	Kranthi	SELECTION FOR VARUNA
7	RNG-73	DRMR
8	Pusa Bold	VARUNA X BIC1780
9	Bhagirathi	SELECTION FROM PUSA JAI KISHAN
10	Durgamahi	DRMR
11	RH-119	DRMR
12	NRCHB-101	BL-4 X PUSA BOLD
13	DRMR-IJ-31	DRMR
14	Pusa Mustard-27	DIVYA/PUSABOLD//PR666EPS///PR704EPS2
15	Pusa Saag-1	DRMR
16	SMR-1	DRMR
17	Urvashi	VARUNA X KRANTI
18	RH-30	SELECTION FROM P26/3-1
19	RH-749	DRMR
20	RB-50	DRMR

**Table 2: Analysis of variance for fourteen characters in Mustard.**

Sr.	Characters	Mean Sum of Squares
-----	------------	---------------------

No.		Source of Variation			
		d.f = 2	d.f = 19	d.f = 38	
		Replication	Genotypes	Error	CV
1	Days to 50% flowering	3.516	25.132**	2.885	3.29
2	Days to maturity	45.033	166.583**	416.966	2.75
3	Plant height (cm)	43.015	328.024**	177.078	7.83
4	Number of primary branches	0.024	4.052**	0.259	7.88
5	Number of secondary branches	0.020	8.649**	0.324	6.67
6	Length of main raceme	6.640	239.879**	22.809	7.32
7	Number of siliquae in main raceme	2.498	65.230**	6.904	6.17
8	Number of siliquae per plant	2325.190	26420.137**	1804.069	10.64
9	Length of siliquae	0.512	0.965	0.534	9.04
10	Number of seeds per siliquae	3.555	16.175**	0.659	5.24
11	1000 seed weight	0.009	0.976	0.043	4.68
12	Biological Yield	5.820	10180.672**	46.409	6.49
13	Seed yield per plant	2.791	274.841**	13.461	7.20
14	Harvesting index	0.695	143.820**	7.751	7.39

**Table 3.: Estimate of coefficient of variation, h<sup>2</sup> (broad sense) and genetic advance in per cent of mean in Mustard.**

S. No.	Characters	Range		Mean	Coefficient of variation		Heritability in broad sense (%)	Genetic advance in per cent of
		min	max					

					PCV	GCV		mean
1	Days to 50% flowering	44	55	49.6167	5.8335%	5.4885 %	88.52	10.6375 %
2	Days to maturity	115	126	118.0833	1.4477 %	0.7261 %	25.15	-0.7501 %
3	Plant height	144	180	165.1100	6.3331 %	4.2961 %	46.02	6.0035 %
4	Number of primary branches	2	7	4.2800	27.1564 %	26.2741 %	93.61	52.3662 %
5	Number of secondary branches	4	11	6.7267	25.2429 %	24.7649 %	96.25	50.0494 %
6	Length of main raceme	41	78	60.5567	14.7664 %	14.0468 %	90.49	27.5263 %
7	Number of siliquae in main raceme	24	48	38.3200	12.1685 %	11.5065 %	89.42	22.4140 %
8	Number of siliquae per plant	222	555	393.8217	23.8291 %	23.0011 %	93.17	45.7360 %
9	Length of siliquae	4.3	6.4	5.1352	11.0482 %	7.3813 %	44.64	10.1586 %
10	Number of seeds per siliquae	11.9	20.4	13.9950	16.5919 %	16.2504 %	95.93	32.7868 %
11	Biological Yield	112	182	140.9333	13.4671 %	13.1747 %	95.71	26.5507 %
12	1000 seed weight	3.2	5.2	3.8358	14.8700 %	14.5317 %	95.50	29.2544 %
13	Seed yield per plant	23	61	46.3007	20.6725 %	20.1599 %	95.10	40.4996 %
14	Harvesting index	20.5	49.5	33.0847	20.9278 %	20.3560 %	94.61	40.7876 %

**Table 4: Summary of phenotypic correlation (above diagonal) coefficient & genotypic correlation (below diagonal) coefficient for yield and yield contributing traits in mustard under study.**

S.N.	Characters	D 50% F	DM	PH	NPB	NSP	LMR	NSMR	NSP	LS	NSPS	TW	BW	HI	SYP
1	DFF	1	***	+	+	-	-	-	+	-	-	-	+	+	+
2	DM	-	1	-	-	-	+	-**	-	***	-	-	-	+	+
3	PH	+	+	1	-	-	+	***	-	-	-	-	+	-	-
4	NPB	+	+	-	1	+	+	+	+	-	+	-	+	-	+
5	NSB	-	+	-	+	1	+	+	+	+	+	+	+	+	+
6	LMR	-	-	+	+	+	1	-	-	+	-	-	+	-	-
7	NSMR	-	+	+	+	+	+	1	+	+	+	+	-	+	+
8	NSP	-	+	-	+	+	-	+	1	+	***	+	+	***	***
9	LS	-	-	-	-	+	+	+	+	1	+	-	+	+	+
10	NSPS	-	+	-	+	+	-	+	***	+	1	+	+	+	***
11	TW	-	-	-	+	+	-	+	+	-	+	1	+	-	+
12	BY	+	+	+	+	+	+	-	+	+	+	+	1	+	+
13	HI	+	-	-	-	+	-	+	***	+	+	-	+	1	***
14	SYPP	+	-	-	+	+	-	+	***	+	***	+	+	***	1

\*\* indicate significant at 5% and 1% probability level, respectively. DF = Days to 50% flowering, DM = Days to maturity, PH = Plant height, NPB = Number of primary branches, NSB = Number of secondary branches, LMR = Length of main raceme, NSMR = Number of siliqua on main raceme, NSP = Number of siliqua/plants, LS = Length of siliqua, NSS = Number of seeds/siliqua, TSW = 1000 seed weight, BY = Biological yield/plant, SY = Seed yield/plant and HI = Harvest index.



**Table 5: Genotypic direct and indirect effect of different characters on seed yield per plant in Mustard**

Characters	D F50%	DM	PH	NPB	NSP	LMR	NSMR	NSP	LS	NSPS	BW	TW	HI
D F50%	-5.5376	-5.0432	-1.0980	-0.3159	0.8396	1.9156	-0.0096	1.0554	1.6396	1.7352	1.0235	-1.3527	-1.9079
DM	0.0498	0.0547	-0.0337	-0.0040	-0.0336	0.0279	-0.0425	-0.0001	0.0325	-0.0228	-0.0018	-0.0175	0.0171
PH	-0.2612	0.8113	-1.3172	0.0076	0.2743	-0.1510	-1.0578	0.0497	0.1204	0.6553	0.2358	-0.0884	0.1768
NPB	-0.2102	0.2674	0.0214	-3.6853	-2.0007	-1.4378	-0.7215	-0.1178	1.5255	-0.4861	-1.4641	-1.2270	0.4990
NSP	0.7439	3.0166	1.0217	-2.6635	-4.9063	-1.9809	-2.1179	-1.2422	-0.0687	-1.6949	-1.3715	-1.2673	-0.9123
LMR	-2.2629	3.3342	0.7499	2.5521	2.6411	6.5415	-0.0956	-0.4678	1.8452	-1.1543	-0.6549	1.1041	-1.1457
NSMR	0.0053	-2.3942	2.4726	0.6028	1.3290	-0.0450	3.0788	1.1274	0.3270	1.2386	0.3596	-0.2355	0.9256
NSP	1.5908	0.0175	0.3151	-0.2669	-2.1133	0.5969	-3.0563	-8.3468	-3.4078	-5.5741	-2.9369	-0.1599	-5.4377
LS	2.3512	-4.7182	0.7258	3.2871	-0.1112	-2.2400	-0.8433	-3.2421	-7.9409	-3.3027	1.2711	-0.7290	-4.2408
NSPS	0.1954	0.2596	0.3103	-0.0823	-0.2154	0.1100	-0.2509	-0.4165	-0.2594	-0.6236	-0.2957	-0.0454	-0.2678
BY	-1.4092	-0.2482	-1.3649	3.0289	2.1314	-0.7632	0.8906	2.6827	-1.2204	3.6158	7.6243	1.3338	-1.6263
TW	-0.2306	0.3024	-0.0634	-0.3143	-0.2438	-0.1593	0.0722	-0.0181	-0.0867	-0.0688	-0.1651	-0.9440	-0.2512
HI	5.1506	4.6634	-2.0068	-2.0242	2.7798	-2.6185	4.4942	9.7393	7.9837	6.4196	-3.1889	3.9776	14.9497
SY	0.1753	0.3235	-0.2672	0.1222	0.3708	-0.2038	0.3403	0.8030	0.4901	0.7374	0.4355	0.3487	0.7786

R SQUARE = 1.0025 RESIDUAL EFFECT = SQRT (1- 1.0025), Bold figures indicate direct effect. DF = Days to 50% flowering, DM = Days to maturity, PH = Plant height, NPB = Number of primary branches, NSB = Number of secondary branches, LMR = Length of main raceme, NSMR = Number of siliqua on main raceme, NSP = Number of siliqua/plants, LS = Length of siliqua, NSS = Number of seeds/siliqua, TSW = 1000 seed weight, BY = Biological yield/plant, SY = Seed yield/plant and HI = Harvest index.

**Table 6: Clustering pattern of 25 Mustard genotypes & based on Mahalanobis's  $D^2$  statistic.**

Cluster No.	No. of Genotypes	Genotypes
Cluster 1	12	Maya, RH 30, Pusa Bold, Laxmi, Geeta, Gujarat Mustard 1, Aravali, Urvashi, SMR 1, DRMRIJ31, Durgamani, RH 749
Cluster 2	5	Kranti, Pusa Mustard 27, RNG73, NRCHB101, RH 119
Cluster 3	1	Pusa saas 1
Cluster 4	1	RB 50
Cluster 5	1	Bhagirathi

**Table 7: Estimates of average intra and inter-cluster distances for the fiveclusters in Mustard**

Clusters	Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5
Cluster 1	11.87	18.71	16.37	18.19	24.02
Cluster 2		12.02	26.71	28.08	16.19
Cluster 3			0.00	23.43	33.97
Cluster 4				0.00	33.28
Cluster 5					0.00

**Table 8: Cluster means and percentage contribution for different characters in mustard**

Characters	DFF	DM	PH	NPBP	NSBP	LMR	NSMR	NSPP	LS	NSPS	BYPP	TSW	HI	SYP
<b>Cluster 1</b>	50.25	117.83	167.87	3.89	6.07	58.21	38.43	384.08	5.02	13.39	138.10	3.63	32.88	45.05
<b>Cluster 2</b>	48.87	118.47	162.40	5.50	8.89	67.73	38.27	398.46	5.20	14.59	150.17	4.41	33.19	49.85
<b>Cluster 3</b>	47.00	118.33	165.00	3.10	4.60	61.40	32.80	222.97	5.63	11.93	115.60	3.36	20.57	23.78
<b>Cluster 4</b>	51.00	118.67	163.13	3.27	5.27	55.20	38.87	520.63	5.79	13.80	112.40	4.47	49.53	55.60
<b>Cluster 5</b>	47.00	118.33	147.60	5.10	7.33	57.33	42.27	531.53	5.01	20.47	182.60	3.30	31.08	56.74

DF = Days to 50% flowering, DM = Days to maturity, PH = Plant height, NPB = Number of primary branches, NSB = Number of secondary branches, LMR = Length of main raceme, NSMR = Number of siliqua on main raceme, NSP = Number of siliqua/plants, LS = Length of siliqua, NSS = Number of seeds/siliqua, TSW = 1000 seed weight, BY = Biological yield/plant, SY = Seed yield/plant and HI = Harvest index.

## CONCLUSION

The conclusion that can be reached from variability, correlations, path coefficient analysis and genetic divergence is that plant height, number of primary branches, number of secondary branches, length of siliqua , number of seeds per siliqua , biological yield per plant, seed yield per plant and harvesting index found the most important component characters. Hence, these traits should be considered as selection criteria for yield improvement in mustard.

## REFERENCES

- [1] Hemingway J. S. (1976). Mustard: *Brassica spp.* and *Sinapis alba (Cruciferae)*. pp 56-9. (in) *Evolution of Crop Plants*. Simmonds (Ed.). Longmans, London.
- [2] USDA, N. (1999). United States department of agriculture. Natural Resources Conservation Service. *Plants Database*. <http://plants.usda.gov> (accessed in 2000).
- [3] Dewey, D. R. and Lu. K. H. (1959). Correlation and path coefficient analysis of seed production, *Agron. J.*, 51: 515-518.
- [4] Panse V. G. and Sukhatme P. V. (1961). Statistical methods for agricultural workers, ICAR, New Delhi.
- [5] Burton, G. W. and De Vane, E. H. (1953). Estimating Heritability in Tall Fescue (*Festuca Arundinacea*) from Replicated Clonal Material. *Agronomy Journal*, 45(10), 478-481.
- [6] Johanson, H. W., Robinson, H. F. and Comstock, R. E. 1955. Estimates of genetic and environmental variability in Soyabean. *Agronomy J.*, 47(7): 314- 315.
- [7] Al-Jibouri, H. and Miller, P. and Robinson, H. (1958). Genotypic and Environmental Variances and Covariances in an Upland Cotton Cross of Interspecific Origin<sup>1</sup>. *Agronomy Journal - AGRON J.*, 50. 10.
- [8] Wright S. (1921). Correlation and causation. *J. agric. Res.*, 20: 557-587.
- [9] Mahalanobis, P. C. (1936). On the generalized distance in statistics. National Institute of Science of India.
- [10] Sohan, R., and Verma, N. (2010). Genetic Variability and yield components in Indian mustard, *Journal of Oilseeds Research*, 27 (2):170-171.
- [11] Akabari, V. R. and Niranjana, M. (2015). Genetic variability and trait association studies in Indian mustard (*Brassica juncea* L.). *International Journal of Agricultural Sciences*, 11. 35-39.

- [12] Devi, T. R., Devi, N. D., Vivekananda, Y. and Sharma, P. R. (2017). Genetic diversity analysis in Indian mustard (*Brassica juncea* L. Czern and Coss) genotypes using agromorphological parameters. *Electronic Journal of Plant Breeding*, 8(3), 749-753.
- [13] Kumar, R., Kaur, S., Bala, K., Kaur, S. and Sharma, L., (2019). Assessment of Genetic Variability, Correlation and Path Analysis for Yield Traits in F1 Hybrids of Indian Mustard [*Brassica juncea* (L.) Czern & Coss.]. *Agriways*, 7(1): 1-7.
- [14] Chowdhury, P. R and Goswami, G. D. (1991). Genetic variability studies in Indian mustard (*Brassica juncea* L. Czern. &Coss.). *Environ. and Ecol.*,9(4):1003- 1006.
- [15] Pant, S. C. and Singh, P. (2001). Genetic variability in Indian mustard. *Agricultural Science Digest.*, 21(1):28-30.
- [16] Singh, Poonam, Singh, D. N., and Chakraborty, M. (2003). Variability, heritability and genetic advance in Indian mustard (*Brassica juncea* L.).
- [17] Rai, S. K., Verma, A. and Pandey, D. D. (2005). Genetic variability and character association analysis in Indian mustard (*Brassica juncea* (L.) Czern and Coss.). *Annals of Agri. Bio. Res.*, 10(1):29-34.
- [18] Singh, S. K., Singh, A. K. and Kumar, K. (2007). Genetic variability for yield and its components in Indian mustard (*Brassica juncea* L. Czern & Coss.). *New Botanist*. 34 (1/4):147-150.
- [19] Yadav, V. K., Srivastava, K. K., Mishra, V. K. and Negi, S. (2021). Studies on Genetic Variability, Character Association and Genetic Divergence in Indian Mustard (*Brassica juncea* L. Czern and Coss.). *Int. J. Curr. Microbiol. App. Sci.*, 10: 132-143.
- [20] Devi, B. (2018). Correlation and path analysis in Indian mustard [*Brassica juncea* (L.)] In agro – climatic conditions of Jhansi (U.P.), *Journal of Pharmacognosy and Phytochemistry*, 7(1): 1678-1681.
- [21] Gangapur, D. R., Prakash, B. G. and Hiremath, C. P. (2011). Genetic variability studies of germplasm accessions in Indian mustard under protected and unprotected conditions at Bijapur, Karnataka. *International Journal of Agricultural Sciences*, 7(1), 54-57.
- [22] Singh, M., Tomar, A., Mishra, C. N. and Srivastava, S. B. L. (2011). Genetic parameters and character association studies in Indian Mustard. *Journal of Oilseed Brassica*, 2(1): 35-38.
- [23] Kumar, H., Srivastava, Ayushi, Vishwakarma, M. K. and Lal, J. P. (2012). Genetic Enhancement of Variability through Induced Mutagenesis in Two Genotypes of (*Brassica napus* L.) *Madras Agric. J.*, 99 (4-6): 228-231.
- [24] Nasim, A., Farhatullah, Iqbal, S., Shah, S., and Azam, S. M. (2013). Genetic variability and correlation studies for morphological traits in (*Brassica napus* L.). *Pak. J. Bot.* 45(4): 1229-1234.
- [25] Singh, Maharaj, and Chauhan. (2013). Genetic variation and correlation for some physiological character and seed yield in Indian Mustard (*Brassica juncea* L.) Under Rain fed Condition. *J. Oilseeds Res.* 30(2): 167-170.

- [26] Tripathi, N., Kumar, K. and Verma, O.P. (2013). Genetic variability, heritability and genetic advance in Indian Mustard (*Brassica juncea* L. Czern and Coss.) for seed yield and its contributing attributes under normal and saline/alkaline condition. *International Journal of Science and Research*. 983-985.
- [27] Meena, P., Chauhan, J.S., Singh, M., Singh, K.H., Rathore, S.S. and Meena, M.L. (2014). Genetic parameters and correlations for seed yield and morphological characters in Indian Mustard [*Brassica juncea* (L.) Czern. & Coss.]. *J. Oilseeds Res.*, 31(2): 114-117.
- [28] Bind, D., Singh, D. and Dwivedi, V. K. (2014). Genetic variability and character association in Indian mustard [*Brassica Juncea* (L.) czerns & coss.], *Agric. Sci. Digest.*, 34 (3): 183 – 188.
- [29] Synrem, G. I., Rangare, N. R., Myrthong, I. and Bahadure, D. M. (2014). Variability studies in intra specific crosses of indian mustard [*Brassica juncea* (L.) Czernand Coss.] Genotypes. *Iosr Journal of Agriculture and Veterinary Science (Iosr-Javs)*. 7 (9): 29-32.
- [30] Rauf, M. A., and Rahim, M. A. (2018). Genetic variability studies among yield and its contributing traits in mustard (*Brassica napus* L.). *Adv. Zool. Bot.*, 6, 101–108.
- [31] Dawar, S., Kumar, N., and Mishra, S. P. (2018). Genetic Variability, Correlation and Path Coefficient Analysis in the Indian Mustard (*Brassica juncea* L. Czern and Coss) Varieties Grown in Chitrakoot, India. *Int. J. Curr. Microbiol. App. Sci.*, 7(3): 883-890.
- [32] Nagda, R., Dubey, D., Avinashe, H. and Tamatam, D. (2018). Assessment of Genetic Diversity in Mustard Genotypes. *Plant Archives*, Vol. 18 No. 2, 2018pp. 2091- 2096.
- [33] Doddabhimappa, R., Gangapur, B., Prakash, G. and Hiremath, C. P. (2010). Genetic diversity analysis of Indian mustard (*Brassica juncea* L.). *Electronic Journal of Plant Breeding*, 1(4), 407-413.
- [34] Chaurasia, R. K, Ram, B. and Gr Chougule. (2014). Genetic divergence for seed yield and component traits in Indian mustard [*Brassica juncea* (L.) Czern & Coss.] G. B. Pant University of Agriculture & Technology, Pantnagar (Uttarakhand). *AGRES-An international e-journal*, 3(3): 260-269.
- [35] Gupta, A., Pant, N. C., Dwivedi, U., Tiwari, S., Pandey, C. S., Dhoundiyal, R. and Verma, O. P. (2018). Studies on correlation and path coefficient analysis for yield and yield related traits in Indian mustard (*Brassica juncea* L. Czern & Coss.) under timely and late sown conditions. *Journal of Pharmacognosy and Phytochemistry*, 7(2), 2545-2551.