

## ABSTRACT

Okra [*Abelmoschus esculentus* L., Moench] a native of South-Africa and commonly known as ‘Bhindi’ is an annual malvaceous vegetable crop, especially grown in tropical and subtropical climates. It is also called “perfect villager’s vegetable”. Root-knot nematodes are considered among the top five major plant pathogens and the first among the ten most important genera of plant parasitic nematodes in the world. Amongst nematodes, root-knot nematode (*Meloidogyne incognita* and *Meloidogyne javanica*) causes severe damage to okra. *Trichoderma viride*, *Pseudomonas fluorescense* and *Rhizobium* were tested under field conditions during Rabi 2020-2021 for their efficacy against the Root knot nematode and growth & nematode population. Among the treatments the maximum plant height(cm) at 60 DAS was recorded in T<sub>6</sub> -Vermicompost+ spent mushroom compost+ *Trichoderma viride*+ *Pseudomonas fluorescense* + *Rhizobium* (51.50 cm) followed by T<sub>5</sub> – Vermicompost +spent mushroom compost+ *Trichoderma viride* + *Rhizobium* (44.45 cm) as compared to untreated check control T<sub>0</sub> – (25.13 cm). The maximum Root weight was recorded in T<sub>3</sub>– Vermicompost +spent mushroom compost + *Rhizobium* (9.90 gm) followed by T<sub>1</sub> - Vermicompost+ spent mushroom compost+ *Trichoderma viride*(9.65gm) as compared to untreated check control T<sub>0</sub>-(10.25gm). The maximum Root length was recorded in T<sub>6</sub> - vermicompost + spent mushroom compost+ *Trichoderma viride* + *Pseudomonas fluorescense* + *Rhizobium* (13.70gm) followed by T<sub>4</sub> –vermicompost + spent mushroom compost + *Trichoderma viride* + *Pseudomonas fluorescense* (13.35gm) as compared to untreated check control T<sub>0</sub> – (10.12). The minimum nematode intensity (%) at 60 DAS was recorded in T<sub>6</sub> – Vermicompost+ spent mushroom compost + *Trichoderma viride* + *pseudomonas fluorescense* + *Rhizobium* (17), followed by T<sub>4</sub> – Vermicompost +spent mushroom compost + *Trichoderma viride* + *Pseudomonas fluorescense* (25.50) as compared to untreated check control T<sub>0</sub>(57).

**Keywords:** *Meloidogyne incognita*, *pseudomonas fluorescense*, *Rhizobium*, Root-knot Nematode of Okra, *Trichoderma viride*.

## 1. INTRODUCTION

Okra [*Abelmoschus esculentus* L. Moench] a native of South- Africa and commonly known as ‘Bhindi’ is an annual malvaceous vegetable crop, especially grown in tropical and subtropical climates [1]. India ranks first in the world with 3.5 million tonnes

(70% of the total world production) of okra produced from over 0.35-million-hectare land[2]. In India, total grown in an area of 511 hectare having total production of 6219 million tons with the productivity of 12.0 million tons per hectare[3]. Dry seed contains 13-22% edible oil; and 20-24% protein [4]. The production of okra also suffers from several pests (fruit borer, white flies, jassids, thrips, mites etc.) and pathogens (yellow vein mosaic, powdery mildew, root rot etc.) including nematodes [5]. Plant parasitic nematodes viz. *Meloidogyne incognita*, *Meloidogyne javanica*, *Rotylenchulus* spp., *Tylenchorhynchus* spp., *Hoplolaimus* spp., *Aphelenchus avenae*, *Nothotylenchus* spp., *Helicotylenchus* spp., *Hemicriconemoides* spp., *Longidorus* spp., *Paralongidorus* spp., *Trichodorus* spp., *Paratrichodorus* spp., *Ditylenchus* spp., *Tylenchus* spp., *Rotylenchulus* spp., and *Xiphinema* spp., were found associated with okra in India [6]. Root-knot nematode (*Meloidogyne incognita* and *Meloidogyne javanica*) causes severe damage to okra [7]. Severe attack of root-knot disease caused by *Meloidogyne* spp[8]. On okra and yield losses up to 27% [9]. Rootknot nematodes (RKN) are sedentary endo-parasite and is among the most damaging agricultural pests, attacking a wide range of crops [10].

## 2. MATERIALS AND METHODS

Keeping in view, the experiment was conducted in nematode infested soil at the courtyard of Department of Plant Pathology, SHUATS, Prayagraj, Uttar Pradesh during Rabi season in the year 2020-2021. The soil sample was collected from the infested field and processed in laboratory by following cobb's decanting and sieving technique followed by modified Baermann funnel technique to estimate the nematode population. Before laying out the experiment it was assured that the experimental field possess 2 larvae/gm of soil.

The selected field was dug up and the soil become pulverized and then whole location was divided into sub-plots and specified in randomized block design with six treatments viz., vermi compost @ 8 t/ha, spent mushroom compost @ 8 t/ha was used as basal application. These were incorporated into the soil by forming specific ridges according to the crop spacing and covered by thin layer of soil. The field was irrigated for fifteen days at regular intervals to enhance decomposition process. After fifteen days, seed treatment was done with *Trichoderma* @10gm/kg, *Pseudomonas* @10gm/kg, *Rhizobium*@5gm/kg of seed where, vermi compost and spent mushroom compost was incorporated initially. Each treatment was replicated four times with plot size of  $2.5 \times 1 \text{ m}^2$  each and local variety seed was sown with a spacing of  $45 \times 30 \text{ cm}$ . Root knots in the root system and plant growth parameters of okra was recorded at 30, 45, 60 days after sowing of the crop. The records have been subjected to the statistical analysis.

At 60 days after sowing the root knots in the roots of okra are identified. The galled roots were removed and washed thoroughly with sterile water. a gall is placed on the sterile slide using sterile forceps and teased using a sterile needle and examined under microscope. Eggs and female *Meloidogyne* were identified when observed under microscope. Mature females are swollen, melon like with elongate neck at anterior end, forms perineal patterns, short stylet with well-developed basal knobs, eggs laid in gelatinous matrix outside the body, tail absent. Males are vermiform, 1.5-2.0 mm long, basal knobs; oesophageal glands overlap intestine ventrally; tail elongate conoid with pointed tip.

### 3. RESULTS

The result presented in table 1 revealed that all the treatments were statistically significant and decreased the number of root knots in the roots of okra as compared to control. Among the bio agents used, the treatment T6- Vermicompost + SMC+ *Trichoderma* spp.+ *Pseudomonas* spp.+ *Rhizobium* (17) significantly decreased the root knots in the root system (57) in okra when compared to other bio agents. The treatments (T<sub>6</sub>, T<sub>4</sub>), (T<sub>2</sub>, T<sub>5</sub>) and (T<sub>1</sub>, T<sub>3</sub>) are not significantly differ from each other.

**Table 1: Effect of compost along with bio-agents on the number of root knot nematodes on the roots of okra at 60 DAS.**

Tr no.	Treatments	Number of root knots
T <sub>0</sub>	Control+ Vermicompost+ SMC	57
T <sub>1</sub>	Vermicompost+ SMC+ <i>Trichoderma</i> spp.	50
T <sub>2</sub>	Vermicompost + SMC+ <i>Pseudomonas</i> spp	31
T <sub>3</sub>	Vermicompost+ SMC+ <i>Rhizobium</i>	51
T <sub>4</sub>	Vermicompost + SMC+ <i>Trichoderma</i> spp.+ <i>Pseudomonas</i> spp.	26
T <sub>5</sub>	Vermicompost + SMC+ <i>Trichoderma</i> spp.+ <i>Rhizobium</i>	34
T <sub>6</sub>	Vermicompost+ SMC+ <i>Trichoderma</i> spp.+ <i>Pseudomonas</i> spp.+ <i>Rhizobium</i>	17
	F- test	S
	S. E. (d) ±	1.91
	C. D. (5%)	3.93



**Plate 1: Root knots in the root system of okra at 60 days after sowing.**

The result presented in table 2 revealed that all the treatments were statistically significant and increased the plant growth parameters of okra. Among the Bio agents, the treatment T<sub>6</sub>- Vermicompost + SMC + *Trichoderma* spp. + *Pseudomonas* spp. + *Rhizobium* (19.01cm) significantly increased the plant height of okra. Among the Bio agents the treatments (T<sub>6</sub>,T<sub>4</sub>), (T<sub>2</sub>,T<sub>4</sub>), (T<sub>2</sub>,T<sub>5</sub>) and (T<sub>1</sub>,T<sub>5</sub>) found non-significant to each other. The treatment T<sub>6</sub>- Vermicompost + SMC + *Trichoderma* spp. + *Pseudomonas* spp. + *Rhizobium* (37.20 cm) significantly increased the plant height of okra. Among the Bio agents the treatment (T<sub>1</sub>,T<sub>3</sub>) found non-significant to each other. The treatment T<sub>6</sub>- Vermicompost + SMC + *Trichoderma* spp. + *Pseudomonas* spp. + *Rhizobium* (51.50 cm) significantly increased the plant height of okra. Among the treatments (T<sub>6</sub>, T<sub>4</sub>), (T<sub>2</sub>,T<sub>4</sub>), (T<sub>2</sub>,T<sub>5</sub>) and (T<sub>1</sub>,T<sub>5</sub>) found non-significant to each other

**Table 2: Effect of bio agents along with bio agents on plant growth parameters of okra:**


Treatments	Plant height (cm)	Root length (cm)	Root weight (gm)
T <sub>0</sub>	25.13	10.12	10.25
T <sub>1</sub>	35.27	11.50	9.65
T <sub>2</sub>	41.85	12.62	8.93
T <sub>3</sub>	32.35	11	9.90
T <sub>4</sub>	44.45	13.35	8.43
T <sub>5</sub>	38.35	12.20	9.13
T <sub>6</sub>	51.50	13.70	8.25
F-test	S	S	S
S. E (d) ±	0.75	0.29	0.40
C.D. (5%)	1.58	0.60	0.84

The result presented in table 2 revealed that all the treatments were statistically significant and decreased the root weight of cowpea as compare to control. The treatments T<sub>6</sub>- Vermicompost + SMC + *Trichoderma* spp.+ *Pseudomonas* spp.+ *Rhizobium* (8.25 gm) significantly decreased the root weight due to the less number of root knots in root system of okra. The treatments (T<sub>3</sub>,T<sub>1</sub>), (T<sub>3</sub>,T<sub>1</sub>,T<sub>5</sub>), (T<sub>1</sub>,T<sub>5</sub>,T<sub>2</sub>), (T<sub>5</sub>, T<sub>2</sub>, T<sub>4</sub>) and (T<sub>2</sub>, T<sub>4</sub>,T<sub>6</sub>) found non-significant to each other. Highest root weight was recorded in control T<sub>0</sub>- (10.25 gm) due to higher number of root knots in the root system of cowpea. The present investigation indicates that Vermicompost +

SMC + *Trichoderma* spp.+ *Pseudomonas* spp.+ *Rhizobium* into the soil used as an effective treatment for root knot nematodes and to develop eco-friendly strategy for the management of root knot nematodes of okra.

#### 4. DISCUSSION

Probable reason for such finding may be due to the inhibitory impact of bio agents because of the nemato-toxic compounds present in the bio agents which help to reduce the severity of the nematodes in the soil and plants [11]. Application of *Trichoderma* had a significant effect on root knot nematode population and found toxic to *Meloidogyne* spp. due to Myco-parasitism mechanisms involved in the antagonisms of *Trichoderma*.



**Fig 1: Effect of bio agents along with bio agents on plant growth parameters of okra.**

#### 5. CONCLUSION

In the present study on the basis of observation, it was found that for managing the root knots in the root system of okra, Vermicompost + SMC+ *Trichoderma* spp.+ *Pseudomonas* spp.+ *Rhizobium* (17) was significant in comparison to control (57). Hence from present study it can be concluded that Vermicompost + SMC+ *Trichoderma* spp.+ *Pseudomonas* spp.+ *Rhizobium* can be used effectively to reduce the root knots and to increase the plant growth of okra.

#### References

- 1.Chin, A.H.G. and Nurshiwan, Z. (1990). Properties of vegetable oils from some unemployed source in Malaysia. M.A.R.D.I. Research. Journal., 18(2);261-265.
- 2.FAOSTAT., (2008). (<http://www.fao.org>)

3. Anonymous (2019 a). Indian Horticulture Data Base At a glance, <http://nhb.gov.in/database>
4. Anonymous (2019b). District wise estimated area & production of horticultural crops 2019b. <https://doh.gujarat.gov.in>.
5. Mishra, J.P. (2002). Okra. In: Handbook of Horticulture (Ed. K.L. Chadha), *India Council of Agricultural Research*, New Delhi, pp 422-427.
6. Bhatti, D.S. and Jain, R.K. (1977). Estimation of losses in okra, tomato and brinjal yield due to *Meloidogyne incognita*. *Indian Journal of Nematology* 7(1): 37-41.
7. Rangasamy, S.D., Parvatha Reddy, P. and Nagesh, M. (2000). Evaluation of biocontrol agents (*Pasteuria penetrans* and *Trichoderma viride*) and botanicals for the management of root-knot nematode, *Meloidogyne. incognita* infecting tomato. *Pest Management in Horticultural Ecosystem*. 6(1): 135-138.
8. Jung C. and Wyss. (1999). New approaches to control plant parasitic nematodes. *Applied Microbiology and Biotechnology*. 51:439-446.
9. Ravichandra, N.G. and Krishnappa, K. (2004). Prevalence and distribution of phyto parasitic nematodes associated with major vegetable crops in Mandya district, Karnataka. *Indian Journal of Nematology*. 34(1): 113-116.
10. Yadav, B.S., Verma, A.C., Mathur, B.N. and Varma, M.K. (1969). Nematodes of cereals and vegetables grown in Rajasthan. *All India Nematology Symposium.*, IARI, New Delhi: 21-22.11.
11. Tariq, M. (2018). Management of Root-Knot Nematode, *Meloidogyne incognita*, in Tomato with Two *Trichoderma Species* *Pakistan Journal of Zoology.*, vol.50(4), pp 1589-1592