

Original Research Article

Impact of different approaches of nutrient recommendations for aerobic on soil fertility of Alfisols of Eastern Dry Zone of Karnataka

Abstract

A field experiment to study the influence of soil test crop response based fertilization on nutrient requirement, nutrient uptake and nutrient use efficiency of aerobic rice was conducted during Kharif 2020 at Zonal Agricultural Research Station (ZARS), GKVK, Bengaluru. The experiment was laid out in RCBD comprising twelve treatments replicated thrice. The results revealed that significantly higher grain yield (68.85 q ha^{-1}) was recorded in treatment receiving fertilizer nutrients based on STCR inorganic approach for the targeted yield of 65 q ha^{-1} based on predicted soil test values which was superior compared to Low-Medium-High approach and Blanket recommendation. The higher post-harvest soil available nutrient status was registered in STCR integrated approach based on predicted soil test values compared to package of practice and LMH approach..

Introduction

The food security in Asia is challenged by increasing food demand and is threatened by declining availability of water with growing population, increased urbanisation and environmental degradation. In India, rice occupies an area of 43.79 million hectare with production of 112.91 million tonnes with an average productivity of 2578 kg per hectare. In Karnataka, rice is being grown in an area of 1.32 m ha with an annual production of 4.24 m t and productivity is 3338 kg ha^{-1} [1]. With emerging water scarcity in many parts of the world, the traditional way of lowland rice cultivation can no longer be sustained. Along with high water requirement, traditional system of rice production in long run leads to destruction of soil aggregates and reduction in macropore volumes [2]. Moreover, lack of rainfall is a major production constraint in rainfed areas. Alternatives to the conventional flooded rice cultivation were developed worldwide to reduce water consumption and to produce more rice with less water. Among the different water saving strategies, “aerobic rice” is considered a promising cultivation system for water scarce areas.

Aerobic rice is broadly defined as “a production system in which, direct seeding of high yielding and input responsive rice cultivars with aerobic adaptation grown in non-puddle, non-flooded and non-saturated soil during the entire growing period” [3]. It is a new concept of reducing water requirement for rice in which rice is grown like an upland crop with high inputs and supplementary irrigations, when rainfall is insufficient. Although India has made considerable advances in agricultural research, still the blanket recommendation of cultivation practices for adoption over larger areas are in vogue.

The current energy crisis prevailing, higher prices and lack of proper supply system of fertilizers, deterioration of soil fertility calls for more effective nutrient management practices using manure and fertilizers judiciously to sustain yield levels [4]. The effective nutrient management involves site specific nutrient recommendations that include timely and balanced fertilizer nutrient application, using appropriate methods and practicing integrated plant nutrient supply system using chemical fertilizers, organic manures, crop residues and biofertilizers.

Considering high cost of fertilizers and their adverse environmental implications, fertilizer recommendations based on soil test values, residual effect and yield targets becomes highly important. This can be achieved by following targeted yield approach involving integrated plant nutrition system (IPNS) for enhancing crop productivity, nutrient use efficiency as well as soil nutrient balance [5].

Material and methods

The field experiment was conducted on aerobic rice during 2020-21 at Zonal Agricultural research station, University of Agricultural Sciences, GKVK, Bangalore (13° 04' 55.2" N latitude, 77° 34' 10.0" E longitude). The experimental soil was sandy loam in texture belonging to vijayapura series of great group *Kandic paleustalfs* and had pH of 5.77, electrical conductivity 0.085 dS m⁻¹ (1: 2.5 soil : water ratio) and organic carbon 4.44 g kg⁻¹. Treatments schedule for aerobic rice comprising of T₁: STCR through inorganics (65 q ha⁻¹) - Actual STV* (*Soil Test Value), T₂: STCR through inorganics (65 q ha⁻¹) - Predicted STV, T₃: STCR through integrated (65 q ha⁻¹) - Actual STV, T₄: STCR through integrated (65 q ha⁻¹) - Predicted STV, T₅: STCR through inorganics (55 q ha⁻¹) - Actual STV, T₆: STCR through inorganics (55 q ha⁻¹) - Predicted STV, T₇: STCR through integrated (55 q ha⁻¹) - Actual

STV, T₈: STCR through integrated (55 q ha⁻¹) - Predicted STV, T₉: Package of practice, T₁₀: LMH (STL) - Actual STV, T₁₁: LMH (STL) - Predicted STV and T₁₂: Absolute control.

The following STCR fertilizer adjustment equation and post harvest soil test value prediction equations developed by AICRP on STCR, UAS, Bengaluru centre under *Alfisols* of Eastern Dry Zone of Karnataka was used for STCR treatments and to predicted the post harvest soil test value for the preceding crop of aerobic rice (dry chilli) which can be used as initial soil test value for the present investigation to prescribe the fertilizer dose. More details are regarding development of targeted yield equations and post harvest soil test value prediction equations are provided in Ph.D. thesis on “Development of targeted yield equation for aerobic rice and its evaluation on *Alfisols* of Eastern dry zone of Karnataka” [6] at the same experimental site. The quantity of fertilizer nutrients (NPK) applied for each treatment is mentioned in Table 2.

Prediction equation	R ² value
Inorganic approach	
PHN = 188.752 + 0.001 ^{**} SN + 0.203 FN - 0.184 UN	0.610 ^{**}
PHP = - 6.133 + 1.089 ^{**} SP + 1.188 ^{**} FP - 1.299 [*] UP	0.965 ^{**}
PHK = 5.075 + 1.138 ^{**} SK + 1.275 ^{**} FK - 0.249 UK	0.925 ^{**}
IPNS approach	
PHN = 191.090 ^{**} - 0.003 SN + 0.087 ^{**} FP - 0.008 UN	0.442 ^{**}
PHP = 7.325 + 0.721 ^{**} SP + 1.167 ^{**} FP + 2.515 ^{**} UP	0.890 ^{**}

Table 1: Post

STCR- Inorganics (NPK alone) equation

$$F.N. = 3.02879 T - 0.20314 STV - N$$

$$F.P_2O_5 = 1.24589 T - 0.07368 STV - P_2O_5$$

$$F.K_2O = 1.51168 T - 0.22617 STV - K_2O$$

STCR- IPNS (Integrated plant nutrient supply) equation

$$F.N. = 2.89282 T - 0.20320 STV - N - 0.72978 OM$$

$$F.P_2O_5 = 1.13206 T - 0.06960 STV - P_2O_5 - 0.48911 OM$$

$$F.K_2O = 1.50402 T - 0.21105 STV - K_2O - 0.42410 OM$$

harvest soil test value prediction equation

Table 2: Quantity of fertilizer nutrients and poultry manure applied through different approaches as per the treatments and soil test values

Treatments	Soil test values			Poultry manure applied	Fertilizer nutrients applied		
	N	P ₂ O ₅	K ₂ O		N	P ₂ O ₅	K ₂ O
	kg ha ⁻¹				t ha ⁻¹	kg ha ⁻¹	
T ₁	260.59	101.05	271.64	0	143.94	73.54	36.82
T ₂	207.10	174.48	298.07	0	154.80	68.13	30.84
T ₃	261.67	106.92	305.92	10	127.56	61.25	28.96
T ₄	205.62	167.75	456.03	10	138.95	57.08	2.12
T ₅	260.21	99.88	221.67	0	113.72	61.16	33.01
T ₆	202.89	149.36	276.69	0	125.37	56.85	20.56
T ₇	262.08	93.65	245.75	10	98.55	50.85	26.62
T ₈	201.60	145.27	340.47	10	110.84	47.26	6.62
T ₉	272.53	115.60	286.76	10	100.00	50.00	50.00
T ₁₀	266.56	98.55	285.69	10	125.00	37.50	50.00
T ₁₁	204.08	165.92	322.27	10	125.00	37.50	45.83
T ₁₂	243.41	56.96	168.13	0	0.00	0.00	0.00

There were twelve treatments replicated three times in a randomized block design.

Aerobic rice seeds (Var. MAS 946-1) were sowed in rows at proper spacing in the first week of July 2020, after basal application of fertilizers as per treatments. The remaining half dose of N was top-dressed in two splits at tillering stage and boot stage. The crop was cultivated adopting proper package of practices. All climatic conditions were favourable for growth and development of the crop. The soil samples were collected before sowing and after the harvest at 0-15 cm soil depth. Basic soil parameters were estimated by using standard laboratory procedures outlined by Jackson [7] and nutrient balance was worked out.

3. Results and Discussion

3.1 Grain yield of aerobic rice

Significantly higher grain yield (Table 3) of 68.85 q ha⁻¹ was recorded with the application of nutrients based on STCR approach for the targeted yield of 65 q ha⁻¹ through

inorganic based on predicted soil test values (T_2) compared to treatment T_8 (60.14 q ha^{-1}) [STCR integrated (55 q ha^{-1}) - Predicted STV], T_7 (57.55 q ha^{-1}) [STCR integrated (55 q ha^{-1}) - Actual STV], T_9 (53.25 q ha^{-1}) (Package of practice), T_{11} (49.15 q ha^{-1}) (LMH - predicted STV), T_{10} (48.76 q ha^{-1}) (LMH - Actual STV), and T_{12} (20.66 q ha^{-1}) (Absolute control). However, it was on par with treatments receiving fertilizers through STCR inorganic approach for the targeted yield of 65 q ha^{-1} based on actual soil test values (T_1 : 65.50 q ha^{-1}); STCR integrated approach for the targeted yield of 65 q ha^{-1} based on predicted soil test values (T_4 : 63.79 q ha^{-1}) and actual test values (T_3 : 61.70 q ha^{-1}); STCR inorganic approach for the targeted yield of 55 q ha^{-1} for predicted soil test values (T_6 : 62.96 q ha^{-1}) and actual soil test values (T_5 : 61.58 q ha^{-1}). The higher yield in STCR treatments could be attributed to the ability of targeted yield approaches to satisfy the nutrient demand of crop more efficiently. These findings are in close accordance with those reported by Kumar and Paramananda, 2018 [8] who opined that application of fertilizers based on STCR approach at critical physiological phases would have supported for better assimilation of photosynthates towards grain.

3.2 Soil pH

After harvesting rice crop, the soil pH was increased compared to the respective initial soil pH but, no significant difference was found with respect to post harvest soil pH among the treatments (Table 3). However, numerically higher pH value (6.02) was recorded in absolute control where no fertilizers were applied (T_{12}) while the lower pH value (5.55) was found in STCR inorganic approach for the targeted yield of 55 q ha^{-1} using actual soil test values (T_5). The soil pH was higher in integrated approach irrespective of target and actual or predicted soil test values compared to inorganics which might be attributed to release of basic cations from poultry manure during its decomposition and source of phosphatic fertilizer used was single super phosphate which contains calcium, which neutralizes soil acidity to some extent [9][10].

3.3 Soil EC

The data clearly indicated that EC values of post harvest soil in all the treatments increased compared to the respective initial treatments. Similarly, the electrical conductivity (EC) values of the post-harvest soil indicated a slight increase in all the treatments as compared to control (0.081 dS m^{-1}) (Table 3). The significantly higher EC (0.104 dS m^{-1}) was found where fertilizers were applied through STCR approach for a targeted yield of 55 q ha^{-1}

through inorganics (T_6) based on predicted soil test values compared to control (T_{12} : 0.081) and the remaining treatments were on par. The results indicates that the EC was higher where fertilizers were applied based on predicted soil test values and application of poultry manure in integrated approach. The increase in soil EC after harvest of aerobic rice in STCR inorganic and integrated approach through predicted soil test values may be due to release of soluble salts from poultry manure upon decomposition in integrated approach and direct application of slightly higher dose of inorganic fertilizers with predicted soil test value that might have caused higher EC values [11].

3.4 Soil organic carbon

The organic carbon content of the post-harvest soil was increased compared to initial and it was found non-significant among the different treatments (Table 3). However, numerically higher organic carbon content (0.52 %) was recorded in STCR target of 65 q ha⁻¹ integrated approach with PM (T_4) based on predicted soil test values, which was on par with LMH approach with predicted soil test value based fertilizer dose whereas, lower value (0.44 %) was recorded in STCR target 55 q ha⁻¹ through inorganics for actual soil test values (T_5). The organic carbon content was found higher in STCR integrated treatments, LMH and RDF approach compared to STCR inorganic approach due to application of poultry manure at 10 t ha⁻¹ in these treatments. This was mainly due to application of poultry manure at the rate of 10 t ha⁻¹ and root biomass of rice crop upon gradual decomposition substantially contributed to pool of soil organic carbon in contrast to STCR inorganic treatments where no poultry manure was added [12]. These increase/ maintenance in organic carbon content due to use of fertilizers can be attributed to contribution of biomass to the soil in the form of crop stubbles and residues.

3.5 Available nitrogen

Significantly higher (280.37 kg ha⁻¹) available nitrogen was recorded in STCR integrated approach for the targeted yield of 65 q ha⁻¹ based on predicted soil test values (T_4) compared to absolute control (T_{12}) where the available nitrogen was 239.12 kg ha⁻¹, but it was found to be on par with all the remaining treatments. Soil available nitrogen after harvest of aerobic rice was improved in all the treatments of fertilizer nutrient application (both integrated and inorganic approach) except in absolute control (T_{12}) where soil available nitrogen was reduced over its initial content (Table 4). The improved available nitrogen after

harvest of rice crop was mainly due to mineralization of applied poultry manure along with direct addition of inorganic nitrogen fertilizers which contributed to the pool of available nitrogen and might have improved water and nutrient holding capacity in integrated approach in contrast with other STCR inorganic treatments [13].

3.6 Available P_2O_5

Available P_2O_5 decreased from the initial level in all the treatments except control and all the STCR treatments of 55 q ha⁻¹ where slightly increased (Table 4). Significantly higher (109.34 kg P_2O_5 ha⁻¹) value was found in STCR yield target 65 q ha⁻¹ through inorganics based on predicted soil test values (T_2) compared to LMH approach where nutrients were applied based on predicted (T_{11} : 93.74 kg P_2O_5 ha⁻¹) and actual (T_{10} : 88.61 kg P_2O_5 ha⁻¹) soil test values and with control plots (T_{12} : 57.51 kg P_2O_5 ha⁻¹) where no fertilizers and manure was applied. Interestingly, no significant difference was noticed among actual and predicted soil test value based fertilizer recommendation under STCR inorganic and integrated approach at both the targets and LMH approach. The significantly higher available phosphorus in STCR inorganic approach compared to integrated approach could be attributed to application of lower dose of inorganic fertilizers and lower initial available phosphorus content in soil. However no significant difference was observed between inorganic and integrated approach. The soils on which the present study was conducted are acidic in reaction (pH 5.77) where Al and Fe ion concentrations may be higher hence, the response to phosphatic fertilizer was high and had higher available phosphorus content in soil. These results are in conformity with findings of Ashwini (2007) [14].

3.7 Available K_2O

Application of nutrient doses as per STCR integrated approach (T_4) for the targeted yield of 65 q ha⁻¹ based on predicted soil test values recorded significantly higher available potassium content (297.52 kg ha⁻¹) (Table 4) in soil after harvest of aerobic rice compared to treatment T_2 [STCR inorganics (65 q ha⁻¹) - Predicted STV] (229.08), T_6 [STCR inorganics (55 q ha⁻¹) - Predicted STV] (222.40 kg ha⁻¹), T_7 [STCR integrated (55 q ha⁻¹) - Actual STV] (221.88 kg ha⁻¹), T_1 [STCR inorganics (65 q ha⁻¹) - Actual STV] (217.36 kg ha⁻¹), T_5 [STCR inorganics (55 q ha⁻¹) - Actual STV] (216.60 kg ha⁻¹) and T_{12} (Absolute control) (166.20 kg ha⁻¹). In all the treatments of different fertilizer nutrient recommendations available potassium content after harvest of the crop was found to be reduced. Higher available

potassium in T₄ treatment even without application of potassium fertilizer might be due to higher potassium content in native soil (316.49 kg ha⁻¹) and also contribution to the pool of available potassium in soil through mineralization of applied poultry manure (Rajput *et al.*, 2016).

3.8 Available sulphur

Significantly higher available sulphur content (35.00 mg kg⁻¹) was recorded in T₈ [STCR integrated (55 q ha⁻¹) - Predicted STV] as compared to STCR target of 65 q ha⁻¹ through inorganic approach using actual soil test values (27.75 mg kg⁻¹) (T₁) Table 5). However, it was found to be on par with all the treatments except in T₁₂ (Absolute control). The sulphur content was higher where poultry manure was applied along with inorganic fertilizers compared to inorganic fertilizers alone. Application of phosphorus through SSP which contain 11 per cent sulphur and mineralization of added poultry manure, which substantially contributed to plant available sulphur. These results are in accordance with Chandrakanth (2015) [15].

3.9 Exchangeable calcium

Significant difference in exchangeable calcium was found between treatments due to different approaches of nutrient recommendations through actual and predicted soil test values (Table 5). Significantly higher [3.33 c mol (p⁺) kg⁻¹] exchangeable calcium was recorded in STCR targeted yield of 55 q ha⁻¹ through integrated approach (T₈) where nutrients were applied using predicted soil test values and lower exchangeable calcium was recorded in absolute control [T₁₂: 2.29 c mol (p⁺) kg⁻¹]. Exchangeable Ca content in soil was decreased from its corresponding initial content after harvest of aerobic rice crop in all the approaches of fertilizer nutrient recommendations including in absolute control due to crop removal. Among inorganic and integrated approach higher calcium content was recorded in integrated approach due to addition of some amount of secondary nutrients from the straight fertilizers particularly SSP which contains 18 per cent of Ca which might have resulted in increase in calcium content and also release of Ca during mineralization of added poultry manure [16].

3.10 Exchangeable magnesium

Perusal of data from Table 5 reveals that there was no significant difference among the treatments with respect to exchangeable magnesium content in post-harvest soils of aerobic rice crop. However, numerically higher value [$1.18 \text{ c mol (p}^+) \text{ kg}^{-1}$] was recorded where fertilizer nutrients were applied through STCR integrated approach based on predicted soil test values for the targeted yield of 65 q ha^{-1} (T_4) and the lower value [$1.00 \text{ c mol (p}^+) \text{ kg}^{-1}$] was recorded in absolute control (T_{12}) where no fertilizers or manures were applied. Exchangeable Mg content in post-harvest soil was decreased from its respective initial content after harvest of rice crop in all the approaches of fertilizer nutrient recommendations due to crop removal.

3.11 Nitrogen balance in soil

The initial available nitrogen in soil ranged from $201.60 \text{ kg N ha}^{-1}$ to $272.53 \text{ kg N ha}^{-1}$ (Table 6) and higher dose of nitrogen was added ($154.80 \text{ kg N ha}^{-1}$) in STCR inorganic approach for the targeted yield of 65 q ha^{-1} based on predicted soil test values (T_2). The maximum uptake of nitrogen ($176.73 \text{ kg N ha}^{-1}$) by aerobic rice was recorded where fertilizer was applied as per STCR inorganic approach based on predicted soil test values for the targeted yield of 65 q ha^{-1} (T_2) followed by STCR target of 55 q ha^{-1} through inorganic approach based on predicted soil test values ($162.21 \text{ kg N ha}^{-1}$) (T_6). Lower uptake of nitrogen ($47.85 \text{ kg N ha}^{-1}$) was recorded in absolute control (T_{12}) where no fertilizers or poultry manure was applied. The higher actual balance ($280.37 \text{ kg N ha}^{-1}$) was recorded in STCR integrated approach for the targeted yield of 65 q ha^{-1} based on predicted soil test values (T_4). However, overall net positive balance ($111.12 \text{ kg N ha}^{-1}$) was higher in T_8 [STCR integrated (55 q ha^{-1}) - Predicted STV] followed by $100.14 \text{ kg N ha}^{-1}$ recorded in T_6 [STCR inorganics (55 q ha^{-1}) - Predicted STV]. The net negative balance of nitrogen ($-12.25 \text{ kg N ha}^{-1}$) was recorded in LMH approach where fertilizer nutrients were applied by considering actual soil test values. The higher actual balance of nitrogen in STCR integrated approaches was due to efficient use of applied nitrogen without wastage. These results were in accordance with the findings of Brar and Singh (1984) [17] who reported that the increase in available N in the post harvest soil might be due to the continuous mineralization of organic sources of N applied along with inorganic fertilizers.

3.12 Phosphorus balance in soil

The initial available phosphorus content in soil ranged from 56.96 kg ha⁻¹ to 174.48 kg ha⁻¹ (Table 7). The higher dose of phosphorus (73.54 kg ha⁻¹) was applied in STCR target of 65 q ha⁻¹ through inorganic approach based on actual soil test values (T₁) where the soil available phosphorus was low (101.05 kg ha⁻¹) and the target was high (65 q ha⁻¹), whereas lower dose was applied in T₁₀ and T₁₁ (LMH approach through actual and predicted soil test values respectively). The maximum uptake of phosphorus (59.43 kg ha⁻¹) by aerobic rice was recorded where NPK fertilizers were applied as per STCR approach through inorganics based on predicted soil test values (T₂), whereas lower uptake was recorded in absolute control (17.48 kg ha⁻¹) where no fertilizers or poultry manure was applied. The higher actual balance (109.34 kg ha⁻¹) was recorded in T₂ [STCR inorganics (65 q ha⁻¹) - Predicted STV]. Interestingly, all the treatments of various approaches of fertilizer recommendations except absolute control and STCR target of 55 q ha⁻¹ through integrated approach based on actual soil test values recorded net negative balance of phosphorus may due to fixation of applied phosphorus. This could be due to conversion of plant available form of phosphorus to plant unavailable form (Al-P, Fe-P and Ca-P). Tomar (2000) [18] reported that nearly 60 to 70 per cent of the applied phosphorus has been found to remain fixed in the form of Al-P, Fe-P and Ca-P after harvest of rice crop, which corroborated with the results of the present study.

3.13 Potassium balance in soil

The initial available potassium content in soil ranged from 168.13 kg ha⁻¹ to 456.03 kg ha⁻¹ (Table 8). The higher dose of applied potassium (50 kg ha⁻¹) was noticed in LMH approach based on both actual and predicted soil test values, whereas lower dose was applied (2.12 kg ha⁻¹) in T₄ (STCR integrated (65 q ha⁻¹) - Predicted STV) where the initial available potassium was very high (456.03 kg ha⁻¹). The maximum uptake of potassium (175.70 kg ha⁻¹) was recorded in STCR target of 65 q ha⁻¹ through integrated approach using predicted soil test values (T₂), whereas, lower uptake of K₂O was recorded in absolute control (43.63 kg ha⁻¹) where no fertilizers or poultry manure was applied. The higher actual balance (297.52 kg ha⁻¹) was recorded in STCR integrated approach for the targeted yield of 65 q ha⁻¹ through predicted soil test values (T₄). Interestingly, all the treatments of various approaches of fertilizer recommendations recorded net positive balance of potassium. The higher actual balance of potassium in these treatments might be due to incorporation of poultry manure or any organic sources along with fertilizer nitrogen (synergistic effect of N on K) which increased the cumulative non exchangeable K release and maintained greater amount of

potassium in solution and on exchange sites, by re-establishing the equilibrium among the different fractions of potassium [19], thereby enhanced the available K_2O in the soil

UNDER PEER REVIEW

Table 3: Influence of different approaches of nutrient application on yield and physico - chemical properties of post harvest soil of aerobic rice

Treatment details		Grain yield	Soil pH (1:2.5)		Electrical Conductivity (dS m ⁻¹)		Organic Carbon (%)	
		q ha ⁻¹	Initial	After harvest	Initial	After harvest	Initial	After harvest
T ₁	STCR inorganics (65 q ha ⁻¹) - Actual STV	65.50	5.75	5.76	0.070	0.087	0.42	0.45
T ₂	STCR inorganics (65 q ha ⁻¹) - Predicted STV	68.85	5.45	5.78	0.088	0.090	0.46	0.48
T ₃	STCR integrated (65 q ha ⁻¹) - Actual STV	61.70	5.93	5.81	0.089	0.099	0.46	0.50
T ₄	STCR integrated (65 q ha ⁻¹) - Predicted STV	63.79	5.85	5.91	0.090	0.100	0.49	0.52
T ₅	STCR inorganics (55 q ha ⁻¹) - Actual STV	61.58	5.54	5.55	0.087	0.102	0.38	0.44
T ₆	STCR inorganics (55 q ha ⁻¹) - Predicted STV	62.96	5.45	5.71	0.088	0.104	0.45	0.45
T ₇	STCR integrated (55 q ha ⁻¹) - Actual STV	57.55	5.84	5.86	0.087	0.101	0.36	0.50
T ₈	STCR integrated (55 q ha ⁻¹) - Predicted STV	60.14	5.81	5.98	0.088	0.103	0.43	0.51
T ₉	Package of practice	53.25	5.76	5.88	0.085	0.096	0.48	0.51
T ₁₀	LMH (STL) - Actual STV	48.76	5.75	5.83	0.091	0.095	0.50	0.51
T ₁₁	LMH (STL) - Predicted STV	49.15	5.82	5.84	0.093	0.100	0.50	0.52
T ₁₂	Absolute control	20.66	5.98	6.02	0.071	0.081	0.45	0.49
S.Em. ±		2.88	-	0.09	-	0.006	-	0.03
C.D. @ 5 %		8.39	-	NS	-	0.018	-	NS

NS: Non significant

Table 4: Influence of difference approaches of fertilizer recommendations on available major nutrients status of post harvest soil of aerobic rice

UNDER PEER REVIEW

Treatment details		Available Nitrogen		Available Phosphorus		Available potassium	
		(kg ha ⁻¹)					
		Initial	After harvest	Initial	After harvest	Initial	After harvest
T ₁	STCR inorganics (65 q ha ⁻¹) - Actual STV	260.59	261.33	101.05	100.81	271.64	217.36
T ₂	STCR inorganics (65 q ha ⁻¹) - Predicted STV	257.23	264.69	115.46	109.34	282.56	229.08
T ₃	STCR integrated (65 q ha ⁻¹) - Actual STV	261.67	267.09	106.92	99.18	305.92	291.20
T ₄	STCR integrated (65 q ha ⁻¹) - Predicted STV	260.32	280.37	113.47	104.30	316.49	297.52
T ₅	STCR inorganics (55 q ha ⁻¹) - Actual STV	260.21	265.07	99.88	103.24	221.67	216.60
T ₆	STCR inorganics (55 q ha ⁻¹) - Predicted STV	261.33	266.20	102.87	105.43	229.44	222.40
T ₇	STCR integrated (55 q ha ⁻¹) - Actual STV	262.08	266.93	93.65	101.47	245.75	221.88
T ₈	STCR integrated (55 q ha ⁻¹) - Predicted STV	260.96	277.39	100.36	103.32	261.11	249.44
T ₉	Package of practice	268.05	268.61	115.60	102.21	286.77	274.30
T ₁₀	LMH (STL) - Actual STV	266.56	268.95	98.55	88.61	285.69	283.72
T ₁₁	LMH (STL) - Predicted STV	269.17	272.16	111.52	93.74	289.36	289.04

T₁₂	Absolute control	243.41	239.12	56.96	57.51	168.13	166.20
S.Em. ±		-	9.59	-	3.53	-	18.14
C.D. @ 5 %		-	28.12	-	10.35	-	53.19

Treatments		S		Ca		Mg	
		(mg kg ⁻¹)		(cmol (p ⁺) kg ⁻¹)			
		Initial	After harvest	Initial	After harvest	Initial	After harvest
T ₁	STCR inorganics (65 q ha ⁻¹) - Actual STV	34.96	27.75	3.12	2.72	1.22	1.12
T ₂	STCR inorganics (65 q ha ⁻¹) - Predicted STV	36.90	30.59	3.05	2.93	1.23	1.10
T ₃	STCR integrated (65 q ha ⁻¹) - Actual STV	36.67	29.84	3.83	2.93	1.22	1.05
T ₄	STCR integrated (65 q ha ⁻¹) - Predicted STV	42.90	33.74	3.13	3.10	1.35	1.18
T ₅	STCR inorganics (55 q ha ⁻¹) - Actual STV	42.73	32.60	3.15	2.62	1.12	1.01
T ₆	STCR inorganics (55 q ha ⁻¹) - Predicted STV	42.88	32.68	3.03	3.03	1.18	1.02
T ₇	STCR integrated (55 q ha ⁻¹) - Actual STV	38.49	32.80	3.33	2.92	1.25	1.10
T ₈	STCR integrated (55 q ha ⁻¹) - Predicted STV	42.94	35.00	3.37	3.33	1.35	1.17
T ₉	Package of practice	42.16	34.40	3.25	2.19	1.17	1.06
T ₁₀	LMH (STL) - Actual STV	43.80	31.68	3.48	2.40	1.15	1.07
T ₁₁	LMH (STL) - Predicted STV	44.55	31.89	3.47	2.45	1.15	1.09
T ₁₂	Absolute control	37.95	27.19	3.10	2.29	1.34	1.00
S.Em. ±			2.10		0.22		0.12
C.D. @ 5 %			6.16		0.64		NS

Table 5: Influence of difference approaches of fertilizer recommendations on secondary nutrients status of post harvest soil of aerobic rice

NS: Non significant

Table 6: Nitrogen balance in soil as influenced by different approaches of nutrient application

Treatment details	IAN	FN	TN	CU	EB	AB	G/L
-------------------	-----	----	----	----	----	----	-----

		1	2	3(1+2)	4	5(3-4)	6	7(6-5)
T₁	STCR inorganics (65 q ha ⁻¹) - Actual STV	260.59	143.94	404.53	161.94	242.59	261.33	18.74
T₂	STCR inorganics (65 q ha ⁻¹) - Predicted STV	207.10	154.80	361.90	176.73	185.17	264.69	79.53
T₃	STCR integrated (65 q ha ⁻¹) - Actual STV	261.67	127.56	389.23	151.50	237.73	267.09	29.36
T₄	STCR integrated (65 q ha ⁻¹) - Predicted STV	205.62	138.95	344.57	159.71	184.86	280.37	95.51
T₅	STCR inorganics (55 q ha ⁻¹) - Actual STV	260.21	113.72	373.93	150.90	223.03	265.07	42.04
T₆	STCR inorganics (55 q ha ⁻¹) - Predicted STV	202.89	125.37	328.26	162.21	166.06	266.20	100.14
T₇	STCR integrated (55 q ha ⁻¹) - Actual STV	262.08	98.55	360.63	143.73	216.90	266.93	50.04
T₈	STCR integrated (55 q ha ⁻¹) - Predicted STV	201.60	110.84	312.44	146.17	166.26	277.39	111.12
T₉	Package of practice	272.53	100.00	372.53	126.61	245.92	268.61	22.69
T₁₀	LMH (STL) - Actual STV	266.56	125.00	391.56	110.37	281.19	268.95	-12.25
T₁₁	LMH (STL) - Predicted STV	204.08	125.00	329.08	114.19	214.89	272.16	57.27
T₁₂	Absolute control	243.41	0.00	243.41	47.85	195.56	239.12	43.56

Legend:

IAN = Initial available nitrogen (kg ha^{-1})	TN = Total nitrogen (kg ha^{-1})	FN = Fertilizer nitrogen (kg ha^{-1})
CU = Crop uptake (kg N ha^{-1})	EB = Expected balance (kg ha^{-1})	
AB = Actual balance (kg ha^{-1})	G/L = Net gain/ net loss (kg ha^{-1})	

Table 7: Phosphorus balance in soil as influenced by different approaches of nutrient application

Treatments		IAP	FP	TP	CU	EB	AB	G/L
		1	2	3(1+2)	4	5(3-4)	6	7(6-5)
T ₁	STCR inorganics (65 q ha ⁻¹) - Actual STV	101.05	73.54	174.59	55.81	118.78	100.81	-17.97
T ₂	STCR inorganics (65 q ha ⁻¹) - Predicted STV	174.48	68.13	242.61	59.43	183.18	109.34	-73.84
T ₃	STCR integrated (65 q ha ⁻¹) - Actual STV	106.92	61.25	168.17	53.85	114.33	99.18	-15.15
T ₄	STCR integrated (65 q ha ⁻¹) - Predicted STV	167.74	57.08	224.82	55.96	168.86	104.30	-64.55
T ₅	STCR inorganics (55 q ha ⁻¹) - Actual STV	99.88	61.16	161.04	53.99	107.05	103.24	-3.81
T ₆	STCR inorganics (55 q ha ⁻¹) - Predicted STV	149.36	56.85	206.21	55.98	150.22	105.43	-44.79
T ₇	STCR integrated (55 q ha ⁻¹) - Actual STV	93.65	50.85	144.50	50.32	94.18	101.47	7.29
T ₈	STCR integrated (55 q ha ⁻¹) - Predicted STV	145.27	47.26	192.53	51.46	141.07	103.32	-37.75
T ₉	Package of practice	115.60	50.00	165.60	42.43	123.17	102.21	-20.96
T ₁₀	LMH (STL) - Actual STV	98.55	37.50	136.05	38.67	97.38	88.61	-8.77
T ₁₁	LMH (STL) - Predicted STV	165.92	37.50	203.42	40.67	162.75	93.74	-69.01
T ₁₂	Absolute control	56.96	0.00	56.96	17.48	39.48	57.51	18.03

Legend: IAP = Initial available phosphorus (kg ha⁻¹), TP = Total phosphorus (kg ha⁻¹) FP = Fertilizer phosphorus (kg P₂O₅ ha⁻¹)
CU = Crop uptake (kg P₂O₅ ha⁻¹), EB = Expected balance (kg ha⁻¹)
AB = Actual balance (kg ha⁻¹) G/L = Net gain/ net loss (kg ha⁻¹)

Table 8: Potassium balance in soil as influenced by different approaches of nutrient application

Treatments		IAK	FK	TK	CU	EB	AB	G/L
		1	2	3(1+2)	4	5(3-4)	6	7(6-5)
T ₁	STCR inorganics (65 q ha ⁻¹) - Actual STV	271.64	36.82	308.46	173.12	135.34	217.36	82.02
T ₂	STCR inorganics (65 q ha ⁻¹) - Predicted STV	298.07	30.84	328.91	175.70	153.21	229.08	75.87
T ₃	STCR integrated (65 q ha ⁻¹) - Actual STV	305.92	28.96	334.88	165.66	169.22	291.20	121.98
T ₄	STCR integrated (65 q ha ⁻¹) - Predicted STV	456.03	2.12	458.15	170.85	287.30	297.52	10.22
T ₅	STCR inorganics (55 q ha ⁻¹) - Actual STV	221.67	33.01	254.68	150.08	104.60	216.60	112.00
T ₆	STCR inorganics (55 q ha ⁻¹) - Predicted STV	276.69	20.56	297.25	153.03	144.22	222.40	78.18
T ₇	STCR integrated (55 q ha ⁻¹) - Actual STV	245.75	26.62	272.37	141.91	130.45	221.88	91.43
T ₈	STCR integrated (55 q ha ⁻¹) - Predicted STV	340.47	6.62	347.09	141.18	205.91	249.44	43.53
T ₉	Package of practice	286.76	50.00	336.76	144.71	192.05	274.30	82.25
T ₁₀	LMH (STL) - Actual STV	285.69	50.00	335.69	120.77	214.93	283.72	68.79
T ₁₁	LMH (STL) - Predicted STV	322.27	45.83	368.10	130.63	237.47	289.04	51.57
T ₁₂	Absolute control	168.13	0.00	168.13	43.63	124.50	166.20	41.70

Legend: IAK = Initial available potassium (kg ha⁻¹), TK = Total potassium (kg ha⁻¹) FK = Fertilizer potassium (kg K₂O ha⁻¹)
CU = Crop uptake (kg K₂O ha⁻¹), EB = Expected balance (kg ha⁻¹)
AB = Actual balance (kg ha⁻¹) G/L = Net gain/ net loss (kg ha⁻¹)

Conclusion

The higher post-harvest soil available nutrient status was registered in STCR integrated approach based on predicted soil test values compared to package of practice and LMH approach. No significant difference was observed between the actual and predicted soil test value based fertilizer recommendation with respect to soil fertility status. Similarly, no significant difference in available nutrient status was noticed between inorganic and integrated approach.

References

1. Anonymous, <https://www.indiaagristat.com> (2019).
2. Shashidhar, H. E., Rice root system under aerobic condition. *Euphytica*, **129**:290-294 (2007).
3. Patil, P.A.S., NAnjappa, H.V., Ramachandrappa. B.K. and Basavaraja, P. K., Quality of aerobic rice as influenced by site specific nutrient management approach. *J. Pharmacogno. Phytochem.*, **9**(3): 1529-1532 (2020).
4. Satish, A., Govind G.V., Chandrappa, H., Nagaraja, K., Long term effect of integrated use of organic and inorganic fertilizers on productivity, soil fertility and uptake of nutrients in rice and maize cropping system. *Int. J. Sci. Nat.*, **2**:84–88 (2011).
5. Ray, P.K., Jana, A.K., Maitra, D.N., Saha, M.N., Chaudhury, J., Saha, A.R., Fertilizer prescription on soil test basis for jute, rice and wheat in *Typic ustochrept*. *J. Indian. Soc. Soil Sci.*, **48**: 79–84 (2000).
6. Bhavya, N., Development of targeted yield equation for aerobic rice and its evaluation on *Alfisols* of Eastern dry zone of Karnataka *Ph.D. Thesis*, (Unpub.) University of Agricultural Sciences, Bangalore. (2021).
7. Jackson, M.L., Soil Chemical Analysis. Prentice Hall of India Pvt. Ltd., New Delhi. (1973).

- 8.Kumar, P. and Parmanand, Evaluation of soil test crop response approach for sustainable production of rice in Balodabazar –Bhatapara District of Chhattisgarh. *Int. J. Curr. Microbiol. App. Sci.*, **7**: 3513-3518. (2018).
- 9.Sarwar, G., Schmeisky, H., Tahir M.A., Iftikhar, Y. and Sabah, N.U., Application of green compost for improving soil chemical properties and fertility status. *J. Animal and Plant Sci.*, **20**: 258-260 (2010).
- 10.Ojo, A.O., Adetunji, M.T., Okeleye, K.A. and Adejuyigbe, C.O., Soil fertility, phosphorus fractions, and maize yield as affected by poultry manure and single superphosphate, *Inter. Scholarly Res. Notices.*, **71**: 325-333 (2015).
- 11.Manish, S., Singh, Y.V., Singh, S.K., Dey, P. and Ram, R.L., Validation of soil test and yield target based fertilizer prescription model for rice on *Inceptisols* of Eastern Zone of Uttar Pradesh, India. *Inter. J. Curr. Microbio. App. Sci.*, **6**(2): 406-415 (2017).
- 12.Mahmood, F., Mohammad, S., Muhammad, A., Sami, U., Imran Khan, Umair, A., Tanvir, S., Sabir, H., Muhammad, S., Muhammad, A., Sami, U., Effects of organic and inorganic manures on maize and their residual impact on soil physico-chemical properties, *J. Soil Sci. Plant Nutri.*, **17**(1): 22-32 (2017).
- 13.Rajput, P.S., Srivastava, S., Sharma, B.L., Sachidanand, B., Dey, P., Aher, S.B. and Yashona, D.S., Effect of soil-test-based long-term fertilization on soil health and performance of rice crop in *Vertisols* of Central India. *Int. J. Agric. Environ. Biotech.*, **9**(5): 801-806 (2016).
- 14.Ashwini, Y., Evaluation of STCR targeted yield approach on ragi crop yield, soil properties, nutrient uptake and nutrient use efficiency. *M.Sc (Agri.) Thesis*, University of Agricultural Sciences, Bangalore (2007).
- 15.Chandrakanth, STCR approach for soil and foliar application of soluble fertilizers and their effect on soil properties, growth and yield of maize. *M.Sc (Agri.) Thesis*, Univ. Agric. Sci., Bangalore (2015).

- 16.Santhosha, V. P., Yield maximization in maize through different forms of fertilizers and approaches of nutrient recommendations. *M.Sc (Agri.) Thesis*, University of Agricultural Sciences, Bangalore (2013).
- 17.Brar, S.P.S. and Singh. B., Effect of long term application of N, P, K and zinc on crop and soil in maize-wheat rotation. *J. Res., Punjab Agril. Uni.*, **26**(4): 572-580 (1984).
- 18.Tomar, N.K., Dynamics of phosphorus in soils. *J. Indian Soc. Soil Sci.*, **48**: 640-672 (2000).
- 19.Sundaresh. R., Development of STCR targeted yield equation for cabbage (*Brassica oleraceae* var. *capitata*) under fertigation with soluble fertilizers and its evaluation. *Ph.D. Thesis*, Univ. Agric. Sci., Bangalore. (2019).