

Zinc Fertilization: Effects on Nutrients Availability and Productivity of Rice (*Oryza sativa* L.)

ABSTRACT

The two years (2020-21 & 2021-22) of on farm trial established at farmers field of district Saran, Bihar to evaluate the effect of zinc fertilization on plant available nutrients and crop productivity. The trial performed under randomised block design involved seven replication of three different treatments viz., T₁: NPK- 130:40:20 kg ha⁻¹ (Farmer's Practice), T₂: RDF (NPK-120:60:40 kg ha⁻¹ + Zn @ 5.0 kg ha⁻¹ and T₃: RDF (NPK-120:60:40 kg ha⁻¹) + foliar spray of 0.5 % ZnSO₄ at 25 DAT. The pooled results of two years trial revealed that basal application of RDF-NPK and Zn (T₂) significantly improved the soil organic carbon by 16.22%, plant available N by 11.96 %, plant available P by 15.32%, plant available K by 10.99% and plant available Zn by 24.00 % as compared to farmers practice. The crop productivity was also improved by 24.22% and 14.43% in treatment having basal application of Zn (T₂) and foliar application of Zn (T₃), respectively over farmers practice (T₁). The two years of investigation can be concluded that in order to maintain the soil nutrients and better crop productivity, the basal application of RDF-NPK and Zn was effective in semi arid region of Bihar.

Key words: Zinc, nutrient availability, productivity and rice

1. INTRODUCTION

Rice (*Oryza sativa* L.) is a staple food for more than 50% of the world's population including regions of high population density and rapid growth. The global population is 7.55 billion presently and is expected to reach 9.10 billion by the year 2050 [1]. Human consumption accounts for 85 percent of total production of rice compared with 72 percent of wheat [2]. At present, rice production alone consumes nearly 24.7 MT. of fertilizer (N+P₂O₅+K₂O) which accounts for approximately 14.0 % of the total global fertilizer consumption in a year and in India it accounts for 31.8% of the total fertilizer consumption [3].

Crops grown in arid or semi-arid regions are mostly exposed to low soil fertility and exhibit multiple nutrient deficiencies due to low soil organic matter and alkaline calcareous nature of soil that limit the crop production [4]. The nutrient deficiency can be corrected by applying appropriate micronutrient containing

fertilizers. The nutrients can be applied to the crop plants by a variety of ways like soil application and foliar spray. Every method of application has its advantages and disadvantages [5], depending upon the soil and climatic conditions of the area.

Micronutrients are essential for increasing crop production and enhancing animal and human health. In India, zinc is considered as the fourth important yield limiting nutrient after nitrogen, phosphorus and potassium [6]. The critical limit of available zinc in the soil, suitable for rice growth is 0.6 mg kg^{-1} . Soil application of zinc increased the grain yield [7], and plant available nutrients [8].

Zinc deficiency is a well known nutritional and health problem in human populations where rice is the dominating staple food crop [9]. Among nutrient deficiencies, Zn deficiency has been identified as a most serious agricultural issue in the world. Forty three percent of Indian soils [10] are identified deficient in zinc. The main reason of deficiency of plant available zinc in soil is the precipitation or adsorption of zinc with various soil components, depending on the soil pH, organic matter, pedogenic oxides and redox potential. Soil zinc found in soil solution as the free ion, Zn^{2+} associated with organic and inorganic ligands on exchanged sites of soil, bound by organic matter and occluded in oxides and hydroxides of Al and Fe [11]. Soil and foliar applications of zinc may increase grain zinc concentration in rice, soil zinc application has been reported to increase grain yield whereas foliar application of zinc increased grain concentration of zinc [12]. Hence, this study was aimed to investigate the effects of zinc fertilization on plant available nutrients and crop productivity of rice.

2. MATERIALS AND METHODS

2.1. Study area

An on-farm trial was established during 2020-21 & 2021-22 at farmer's fields of district Saran, Bihar, under the supervision of Krishi Vigyan Kendra, Manjhi, Saran, Dr. Rajendra Prasad Central Agricultural University, Pusa, Samastipur, Bihar. The area falls in the subtropical, humid agro-climatic zone of Bihar. The average annual rainfall of the area is about 800-1100 mm and mean monthly maximum & minimum temperature varies from $24-33^{\circ}\text{C}$ and $16-23^{\circ}\text{C}$, respectively. The soil of the experimental site was sandy loam in texture. The initial properties of experimental soil were given in table 1.

Table 1. Initial fertility status of soil during two years (2020-21 & 2021-22) of study.

Soil parameters	Years	Average value
-----------------	-------	---------------

	2020-21	2021-22	
pH (1:2.5)	8.44	8.42	8.43
EC (dS/m)	0.32	0.33	0.33
SOC (%)	0.37	0.36	0.37
Available N (kg/ha)	214.20	213.00	213.60
Available P(kg/ha)	11.50	12.40	11.95
Available K (kg/ha)	129.00	130.10	129.55
Available Zn (mg/kg)	0.48	0.47	0.48

2.2. Experimental Design

An experiment on Zn fertilization was established at seven farmer fields of Saran district of Bihar under supervision of Krishi Vigyan Kendra, Manjhi, Saran (Dr. Rajendra Prasad Central Agricultural University, Pusa Samastipur) during *kharif*, 2020-21 & 2021-22 (Two season). The experiment was laid out in Randomized Block Design and replicated seven times involved three Zn fertilization treatments *i.e.*, T₁: NPK- 130:40:20 kg ha⁻¹ (Farmer's Practice), T₂: RDF (NPK-120:60:40 kg ha⁻¹ + Zn @ 5.0 kg ha⁻¹ and T₃: RDF (NPK-120:60:40 kg ha⁻¹) + foliar spray of 0.5 % ZnSO₄ at 25 DAT. A total of twenty one plots were established with each plot sized of 180.0 m × 22.2 m.

2.3. Nursery raising

The seed was treated with fungicide SAAF (Carbendazim + Mencozeb) @ 3g/kg seed before sowing to protect the crops from seed borne diseases. Seed of rice variety Sahbhagi was raised in nursery by "Wet bed method". Seed beds of 8 × 1.25 m size were prepared in dry condition. In addition 1 kg of nitrogen, 1 kg of phosphorus and 0.5 kg of potash were also applied @ 1000 sq. m through Urea, DAP and MOP, respectively at the time of last ploughing. Further, top dressing was done with @ 1.0 kg N/1000 sq. m in the form of urea at 10 days after sowing. Need based irrigation and weeding was also done.

2.4. Field preparation

The experimental field was ploughed immediately after the harvest of previous wheat crop by a tractor drawn harrow in summer to expose weeds and the eggs of harmful insects. The field was prepared by following two cross disc harrowing and two cross tiller operations and finally the field was levelled by planking. Thereafter, the field was flooded with water and puddled by tractor. After puddling field was levelled finally.

2.5. Nutrients application

Irrespective of treatments, the recommended doses of nitrogen (N), phosphorus (P_2O_5) and potassium (K_2O) were used 120 kg, 60 and 40 kg per hectare, respectively. N, P, K were applied in the form of urea, di-ammonium phosphate and muriate of potash. Whole dose of P and K fertilizers was applied at the time of field preparation. The nitrogen fertilizer was applied in four equal splits *i.e.*, 1/4 at basal dressing, 1/4 at mid-tillering, 1/4 at active tillering and 1/4 at panicle initiation in rice cropping. Zinc fertilizers were applied in the form of zinc sulphate @ 5.0 kg ha⁻¹ as basal dose. The foliar application of Zn was applied @ 0.5% at 25 days after transplanting of paddy.

2.6. Data collections

2.6.1. Soil properties

Soil analysis was done before the commencement of experiment and after harvesting of crop by standard procedure. Firstly, three representative soil samples from 0-15 cm depth were collected for analysis of physical and chemical properties. The pH of soil was measured with the help of a pH meter, maintaining the soil-water ratio of 1:2.5 as described by Jackson [13]. The electrical conductivity (dS m⁻¹) in the clear extract of soil-water ratio of 1:2.5 was determined with the help of conductivity meter [13]. The organic carbon content in soil samples was estimated by Walkely and Black [14] method as suggested by Jackson [13]. Available N of soil was measured by Modified macro-Kjeldahl method [15]. Available P was determined by the ascorbic acid procedure using a blue filter (660 mμ) as suggested by Olsen et al. [16]. Available potassium were analysed after extraction of soil with neutral normal 1N NH_4OAc (pH 7.0) as suggested by Tondon [17]. Available zinc (Zn) was estimated by Diethylene Triamine Penta Acetic Acid solution (0.005 M DTPA + 0.1 M TEA + 0.01 M $CaCl_2$; pH 7.3) as suggested by Lindsay and Norvell [18].

2.6.2. Crop productivity

The crops were harvested manually from 1 m² randomly selected 3 quadrates from each plot and recorded the crop productivity in terms of quintal per hectare.

2.7. Data Analysis

The data generated from the present investigation were subjected to statistical analysis using the statistical package SPSS 13.0. The least significant difference (LSD) at 5% for testing the significant difference among the treatment means [20].

3. RESULTS AND DISCUSSION

3.1. Soil pH, EC and organic carbon

A significant effect of zinc fertilization on soil pH, EC and soil organic carbon was observed in two years of investigation of rice (Table 2). The pooled value of pH revealed that the basal application of Zn fertilization produced significant effect whereas, the foliar application of Zn and farmers practice produced almost similar effect on soil pH. The treatment T₂ (basal application of RDF- NPK and Zn) and T₃ (basal application of RDF-NPK + foliar application of Zn @ 0.5% at 25 DAT) lowered the pH values by 0.97 & 0.99 times as compared to farmers practice (T₁). When compared with initial value (average of two years), the pH value was lowered by 0.12 times in both the treatment T₂ and T₃. Basal application of Zn fertilization (T₂) lowered the pH value by 0.98 times as compared with foliar application of Zn fertilization (T₃). In contrast, the EC value was raised by 1.20 & 1.17 times in treatment T₂ & T₃, respectively as compared to T₁ but statistically similar to each other. When compared with initial value, the EC value was raised by 1.27 & 1.24 times in treatment T₂ & T₃, respectively.

Soil organic carbon (SOC) was significantly affected by zinc fertilization in two years of study (Table 2). The basal application of zinc (T₂) maintained the highest per cent of increment i.e., 16.22 over farmers practice which was statistically at par with foliar application of zinc (T₃). In comparison with initial value, the similar per cent increment in SOC was recorded in treatment T₂ and T₃. When comparisons with basal and foliar application zinc, the basal application of zinc was improved the SOC by 4.88% over foliar application of zinc. Comparatively the higher production of carbonic acid in zinc treated plots after decomposition of root biomass that hastens the soil organic carbon and lowering the soil pH.

Table 2. Zn fertilization: effect on availability of nutrients in soil after harvesting of rice.

Treatments	Years		Pooled average
	2020-21	2021-22	
pH (1:2.5)			
T ₁	8.39b	8.40b	8.40b
T ₂	8.10a	8.26a	8.18a
T ₃	8.28b	8.37b	8.33b
EC (dS/m)			
T ₁	0.35b	0.35b	0.35b
T ₂	0.41a	0.42a	0.42a
T ₃	0.40a	0.41a	0.41a
SOC (%)			
T ₁	0.38b	0.37b	0.37b
T ₂	0.42a	0.43a	0.43a

T ₃	0.40ab	0.41ab	0.41ab
Available N (kg/ha)			
T ₁	216.70b	215.50b	216.10b
T ₂	241.60a	242.30a	241.95a
T ₃	231.50a	233.10a	232.30a
Available P (kg/ha)			
T ₁	11.90b	12.90b	12.40b
T ₂	14.30a	14.30a	14.30a
T ₃	14.20a	14.10a	14.15a
Available K (kg/ha)			
T ₁	136.20b	135.80b	136.00b
T ₂	151.50a	150.40a	150.95a
T ₃	147.80a	147.90a	147.85a
Available Zn (mg/kg)			
T ₁	0.50b	0.49b	0.50b
T ₂	0.62a	0.61a	0.62a
T ₃	0.53b	0.53b	0.53b

* Within variable means in the same column followed by different letters are significantly different from each other

3.2. Plant available nutrients (N, P, K and Zn)

Availability of primary soil nutrients was also significantly influenced by different methods of zinc fertilization in rice (Table 2), the maximum availability of nitrogen (N), phosphorus (P) and potassium was estimated with treatment having basal application of zinc fertilization (T₂) which was statistically at par with foliar application of zinc (T₃). An improvement in available N, P and K was recorded by 11.96, 15.32 and 10.99% in treatment having basal application of Zn (T₂) over farmers practice (T₁). When compared with initial value of plant available N, P and K the highest increment in availability of N, P and K was estimated in treatment T₂ by 13.27, 19.67 and 16.52%. Among the zinc fertilization, the basal application of zinc was considered superior treatment in respect of plant available nutrients. These results were confirmed with the findings of Ghoneim [8] and Haldar et al. [20] for available N, Ghoneim [8] and Lonergan et al. [21] for available P, and Ghoneim [8] and Wu et al. [22] for available potassium.

The basal application of zinc also maintained the highest availability of zinc as compared to rest of the treatments, and the improvement was recorded by 16.04 and 24.24% as compared to foliar application of zinc and farmer practice, respectively. As compared with initial value, an increment was recorded by 28.13 & 1.42% in treatment T₂ & T₃, respectively. A positive correlation (Figure 1) was observed between SOC and available Zn due to that reason the available Zn was improved under treatment having basal

application Zn. The availability of zinc increased significantly with increase in soil organic carbon because zinc forms soluble complexes with soil organic matter. Sharma et al. [23] reported that the positive correlations of soil zinc and organic carbon. Sidhu and Sharma [24] also reported that the available micronutrients including Zn increased with increase in organic carbon.




Figure 1. Polynomial correlation between SOC and available Zn.

3.3. Crop productivity

The zinc fertilization well reflected to satisfactory crop productivity (Figure 2) of rice. The treatment having basal application of zinc (T_2) recorded the highest productivity of rice which was statistically followed by foliar application (T_3) and farmers practices (T_1). An improvement in rice yield with the basal application of Zn and RDF-NPK by 7.74% & 24.22% over treatment T_3 & T_1 , respectively. Similar results were reported by Shamim et al. [23] who reported greater grain yield with soil application of Zn than foliar application. Kumar et al. [24], and Savithri et al. [19] also reported somewhat similar results.




Figure 2. Zn fertilization: effects on productivity of rice. Vertical bars indicate $\pm$ S.E. of mean of the observed values.

4. CONCLUSION

On the basis of two years on farm trial data, we conclude that the basal application of zinc along with RDF-NPK improved the soil organic carbon and plant available nutrients like nitrogen, phosphorus, potassium and zinc and also produced satisfactory crop productivity. Therefore, in order to maintain the soil nutrients and better crop productivity, the basal application of RDF-NPK and Zn was effective in semi arid region of Bihar.

REFERENCES

1. FAO. Global Agriculture Towards 2050: FAO, Rome; 2009.
2. Prakash P, Hemalatha M, Joseph M. Methods of Zinc Application with Green Leaf Manuring. *Advances in Crop Science and Technology*. 2018; 6:3.
3. Nayak AK, Khanam R, Mohanty S, Chatterjee D, Bhaduri D. Changing dimension of nutrient management in rice. 3rd ARRW International Symposium, Feb. 6-9, Cuttack, India. 2018; pp: 40-44.
4. Rafique E, Rashid A, Rayan J, Bhati AU. Zinc deficiency in rainfed wheat in Pakistan: magnitude spatial variability, management and plant analysis diagnostic norms. *Communications in Soil Science and Plant Analysis*. 2006; 37:181-197.
5. Rehim A, Farooq M, Ahmad F, Hussain M. Band placement of phosphorus improves the phosphorus use efficiency and wheat productivity under different irrigation regimes. *International Journal of Agriculture and Biology*. 2012; 14: 727-733.

6. Singh V, Singh S. Organic rice production technology. Research Trends in Agriculture. AkiNik Publications, New Delhi. 2019;16: 99-129.
7. Singh K, Verma G, Manchanda JS. Soil and foliar zinc application for enhancing grain zinc content of aromatic rice genotypes grown in zinc deficient and sufficient soil. Journal of Soil and Water Conservation. 2020; 19(2): 0-0.
8. Ghoneim A M. Effect of different methods of Zn application on rice growth, yield and nutrients dynamics in plant and soil. Journal of Agriculture and Ecology Research International. 2016; 6(2): 1-9.
9. Stein AJ. Global impacts of human mineral malnutrition. Plant Soil. 2010;335: 133–154.
10. Shukla, A.K., Tiwari, P.K. and Chandra, Prakash. Micronutrient deficiencies vis a vis food and nutritional security of India. Indian Journal of fertilizers. 2014; 10(12): 94-112.
11. Havlin JL, Beaton JD, Tisdale SL, Nelson WL. Soil fertility and fertilizer – an introduction to soil nutrient management. Pearson Education Inc. 2005.
12. Wissuwa M, Ismail A, Graham R. Rice grain zinc concentrations as affected by genotype, native soil-zinc availability, and zinc fertilization. Plant and Soil. 2008; 306:37-48.
13. Jackson M L. Soil Chemical Analysis. Printice Hall, India Pvt. Ltd., New Delhi; 1973.
14. Walkely AJ, Black CA. Estimation of soil organic carbon by the chromic acid titration method. Soil Science. 1934;37: 259-260.
15. Bremner JM, Mulvaney CS. Nitrogen-total. In Page *et al.* (Eds). Method of soil analysis, part 2, chemical and microbiological properties. Agronomy monograph No, 9. American Society of Agronomy: Inc. Madison, Wisconsin, USA;1982.
16. Olsen SR, Culs CV, Wortanade FS, Deam LA. Estimation of available phosphorus by extraction with sodium bicarbonate. United States Department of Agriculture. 1954; 939: 19-23.
17. Tandon H L S. Methods of analysis of soils, plants, waters, fertilizers and organic manures (2nd Eds): Fertilizer Development and Consultation Organization, New Delhi, India; 2005.
18. Lindsay WL, Norvell WA. Development of a DTPA soil test for zinc, iron, manganese and copper. Soil Science Society of American Journal. 1978; 42: 421-428.
19. Gomez K, Gomez A. Statistical procedures for agricultural research: John Wiley & Sons New York, USA;1984.
20. Haldar M, Mandal LN. Influence of soil moisture regimes and organic matter application on the extractable Zn and Cu content in rice soils. Plant Soil. 1979; 53: 203-213.

21. Lonergan PF, Pallotta MA, Lorimer M. Multiple genetic loci for zinc uptake and distribution in barley (*Hordeum vulgare*). *New Phytol.* 2009;184(1):168-179.
22. Wu CY, Feng Y, Shohag MJ. Characterization of Zn uptake, translocation, and accumulation into developing grains and young leaves of high Zn-density rice genotype. *J. Zhejiang Univ.-Sci. B (Biomed. & Biotechnol)*. 2011;12(5):408-418.
23. Sharma RP, Singh M, Sharma JP. Correlation studies on micronutrients vis-à-vis soil properties in some soils of Nagaur district in semi-arid region of Rajasthan. *Journal of Indian Society of Soil Science*. 2003; 51(4): 522-527.
24. Sidhu GS, Sharma BD. Diethylenetriaminepenta acetic Acid-Extractable Micronutrients Status in Soil under a Rice-Wheat System and Their Relationship with Soil Properties in Different Agro-climatic Zones of Indo-Gangetic Plains of India. *Communications in Soil Science and Plant Analysis*. 2010; 41(1): 29-51.