

Influence of 2-(3- Chlorophenoxy) propionic acid on Crown size, Yield and Quality of Pineapple cv. Giant Kew

ABSTRACT

Aims: The present study was carried out to determine the effect of 2-(3- chlorophenoxy) propionic acid (3- CPA) on crown size, yield and quality of pineapple cv. Giant Kew.

Study design: Completely Randomised Design

Place and Duration of Study: Department of Horticulture, School of Agriculture Sciences and Rural Development, Nagaland University, Medziphema campus.

Methodology: Five treatments consisted of different concentrations of the growth regulator viz., C₁ (Control), C₂ (50 ppm), C₃ (100 ppm), C₄ (150 ppm) and C₅ (200 ppm) were evaluated with four replications each. Treatment was applied as foliar spray on pineapple crowns when apical flowers were in final stage of dehiscence. Growth and yield attributes viz., length and circumference of crown, crown to fruit length ratio, length of fruit with and without crown, pulp weight, peel weight, pulp to peel ratio, core weight, days taken to mature and dry matter content were recorded. Furthermore, biochemical parameters viz., TSS, titratable acidity, TSS: acid ratio, ascorbic acid, total sugar, reducing sugar and non- reducing sugar were analyzed, along with shelf life and Physiological loss in weight (PLW).

Results: The experiment revealed C₃ (100 ppm) as optimum concentration for reduction of crown size with minimum crown length (14.28 cm), circumference (49.72 cm) and crown to fruit ratio (0.83) and showed 48% reduction in crown length over control. Also, maximum values for length of fruit (17.22 cm), pulp weight (1080 g), peel weight (403 g), core weight (200 g), shelf life (19.33 days) and minimum PLW were observed in C₃ (100 ppm). It was found that higher concentration of 3- CPA reduced TSS and increased titratable acidity. Incidence of burns on lower leaves of crown was observed in 150 and 200 ppm treatments.

Conclusion: 2-(3- chlorophenoxy) propionic acid is effective in altering the plant physiological processes and from the present experiment, treatment with 100 ppm showed effective result in terms of crown attributes, quality and shelf life.

Keywords: Pineapple crown, 2-3(chlorophenoxy) propionic acid, Fruitone CPA, 3-CPA.

1. INTRODUCTION

Pineapple (*Ananas comosus* L.) is a unique fruit of the tropics with a beautiful crown, however this crown becomes a burden to farmers and traders due to its big size and heavy weight which increases logistical expenses during storage and transport. Also, during the period of growth, the crown becomes so large and heavy that the fruit and peduncle are lodged and bends or falls over, leading to spoilage of fruits due to contact with soil. The complete removal of crown does not result in any increase in fruit size or yield, rather leads to injury of the central core and fruit becomes prone to pathogenic infection and sun burning (Dalldorf, 1975). Furthermore, according to Liu *et al.* (2017), postharvest detachment of pineapple crown significantly aggravated internal browning and increased yellowing or ripening of pineapple fruit, suggesting that the crown is necessary for extending shelf-life of pineapple.

Manual removal of crown becomes difficult and expensive when cultivated in large scale. Spraying of growth hormones to inhibit vegetative growth tends to be a practical approach in achieving small crown pineapples with enhancement in fruit quality and yield. The auxin 2 – (3- chlorophenoxy) propionic acid has been used in thinning peach and nectarines thus, Inhibitory effect on vegetative growth has been reported along with fruit thinning effect (Retamales & Copper, 1990). This growth hormone consists of synonyms such as Cloprop, 3- CPA, 3- CP, Propanoic acid, Fruitone CPA, α- (3- Chlorophenoxy) propionic acid etc (Anon., 2017). Findings have been reported

on the effect of this chemical on pineapple crown size, fruit size (Dalldorf, 1978), flowering time (Barbosa *et al.*, 2003) and delay in harvest (Rocha *et al.*, 1982). In plants, the chemical mimic the action of auxins, hormones chemically related to Indoleacetic acid that stimulates growth.

Taking into account the commercial importance of pineapple, a need was perceived to study the effect of the growth regulator 2- (3-chlorophenoxy propionic acid) on pineapple crown size, yield and quality of the fruits.

2. MATERIALS AND METHODS

The present study was conducted in the experimental field of the Department of Horticulture, SASRD, Nagaland University, Medziphema campus. The experimental site was located in the foothill of Nagaland at an altitude of 305 meters above mean sea level (MSL) with geographical location of 25°45'43" N latitude and 93°53'04" E. The experimental plot was situated at the subtropical and sub-humid climatic condition in foothills of Nagaland. Fruit quality parameters were analysed in the laboratory of the Department of Horticulture, SASRD, Nagaland University.

For enhancing growth of the ratoon pineapple plants, intercultural operations such as fertilization, weeding and earthing up were done at regular intervals. NPK dose of 12:4:12g per plant was applied in the month of July after harvest. Three times manual weeding was done during the period of investigation followed by earthing up of the plants.

The experiment was laid out in Completely Randomised Design with four replications comprising of five pineapple plants per replication. The treatments consisted of control and four concentrations: C₁ (Control), C₂ (50 ppm), C₃ (100 ppm), C₄ (150 ppm) and C₅ (200 ppm). The acid was dissolved in methanol and then added to water. The solution was applied as foliar spray on the crowns when the apical flowers were in the final stage of dehiscence or when the last flowers were about to dry out. Fruits were harvested when there was visible colour change at the base of developing fruit.

Growth and yield attributes *viz.*, length and circumference of crown, crown to fruit length ratio, length of fruit with and without crown, width of fruit, size of fruit, crown weight, weight of fruit with and without crown, pulp weight, peel weight, pulp to peel ratio, core weight, days taken to mature and dry matter content were recorded. Different bio-chemical parameters *viz.*, Total Soluble Solids, titratable acidity, TSS: acid ratio, ascorbic acid, total sugars, reducing sugars and non-reducing sugars were analysed according to the procedure reported by A.O.A.C (1995). Shelf life and Physiological Loss in Weight (PLW) during storage at ambient temperature were also recorded. The data obtained was statistically analyzed using method of Completely Randomised Design (CRD) with 5% level of significance as described by Gomez and Gomez (2012) and F- test was used to determine the significant difference between two means and critical difference (CD) was calculated for comparison in those cases where F was significant at 5% level of significance.

3. RESULTS AND DISCUSSION

3.1. Growth Parameters

Crown size showed significant decrease in terms of length, circumference, crown to fruit length ratio and weight of treated pineapple crowns in comparison to control pineapple crowns. The lowest crown length (14.28 cm), circumference (49.72 cm) and crown to fruit length ratio (0.83) were reported in C₃ (100 ppm) followed by C₄ (150 ppm). The highest values were observed in control (C₁). The highest crown weight (712 g) was observed in control and the lowest crown weight (163 g) was observed in C₅ (200 ppm) followed by C₄ (150 ppm) with 175 g. Incidence of burns on the lower leaves of crown at 150 and 200 ppm was observed which was also reported by Vieira *et al.* (1982). Similar findings have been reported by Fahl and Franco (1981) who reported the least crown length obtained from 112.5 ppm application. Cooke (1975), Selamat and Rahim (1997), Dalldorf (1978), Fahl and Franco (1981), Rocha *et al.* (1982), Vieira *et al.* (1982), Vieira and Gadelha (1987), Soler (1990), Rebolledo *et al.* (2002) and Othman *et al.* (2006) confirms reduction in crown size using the growth regulator 2- (3- Chlorophenoxy) propionic acid at various concentrations. The reduction in crown size may be due to the ability of 2- (3- chlorophenoxy) propionic acid to influence physiological processes (Vieira *et al.*, 1982) and alter growth cycle of pineapple, which can reduce crown development (Othman *et al.*, 2006).

Maximum fruit length (17.22 cm) was recorded in C₃ (100 ppm) followed by C₄ (150 ppm) with 16.94 cm and maximum fruit width (13.42 cm) was observed in C₄ (150 ppm) followed by C₃ (100 ppm). Subsequently, maximum fruit size was observed in C₄ (150 ppm) with 221 cm² followed by C₃ (100 ppm). The least values were observed in control. In the present finding, delay in maturity of

treated fruits was also observed with maximum days taken to mature (125 days) recorded in C₄ (150 ppm) followed by C₅ (200 ppm) with 120 days.

These findings are in partial conformity with Vieira *et al.* (1982) who reported maximum fruit length and fruit size in 200 ppm and maximum fruit width in 150 ppm concentration. Cantliffe (1982) has also reported increase in fruit circumference in pumpkin at 250 ppm and muskmelon at 500 ppm concentration. Increase in fruit diameter in peach was also reported by Shaybany (1979). Soler (1985) reported four weeks delay in harvest with 197 ppm concentration. Fahl and Franco (1981) also reported delay in maturation of fruits where 82% of control plant pineapples have already matured when the maturation of treated fruits begun. Rebolledo *et al.* (2002) reported 19 days delay in harvest over control when 1500 ml per ha of 3- CPA was applied in two equal applications. Selamat and Rahim (1997) also reported delay in shell colouring or ripening process by the use of the growth regulator.

3- CPA can modify the growth cycle of pineapple plant, which may delay maturation, and this allows for an extended assimilation of plant nutrients by the fruit, producing larger and heavier fruits and reducing the air space in the cavities, giving a more cylindrical shaped fruit, and delaying maturity (Othman *et al.*, 2006). Increase in weight of fruit was observed in the treated fruits. Maximum weight (1.83 kg) was recorded in C₅ (200 ppm) followed by 100 ppm (C₃) with 1.79 kg and the lowest weight of fruit was observed in C₁ (1.09 kg).

Table 1: Effect of 2- (3- chlorophenoxy) propionic acid on growth attributes of pineapple at harvest

TREATMENTS	Crown length (cm)	Crown circumference (cm)	Crown to fruit length ratio	Length of fruit with crown (cm)	Length of fruit without crown (cm)	Width of fruit (cm)
C ₁ (Control)	27.50	101.00	1.73	44.20	15.89	11.90
C ₂ (50 ppm)	25.10	80.59	1.55	42.30	16.23	12.93
C ₃ (100 ppm)	14.28	49.72	0.83	31.50	17.22	13.29
C ₄ (150 ppm)	16.72	58.09	0.99	33.67	16.94	13.42
C ₅ (200 ppm)	17.84	66.99	1.08	34.43	16.59	13.26
SEm±	0.79	1.63	0.05	0.63	0.24	0.17
CD (P=0.05)	2.39	4.91	0.16	1.90	0.72	0.52

Table 2: Effect of 2-(3-chlorophenoxy) propionic acid on yield attributes of pineapple at harvest

TREATMENTS	Size of fruit (cm ²)	Crown weight (g)	Weight of fruit with crown (kg)	Weight of fruit without crown (kg)	Pulp weight (g)	Peel weight (g)	Pulp to peel weight	Core weight (g)
C ₁ (Control)	169	1.81	1.09	712	731	322	2.27	158
C ₂ (50 ppm)	203	1.56	1.25	308	945	339	2.78	191
C ₃ (100 ppm)	210	2.01	1.79	215	1080	403	2.71	200
C ₄ (150 ppm)	221	1.99	1.83	163	1017	393	2.58	194
C ₅ (200 ppm)	213	1.88	1.75	134	1058	398	2.62	197
SEm±	3.67	0.05	0.04	24.87	26.62	6.53	0.06	6.46
CD (P=0.05)	11.06	0.15	0.13	74.98	80.23	19.69	0.19	19.48

3.2. Biochemical attributes

The treatment significantly reduced TSS and increased titratable acidity in pineapples whereby highest TSS (15.29°B) and lowest acidity (1.02%) was recorded in control (C₁) and the minimum TSS (13.28°B) and maximum titratable acidity (1.39%) was observed in C₄ (150 ppm).

This finding is in conformity with the findings of Othman *et al* (2006), Selamat and Rahim (1997), Soler (1985), Rebolledo and Uriza (1994), and Vieira *et al.* (1982) who reported decrease in TSS and increase in titratable acidity over control pineapples. However, in contrary, Vieira and Gadelha (1987)

reported no reduction in TSS content with application of 100 ppm at different stages of growth. Rebolledo *et al.* (2002) also reported Fruitone CPA did not lower the brix degrees at any concentrations, for which he concludes, reduction in TSS occurred in November and December application or in winter harvest.

Highest ascorbic acid value (8.58 mg/100g) was observed in C₅ (200 ppm) followed by C₃ (100 ppm) with 8.46 mg/ 100g and the lowest value was observed in C₂ (50 ppm) with 7.92 mg/100 g. Similar increase in ascorbic acid content due to application of Fruitone CPA was reported by Othman *et al.* (2006) and Soler (1985).

Decrease in total sugar content was recorded where the highest value (9.16%) was observed in control followed by C₃ (100 ppm) with 8.98% and the lowest content was reported in C₅ (8.46%). Minimum sugar content at harvest may be due to delay in ripening, thus delaying the conversion of polysaccharides to free sugars. Selamat and Rahim (1997) reported that with increase in Fruitone concentration, there was reduction in sugar content. Othman *et al.* (2006) also reported higher total sugar content in control pineapples over the treated ones at harvest.

Highest value for reducing sugar (4.40%) was observed in C₃ (100 ppm) followed by C₁ (3.95%) and the lowest value was observed in C₅ (3.35%). Highest non reducing sugar (5.17%) was observed in 50 ppm (C₂) followed by C₁ (4.95%) and the lowest ratio was reported in C₃ (4.35%).

Significant difference in dry matter content was observed. Highest value (3.89%) was observed in control followed by 50 ppm (3.83%) and the lowest (2.70%) was reported in C₅ (200 ppm). Nardoza *et al.* (2011) reported high dry matter genotypes in Kiwi fruit had higher TSS and were perceived as sweeter than low dry matter genotypes. The same trend between TSS and dry matter content has been observed in the findings.

Prolonged shelf life and minimal PLW of treated pineapples at ambient temperature was observed. Longest shelf life (19.33 days) and lowest PLW was observed in C₃ (100 ppm) followed by C₄ and the lowest shelf life and highest PLW was observed in C₁. The reduced weight loss may be due to increased fruit size resulting in reduced surface area to volume ratio, better filling of fruitlets, narrower locules and reduced air spaces which modified water deficit stress and the effect of the growth regulator on crown size reduction may also reduce weight loss because most of the fruit's initial water loss is through their stomata found substantially in the crown. The reduction in weight loss during storage, subsequently reduced the texture loss, maintained the freshness and delayed the ripening, hence, increased shelf life of treated pineapples. Delay in ripening in treated pineapples may be related to alterations in fruitlets air space which may have altered the internal atmosphere, retention of chlorophyll, photosynthesis, and respiratory rates (Othman *et al.*, 2006, 2007).

Table 3: Effect of 2-(3-chlorophenoxy) propionic acid on quality attributes of pineapple at harvest

Treatments	TSS (°B)	Titrateable acidity	TSS:Acid ratio	Ascorbic acid (mg/100g pulp)	Total sugar	Reducing sugar	Non reducing sugar	Dry mater content	Shelf-life days
C ₁ (Control)	15.29	1.02	15.13	8.20	9.16	3.95	4.95	3.89	14.67
C ₂ (50 ppm)	14.94	1.19	12.53	7.92	8.84	3.40	5.17	3.83	16.00
C ₃ (100 ppm)	14.80	1.22	12.15	8.46	8.98	4.40	4.46	3.61	19.33
C ₄ (150 ppm)	13.33	1.39	9.62	8.24	8.56	3.55	4.76	2.70	19.00
C ₅ (200 ppm)	13.28	1.23	10.78	8.58	8.46	3.35	4.85	2.92	18.33
SEm±	0.18	0.04	0.34	0.10	0.09	0.10	0.28	0.22	0.28
CD (P=0.05)	0.55	0.11	1.02	0.32	0.26	0.32	0.85	0.67	0.84

Table 4: Effect of 2-(3-chlorophenoxy) propionic acid on Physiological Loss of Weight (PLW) of pineapple during storage

Treatments	PLW (%)
------------	---------

	Initial weight of fruit (kg)	2 DAS	4 DAS	6 DAS	8 DAS	10 DAS	12 DAS	14 DAS
C₁ (Control)	1.68	1.44	2.39	2.84	5.88	7.84	9.48	14.04
C₂ (50 ppm)	1.80	1.30	2.22	3.04	4.19	5.40	6.03	6.85
C₃ (100 ppm)	1.94	1.51	1.76	2.97	3.53	4.32	4.54	4.85
C₄ (150 ppm)	1.87	1.58	1.17	2.07	2.87	3.44	4.05	4.97
C₅ (200 ppm)	1.64	1.75	1.38	2.31	3.37	4.04	5.24	6.28
SEm±	0.01	0.21	0.33	0.34	0.28	0.23	0.32	0.64
CD (P=0.05)	0.02	0.65	1.00	NS	0.85	0.69	0.96	1.92

4. CONCLUSION

From the present study, it was observed that treatment of pineapple with 100 ppm 2- (3-Chlorophenoxy) propionic acid (C₃) gave the most effective result in terms of crown attributes (minimum length, circumference and crown to fruit ratio), quality (maximum length of fruit, pulp weight, peel weight, core weight, reducing sugar), minimum PLW, longest shelf life and resulted in 48% reduction in crown length of treated pineapples over control pineapples.

5. REFERENCES

- A.O.A.C. 1995. Official method of Analysis of the Association of Official Agricultural Chemists. 16th Ed. Washington D.C. USA.
- Anonymous. 2017. 2- (3- Chlorophenoxy) propionic acid. National Institute of Standards and Technology, U.S. Department of Commerce.
- Barbosa, N. M. L., Cunha, G. A. P., Reindardt, D. H., Barros, P. G., Santos, A. R. L. and Do, S. 2003. Induction of leaf morphological and anatomical alterations on 'Perola' pineapple leaves by the 2- (3-Chlorophenoxy) propionic acid. *Revista Brasileira de Fruticultura*. **25**(3): 386-389.
- Cantliffe, D. J. 1982. Increased fruit size of several vegetable species by application of a growth regulator. In: Proceedings of the Florida State Horticultural Society. **95**: 351-353
- Cooke, A. R. 1975. Inhibiting crown growth on pineapple fruit. United States patent 3,894,860.
- Dalldorf, D. B. 1978. The effect of Chlorophenoxy propionamide (Fruitone CPA) on the fruit of the smooth cayenne pineapple. *The Citrus and Subtropical Fruit Journal*. (534): 17-18.
- Dalldorf, E. R. 1975. Removal of tops from Cayenne Pineapple fruit. *Citrus and Subtropical Fruit Journal*. **494**: 9- 10.
- Fahl, J. I. and Franco, J. F. 1981. Effects of 2-(3-chlorophenoxy) propionic acid (3-CPA) on fruits of 'cayenne' pineapple. *Planta daninha*. **4**(1): 17-20.
- Gomez, K. A. and Gomez, A. A. 2012. Statistical procedures for agricultural research. 2nd Ed. Ar. Emm. International, Delhi, ISBN: 978-81-65-2395-5.
- Liu, J., Congcong, H., Shen, F. S., Zhang, K. and Zhu, S. 2017. The crown plays an important role in maintaining quality harvested pineapple, *Post harvest biology and technology* **124**: 18-24.
- Nardoza, S., Gamble, J., Axten, L. G., Wohlers, M. W., Clearwater, M. J., Feng, J. and Harker, F. R. 2011. Dry matter content and fruit size affect flavor and texture of novel *Actinidia deliciosa* genotypes. *Journal of Science of Food Agriculture*. **91**(4): 742- 748
- Othman, Z., Mohamed, S., Madom, M. S. and Osman, A. 2007. Effect of different Fruitone concentrations on the physical characteristics and postharvest physiological disorder of cold stored pineapple. *Journal of Tropical Agriculture and Food Science*. **35**(1): 59- 69.

- Othman, Z., Madom, M. S., Osman, A. and Mohamed, S. 2006. The post harvest sensory and chemical characteristics of cold- stored pineapples pretreated with different Fruitone concentrations. *Journal of the Science of Food and Agriculture*. **86**: 2067-2075.
- Retamales, J. and Cooper, T. 1990. Effects of CGA 15281 and 3- CPA as thinning agents in nectarines. *Journal of Horticultural Science*. **65**(6): 639- 647.
- Rocha, A. D. D., Ruggiero, C., Yoshiura, A. Y. and Banzatto, D. A. 1982. Influence of rates and split applications of 2- (3- Chlorophenoxy) propionic aci American Society for Horticultural Science. **25**: 41-45.
- Rebolledo, M. A. and Uriza, A. D. 1994. Application of Fruitone and Gibberellins to delay ripening in pineapple in winter. In: Meeting of the InterAmerican Society for Tropical Horticulture. **10**: 51
- Rebolledo, M. A., Uriza, D. E. A., Rebolledo, M. L., Becerril, R. A. E. and Ruiz, P. Y. L. D. 2002. Fruitone CPA for delay the fruit maturity of pineapple cv. Smooth Cayenne, harvested in spring. *Brazilian Journal of Horticulture*. **24**(2): 354- 358
- Selamat, M. M. and Rahim, A. M. 1997. The effect of Fruitone® application on the yield and fruit quality of pineapple cv. Gandul grown on deep peat soil in Malaysia. *Pineapple news*. **3**(1): 11-12
- Shaybany, B., Costa, G., Brown, S. S., Obernauff, G., Martin, G. C. and Gerdts, M. 1979. Effect of 2- (3- Chlorophenoxy) propionic acid and 2- (3-chlorophenoxy) propionamide applications on fruit size and maturity of peach. *Journal of the American Society for Horticultural Science*. **104**(1): 34- 36.
- Soler, A. 1990. Advantages and limitations of the use of 3CPA (2-3-chlorophenoxy propionic acid) in d on agronomic characteristics of pineapple, cv. Smoth Cayenne. *Proceedings of the Tropical pineapple growing in Cote d'Ivoire*. *Fruits*. **45**(4): 357-365
- Soler, A. 1985. Use of Fruitone 3CPA as a growth regulator on pineapple cv. Smooth Cayenne in Ivory Coast. *Fruits*. **40**(1): 31- 38.
- Vieira, A., Gadelha, R. S. D. S. and Santos, A. C. D. 1982. Application of Fruitone CPA on pineapple cv. Smooth Cayenne. *Pesq. Agropec. Bras. Brasilia*. **17**(11): 1599- 1601.
- Vieira, A. and Gadelha, R. S. S. 1987. Effect of Chlorophenoxy propionic acid in Abacaxi fruit. *Pesq. Agropec. Bras. Brasilia*. **22**(7): 725- 727.