

Effect of Soil Test Crop Response Approach on Yield, Nutrient Content and Uptake by Wheat (*Triticum aestivum* L.)

ABSTRACT

A field experiment was carried out during *Rabi* season of 2018-19 under AICRP project on Soil Test Crop Response (STCR) at Agricultural Research Station, Swami Keshwanand Rajasthan Agricultural University, Bikaner to find out the effect of soil test crop response approach on yield, nutrient content and uptake of wheat. The experiment was laid out in randomized block design with five treatment combinations viz. general recommendation dose (T_1), STCR recommendation for target yield 30 q ha⁻¹ (T_2), STCR recommendation for target yield 30 q ha⁻¹ with IPNS (T_3), STCR recommendation for target yield 40 q ha⁻¹ (T_4) and STCR recommendation for target yield 40 q ha⁻¹ with IPNS (T_5) with four replications. Research finding of long term field experiment revealed that application of STCR recommendation for target yield 40 q ha⁻¹ with IPNS (T_5) gave maximum number of effective tillers m⁻¹ row length, spike length, number of grains spike⁻¹, weight of grains spike⁻¹, test weight, grain yield (4026 kg ha⁻¹) and straw yield (5629 kg ha⁻¹) of wheat which was found statistically at par with STCR recommendation for target yield 30 q ha⁻¹ with IPNS (T_3). Application of STCR recommendation for target yield 40 q ha⁻¹ with IPNS (T_5) also improved the N, P and K content and their uptake by grain and straw of wheat. These study results proved that STCR recommendation with IPNS gave better results of all parameters in the comparison of general recommendation dose and STCR without IPNS.

Key words: Growth, IPNS, Nutrient, STCR, Wheat, Yield

1. INTRODUCTION

Among the different food crops, role of cereals are highest towards transformation of Indian agriculture from deficit to sufficiency. Wheat (*Triticum aestivum* L.) is an important crop cultivated for food and feed under a wide range of agro-climatic

conditions in India. India stands in second position next to china in the world with regard to area and production of wheat. In India, wheat is grown on 29.14 million hectares with total production of 102.19 million tonnes with average productivity of 3507 kg ha⁻¹ (Anonymous, 2020). Rajasthan contributed 10.49 million tonnes of wheat from 3.0 million hectares area with productivity of 3501 kg ha⁻¹ (Anonymous 2020) to the national pool. Wheat productivity is highly variable within different agro-ecologies of India, due to variable climatic conditions, genotypes, seeding time and practices and other management practices (Kantwa *et al.*, 2015)

The wheat crop contributed significantly in food grain production for ensuring nations food security. There are reports of decreasing crop yields which raised questions on sustainability of the wheat crop. The reasons behind stability of crop yields are not exactly known, though it has been found that changes in quantity and quality of soil organic matter (SOM) and a gradual decline in the supply of soil nutrients are major responsible factors (Ladha *et al.* 2000). Long-term studies indicated that soil fertility is decreased mostly due to excessive removal of nutrients and inadequate refill through manures and fertilizers. Appropriate application of fertilizers based on soil test nutrient availability and crop response to fertilizer applied for specific target yield seems to be a good option to achieve targeted food grain production as well as reducing environmental degradation. STCR approach of fertilizer application involved both soil and plant analysis in a scientific basis that proved to be a refined and unique technique for most efficient use of fertilizer and soil nutrients. Several studies have documented effects of STCR based fertilizer recommendation with IPNS on soil nutrient status, soil organic carbon pools and potassium dynamics in soils. STCR approach is aiming at obtaining a basis for precise quantitative adjustment of fertilizer doses under varying soil test values and response conditions of the farmers and for targeted levels of crop production. According to STCR based research experiment results shown very close correlation between targeted yield and yield actually obtained and an evidence of the usefulness of soil testing within limit of variation under field conditions (Gulati *et al.*, 2005). Keeping all this point in mind the present investigation was conducted to determine the effect of Soil Test Crop Response approach on yield, nutrient content and uptake by wheat (*Triticum aestivum* L.).

2. MATERIALS AND METHOD

Experimental site: - The field experiment was carried out during *Rabi* season of 2018-19 under AICRP project on Soil Test Crop Response (STCR) at Agricultural Research Station, Swami Keshwanand Rajasthan Agricultural University, Bikaner. Geographically, the study area is located at 28° 10' N latitude, 73° 18' E longitude and 223.88 meters above mean sea level and this region falls under agro-climatic zone I C of Rajasthan. The climate of this tract is typically arid nature by aridity of the atmosphere and some salinity in the rhizosphere with extremes of temperature in summers and winters. There is wide range of temperature in summer (40° to 48°C) and in winter (-1° to 10°C) with annual rainfall of 250-270 mm. The soil of experimental field was loamy sand in texture, slightly alkaline in reaction.

Experimentation and crop husbandry: - The experiment was laid out in Randomized Block Design with five treatment combinations viz. general recommendation dose (T_1), STCR recommendation for target yield 30 q ha⁻¹ (T_2), STCR recommendation for target yield 30 q ha⁻¹ with IPNS (T_3), STCR recommendation for target yield 40 q ha⁻¹ (T_4) and STCR recommendation for target yield 40 q ha⁻¹ with IPNS (T_5) and replicated four times. Standard crop production practice and methods were followed for weeding, fertilizer application and crop protection management to grow the crop.

Data collection: - Five plants were selected from net plot and tagged for measurement of number of effective tillers m⁻¹ row length, spike length, number of grains spike⁻¹, weight of grains spike⁻¹, test weight, grain yield, straw yield and biological yield of wheat. Numbers of effective tillers were counted in five randomly selected per meter row lengths in each plot at physiological maturity of the crop. These were averaged to record and express effective tillers per meter row length. Five spikes were taken randomly from the observational plants in each plot and their length (cm) was measured with meter scale from neck node to tip and their average was computed. Randomly selected five productive spike heads plot⁻¹ was threshed separately and grains were counted and average number of grains spikes⁻¹ was worked out. The grains from five spikes of the plants tagged already were collected, weighed and average weight of grains spike⁻¹ (g) was worked out. A seed sample was select from the produce of each of the net plot harvested and 1000 seeds were counted and weighed to record as test weight in grams. The grain yield of each net plot (inclusive of tagged plants) was recorded in kg plot⁻¹ and the threshed produce was converted as q ha⁻¹. Straw yield in kg plot⁻¹ was calculated by deducting the grain yield from biological yield per plot

recorded earlier and expressed in $q\ ha^{-1}$. Grain and straw sample of wheat from individual plot was taken at the time of threshing for estimation of nutrients content. The samples were dried in oven and grind separately using grinder and N, P and K contents were determined by using standard methods. Nitrogen content was determined by colorimetric method using Nessler's reagent (Snell and Snell, 1939). Nitrogen content was expressed as percentage. Phosphorus content was estimated by Vanadomolybdate phosphoric yellow colour method (Jackson, 1973) and was expressed as percentage. Potassium content was obtained by flame photometer (Richards, 1954) and expressed as percentage. The uptake of nitrogen, phosphorus and potassium by grain, straw and total plant of wheat was calculated by following formula.

$$\text{Nutrient uptake = } \frac{\text{Per cent nutrient content in grain or straw} \times \text{Seed or straw yield (kg ha}^{-1}\text{)}}{100}$$

(kg ha⁻¹)

Statistical analysis: - The experimental data were subjected to statistical analysis by adopting appropriate method of analysis of variance assuming homogeneity, analysis of the data was carried out to establish the trend of treatments applied as per Gomez and Gomez [6]. Wherever, the F values were found significant at 5% level of probability, the critical difference (CD) values were computed for making comparison among the treatment means.

3. RESULTS AND DISCUSSION

Yield attributes and yield: -

Analysis of data stated that different treatments bring significant effect on yield attributes viz. effective tillers m^{-1} row length, spike length, number of grains spike⁻¹, weight of grains spike⁻¹ and yields i.e. (grain and straw) of wheat. The maximum values of effective tillers m^{-1} row length, spike length, number of grains spike⁻¹, weight of grains spike⁻¹ and yields i.e. (grain and straw) was obtained by treatment T₅ (STCR recommended for target yield 40 $q\ ha^{-1}$ with IPNS). The data related to correlation study between grain yield and yield attributes showed that relationship between grain yield and yield attributes of wheat was also found positive and significant. The reason behind improvement in yield parameters and yield of wheat with application of STCR based fertilizer recommendation for yield of the crop along with IPNS treatments in the

comparison of normal recommendation and soil test-based treatments because steady and justifiable use of the inorganic fertilizers with FYM. The beneficial impact of fertilizers and organic manure (STCR recommended for target yield with IPNS) on wheat might be the availability of sufficient amount of plant nutrients to crop which resulted better uptake of nutrients, plant vigour and improved yield. By virtue of increased availability of nutrients, there might have been increment in dry matter with increasing fertilizer application and increased growth components due to improved fertilizer levels gave stability in higher supply of photosynthates towards the sink (grains spike⁻¹). As a result, all yields attributes of wheat increased significantly. The improvement in yield (grain and straw) might be due to improved nutritional environment in low status of N and P₂O₅ soil through increased supply of major nutrients which increase their uptake by plants led to enhanced growth and yield attributes, which ultimately resulted in increased yield (grain and straw) of the crop. Tiwari *et al.* (2013), Sellamuthu *et al.* (2015) and Singh *et al.* (2016) were also noted similar results which support the results of present experiment.

Nutrient content and uptake: - The significantly highest nitrogen, phosphorus and potassium content in grain, straw and root and their uptake in grain, straw, root and total plant of wheat was obtained with STCR recommended for target yield 40 q ha⁻¹ with IPNS (T₅) which was superior to T₂ (STCR recommended for target yield 30 q ha⁻¹), and T₄ (STCR recommended for target yield 40 q ha⁻¹) but found at par with T₁ (general recommended dose) and T₃ (application of STCR recommended for target yield 30 q ha⁻¹ with IPNS). The significant increase in nitrogen, phosphorus and potassium content in grain, straw and root and their uptake in grain, straw, root and total plant of wheat with the application of STCR recommended for target yield (40 q ha⁻¹) with IPNS could be due to improved nutrient supplying capacity of nutrients and flush of available nutrients on autolysis of microbial cells besides, improvement in biochemical parameters of soil. When organic manure is added to soil, complex nitrogenous compounds slowly break-down and make steady nitrogen, phosphorus and potassium supply throughout growth period of crop which might be attributed to more nutrient availability. The increase in nitrogen, phosphorus and potassium content in grain, straw and root of wheat with the application of fertilizers might be due to improved nutritional environment in the rhizosphere as well as in the plant system leading to enhanced translocation of N, P and K in plant parts which ultimately enhance the uptake of

nutrient in all parts of plant. Similar finding has also been reported by Mali *et al.* (2015) who reported positive influence of fertilizers and FYM on nitrogen content in grain and straw of wheat. Sawarkar *et al.* (2013), Jat *et al.* (2014) and Sharma *et al.* (2016) has also been reported similar positive influence of inorganic fertilizers alone or in combination with organic manure.

4. CONCLUSION

Results of a long-term experiment concluded that STCR recommendation with IPNS on wheat brought an additive effect in increasing yield attributes, yield, nutrient content and nutrient uptake by wheat crop. Application of STCR recommended for target yield 40 q ha⁻¹ with IPNS (T₅) gave maximum yield attributes, grain yield (4026 kg ha⁻¹), straw yield (5629 kg ha⁻¹), N, P and K content in grain and straw and their uptake in by grain, straw and total plant which was statistically found at par with treatment T₃ (STCR recommended for target yield 30 q ha⁻¹ with IPNS). These study results proved that STCR recommended with IPNS gave better results of all parameters in the comparison of general recommended dose and STCR without IPNS.

Table 1. Effect of STCR approach on yield attributes and yield of wheat

Treatments	Yield attributes					Yield (kg ha ⁻¹)		
	No. of effective tiller m ⁻¹ row length	No. of non-effective tiller m ⁻¹ row length	Spike length (cm)	No. of grains spike ⁻¹	Grain weight spike ⁻¹ (g)	Test weight (g)	Grain yield	Straw yield
T ₁	72.1	13.0	7.96	30.8	2.71	31.26	2711	3852
T ₂	73.4	12.7	8.13	31.1	2.76	32.10	3314	4668
T ₃	85.8	11.2	9.63	37.7	3.44	37.83	3980	5580
T ₄	75.1	11.4	8.27	31.4	2.84	32.80	3397	4770
T ₅	87.2	11.0	9.77	38.7	3.50	38.44	4026	5629
SEm±	2.53	0.36	0.36	0.36	0.15	1.76	164	212
CD (P = 0.05)	7.8	1.11	1.12	1.1	0.45	5.4	505	652

Table 2. Relationship between grain yield and yield attributes of wheat

	Grain yield	No. of effective tiller	No. of non-effective tiller	Spike length	No. of grains spike ⁻¹	Grain weight spike ⁻¹	Test weight
Grain yield	1						
No. of effective tiller	0.927*	1					
No. of non-effective tiller	0.932*	0.999**	1				
Spike length	0.923*	1	0.999**	1			
No. of grains spike ⁻¹	0.894*	0.995**	0.995**	0.997**	1		
Grain weight spike ⁻¹	0.916*	0.999**	0.998**	1	0.997**	1	
Test weight	0.980**	0.982**	0.984**	0.981**	0.965**	0.977**	1

Table 3. Effect of STCR approach on N, P and K content in grain, straw and their uptake by grain, straw and total plant of wheat

Treatments	N content (%)		N uptake (kg ha ⁻¹)			P content (%)		P uptake (kg ha ⁻¹)			K content (%)		K uptake (kg ha ⁻¹)		
	In grain	In straw	By grain	By straw	Total	In grain	In straw	By grain	By straw	Total	In grain	In straw	By grain	By straw	Total
T ₁	1.64	0.67	35.4	25.7	61.1	0.542	0.225	14.6	8.7	23.3	0.442	1.09	12.0	42.1	54.1
T ₂	1.53	0.61	50.8	28.4	79.2	0.492	0.202	16.3	9.5	25.7	0.394	1.01	13.0	46.9	59.9
T ₃	1.68	0.69	66.8	38.3	105.0	0.588	0.252	23.4	14.0	37.4	0.447	1.15	17.8	64.2	82.0
T ₄	1.60	0.63	54.3	29.9	84.2	0.546	0.230	18.6	11.0	29.5	0.407	1.06	13.8	50.8	64.6
T ₅	1.72	0.71	69.3	40.0	109.2	0.593	0.256	23.9	14.4	38.3	0.458	1.17	18.4	65.9	84.4
SEm+	0.05	0.02	4.2	1.7	4.8	0.013	0.006	0.9	0.7	1.4	0.012	0.03	0.7	2.7	3.0
CD (P = 0.05)	0.14	0.06	12.9	5.3	14.9	0.040	0.018	2.6	2.1	4.3	0.037	0.10	2.1	8.2	9.4

REFERENCES

1. Anonymous, Government of India, Ministry of Agriculture and Farmers Welfare, Department of Agriculture Cooperation and Farmers Welfare, Directorate of Economics and Statistics, 2020, Page- 51.
2. Gulati, R.; Lawrence, P.R.; Puranam, P. Adaptation in vertical relationships: beyond incentive conflict. *Strategic Management Journal*, 2005, **26**: 415-440.
3. Jackson, M.L. Soil chemical analysis, Prentice Hall of India Pvt. Ltd., New Delhi, 1973.
4. Jat, G.; Majumdar, S.P.; Jat, N.K.; Mazumdar, S.P. Effect of potassium and zinc fertilizer on crop yield, nutrient uptake and distribution of potassium and zinc fractions in Typic Ustipsamment. *Indian Journal of Agricultural Sciences* 2014, **84**: 832–838
5. Kantwa, S.R.; Choudhary, U.; Sai Prasad, S.V. Tillage x early sown wheat genotypes interaction effect on nutritional quality, productivity and profitability in central India. *Green farming*. 2015, **6**: 1098-1101.
6. Ladha J.K.; Fischer, K.S., Hossain, M., Hobbs, P.R., Hardy B. Improving the productivity and sustainability of rice-wheat systems of the Indo-Gangetic Plains: a synthesis of NARS-IRRI partnership research. 2000, **40**: 31-35.
7. Mali, D.V.; Kharche, V.K.; Jadhao, S.D.; Jadhao, S.M.; Saoji, B.V.; Gite, P.A.; Age, A.B. Soil biological health under long-term fertilization in sorghum- wheat sequence on swell-shrink soils of central India. *Journal of the Indian Society of Soil Science* 2015, **63**: 423-428.
8. Richards, L.A. Diagnosis and improvement of saline and alkaline soils. USDA Hand Book No. 60. *Oxford and IBH Publication*, New Delhi, 1954.
9. Sawarkar, S.D.; Khamparia, N.K.; Thakur, R.; Dewda, M.S.; Singh, M. Effect of long-term application of inorganic fertilizers and organic manure on yield, potassium uptake and profile distribution of potassium fractions in Vertisol under soybean-wheat cropping system. *Journal of the Indian Society of Soil Science*, 2013, **61**: 94-98.
10. Sellamuthu, K.M.; Santhi, R.; Maragatham, S.; Dey, P. Validation of soil test and yield target-based fertilizer prescription model for wheat on Inceptisol. *Research on Crops* 2015, **16**: 53-58.

11. Sharma, V.K.; Pandey, R.N.; Kumar, S.; Chobhe, K.A.; Chandra, S. Soil test crop response-based fertilizer recommendations under integrated nutrient management for higher productivity of pearl millet (*Pennisetum glaucum*) and (*Triticum aestivum*) wheat under long term experiment. *Indian Journal of Agricultural Sciences*, 2016, **86**: 1076–81.
12. Singh, Y.V.; Singh, S.K.; Sharma, P.K.; Singh, P. Soil test based integrated fertilizer recommendation for wheat (*Triticum aestivum*) in an Inceptisol of eastern plain zone of Uttar Pradesh. *Journal of the Indian Society of Soil Science* 2014, **62**: 255-258.
13. Snell, F.D.; Snell, C.P. Colorimetric Methods of Analysis, 3rd Ed. Vol. 2nd D. Van Nostrand Inc. New York, 1949.
14. Tiwari, L.B.; Mishra, D.D.; Gupta, J. Evaluation of integrated nutrient management options in rice (*Oryza sativa*)- wheat (*Triticum aestivum*) cropping system in reclaimed sodic land. *Indian Journal of Agronomy* 2013, **58**: 137-14.