

**EVALUATION OF SOIL CHEMICAL PROPERTIES AND ORGANIC CARBON ALONG THE DISTANCE GRADIENT FROM SEA COAST IN COASTAL SOILS OF NORTHERN SAURASHTRA REGION OF GUJARAT**

**ABSTRACT**

**Background-** In present time, salinity and alkalinity of the soils are serious problems in India as well as in Gujarat. These soils are usually supposed to be originated as a result of high water table, arid and semi-arid weather, ingress the sea, water, saline nature of barren materials, poor drainage and salt deposition through wind-blown particles. The current study aimed to study the soil chemical properties and organic carbon in soils of Northern Saurashtra coastal region (Jamnagar, Devbhumi Dwarka and Porbandar district) of Gujarat.

**Methods-** In this field-laboratory investigation, different talukas of Northern Saurashtra coastal region were surveyed during May, 2019, to study the soil properties by collecting 141 grid based surface (0-15 cm) soil samples from farmer's cultivated field, through the use of GPS at distance demarcation of 0-5, 5-10, 10-15 and 15-20 km from sea coast, during May, 2019 and were analyzed for different chemical properties. Sampling localities were selected based on geographical situation, climate and local data. In the laboratory, the collected samples were analyzed for SOC, EC<sub>2.5</sub>, CaCO<sub>3</sub>, CEC and water soluble ions.

**Result-** On the basis of analyzed data, it can be concluded that EC<sub>2.5</sub>, CaCO<sub>3</sub>, CEC and water soluble ions except K<sup>+</sup> were decreased, while pH<sub>2.5</sub> and SOC was slightly increased with increasing the distance from sea coast. Soil EC<sub>2.5</sub> was found beyond to its critical or marginal limit up to 0 to 5 km distance from sea coast. The soil organic carbon status of Northern Saurashtra coastal region was found in medium (5.14 g kg<sup>-1</sup>) category. The soils are calcareous in nature (CaCO<sub>3</sub> 121.20 g kg<sup>-1</sup>) with alkaline in reaction (pH<sub>2.5</sub> 7.58).

**Keywords:** Northern, CEC, CaCO<sub>3</sub>, Saurashtra, Coastal

**1. INTRODUCTION**

Soil survey provides useful information for planning proper soil management practices, with play an important part in augmenting crop production. Poor performance of crops in salts affected soil may be due to excessive quantities of soluble salts, which


consequently resulted in nutritional disorders in plants. In present time, salinity and alkalinity of the soils are serious problems in India as well as in Gujarat. These soils are usually supposed to be originated as a result of high water table, arid and semi-arid weather, ingress the sea, water, saline nature of barren materials, poor drainage and salt deposition through wind-blown particles.

Around 6.72 million ha area in India is salt-affected, of which 2.95 million ha is saline and the rest 3.77 million ha is sodic [1]. Nearly 75% of salt-affected soils in the country exist in the states of Gujarat (2.23 million ha), Uttar Pradesh (1.37 million ha), Maharashtra (0.61 million ha), West Bengal (0.44 million ha) and Rajasthan (0.38 million ha) [2]. Salt build-up in soils and water are major constraints for human habitat, sustainable development, soil health and crop productivity [3] due to severe problem of water and soil salinity in the coastal regions [4]. Since soils in general are degrading due to poor management and faulty land use at a rate faster than their natural degeneration, it becomes imperative to protect them from further degradation; as there is a concomitant decline in the quality of soil to produce healthy crops. Soil reaction (pH), electrical conductivity (EC) and other properties of soil have marked effects on plant growth. The most important constituent in soil is organic matter, an appreciable amount of it in the soil tremendously increase soil fertility. Decay of organic matter releases nitrogen, phosphorus and mineral nutrient in available form to plants. Organic carbon is also positively correlated with total and available nitrogen in all soil groups [5]. Better crop production in salt affected soils can be attained by if the nature and extent of salinity problems are correctly diagnosed and appropriate reclamation and management practices are adopted.

## **2. MATERIAL AND METHODS**

### **2.1.1 Collection and preparation of soil samples**

Study had been performed in Northern Saurashtra coastal region of Gujarat. Soil samples were collected from distance demarcation of 0-5, 5-10, 10-15 and 15-20 km from sea coast through use of GPS.



**Fig. 1: Map of survey talukas of Jamnagar, Devbhumi Dwarka and Porbandar district**

The twenty surface soil samples were collected from each taluka viz. Jodiya, Jamnagar and Lalpur talukas of Jamnagar district, Khambhaliya, Dwarka and Kalyanpur talukas of Devbhumi Dwarka district and Porbandar taluka of Porbandar district of Northern Saurashtra Coastal region of Gujarat during the summer season of year 2019 (Fig. 1).

Two kilograms of soil sample were collected in plastic bag and transfer to cotton bag from the surface soils of study area and it was label properly. The collected soil samples were air dried and gently crushed (ground) with a wooden mortar with pestle and passed though 2 mm sieve and chemically analyzed.

### 2.1.2 Methods for analysis

Soil samples were analyzed for chemical properties viz;  $EC_{2.5}$ ,  $pH_{2.5}$ , free lime, organic carbon, CEC and water soluble ions by using standard methods. The  $EC_{2.5}$ ,  $pH_{2.5}$  and water soluble ions of soil samples were determined by use of 1:2.5 soil-water ratio, free lime (equivalent) content was estimated by the rapid acid neutralization method, SOC was determined by 1 N  $K_2Cr_2O_7$  method as described by Walkley and Black [6] and CEC was worked out by Ammonium acetate extractable method as described by Chapman [7].

## 3. RESULTS AND DISCUSSION

### 3.1 EC<sub>2.5</sub>

The EC<sub>2.5</sub> of soil samples were determined by use of 1:2.5 soil-water ratio. Overall, it was varied widely ranging from 0.23 to 5.97 dS m<sup>-1</sup> with a mean value of 1.23 dS m<sup>-1</sup> (Table 1). The lowest (0.23 dS m<sup>-1</sup>) EC<sub>2.5</sub> value was recorded in the soil sample collected from Kalyanpur taluka at the distance of 15 to 20 km from the sea coast, whereas highest value of (5.97 dS m<sup>-1</sup>) was recorded in the soil sample collected from Dwarka taluka at the distance of 0 to 5 km from the sea coast. The data further revealed that the lowest mean value of 0.86 dS m<sup>-1</sup> was obtained in the soils of Lalpur taluka and the highest mean value of 1.39 dS m<sup>-1</sup> was registered in the soils of Porbandar taluka. In Jamnagar district, overall mean value of EC<sub>2.5</sub> was 1.09 dS m<sup>-1</sup>, maximum EC<sub>2.5</sub> (3.67 dS m<sup>-1</sup>) was found at 0 to 5 km distance from the sea coast and minimum EC<sub>2.5</sub> (0.29 dS m<sup>-1</sup>) was found at 15 to 20 km distance from the sea coast. In Devbhumi Dwarka district, maximum EC<sub>2.5</sub> (5.97 dS m<sup>-1</sup>) was found at 0 to 5 km distance from the sea coast and minimum EC<sub>2.5</sub> (0.23 dS m<sup>-1</sup>) was found at 15 to 20 km distance from the sea coast, while overall mean value of EC<sub>2.5</sub> was 1.32 dS m<sup>-1</sup>. In Porbandar district, overall mean value of EC<sub>2.5</sub> was 1.39 dS m<sup>-1</sup>, maximum EC<sub>2.5</sub> (4.80 dS m<sup>-1</sup>) was found at 0 to 5 km distance from the sea coast and minimum EC<sub>2.5</sub> (0.30 dS m<sup>-1</sup>) was found at 15 to 20 km distance from the sea coast.

The wide variation in EC<sub>2.5</sub> could be due to the accumulation of salts in underground water or influence of sea water or shallow water table or poor quality of ground water prevailing in a particular location. Similar results were also obtained for Girnar toposequence [8], for Patan district [9], for Kapada district of Andhra Pradesh [10], for Nagpur district of Maharashtra [11], for Nagaur district of Rajasthan [12] and for Southern saurashtra of Gujarat [13].

**Table 1: Talukawise range and mean values of EC<sub>2.5</sub> (dS m<sup>-1</sup>) in different districts of Northern Saurashtra coastal region**

| Distance (km)<br>Talukas | 0 to 5           |             | 5 to 10          |             | 10 to 15         |             | 15 to 20         |             | Overall          |             |
|--------------------------|------------------|-------------|------------------|-------------|------------------|-------------|------------------|-------------|------------------|-------------|
|                          | Range            | Mean        | Range            | Mean        | Range            | Mean        | Range            | Mean        | Range            | Mean        |
| Jamnagar                 | 2.78-3.56        | 3.17        | 0.55-1.84        | 1.07        | 0.46-1.13        | 0.76        | 0.37-0.84        | 0.55        | 0.37-3.56        | 1.08        |
| Jodiya                   | 1.37-3.67        | 2.53        | 0.76-1.74        | 1.23        | 0.48-1.20        | 0.81        | 0.32-0.89        | 0.58        | 0.32-3.67        | 1.34        |
| Lalpur                   | 1.19-3.13        | 2.01        | 0.53-1.32        | 0.85        | 0.36-0.92        | 0.59        | 0.29-0.47        | 0.36        | 0.29-3.13        | 0.86        |
| Jamnagar District        | <b>1.19-3.67</b> | <b>2.41</b> | <b>0.53-1.84</b> | <b>1.05</b> | <b>0.36-1.20</b> | <b>0.70</b> | <b>0.29-0.89</b> | <b>0.51</b> | <b>0.29-3.67</b> | <b>1.09</b> |
| Kalyanpur                | 1.31-5.60        | 2.58        | 0.58-1.33        | 0.86        | 0.38-1.05        | 0.67        | 0.23-0.56        | 0.36        | 0.23-5.60        | 1.13        |
| Khambhalia               | 1.17-4.33        | 2.59        | 0.54-1.64        | 0.90        | 0.45-0.97        | 0.65        | 0.24-0.46        | 0.36        | 0.24-4.33        | 1.05        |
| Dwarka                   | 1.44-5.97        | 2.67        | 0.80-2.31        | 1.45        | 0.57-1.13        | 0.82        | 0.36-0.69        | 0.53        | 0.36-5.97        | 1.78        |

|                          |           |      |           |      |           |      |           |      |           |      |
|--------------------------|-----------|------|-----------|------|-----------|------|-----------|------|-----------|------|
| Devbhumi Dwarka District | 1.17-5.97 | 2.63 | 0.54-2.31 | 1.05 | 0.38-1.13 | 0.71 | 0.23-0.69 | 0.39 | 0.23-5.97 | 1.32 |
| Porbandar                | 1.44-4.80 | 2.73 | 0.82-1.74 | 1.18 | 0.58-1.11 | 0.86 | 0.30-0.78 | 0.51 | 0.30-4.80 | 1.39 |
| Overall                  | 1.17-5.97 | 2.58 | 0.53-2.31 | 1.06 | 0.36-1.20 | 0.73 | 0.23-0.89 | 0.46 | 0.23-5.97 | 1.23 |

### 3.2 pH<sub>2.5</sub>

In general, the soil samples of Northern Saurashtra coastal region were slightly alkaline in reaction. The overall pH<sub>2.5</sub> values of Northern Saurashtra's soil samples were ranged from 7.20 to 8.59 with mean value of 7.78. The data (Table 2) revealed that the lowest mean value of pH<sub>2.5</sub> 7.66 was obtained from the samples of Kalyanpur taluka of Devbhumi Dwarka district and the highest mean value of pH<sub>2.5</sub> 7.96 was found in the samples of Dwarka taluka of Devbhumi Dwarka district. In Jamnagar district, maximum pH<sub>2.5</sub> (8.21) was found at 5 to 10 km distance from the sea coast and minimum pH<sub>2.5</sub> (7.20) was found at 0 to 5 km distance from the sea coast, whereas overall mean value of pH<sub>2.5</sub> was 7.75.

**Table 2: Talukawise range and mean values of pH<sub>2.5</sub> in different districts of Northern Saurashtra coastal region**

| Distance (km)<br>Talukas | 0 to 5           |             | 5 to 10          |             | 10 to 15         |             | 15 to 20         |             | Overall          |             |
|--------------------------|------------------|-------------|------------------|-------------|------------------|-------------|------------------|-------------|------------------|-------------|
|                          | Range            | Mean        | Range            | Mean        | Range            | Mean        | Range            | Mean        | Range            | Mean        |
| Jamnagar                 | 7.20-7.64        | 7.42        | 7.51-8.21        | 7.84        | 7.41-8.20        | 7.78        | 7.51-8.18        | 7.77        | 7.20-8.21        | 7.76        |
| Jodiya                   | 7.22-7.66        | 7.43        | 7.37-7.95        | 7.62        | 7.68-8.16        | 7.91        | 7.74-8.20        | 7.97        | 7.22-8.20        | 7.70        |
| Lalpur                   | 7.51-7.83        | 7.70        | 7.37-8.02        | 7.79        | 7.42-7.90        | 7.71        | 7.91-8.07        | 7.97        | 7.37-8.07        | 7.78        |
| Jamnagar District        | <b>7.20-7.83</b> | <b>7.51</b> | <b>7.37-8.21</b> | <b>7.75</b> | <b>7.42-8.20</b> | <b>7.78</b> | <b>7.51-8.20</b> | <b>7.90</b> | <b>7.20-8.21</b> | <b>7.75</b> |
| Kalyanpur                | 7.28-7.74        | 7.48        | 7.60-8.14        | 7.77        | 7.64-7.84        | 7.71        | 7.51-7.87        | 7.70        | 7.28-8.14        | 7.66        |
| Khambhalia               | 7.21-8.09        | 7.63        | 7.50-7.83        | 7.70        | 7.62-8.12        | 7.87        | 7.70-8.02        | 7.87        | 7.21-8.12        | 7.76        |
| Dwarka                   | 7.44-8.04        | 7.81        | 7.65-8.12        | 7.82        | 7.93-8.59        | 8.19        | 8.52-8.55        | 8.54        | 7.44-8.59        | 7.96        |
| Devbhumi Dwarka District | <b>7.21-8.09</b> | <b>7.68</b> | <b>7.50-8.14</b> | <b>7.76</b> | <b>7.62-8.59</b> | <b>7.89</b> | <b>7.51-8.55</b> | <b>7.93</b> | <b>7.21-8.59</b> | <b>7.80</b> |
| Porbandar                | 7.36-8.10        | 7.74        | 7.50-8.12        | 7.75        | 7.55-8.15        | 7.83        | 7.71-8.19        | 7.90        | 7.36-8.19        | 7.80        |
| Overall                  | <b>7.20-8.10</b> | <b>7.64</b> | <b>7.37-8.21</b> | <b>7.75</b> | <b>7.42-8.59</b> | <b>7.83</b> | <b>7.51-8.55</b> | <b>7.91</b> | <b>7.20-8.59</b> | <b>7.78</b> |

In Devbhumi Dwarka district, maximum pH<sub>2.5</sub> (8.59) was found at 10 to 15 km distance from the sea coast and minimum pH<sub>2.5</sub> (7.21) was found at 0 to 5 km distance from the sea coast, while overall mean value of pH<sub>2.5</sub> was 7.80. In Porbandar district, overall mean value of pH<sub>2.5</sub> was 7.80, maximum pH<sub>2.5</sub> (8.19) was found at 15 to 20 km distance from the sea coast and minimum pH<sub>2.5</sub> (7.36) was found at 0 to 5 km distance from the sea coast. Overall, soil pH<sub>2.5</sub> was below its critical limit in coastal region of North Saurashtra. Similar results

were also obtained for Pedapuluguvaripalem village of Guntur district [14], for Patan district [9], for Kapada district of Andhra Pradesh [10], for Nagpur district of Maharashtra [11], for Nagaur district of Rajasthan [12] and for Southern saurashtra of Gujarat [13].

### 3.3 Free Lime ( $\text{CaCO}_3$ )

The overall free lime content was ranging from 12.34 to 420.31  $\text{g kg}^{-1}$  with mean value of 121.20  $\text{g kg}^{-1}$  (Table 3) indicating the calcareous nature of the soil. This might be due to impregnation of lime in the transported materials and accumulation of shells in the marine alluvial soils particularly in Northern Saurashtra coastal region.

**Table 3: Talukawise range and mean values of  $\text{CaCO}_3$  ( $\text{g kg}^{-1}$ ) in different districts of Northern Saurashtra coastal region**

| Distance (km)<br>Talukas    | 0 to 5                    |                    | 5 to 10                  |                    | 10 to 15                 |              | 15 to 20                |              | Overall                  |                    |
|-----------------------------|---------------------------|--------------------|--------------------------|--------------------|--------------------------|--------------|-------------------------|--------------|--------------------------|--------------------|
|                             | Range                     | Mean               | Range                    | Mean               | Range                    | Mean         | Range                   | Mean         | Range                    | Mean               |
| Jamnagar                    | 103.24-<br>132.54         | 117.8<br>9         | 63.12-<br>90.00          | 74.05              | 42.58-<br>60.23          | 50.01        | 19.99-<br>40.67         | 28.95        | 19.99-<br>132.54         | 62.20              |
| Jodiya                      | 174.07-<br>231.04         | 194.7<br>8         | 114.20-<br>170.00        | 136.6<br>4         | 60.12-<br>95.37          | 72.16        | 42.58-<br>52.02         | 47.45        | 42.58-<br>231.04         | 120.4<br>4         |
| Lalpur                      | 122.08-<br>152.47         | 138.5<br>2         | 50.00-<br>102.03         | 67.90              | 23.21-<br>50.00          | 36.65        | 13.02-<br>34.06         | 23.73        | 13.02-<br>152.47         | 60.93              |
| Jamnagar District           | <b>103.24-<br/>231.04</b> | <b>162.5<br/>3</b> | <b>50.00-<br/>90.00</b>  | <b>92.01</b>       | <b>23.21-<br/>95.37</b>  | <b>49.72</b> | <b>13.02-<br/>52.02</b> | <b>34.25</b> | <b>13.02-<br/>231.04</b> | <b>81.19</b>       |
| Kalyanpur                   | 224.10-<br>360.12         | 269.4<br>2         | 56.24-<br>163.28         | 116.6<br>6         | 12.34-<br>80.00          | 50.43        | 12.34-<br>41.03         | 25.85        | 12.34-<br>360.12         | 116.8<br>2         |
| Khambhalia                  | 114.02-<br>241.03         | 186.3<br>4         | 68.17-<br>114.00         | 93.74              | 42.74-<br>90.00          | 65.90        | 18.24-<br>52.64         | 31.01        | 18.24-<br>241.03         | 91.01              |
| Dwarka                      | 190.00-<br>400.28         | 285.5<br>4         | 100.04-<br>187.28        | 136.5<br>1         | 88.17-<br>116.26         | 97.20        | 32.17-<br>68.02         | 50.10        | 32.17-<br>400.28         | 187.0<br>7         |
| Devbhumi<br>Dwarka District | <b>114.02-<br/>400.28</b> | <b>259.0<br/>2</b> | <b>56.24-<br/>187.28</b> | <b>113.0<br/>6</b> | <b>12.34-<br/>116.26</b> | <b>68.21</b> | <b>12.34-<br/>41.03</b> | <b>32.60</b> | <b>12.34-<br/>400.28</b> | <b>131.6<br/>3</b> |
| Porbandar                   | 241.25-<br>420.31         | 348.3<br>2         | 192.03-<br>292.57        | 232.9<br>2         | 91.47-<br>190.36         | 138.6<br>3   | 45.05-<br>92.14         | 74.29        | 45.05-<br>420.31         | 205.6<br>7         |
| Overall                     | <b>103.24-<br/>420.31</b> | <b>246.4<br/>0</b> | <b>50.00-<br/>292.57</b> | <b>116.1<br/>5</b> | <b>12.34-<br/>190.36</b> | <b>69.26</b> | <b>12.34-<br/>92.14</b> | <b>41.00</b> | <b>12.34-<br/>420.31</b> | <b>121.2<br/>0</b> |

In Jamnagar district, overall mean value of free lime was 81.19  $\text{g kg}^{-1}$ , maximum free lime (231.04  $\text{g kg}^{-1}$ ) was found at 0 to 5 km distance from the sea coast and minimum free lime (13.02  $\text{g kg}^{-1}$ ) was found at 15 to 20 km distance from the sea coast. In Devbhumi

Dwarka district, maximum free lime ( $400.28 \text{ g kg}^{-1}$ ) was found at 0 to 5 km distance from the sea coast and minimum free lime ( $12.34 \text{ g kg}^{-1}$ ) was found at 15 to 20 km distance from the sea coast, while overall mean value of free lime was  $131.63 \text{ g kg}^{-1}$ . In Porbandar district, overall mean value of free lime was  $205.67 \text{ g kg}^{-1}$ , maximum free lime ( $420.31 \text{ g kg}^{-1}$ ) was found at 0 to 5 km distance from the sea coast and minimum free lime ( $45.05 \text{ g kg}^{-1}$ ) was found at 15 to 20 km distance from the sea coast. The lowest value of free lime  $12.34 \text{ g kg}^{-1}$  was recorded in the samples collected from Kalyanpur taluka in Devbhumi Dwarka district, whereas the highest value of free lime  $420.31 \text{ g kg}^{-1}$  was found in Porbandar taluka of Porbandar district.


Similar results were also observed for the soils of north-west agro climatic zone of Gujarat [15], for Bhavnagar district [16], for Latur district [17] and for Southern saurashtra of Gujarat [13].

### 3.4 Soil Organic Carbon

Organic carbon content in the soils is important parameter from the fertility and physical properties points of view. Hence, the samples were analyzed for soil organic carbon content in Northern Saurashtra coastal region (Table 4, Fig. 2). In general, the soils of Northern Saurashtra coastal region are medium in O.C. status. The overall organic carbon content in the soils of Northern Saurashtra coastal region were ranged from  $1.40$  to  $9.40 \text{ g kg}^{-1}$  having mean value of  $5.14 \text{ g kg}^{-1}$ . In Jamnagar district, maximum soil organic carbon ( $9.00 \text{ g kg}^{-1}$ ) was found at 15 to 20 km distance from the sea coast and minimum organic carbon ( $1.50 \text{ g kg}^{-1}$ ) was found at 0 to 5 km distance from the sea coast, whereas overall mean value of soil organic carbon was  $4.85 \text{ g kg}^{-1}$ . In Devbhumi Dwarka district, maximum soil organic carbon ( $9.70 \text{ g kg}^{-1}$ ) was found at 15 to 20 km distance from the sea coast and minimum soil organic carbon ( $1.40 \text{ g kg}^{-1}$ ) was found at 0 to 5 km distance from the sea coast, while overall mean value of soil organic carbon was  $5.17 \text{ g kg}^{-1}$ . In Porbandar district, overall mean value of organic carbon was  $5.86 \text{ g kg}^{-1}$ , maximum soil organic carbon ( $9.40 \text{ g kg}^{-1}$ ) was found at 15 to 20 km distance from the sea coast and minimum organic carbon ( $2.00 \text{ g kg}^{-1}$ ) was found at 0 to 5 km distance from the sea coast. The lowest value of organic carbon ( $1.40 \text{ g kg}^{-1}$ ) was recorded in the samples collected from Dwarka taluka in Devbhumi Dwarka district, whereas the highest value of organic carbon ( $9.40 \text{ g kg}^{-1}$ ) was found in Porbandar taluka of Porbandar district. The content of organic carbon in soil was noted below to its critical limit upto 10 km distance from sea coast in North Saurashtra which might be due to salinity hazard and shallow light texture.

**Table 4: Talukawise range and mean values of SOC ( $\text{g kg}^{-1}$ ) in different districts of Northern Saurashtra coastal region**

| Distance (km)<br>Talukas    | 0 to 5           |             | 5 to 10          |             | 10 to 15         |             | 15 to 20         |             | Overall          |             |
|-----------------------------|------------------|-------------|------------------|-------------|------------------|-------------|------------------|-------------|------------------|-------------|
|                             | Range            | Mean        | Range            | Mean        | Range            | Mean        | Range            | Mean        | Range            | Mean        |
| Jamnagar                    | 1.90-2.80        | 2.35        | 3.50-4.80        | 4.19        | 5.10-6.90        | 6.07        | 7.20-9.00        | 7.80        | 1.90-9.00        | 5.29        |
| Jodiya                      | 1.80-3.00        | 2.46        | 3.10-4.90        | 4.06        | 5.00-6.10        | 5.55        | 6.40-8.10        | 7.33        | 1.80-8.10        | 4.61        |
| Lalpur                      | 1.50-1.90        | 1.70        | 2.40-4.80        | 3.73        | 4.50-6.60        | 5.60        | 6.20-8.70        | 7.47        | 1.50-8.70        | 4.64        |
| Jamnagar District           | <b>1.50-3.00</b> | <b>2.21</b> | <b>2.40-4.90</b> | <b>4.00</b> | <b>4.50-6.90</b> | <b>5.75</b> | <b>6.20-9.00</b> | <b>7.54</b> | <b>1.50-9.00</b> | <b>4.85</b> |
| Kalyanpur                   | 3.00-4.10        | 3.56        | 4.20-5.40        | 4.84        | 5.30-6.70        | 6.15        | 6.50-8.30        | 7.28        | 3.00-8.30        | 5.40        |
| Khambhalia                  | 1.50-3.10        | 2.35        | 3.70-5.50        | 4.57        | 5.30-6.70        | 5.98        | 6.20-7.60        | 6.90        | 1.50-7.60        | 4.99        |
| Dwarka                      | 1.40-3.60        | 2.44        | 5.20-6.80        | 6.16        | 7.20-8.40        | 7.75        | 9.10-9.70        | 9.40        | 1.40-9.70        | 5.13        |
| Devbhumi<br>Dwarka District | <b>1.40-4.10</b> | <b>2.73</b> | <b>3.70-6.80</b> | <b>5.12</b> | <b>5.30-8.40</b> | <b>6.56</b> | <b>6.20-8.30</b> | <b>7.49</b> | <b>1.40-9.70</b> | <b>5.17</b> |
| Porbandar                   | 2.00-4.70        | 3.48        | 4.90-6.00        | 5.44        | 6.10-7.20        | 6.66        | 7.10-9.40        | 8.34        | 2.00-9.40        | 5.86        |
| Overall                     | <b>1.40-4.70</b> | <b>2.71</b> | <b>2.40-6.80</b> | <b>4.60</b> | <b>4.50-8.40</b> | <b>6.19</b> | <b>6.20-9.40</b> | <b>7.67</b> | <b>1.40-9.40</b> | <b>5.14</b> |



**Fig. 2: Map of overall SOC status in coastal soils of Northern Saurashtra region**

Similar findings were made for the north west agro climatic zone of Gujarat [15], for Hassan district of Karnataka [18], for Degana tahsil, Nagaur district of Rajasthan [19], for



Kishtwar district of Jammu and Kashmir [20], for Veeranam command area of Cuddalore district of Tamil Nadu [21], for Gir Somnath district [22], for Haveri, Gadag and Dharwad districts of Karnataka [23] and for Southern saurashtra of Gujarat [13].

### 3.5 Cation Exchange Capacity (CEC)

The CEC was worked out from the values of exchangeable cations (Ammonium acetate extractable) and are given in table 5. The overall range of CEC in Northern Saurashtra coastal region was 19.49 to 59.72 (cmol (p+) kg<sup>-1</sup>) with the mean value of 36.99 (cmol (p+) kg<sup>-1</sup>). In Jamnagar district, maximum CEC [56.25 (cmol (p+) kg<sup>-1</sup>)] was found at 0 to 5 km distance from the sea coast and minimum CEC [23.29 (cmol (p+) kg<sup>-1</sup>)] was found at 5 to 10 km distance from the sea coast, whereas overall mean value of CEC was 38.82 (cmol (p+) kg<sup>-1</sup>). In Devbhumi Dwarka district, maximum CEC [47.91 (cmol (p+) kg<sup>-1</sup>)] was found at 5 to 10 km distance from the sea coast and minimum CEC [19.49 (cmol (p+) kg<sup>-1</sup>)] was found at 5 to 10 km distance from the sea coast, while overall mean value of CEC was 34.81 (cmol (p+) kg<sup>-1</sup>).

**Table 5: Talukawise range and mean values of CEC (cmol (p+) kg<sup>-1</sup>) in different districts of Northern Saurashtra region**

| Distance (km)<br>Talukas    | 0 to 5                  |              | 5 to 10                 |              | 10 to 15                |              | 15 to 20                |              | Overall                 |              |
|-----------------------------|-------------------------|--------------|-------------------------|--------------|-------------------------|--------------|-------------------------|--------------|-------------------------|--------------|
|                             | Range                   | Mean         | Range                   | Mean         | Range                   | Mean         | Range                   | Mean         | Range                   | Mean         |
| Jamnagar                    | 41.02-<br>42.74         | 41.88        | 23.29-<br>52.83         | 38.42        | 30.14-<br>47.81         | 42.52        | 34.63-<br>42.86         | 38.50        | 23.29-<br>52.83         | 40.01        |
| Jodiya                      | 44.33-<br>56.25         | 48.49        | 32.59-<br>49.96         | 44.70        | 30.93-<br>39.36         | 34.88        | 23.59-<br>28.49         | 26.77        | 23.59-<br>56.25         | 40.10        |
| Lalpur                      | 32.50-<br>44.87         | 39.62        | 30.97-<br>50.54         | 39.87        | 29.27-<br>37.51         | 33.60        | 27.92-<br>33.14         | 31.38        | 27.92-<br>50.54         | 36.36        |
| Jamnagar District           | <b>32.50-<br/>56.25</b> | <b>44.51</b> | <b>23.29-<br/>52.83</b> | <b>40.88</b> | <b>29.27-<br/>47.81</b> | <b>37.05</b> | <b>23.59-<br/>42.86</b> | <b>32.29</b> | <b>23.29-<br/>56.25</b> | <b>38.82</b> |
| Kalyanpur                   | 23.70-<br>45.79         | 35.95        | 35.66-<br>47.91         | 40.04        | 34.67-<br>44.02         | 40.19        | 33.27-<br>43.14         | 39.35        | 23.70-<br>47.91         | 38.92        |
| Khambhalia                  | 32.45-<br>44.58         | 40.21        | 28.47-<br>45.42         | 37.82        | 27.16-<br>35.03         | 31.82        | 26.38-<br>47.13         | 34.54        | 26.38-<br>47.13         | 36.28        |
| Dwarka                      | 19.88-<br>40.73         | 29.37        | 19.49-<br>42.04         | 30.27        | 22.53-<br>42.31         | 30.48        | 23.20-<br>24.01         | 23.61        | 19.49-<br>42.31         | 29.24        |
| Devbhumi<br>Dwarka District | <b>19.88-<br/>45.79</b> | <b>33.61</b> | <b>19.49-<br/>47.91</b> | <b>36.25</b> | <b>22.53-<br/>44.02</b> | <b>35.02</b> | <b>23.20-<br/>47.13</b> | <b>34.30</b> | <b>19.49-<br/>47.91</b> | <b>34.81</b> |
| Porbandar                   | <b>39.64-<br/>48.78</b> | <b>48.78</b> | <b>31.04-<br/>37.58</b> | <b>37.58</b> | <b>29.23-<br/>34.04</b> | <b>34.04</b> | <b>27.06-<br/>29.39</b> | <b>29.39</b> | <b>27.06-<br/>37.99</b> | <b>37.99</b> |

|         |           |      |           |      |           |      |           |      |           |      |
|---------|-----------|------|-----------|------|-----------|------|-----------|------|-----------|------|
|         | 59.72     |      | 42.32     |      | 39.72     |      | 32.11     |      | 59.72     |      |
| Overall | 1.40-4.70 | 2.71 | 2.40-6.80 | 4.60 | 4.50-8.40 | 6.19 | 6.20-9.40 | 7.67 | 1.40-9.40 | 5.14 |

In Porbandar district, overall mean value of CEC was 37.99 (cmol (p+) kg<sup>-1</sup>), maximum CEC [59.72 (cmol (p+) kg<sup>-1</sup>)] was found at 0 to 5 km distance from the sea coast and minimum CEC [27.06 (cmol (p+) kg<sup>-1</sup>)] was found at 15 to 20 km distance from the sea coast.

Similar findings were also recorded for Bhatinda district of Punjab [24], for Sivaganga district of Tamil Nadu [25], for Gir Somnath district [26], for Nagaur district of Rajasthan [12] and for Southern saurashtra of Gujarat [13].

### 3.6 Water soluble ions

Water soluble cations (Ca<sup>++</sup>, Mg<sup>++</sup>, Na<sup>+</sup> and K<sup>+</sup>) and anions (CO<sub>3</sub><sup>-</sup>, HCO<sub>3</sub><sup>-</sup>, Cl<sup>-</sup> and SO<sub>4</sub><sup>-</sup>) were estimated from the soil:water (1:2.5) extract. Sample wise values of water soluble cations and anions for all the 141 samples are given in tables 6 and 7.

**Table 6: Mean values of water soluble cations (me L<sup>-1</sup>) in soils of Northern Saurashtra coastal region**

| Distance (km) | Water soluble cations (me L <sup>-1</sup> ) |                  |                 |                |
|---------------|---|------------------|-----------------|----------------|
|               | Ca <sup>++</sup>                            | Mg <sup>++</sup> | Na <sup>+</sup> | K <sup>+</sup> |
| 0 to 5        | 3.97  | 3.83             | 17.89           | 0.05           |
| 5 to 10       | 1.91  | 1.39             | 7.26            | 0.08           |
| 10 to 15      | 1.33  | 0.86             | 4.89            | 0.18           |
| 15 to 20      | 0.83  | 0.59             | 2.71            | 0.52           |
| Overall       | 2.05  | 1.69             | 8.35            | 0.18           |

**Table 7: Mean values of water soluble anions (me L<sup>-1</sup>) in soils of Northern Saurashtra coastal region**

| Distance (km) | Water soluble anions (me L <sup>-1</sup> ) |                               |                 |                              |
|---------------|--|-------------------------------|-----------------|------------------------------|
|               | CO <sub>3</sub> <sup>-</sup>               | HCO <sub>3</sub> <sup>-</sup> | Cl <sup>-</sup> | SO <sub>4</sub> <sup>-</sup> |
| 0 to 5        | 0.32                                       | 2.96                          | 22.67           | 0.18                         |
| 5 to 10       | 0.26                                       | 1.65                          | 8.61            | 0.06                         |
| 10 to 15      | 0.18                                       | 1.17                          | 5.74            | 0.04                         |
| 15 to 20      | 0.08                                       | 0.79                          | 3.55            | 0.03                         |
| Overall       | 0.22                                       | 1.68                          | 10.30           | 0.08                         |

Among the water soluble cations,  $\text{Na}^+$  was predominant one followed by  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$  and  $\text{K}^+$ . The mean values of water soluble  $\text{Ca}^{++}$ ,  $\text{Mg}^{++}$ ,  $\text{Na}^+$  and  $\text{K}^+$  were 2.05, 1.69, 8.35 and 0.18 me  $\text{L}^{-1}$ , respectively. Among the water soluble anions,  $\text{Cl}^-$  was predominant one followed by  $\text{HCO}_3^-$ ,  $\text{CO}_3^{--}$  and  $\text{SO}_4^{--}$ . The mean values of water soluble  $\text{CO}_3^{--}$ ,  $\text{HCO}_3^-$ ,  $\text{Cl}^-$  and  $\text{SO}_4^{--}$  were 0.22, 1.68, 10.30 and 0.08 me  $\text{L}^{-1}$ , respectively. The results of water soluble ions revealed that  $\text{Na}^+$  and  $\text{Cl}^-$  were predominant cation and anion in soil solution, respectively. Similar findings were also recorded for Sivaganga district of Tamil Nadu [25], for Gir Somnath district [26] and for Southern saurashtra of Gujarat [13].

#### 4. CONCLUSION

On the basis of analyzed data of soil samples, collected from different districts (Jamnagar, Devbhumi Dwarka and Porbandar) of Northern Saurashtra coastal region of Gujarat, it can be concluded that  $\text{EC}_{2.5}$ ,  $\text{CaCO}_3$ , CEC and water soluble ions except  $\text{K}^+$  were decreased, while  $\text{pH}_{2.5}$  and SOC was slightly increased with increasing the distance from sea coast. Soil  $\text{EC}_{2.5}$  was found beyond to its critical or marginal limit up to 0 to 5 km distance from sea coast. The content of organic carbon in soil was noted below to its critical limit upto 10 km distance from sea coast in Northern Saurashtra which might be due to salinity hazard and shallow light texture.

#### REFERENCES

1. Arora S, Singh Y P, Vanza M and Sahni D. Bioremediation of saline and sodic soils through halophilic bacteria to enhance agricultural production. *Journal of Soil and Water Conservation*. 2016;15:302-305.
2. Mandal S, Raju R, Kumar A, Kumar P and Sharma P C. Current status of research, technology response and policy needs of salt-affected soils in India – a review. *Journal of Indian Society of Coastal Agricultural Research*. 2018;36:40-53.
3. Mimura N. Sea-level rise caused by climate change and its implications for society. *Proceedings of the Japan Academy, Ser. B, Physical and Biological Sciences*. 2013;89(7):281–301.
4. Rao G, Kanani A D, Purohit D and Waghela D. Coastal Saline Soils of Gujarat (India): Problems, Reclamation Measures and Management Strategies. In: Dagar J., Yadav R., Sharma P. (eds) Research Developments in Saline Agriculture. Springer, Singapore, 2019;629-651.

5. Verma B L, Sharma Y and Sighania R A. Quality of underground irrigation waters of Charu District in Rajasthan. *Journal of Indian Society of Soil Science*. 2003;**51**(2):214-216.
6. \* Walkley A and Black I A. An examination of methods for determination of organic carbon and nitrogen in soils. *Journal of Agricultural Science*. 1935;**25**: 589-609.
7. \*Chapman H D. Cation exchange capacity, in C.A. Black (ed). Method of soil analysis. *American Society of Agronomy madisson*. Washington, D.C. USA. 1965;P-891.
8. Gandhi G. Characterization, classification and evaluation of soil and water resources of the soils of Girnar toposequence of South Saurashtra region. M. Sc. (Agri.) Thesis (Unpublished). Junagadh Agricultural University, Junagadh, 2013.
9. Patel J M, Patel B T and Patel I M. Fertility status of cultivated soils in Patan district of North Gujarat. *Gujarat Agricultural Universities Research Journal*. 2016;**41**(1): 23-27.
10. Reddy K S and Naidu M V S. Characterization and Classification of Soils in Semi-arid Region of Chennur Mandal in Kadapa District, Andhra Pradesh. *Journal of the Indian Society of Soil Science*. 2016;**64**(3): 207-217.
11. Wagh N S, Mandaland D K and Sadanshiv N S. Available micronutrient status of sunflower growing soils of Nagpur district (Maharashtra). *An Asian Journal of Soil Science*. 2016;**11**(1): 225-229.
12. Singh P, Sharma K K, Legese B and Godana G. Effect of Irrigation Salinity Water on Soil Properties of Nagaur Region, Rajasthan, India. *Annals of Romania Society for Cell Biology*. 2021;**25**(5): 470-476.
13. Bhorania N C, Savalia S G and Sakarvadia H L. Soil salinity Patten along the distance gradient in coastal region soils of southern Saurashtra of Gujarat. *The Pharma Innovation Journal*. 2021;**10**(8): 1753-1759.
14. Nandy T, Prasuna R P and Madhuvani P. Characterization and classification of some coastal soils of Guntur district, Andhra Pradesh. *Journal of the Indian Society of Coastal Agricultural research*. 2013;**31**(1):1-7.
15. Polara K B, Patel M S and Kalynsundram N K. Salt affected soils of north-west agroclimatic zone of Gujarat: Their characterization and categorization. *Journal of the Indian Society of Coastal Agricultural Research*. 2006;**26**(1): 52-55.

16. Rajput S G and Polara K B. Fertility status of cultivated soils in coastal Bhavnagar district of Saurashtra region of Gujarat. *Journal of the Indian Society of Soil Science*. 2012;**60**(4): 317-320.
17. Gajare A S, Dhawan A S, Ghodke S K and Bhor S D. Available sulphur and phosphorus status of soybean growing soils of Latur district. *An Asian Journal of Soil Science*, 2013;**8**(1): 94-97.
18. Punithraj T S, Nagaraja M S, Dhumgond P, Bhoopal S and Shivakumar KM. Soil fertility status of tomato (*Lycopersicon esculentum*, Mill) grown in areas of Hassan district, Karnataka. *An Asian Journal of Soil Science*. 2012;**7**(2): 288-291.
19. Singh B. Appraisal of fertility indices of the soils of Degana tahsil, Nagaur district (Rajasthan). *An Asian Journal of Soil Science*. 2014;**9**(1): 100-102.
20. Tundup P and Akbar A. Distribution of micronutrient cations and their relationship with soil properties of saffron growing soils of district Kishtwar in Jammu and Kashmir. *An Asian Journal of Soil Science*. 2014;**9**(1): 59-62.
21. Arunkumar V and Paramasivan M. Spatial variability and geostatistics application for mapping of soil properties and nutrients in intensively cultivated village of Veeranam Command Area, Tamil Nadu. *An Asian Journal of Soil Science*. 2015;**10**(2): 299-305.
22. Hadiyal S T, Rathod A D and Rajawat B S. Fertility status of soils of Girgadhda and Una taluka of Gir Somnath district. *An Asian Journal of Soil Science*. 2016;**11**(1): 184-186.
23. Nagaral I N, Kuligod V B and Singh V P. Soil nutrient status of chilli growing area of Northern Transitional Zone of Karnataka. *An Asian Journal of Soil Science*. 2016;**11**(1): 140-145.
24. Singh K B and Sharma B D. Morphological, physical and chemical properties of arid soils of Bhatinda district of Punjab. *An Asian Journal of Soil Science*. 2013;**8**(1): 48-52.
25. Malavath R and Mani S. Nutrients status in the surface and subsurface soils of dryland Agricultural Research Station at Chettinad in Sivaganga district of Tamil Nadu. *An Asian Journal of Soil Science*. 2014;**9**(2): 169-175.
26. Chauhan R B and Polara J V. Characterization and classification of cultivated soils of coastal Gir Somnath district of Gujarat in relation to salinity. *Journal of the Indian Society of Coastal Agricultural Research*. 2015a;**33**(2): 12-15.

**NOTE:** \* Original not seen