

## Original Research Article

### **“Response of different pre-sowing seed treatments with Biofertilizers, Biostimulants and Botanicals on growth, yield and yield attributing traits in chickpea (*Cicer arietinum* L.) var. desi himmat”**

#### **ABSTRACT**

The present study entitled, “Response of different pre-sowing seed treatments with Biofertilizers, Biostimulants and Botanicals on growth, yield and yield attributing traits in chickpea (*Cicer arietinum* L.) var. desi himmat”, was carried out in rabi 2021-2022 at crop research farm, SHUATS, Prayagraj with rhizobium, vermiwash, neem and tulsi leaf extracts of varying concentrates at duration of 12hrs. with an objective to assess the effect of these priming treatments on chickpea and to find the promising pre-sowing treatment. The results depicted that seed priming showed significant variation for growth and yield characters that were recorded and performed well on comparison to the untreated control (T<sub>0</sub>). Seed priming with T<sub>5</sub>-Vermiwash at 10% for 12 hrs. recorded field emergence of 90% and 35.00 pods per plant, 1.66 seeds per pod, 14.70 g seed yield per plant, 24.79 g of 100 seed weight, 37.65 g of biological yield per plant and 39.08 % of harvest index. Thus, priming with T<sub>5</sub>-Vermiwash at 10% for 12 hrs. was found so productive on chickpea and can be suggested for cultivation on commercial scale.

**Keywords:** Neem leaf extract, rhizobium, tulsi leaf extract and vermiwash.

#### **INTRODUCTION**

Legumes are the richest protein source of human diet and livestock in poor areas besides they are used as green manures and green fodder to animals. Chickpea (*Cicer arietinum* L.) is the most important Rabi season self-pollinated pulse crop, with the chromosomal number ( $2n = 2x = 16$ ) and it is the third most widely cultivated legume crop after dry bean (*Phaseolus vulgaris* L.) and field pea (*Pisum sativum* L.) belonging to the family Fabaceae.

The chick pea production in India has gone up from 38.55 to 122.29 lakh tonnes during 2000-01 to 2018-19, while the area has also gone up from 53.85 to 109.21 lakh ha, whereas, the yield has steadily increased from 744 kg/ha to 1077kg/ha during the same period. In India, Madhya Pradesh holds the highest share of 39% chickpea production followed by Maharastra (14%), Rajasthan (14%), Andhra Pradesh (10%), Uttar Pradesh (7%) and Karnataka (6%) (**Preeti varma et al., 2021**). Seed is the main edible part of the plant and is a rich source of protein (23.3- 28.9%), carbohydrates (61.5%), fats (4.5%) and minerals (phosphorus, calcium, magnesium, iron, zinc).

Seed enhancement includes priming, hardening, pelleting, encrusting, film coating *etc.*, but excludes the treatment for the control of seed-borne pathogens. Organic seed priming *viz., treatment* with Panchagavya, Beejamrutha, Jeevamrutha and botanicals provides resistant to high temperature and low moisture especially in semi-arid tropics. It encourages quicker germination, greater seedling vigour resulting in increased crop productivity. Priming helps the plants to cope with the adverse effects of unfavorable environmental conditions (**Muhammad Moaaz Ali *et al.*, 2020**).

Botanical seed treatment is extracted from naturally occurring sources based on the botanical ingredients and have synergistic effect on early and uniform seed germination and enhances tolerance to the pest and diseases during the early crop growth stages. *Ocimum sanctum* is commonly known as tulsi or Holy basil comprises essential bioactive or phytochemical components such as oleanolic acid, rosmarinic, ursolic acid, eugenol, carvacrol, linalool, and  $\beta$ -caryophyllene essential for plant growth and yield. *Azadirachtin* is the main pesticidal component of the neem. Neem products are naturally available materials, cheaper and also safe for microorganisms; its usage in agriculture is increasing extensively because of beneficial effects on plants (**Nelakurthi Venkata Praveen *et al.*, 2020**).

Earthworms play a vital role in plant growth and productivity. It contains several enzymes, vitamins, auxins and cytokinins along with nutrients phosphorous, calcium and potassium that enhances resistivity of crops and increases productivity (**Suchit Tiwari *et al.*, 2018**). It plays an important role in growth of plants and its development contribute to initiation of rooting, root growth, plant development, promotion of growth rate of the plants.

Application of PGPR to seed through seed bio-priming enhances plant performance under stress environments and consequently enhances plant yield both directly and indirectly. Helpful bacteria in rhizosphere are of two types: (a) bacteria forming symbiotic association through particular assemblies and (b) free-living bacteria present in the neighbourhood of plant province which are mainly known as plant growth-promoting rhizobacteria (PGPR). **Belal *et al.*, (2013)** reported that strains of rhizobia are capable of tolerating abiotic factors like salinity and drought by preserving its capability to nodulate and fix atmospheric nitrogen.

## MATERIALS AND METHODS

The present study was conducted during Rabi 2021-2022 at Crop Research Farm, Naini Agriculture Institute, SHUATS, Prayagraj. Thirteen treatment combinations of rhizobium, vermiwash, neem and tulsi leaf extracts of varying concentrations viz., rhizobium at 8%, 10% and 15% for 12 hrs., vermiwash at 8%, 10% and 15% for 12 hrs., neem leaf extract at 5%, 10% and 15% for 12 hrs., and tulsi leaf extract at 5%, 10% and 15% for 12 hrs. were prepared and treated with seeds of desi himmat chickpea variety for a duration of 15 mins.

## RESULTS AND DISCUSSION

The results on the effect of rhizobium, vermiwash, neem and tulsi extract solutions on plant growth and yield attributes were represented in Table 01.

### Field emergence (%)

Seed priming with Vermiwash at 10% for 12 hrs. (T<sub>5</sub>) recorded maximum emergence of 20.00% on 4<sup>th</sup> day after sowing; The treatment T<sub>7</sub>- Neem leaf extract at 5% for 12 hrs. significantly recorded maximum of 54.00% and 91.00% at 7 DAS and 10 DAS. The untreated control (T<sub>0</sub>) recorded least emergence of 13.00%, 42.00% and 84.00% at 4, 7 and 10 DAS respectively. According to (K. Sundararasuet *et al.*, 2016) vermiwash contains nutrients like auxins, cytokinins, bacteria, fungi and nutrients like calcium, phosphorous that enhance germination. Similar trends of increased emergence % due to organic priming were in harmony with the earlier research findings of (Pandit S. Rathodeet *et al.*, 2016).

### Plant height (cm)

Effect of the treatment T<sub>2</sub>- Rhizobium at 10% for 12 hrs. was found significant and recorded 23.11 cm plant height at 30 DAS, Priming with Vermiwash at 10% for 12 hrs. (T<sub>5</sub>) shown maximum of 37.32 cm and 71.16 cm height at 60 DAS and at time of harvest. The untreated control (T<sub>0</sub>) recorded the minimum height of 16.29 cm, 27.99 cm and 62.39 cm at 30 DAS, 60 DAS and at harvest respectively. According to (Tripathi and Bhardwaj 2014) vermiwash contains nitrogen in the form of mucus and nitrogen excretory substances that stimulates hormones and enzymes for plant growth. These results are comparable with the earlier research findings of (H.G. Harshitha *et al.*, 2021)

### Days to 50% flowering

Minimum number of 63.33 days was taken by treatment T<sub>2</sub>-Rhizobium at 10% for 12 hrs. for which in at least 50% of flowers get opened while priming with T<sub>6</sub>- Vermiwash at 15% for 12 hrs. took 70.67 days

for the same. Similar trends for days to 50% flowering were found by earlier research works of (**S. Jaffar Basha *et al.*, 2021**).

### **Days to 50% pod setting**

The least number of days for 50% pod setting was shown by the treatment T<sub>2</sub>-Rhizobium at 8% for 10 hrs. of 71.33 days and treatment T<sub>6</sub>-Vermiwash at 15% for 8 hrs. took 78.33 days for 50% pod setting which was maximum amongst the treatments. The days taken for 50% pod setting are in compatible with the earlier findings of (**Dan Singh Jakkar *et al.*, 2016**).

### **Days to maturity**

Priming with T<sub>8</sub>- Neem leaf extract at 10% for 12 hrs. and T<sub>11</sub>- Tulsi leaf extract at 10% for 12 hrs. recorded the minimum days for maturity of 106.33 days and 106.67 days respectively; T<sub>6</sub>-Vermiwash at 15% for 12hrs. recorded 113.00 days which was maximum amongst treatments. Vermiwash being enriched with NPK, micro nutrients, plant growth hormones, microbes and enzymes enhances the growth and thereby shows effect on maturity (**Vijayantie R.R. Awadhpersadet *et al.*, 2021**). These results are in agreement with earlier findings of (**Dikhshatkumaret *et al.*, 2020**).

### **Number of pods per plant**

Effect of priming with T<sub>5</sub>-Vermiwash at 10% for 12 hrs. significantly recorded maximum of 35.76 pods per plant amongst treatments and untreated control (T<sub>0</sub>) shown minimum of 25.20 pods plant<sup>-1</sup> respectively. Timely supply of nutrients with effect of vermiwash enhances the yield parameters *viz.*, number of pods, pod length and seed-bearing capacity of the pods (**Jijo George *et al.*, 2019**).

### **Number of seeds per pod**

Soaking of chickpea seeds with Vermiwash at 10% at 12 hrs. (T<sub>5</sub>) shown maximum of 1.66 seeds in each pod and the untreated control shown average of 1.33 seeds in each single pod respectively. These results concur with the previous research findings of (**Mebrahtu Gebremariumet *et al.*, 2021**).

### **Seed yield per plant (g)**

Seed priming with T<sub>5</sub>-Vermiwash at 10% for 12 hrs. significantly recorded the maximum of 14.70g of chickpea seeds per plant and minimum yield of 7.08 g was recorded by the untreated control (T<sub>0</sub>) respectively. Organic priming enhances the growth and yield of the pulses; which are highly sensitive to the salinity and alkalinity factors through timely supply of nutrients and proper hormonal balance (**Jalla Manjunath *et al.*, 2021**).

### **Seed index (g)**

Significantly, the maximum hundred seed weight of 24.79 g was recorded with T<sub>5</sub>- Vermiwash at 10% for 12 hrs. while the minimum of 19.50 g was shown by untreated (T<sub>0</sub>) Control. Similar trends of 100 seed weight were recorded by earlier research works of (**Dikhshatkumaret al., 2020**).

### **Biological yield per plant (g)**

Seed priming enhanced the biological yield per plant amongst the treatments; the maximum was recorded of 37.65 g was with T<sub>5</sub>- Vermiwash at 10% for 12 hrs. while the minimum of 22.15 g was shown by the treatment T<sub>0</sub>- Control. Seed priming enhances the metabolic activities of the seed and resistance against to the diseases and pests, hence increases the dry matter content of the plants (**Arjun Sharma et al., 2006**).

### **Harvest index (%)**

Maximum harvest index of 39.08% was recorded by priming with T<sub>5</sub>- Vermiwash at 10% for 12 hrs. and minimum of 32.02% was shown by the untreated control (T<sub>0</sub>) respectively. These results were in conformity with the earlier research findings of (**S. Jaffar Basha et al., 2018**).

## **CONCLUSION**

The results depicted that priming with organics, botanicals and biostimulants shown significant variation with the untreated control for all the growth and yield parameters that were recorded and further analyzed. Effect of the treatment T<sub>5</sub>-Vermiwash at 10% for 12 hrs. found to be promising with 14.70 g seed yield per plant, 24.79 g of 100 seed weight, 37.65 g of biological yield per plant and 39.08 % of harvest index and can be recommended for cultivation on commercial scale.

Table 01. Response of Chickpea *var.* desi himmat to rhizobium, vermiwash, neem and tulasi leaf extract solutions.

TREATMENTS	FIELD EMERGENCE (%) AT 4 DAS	FIELD EMERGENCE (%) AT 7 DAS	FIELD EMERGENCE (%) AT 10 DAS	PLANT HEIGHT AT 30 DAS (cm)	PLANT HEIGHT AT 60 DAS (cm)	PLANT HEIGHT AT MATURITY (cm)	DAYS TO 50% FLOWERING	DAYS TO 50% POD SETTING
T <sub>0</sub>	13	42	84	16.29	28.11	62.39	65.00	73.00
T <sub>1</sub>	16	46	86	19.68	31.44	64.92	67.00	75.00
T <sub>2</sub>	17	48	87	23.11	36.77	67.24	63.33	71.33
T <sub>3</sub>	15	45	84	17.22	30.75	65.99	67.67	74.67
T <sub>4</sub>	16	46	83	18.10	33.61	68.38	66.33	73.00
T <sub>5</sub>	20	52	87	22.56	37.32	71.16	68.67	75.33
T <sub>6</sub>	17	47	83	21.16	35.18	69.01	70.67	78.33
T <sub>7</sub>	19	54	91	22.33	36.78	69.43	66.33	74.67
T <sub>8</sub>	18	48	86	16.84	27.99	63.98	63.67	74.00
T <sub>9</sub>	19	49	87	20.69	32.22	66.26	64.67	72.00
T <sub>10</sub>	17	49	87	20.97	31.38	66.66	66.33	72.67
T <sub>11</sub>	20	52	89	21.29	32.96	68.35	63.33	71.67
T <sub>12</sub>	17	47	85	19.89	31.26	66.43	67.00	73.67
<b>GRAND MEAN</b>	17	48	86	20.01	32.75	66.94	66.15	73.79
<b>F TEST</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>
<b>S. Em</b>	0.73	0.84	0.53	0.27	0.37	0.90	0.88	0.89
<b>S. Ed</b>	1.03	1.20	0.76	0.39	0.52	1.27	1.24	1.25
<b>C.V.</b>	7.36	3.05	1.08	2.39	1.96	2.33	2.31	2.09
<b>T. value</b>	2.06	2.06	2.06	2.06	2.06	2.06	2.06	2.06
<b>C.D.</b>	2.13	2.47	1.56	0.80	1.08	2.63	2.57	2.59

Cont..

TREATMENTS	DAYS TO MATURITY	NUMBER OF PODS PER PLANT	NUMBER OF SEEDS PER POD	SEED YIELD PER PLANT (g)	SEED INDEX (g)	BIOLOGICAL YIELD (g/plant)	HARVEST INDEX (%)
T <sub>0</sub>	109.33	25.00	1.44	7.08	19.50	22.15	32.02
T <sub>1</sub>	107.33	28.00	1.48	8.53	20.28	25.80	33.05
T <sub>2</sub>	108.33	31.00	1.31	9.78	23.85	27.79	35.20
T <sub>3</sub>	108.00	28.00	1.35	9.05	23.23	27.42	33.00
T <sub>4</sub>	109.33	30.00	1.37	9.53	22.86	31.45	30.29
T <sub>5</sub>	109.67	35.00	1.66	14.70	24.79	37.65	39.08
T <sub>6</sub>	113.00	32.00	1.48	11.28	23.72	32.03	35.22
T <sub>7</sub>	110.67	35.00	1.60	13.79	24.55	36.08	38.22
T <sub>8</sub>	106.33	27.00	1.41	9.02	22.84	26.55	33.98
T <sub>9</sub>	109.67	28.00	1.44	9.36	23.11	31.48	29.72
T <sub>10</sub>	108.67	31.00	1.33	8.99	21.33	27.31	32.97
T <sub>11</sub>	106.67	32.00	1.39	9.91	22.06	26.94	36.79
T <sub>12</sub>	109.67	30.00	1.31	8.31	20.77	25.51	32.54
GRAND MEAN	108.97	32.66	1.43	9.95	22.53	29.09	34.01
F TEST	S	S	S	S	S	S	S
S. Em	0.88	0.42	0.03	0.27	0.23	0.64	0.71
S. Ed	1.25	0.59	0.05	0.38	0.33	0.90	1.01
C.V.	1.41	2.38	4.58	4.76	1.83	3.82	3.66
T. value	2.06	2.06	2.06	2.06	2.06	2.06	2.06
C.D.	2.58	1.23	0.11	0.79	0.69	1.87	2.09

## REFERENCES

- K. Sundararasu (2016).** Effect of vermiwash on growth and yielding pattern of selected vegetable crop Chilli (*Capsicum annum* L.). *International journal of Advanced Research on biological science*. 3 (9): 155-160.
- Pandit S. Rathode., S.B. Bellad., D.H. Patil and B.M. Doddamani (2016).** Effect of seed priming on growth and productivity of chickpea (*Cicer arietinum* L.) under rainfed conditions of Karnataka. *The Bioscan*. 11(4): 2695-2698.
- H. G. Harshitha., Abhinav Dayal., Prashanth Kumar Rai and Neha Thomas (2021).** Pre-sowing seed treatment with organic and inorganic treatments on growth, yield and yield attributes of chickpea (*Cicer arietinum* L.) variety Pusa-32. *International Journal of Plant and soil science*. 33 (20):71-78.
- S.Jaffar Basha., V. Jaya Lakshmi., S. KhayumAhmadh and N. Kamakshi(2018).** Studies on growth and yield characters of chickpea (*Cicer arietinum* L.) varieties suitable for mechanical harvesting. *Tropical Plant Research*. 7(3): 634-637.
- Dikshat Kumar., Sunil Kumar., and Harbhinder Singh (2020).** Growth and yield of Chickpea (*Cicer arietinum* L.) as influenced by seed rate and seed priming. *International Journal of Current microbiology and applied sciences*. 9(10): 1011-1016.
- Jijo George., Deepthi, M.P., Rini Joseph., and Kathireswari, P. (2019).** Beneficial utilization of vermiwash in sustainable organic priming. *International Journal of Recent scientific research*. 10 (10): 35403-35406.
- JallaMnjunath., Abhinav Dayal., Sasya Nagar and Prashanth Kumar Rai. (2021).** Influence of bio-fertilizer and organic seed treatment on growth and yield attributing traits of Mustard (*Brassica nigra* L.) variety Pusa 21.
- Dan singhJakhar., M. S. Kamble., Abishek singh and Saket Kumar (2016).** Performance of chickpea (*Cicer arietinum* L.) genotypes in Kohlapur region. *International Journal of Agriculture, Environment and Biotechnology*. 9 (2): 283-289.
- MebratuGebremariam and Teklay Tesfay (2021).**Effect of P application rate and rhizobium inoculation on nodulation, growth and yield performance of chickpea (*Cicer arietinum* L.). *International Journal of Agronomy*. 2021. 01-14.
- Arjun Sharma and Deshpande V. K. (2006).** Effect of pre-soaking of pigeonpea seeds with organics on seed quality. *Karnataka Journal of Agricultural Sciences*. 19: 396-399.



**Preeti varma., Rajesh Kumar., R. K. Solanki, Chaman Jodan and Pradeep Kumar (2021).** Chickpea(*Cicer arietinum*L.)scenario in India and south eastern Rajasthan. *International Journal of Current microbiology and applied sciences*.10 (1): 1057-1067.

**Subha, S., Rehman G. L., Usha, O. P. and Sirisha, A. (2019).** Effect of seed treatment with rhizobia on plant growth, seed yield and seed quality parameters in Pigeonpea (*Cajanus cajan*L.). *International Journal of current microbiology and Applied sciences*. 24 (3): 564-56.

**Muhammad Mooaz Ali., Talha Jawad., Rosario Paolo Mauro., Rubab Shabir., Irfan Afzal and AhmadhFathey Yousuf. (2020).** Effect of seed priming with Potassium Nitrate on the performance of Tomato. 10 (11):498; <https://doi.org/10.3390/agriculture> 10110498.

**Nelakurthi Venkata Praveen., Prashanth Kumar Rai., Rupesh Kumar and Ruksana. (2020).** Effect of priming on growth, seedling, yield and its attributing characters of desi chickpea (*Cicer arietinum* L.).*International Journal of current microbiology and Applied sciences*.9 (9): 3162-3171.

**Suchith Tiwari., A. K. Chaurasia., Nithyananda N and Bineeta Bara.** Effect of organic priming on seed germination behaviour and vigour of chickpea. (*Cicer arietinum* L.). *Journal of Pharmacognosy and phytochemistry*. 7 (4): 1064-1067.

**Belal, E. B., Hassan, M. M., El- Rammady, H. R. (2013).**Phylogenic and charecterization of salt tolerant rhizobial strain nodulatingfaba bean plants. *African Journal of biotechnology*. 12: 4324-4337.