

Original Research Article

Mycobiota of *Cnidoscolus aconitifolius* (Mill.) I.

M. Johnston Phyllosphere

ABSTRACT

Aims: The phyllosphere, which is the above-ground (aerial) part of plants, is colonized by different microorganisms some of which may be pathogenic to plants and also to humans and animals.

Methodology: This study was conducted to determine the mycobiota of *Cnidoscolus aconitifolius* phyllosphere using metagenomics. The mycobiota was determined by sequencing the 18S rRNA gene on Illumina MiSeq platform.

Results: A total of 107 Operational Taxonomic Units (OTUs) were obtained. The mycobiota of *C. aconitifolius* had 100% Ascomycota classified into Dothideomycetes (84.15%), Eurotiomycetes (2.26%) and Sordariomycetes (12.45%). Only 1.13% of the fungi were unassigned at the class level. The core mycobiota of chaya consisted of the genera *Cladosporium* (51.70%), *Lasiodiplodia* (18.11%), *Allophoma* (6.79%), *Stagonosporosis* (2.26%) and *Aspergillus* (2.26%).

Conclusion: The economic importance of the organisms obtained were highlighted. The result from this study shows that *C. aconitifolius* phyllosphere harbors diverse fungi some of which may promote plant growth or are pathogenic to plants and/or humans.

Keywords: Fungi, Illumina next-generation sequencing, Mycobiota, Phyllosphere,

1. INTRODUCTION

Cnidoscolus aconitifolius (Mill.) I. M. Johnston commonly called “chaya” or “tree spinach” is an economic plant that has both food and medicinal uses. *C. aconitifolius* leaves are used as vegetable to cook soup and yam portage in Nigeria and other parts of the world. The leaves have been reported to contain more nutrients than every other land-based leafy vegetable by two to three folds [1]. It is eaten in Mexico as a vegetable and also used as feed for domestic animals. It also has the ability to darken grey hair and strengthen fingernails. In Nigeria, it is used locally as a blood-booster in the rural areas especially in the Southern part of the country. Hamid *et al.* [2] reported that the above-ground parts of *Cnidoscolus aconitifolius* exhibit antibacterial and antifungal activities. Mordi and Akanji [3] also reported that *C. aconitifolius* have antihaemorrhagic, antihypertensive and cardiac depressant properties.

The phyllosphere is a microbial habitat that exists in the aerial parts of vascular plants. It is regarded as a hostile environment for the survival of microorganisms as microorganisms in this sphere need to have the ability to tolerate high ultraviolet (UV) radiations, rapid variations in humidity, temperature and heterogeneous availability of nutrients [4,5]. The phyllosphere is largely colonized through the migration of bacteria, fungi, and other microorganisms from soil, seed, air, water, or through animal-borne sources [5]. Some plant pathogens can inhabit the phyllosphere prior to an infection or in the apparent absence of an infection [5]. Filamentous fungus population in the phyllosphere ranges between 10^2 and 10^8 CFU g⁻¹ leaf [6].

Understanding the modalities of the structure of microbial communities in the phyllosphere is vital for developing bio-control strategies for both plant and human pathogens [7]. Phyllosphere microbiota plays an important role in promoting plant growth through various mechanisms and also protecting plants from various diseases [8]. Microbial communities of the phyllosphere are believed to play a significant role in remediating atmospheric hydrocarbon pollutants and residual pesticides [9,10]. Phyllosphere fungi can increase drought tolerance of plants and confer protection against plant pathogens [11,12]. Leafy vegetables significantly inhibit human pathogens in the phyllosphere which may lead to food-borne diseases [13]. It has been established that host genotype, site characteristics and seasonal changes are some of the factors that determine the structure of microbial communities in the phyllosphere [14,15]. Cordier *et al.* [16] investigated the variations in fungal communities of *Fagus sylvatica* (European beech) leaf surfaces using ITS1 gene sequencing and observed a relationship between the genetic distance of beech trees and differences in fungal community structure, signifying that host genetics determines the fungal community structure on beech leaves. Plant genotype has also been identified as a determinant of fungal community structure on balsam poplar (*Populus balsamifera* L.) phyllosphere using ITS pyrosequencing [17].

Earlier studies have employed the use of culture-dependent method to study and describe community structure and function. These methods are classified as low-through-put molecular techniques as they are believed to underestimate the diversity of microorganisms. All taxa present in a sample cannot be fully represented or identified using these methods. Only culturable members of the microbial communities are represented while majority of the organisms in the habitat are left out. The advent of high-throughput molecular technologies which are also cost-effective has led to remarkable improvements in the field of phyllosphere microbiology as researchers have been able to analyse large number of samples at ease with in-depth coverage of phyllosphere microbiota present [8]. There is more information on the structure and composition of bacteria in the phyllosphere than there is for fungi [8,18]. In this study, we conducted ITS gene-based sequencing on Illumina Miseq platform to determine the mycobiota of *Cnidioscolus aconitifolius* phyllosphere. The results from this study would widen our knowledge on the fungal community associated with *Cnidioscolus aconitifolius* phyllosphere.

2. MATERIAL AND METHODS

2.1 Sample Collection

Cnidioscolus aconitifolius leaves were obtained from Choba, Rivers State, Nigeria in April 2018. The coordinates of the sample collection location is 4.89°N and 6.91°E. The leaves were transferred to the lab in a zip bag prior to DNA extraction.

2.2 DNA Extraction and Illumina Next Generation Sequencing

DNA was extracted from *C. aconitifolius* leaves using Zymo Fungal/Bacterial DNA Extraction kit (Zymo Research Group, California, USA) with a slight modification as described in a previous study [19]. To analyze the total fungal community of the phyllosphere, 0.50g of leaves was used.

The forward primer, ITS1F (5'-CTTGGTCATTTAGAGGAAGTAAT-3') and reverse primer, ITS4 (5'-TCCTCCGCTTATTGACATGS-3') were used to target the ITS region 1 between

the 18S and 5.8S rDNAs, ITS region II and a portion of 28S rDNA. The samples were analyzed with 300 bp paired-end read, Illumina MiSeq, at Inqaba Biotechnology Limited, South Africa. The resulting amplicon was gel purified, end repaired and Illumina specific adapter sequence added to the 5' end of each primer.

2.3 Processing of Sequence Reads

The reads obtained were preprocessed to check sequencing errors. Sequences that did not contain the exact match for both forward and reverse reactions were eliminated from the analysis. Sequences were trimmed with Next-generation sequencing Short Reads (*ngsShoRT*) trimmer as described by [20]. The ITS gene sequences were processed using QIIME v.1.9.0 (Quantitative Insights Into Microbial Ecology) pipeline as described by Caporaso *et al.* [21]. Sequences with less than 200 base pairs and reads with more than 2% of ambiguities were excluded from the final analysis. The UCLUST algorithm [22] was used to cluster sequences into operational taxonomic units (OTUs) at a 97% identity threshold. Each OTU sequence was represented by the most abundant read. The UNITE reference database [23] was used for both open reference OTU picking and taxonomic assignment for the sequences. Raw sequences of *Cnidoscolus aconitifolius* microbiota were deposited on NCBI (National Centre for Biotechnology Information) database under Sequence Read Archive (SRA) in GenBank as BioProject ID PRJNA592288.

3. RESULTS

3.1 *Cnidoscolus aconitifolius* Mycobiota at the Division and Class Levels

Sequences were assembled and a total of One hundred and seven (107) OTUs were successfully characterized and grouped into twenty-eight (28) taxa (genera). The fungal microbiome of *C. aconitifolius* had 100% Ascomycota classified into Dothideomycetes (84.15%), Eurotiomycetes (2.26%) and Sordariomycetes (12.45%). Only 1.13% of the sequences were unassigned at the class level. The fungal community of *C. aconitifolius* at the division and class levels is presented in Figures 1 and 2.




Figure 1: Fungal division obtained from *Cnidoscopus aconitifolius* phyllosphere




Figure 2: Fungal classes obtained from *Cnidoscopus aconitifolius* phyllosphere

3.2 Distribution of Fungi at the Genus and Species Levels

The most represented sequences (each representing more than 1% of the total classified fungi) in *C. aconitifolius* phyllosphere belonged to the families Cladosporiaceae (52.83%), Botryosphaeriaceae (18.11%) and Didymellaceae (9.06%).

Out of the 107 OTUs obtained, only thirty-five (35) belonging to eight (8) taxa were successfully identified to the species level on UNITE database. The other remaining OTUs

were blasted on NCBI database for species identification. This is because NCBI is a constantly updated “gene-house” database as thousands of sequences are deposited on GenBank on a daily bases. The BLAST searches revealed the match sequences of the clones against known sequences on NCBI with 86 to 100% identity. The OTU number and GenBank accession number of match sequences are listed in Table 1.

UNDER PEER REVIEW

Table 1: Taxonomic affinities of OTUs with BLAST searches from NCBI Database based on their ITS sequences

OTU number	Taxonomic affinity (GeneBank Accession no.)	Percentage Similarity (%)
1	<i>Penicillium citrinum</i> (MF476066.1)	96
11	<i>Lasiodiplodia theobromae</i> (GQ469915.1)	100
27	<i>Aspergillus nomius</i> (MK841463.1)	99
39	<i>Phoma eupyrena</i> (KY765281.1)	89
42	<i>Lasiodiplodia theobromae</i> (MH251950.1)	99
50	<i>Penicillium citrinum</i> (MK852473.1)	100
57	<i>Allophoma minor</i> (MF380953.1)	100
70	<i>Phoma eupyrena</i> (KX610328.1)	99
84	<i>Periconia pseudobyssoides</i> (KU214550.1)	99
108	<i>Cladosporium cladosporioides</i> (KU182497.1)	90
112	<i>Acremonium charticola</i> (KT878345.1)	95
114	<i>Aspergillus flavus</i> (MG976497.1)	98
118	<i>Corynespora asiicola</i> (AY238605.1)	99
119	<i>Lasiodiplodia theobromae</i> (GQ469915.1)	97
127	<i>Cladosporium cladosporioides</i> (MH535968.1)	98
137	<i>Cladosporium tenuissimum</i> (MF473305.1)	89
138	<i>Aspergillus flavus</i> (MG976497.1)	100
142	<i>Lasiodiplodia theobromae</i> (GQ469915.1)	100
147	<i>Corynespora asiicola</i> (AY238605.1)	96
159	<i>Nigrospora sphaerica</i> (JQ936184.1)	96
162	<i>Periconia byssoides</i> (KC954157.1)	99
164	<i>Lasiodiplodia theobromae</i> (MH251950)	98
185	<i>Allophoma minor</i> (MF380953.1)	97
193	<i>Corynespora asiicola</i> (AY238605.1)	97

OTU number	Taxonomic affinity (GeneBank Accession no.)	Percentage Similarity (%)
214	<i>Lasiodiplodia theobromae</i> (MH251950.1)	100
217	<i>Cladosporium cladosporioides</i> (MH535968.1)	98
224	<i>Postia placenta</i> (KJ995944.1)	91
227	<i>Aspergillus violaceofuscus</i> (MG682503.1)	99
232	<i>Cladosporium xanthochromaticum</i> (MF473323.1)	97
235	<i>Lasiodiplodia theobromae</i> (EJ904912.1)	89
245	<i>Corynespora casiicola</i> (MH864416.1)	93
254	<i>Nigrospora oryzae</i> (MH619723.1)	87
259	<i>Corynespora casiicola</i> (AY238605.1)	89
268	<i>Colletotrichum gleosporioides</i> (MH392749)	100
272	<i>Corynespora casiicola</i> (AY238605.1)	98
274	<i>Schizothyrium pomi</i> (EF134949.1)	98
283	<i>Cladosporium cladosporioides</i> (MH535968.1)	100
287	<i>Aspergillus versicolor</i> (JQ717322.1)	99
324	<i>Corynespora casiicola</i> (MK685154.1)	99
325	<i>Leptosphaerulina cartarum</i> (KC879283.1)	99
326	<i>Helminthosporium asterinum</i> (MH178554.1)	99
327	<i>Cladosporium cladosporioides</i> (MH790419.1)	100
335	<i>Periconia pseudobyssoides</i> (KU214550.1)	99
338	<i>Lasiodiplodia theobromae</i> (GQ469915.1)	97
343	<i>Schizothyrium pomi</i> (EF134949.1)	98
345	<i>Lasiodiplodia theobromae</i> (MH251950.1)	99
349	<i>Cladosporium colombiae</i> (MH244376.1)	99
364	<i>Corynespora casiicola</i> (MK685154.1)	100
366	<i>Corynespora casiicola</i> (AY238605.1)	97

OTU number	Taxonomic affinity (GeneBank Accession no.)	Percentage Similarity (%)
367	<i>Zygosporium oscheoides</i> (MH861194.1)	96
368	<i>Devriesia Lagerstroemiae</i> (KP197670.1)	94
369	<i>Cladosporium tenuissimum</i> (MG569541.1)	100
394	<i>Aspergillus amstelodami</i> (MK267406.1)	99
395	<i>Pithomyces chartarum</i> (MH859914.1)	99
410	<i>Corynespora casiicola</i> (MK685154.1)	100
420	<i>Allophoma minor</i> (MN380953.1)	100
428	<i>Acremonium charticola</i> (KT878345.1)	94
431	<i>Lasiodiplodia pseudotheobromae</i> (FT904912.2)	89
434	<i>Aspergillus penicillioides</i> (Mh86439.1)	100
439	<i>Colletotrichum siamense</i> (MK471371.1)	100
442	<i>Nigrospora oryzae</i> (MK429852.1)	100
443	<i>Penicillium paxilli</i> (JN617709.1)	98
454	<i>Trichoderma harzianum</i> (MN046978.1)	100
457	<i>Helminthosporium asterinum</i> (AF073918.1)	96
475	<i>Rombousta ilealis</i> (LN555523.1)	99
479	<i>Lasiodiplodia pseudotheobromae</i> (FJ904834.1)	89
480	<i>Spegazzinia tessarthra</i> (JQ673429.1)	94
483	<i>Periconia byssoides</i> (MK370654)	98
506	<i>Simplicillium lanosoniveum</i> (KT878334.1)	97
510	<i>Periconia byssoides</i> (MK907734)	99
523	<i>Nigrospora sphaerica</i> (KT259476.1)	98
529	<i>Aspergillus gracillis</i> (MH858708.1)	99

The core mycobiota of chaya consisted of the genera *Cladosporium* (51.70%), *Lasiodiplodia* (18.11%), *Allophoma* (6.79%), *Stagonosporosis* (2.26%) and *Aspergillus* (2.26%). At the

genus level, the unassigned OTUs obtained 15.47% of the total OTUs. The most predominant species were: *Lasiodiplodia theobromae* (14.98%), *Corynespora casiicola* (14.98%), *Cladosporium tenuissimum* (14.98%), *Cladosporium xanthochromaticum* (14.98%), *Cladosporium cladosporioides* (14.98%), *Nigrospora oryzae* (14.98%), *Nigrospora sphaerica* (14.98%), *Stagonosporopsis curcubitacearum* (14.98%), *Allophoma minor* (14.98%), *Aspergillus flavus* (14.98%) and *Aspergillus chavelaeri* (14.98%). The fungal community of *Cnidoscopus aconitifolius* at the genus and species levels is presented in Figures 3 and 4 respectively. A phylogenetic tree was constructed to show the relationship between the genera obtained (Figure 5).




Figure 3: Fungal genera obtained from *Cnidoscopus aconitifolius* phyllosphere




Figure 4: Fungal species obtained from *Cnidoscolus aconitifolius* phyllosphere




Figure 5: Phylogenetic tree generated by maximum composite likelihood analysis based on the ITS sequences of the OTUs

4. DISCUSSION

Many authors have reported that the majority of taxa obtained from plant leaves belong to the division, Ascomycota [24,25]. Ascomycota, Basidiomycota and Chytridiomycota have also been reported as the dominant fungal divisions on plants [26,27].

Cladosporium belongs to the class Dothideomycetes, order Capnodiales and family Cladosporiaceae. Species occur in clusters of black, green or yellow spots. Different species of *Cladosporium* has been found on a variety of plants including *Phaseolus vulgaris*, *Alium porrum*, *Ananas comosus*, *Pinus ponderosa* etc; and on different parts of human samples including sputum, toe nail, lung, foot, skin, scalp, etc [28]. Ten new species belonging to the genus *Cladosporium* were reported by Sandoval-Denis *et al.* [28] and these species are associated with animal and human infections. *C. cladosporioides* and *C. tenuissimum* were

detected in air conditioning, ventilation and heating equipments in China with *C. cladosporioides* having the highest frequency and concentration [29]. Long term exposure to *Cladosporium* is associated with allergies, asthma symptoms, and eye, ear and skin infections.

Stagonosporopsis belongs to the class Dothideomycetes, order Pleosporales and family Didymellaceae. *Stagonosporopsis cucurbitacearum* is a fungal parasitic pathogen reported by Zhao *et al.* [30] to be responsible for gummy stem blight disease of pumpkin in North-east China. The disease stands as one of the major severe pumpkin diseases in North-eastern region of China and leads to huge crop losses. *S. cucurbitacearum* as a parasite inhabits its host, causing serious damage to crops but does not directly kill its host. The pathogen makes use of penetration pegs (conidial structure) at the early stage of infection, to absorb nutrients necessary for its survival [30]. Wipornpan *et al.* [31] also reported gummy stem blight of cantaloupe caused by *Stagonosporopsis cucurbitacearum* in Thailand. Bhuiyan *et al.* [32] and Vaghefi *et al.* [33] reported ray blight disease of pyrethrum (*Tanacetum cinerariifolium*) caused by *Stagonosporopsis tanacetii*. This is considered as a limiting factor to pyrethrum cultivation in Australia.

Corynespora belongs to the class Dothideomycetes, order Pleosporales and family Corynesporaceae. The genus *Corynespora* consists of mostly plant pathogenic species causing diseases on various plants, world-wide. Some of the species of this genus occur as saprobes and endophytes. About 200 species of *Corynespora* have been recorded and several novel species have been reported [34-37]. *Corynespora* is synonymous with the genera, *Cercospora* and *Helminthosporium*. *Corynespora cassiicola*, a phytopathogen reduces yield of natural rubber latex in African and Asian countries. *Corynespora cassiicola* has 375 host species. Symptoms occur as leaf and fruit decay.

The genus *Allophoma* belongs to the class Dothideomycetes, order Pleosporales and family Didymellaceae. *Allophoma* causes lesion and leaf spots on plants. *Allophoma tropica* was first observed on lettuce in Northern Italy in 2011 [38]. *Allophoma tropica* causes leaf spot of lettuce [39]. *Allophoma zantedeschiae* caused symptoms of small yellow lesions on lower leaves of *Papaver dubium* which eventually spread to the whole plant in Iran [40]. *Allophoma minor* is a phytopathogenic fungus [41,42]. It was isolated as one of the pathogens responsible for lesion blight of *Papaver dubium* in Iran.

Aspergillus belongs to the class Eurotiomycetes, order Eurotiales and family Trichocomaceae. *Aspergillus chevalieri* was reported on peanuts in Malaysia [43]. *A. chevalieri* is a xerophilic organism which provides a favorable growth condition for other spoilage-related fungal organisms. This organism might affect the quality of plant products and lead to reduced shelf life. A study conducted by Chukwu *et al.* [44] indicated that *Aspergillus niger*, *A. flavus* and *A. terreus* were associated with both fresh and dry tiger nuts and they can possibly endure processing treatment. The occurrence of these fungi may cause diverse effects on human health as they have the potential of producing mycotoxins [45]. *A. flavus* produces two most common aflatoxins; aflatoxins B1 and B2 [46].

Nigrospora species exist as endophytes on stems and leaves of different plant species [47] or as saprobes from leaf litter or dead larvae [48,49]. The genus consists of plant pathogenic species infecting various fruits, economic crops and ornamentals. Zhai *et al.* [50] reported the occurrence of *Nigrospora oryzae* on *Aloe vera* in China where it caused leaf spots. *N. oryzae* was also reported in India on *Brassica juncea* where it caused stem blight [51].

Nigrospora sphaerica has been isolated from different plants where it caused leaf spots, rots, blight and lesions. *N. sphaerica* was reported in China as the causal agent of leaf blight

of *Camellia sinensis* [52]. Alam *et al.* [53] reported that *N. sphaerica* caused fatal leaf spot on Kinnow mandarin (*Citrus reticulata*). *Nigrosora sphaerica* has also been reported to be associated with a disease in human. Ananya *et al.* [54] reported that *N. sphaerica* caused corneal ulcer in an immunocompetent woman. *N. oryzae* and *N. sphaerica* were found to cause leaf spot on date palms (*Phoenix dactylifera* L.) [55,56].

5. CONCLUSION

The use of next-generation molecular techniques has led to advances in phyllosphere microbiology. These techniques have helped researchers to know the structure of plant microbial communities, the organisms present on leaf surfaces, and what these organisms do in plants. Most of the organisms obtained in this study are plant pathogens, causing deterioration on plants, reduced quality of plant products and devastating decrease in the quantity of agricultural produces recovered after harvest. *Cnidoscopus aconitifolius* is a highly neglected and underutilized plant. Insight into the mycobiota of *C. aconitifolius* phyllosphere is the starting point for devising ways of combating the pathogenic species and increasing the yield of this plant thereby making it more available to the fast-growing population. The correct identification of microorganisms in the phyllosphere and the in-depth understanding of the interaction that exists among these organisms will help in protecting plants against pre- and post-harvest diseases.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. Ross-Ibarra J and Molina-Cruz A. The ethnobotany of chaya (*Cnidoscopus aconitifolius*): A nutritious maya vegetable. *Journal of Ethnobotany*. 2002;56(4):350-364.
2. Hamid AA, Oguntoye SO, Negi AS, Ajao AO. Chemical constituents, antibacterial, antifungal and antioxidant activities of the aerial parts of *Cnidoscopus aconitifolius*. *Ife Journal of Science*. 2016;18(2):561-571.
3. Mordi JC and Akanji MA. Phytochemical screening of the dried leaf extract of *Cnidoscopus aconitifolius* and associated changes in liver enzymes induced by its administration in wistar rats. *Current Research Journal of Biological Sciences*. 2012;4(2):153-158.
4. Lindow SE and Brandl MT. Microbiology of the phyllosphere. *Applied and Environmental Microbiology*. 2003;69:1875-1883.
5. Vorholt JA. Microbial life in the phyllosphere. *Nature Reviews Microbiology*. 2012;10:828-840.

6. Inacio J, Pereira P, De Carvalho M, Fonseca A, Amaral-Collaco MT and Spencer-Martins I. Estimation and diversity of phylloplane mycobiota on selected plants in a mediterranean-type ecosystem in Portugal. *Microbial Ecology*. 2002;44:344-353.
7. Lopez-Velasc G, Carder PA, Welbaum GE and Ponder MA. Diversity of the spinach (*Spinacia oleracea*) spermosphere and phyllosphere bacterial communities. *FEMS Microbiology Letters*. 2013;346:146–154.
8. Rastogi G, Coaker GL and Leveau JH. New insights into the structure and function of phyllosphere microbiota through high-throughput molecular approaches. *FEMS Microbiology Letters*. 2013;348:1–10.
9. Zhou Y, Qiao X, Li W, Xu J, Wang W and Chen X. Phyllosphere bacterial communities associated with the degradation of acetamiprid in *Phaseolus vulgaris*. *African Journal of Biotechnology*. 2011;10:3809-3817.
10. Ali N, Sorkhoh N, Salamah S, Eliyas M and Radwan S. The potential of epiphytic hydrocarbon-utilizing bacteria on legume leaves for attenuation of atmospheric hydrocarbon pollutants. *Journal of Environmental Management*. 2012;93:113-120.
11. Arnold AE, Mejia LC, Kylo D, Rojas EI, Maynard Z, Robbins N and Herre EA. Fungal endophytes limit pathogen damage in a tropical tree. *Proceedings of the National Academy of Sciences of the United States of America*. 2003;100: 5649-15654.
12. Schweitzer JA, Bailey JK, Bangert RK, Hart SC and Whitham TG. The role of plant genetics in determining above- and below-ground microbial communities. In: MJ Bailey, AK Lilley, PT Timms-Wilson and PT Spencer-Phillips (eds.). *Microbial Ecology of the Aerial Plant Surface*. CABI International, Wallingford, UK; 2006.
13. Teplitski M, Warriner K, Bartz J and Schneider KR. Untangling metabolic and communication networks: interactions of enteric with phyto bacteria and their implications in produce safety. *Trends in Microbiology*. 2011;19(3):121-127.
14. Dees MW, Lysoe E, Nordskog B, Brurberg MB. Bacterial communities associated with surfaces of leafy greens: shift in composition and decrease in richness over time. *Applied and Environmental Microbiology*. 2015;81:1530-1539. <https://doi.org/10.1128/aem.03470-14>.
15. Ding T and Melcher U. Influences of plant species, season and location on leaf endophytic bacterial communities of non-cultivated plants. *PLoS One*. 2016;11(3):1-13. <https://doi.org/10.1371/journal.pone.0150895>.
16. Cordier, T., Robin, C., Capdevielle, X., Fabreguettes, O., Desprez-Loustau, M. L. and Vacher, C. (2012). The composition of phyllosphere fungal assemblages of European beech (*Fagus sylvatica*) varies significantly along an elevation gradient. *New Phytologist*, 196: 510–519.
17. Bálint M, Tiffin P, Hallström B, O'Hara RB, Olson MS, Fankhauser JD, Piepenbring M and Schmitt I. Host genotype shapes the foliar fungal microbiome of balsam poplar (*Populus balsamifera*). *PLoS One*. 2013;8(1):1-9.
18. Vacher C, Hampe A, Porté AJ, Sauer U, Compant S and Morris CE. The phyllosphere: microbial jungle at the plant-climate Interface. *Annual Review of Ecology, Evolution, and Systematics*. 2016;47(1):1-24. <https://doi.org/10.1146/annurev-ecolsys-121415-032238>.

19. Iyanyi NG, Ataga AE and Nwachukwu EO. Molecular characterization of fungi associated with *Cnidioscolus aconitifolius* (Mill.) I. M. Johnston. Researchjournal's Journal of Agriculture. 2019;6(4):1-12.
20. Chen C, Khaleel SS, Huang H and Wu CH. Software for pre-processing Illumina next-generation sequencing short read sequences. Source Code for Biology and Medicine. 2014;9(8):1-11.
21. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Buschman FD and Costello EK. QIIME allows for high-throughput community sequencing data. Nature Methods. 2011;7:335-336.
22. Edgar RC, Haas BJ, Clemente JC, Quince C and Knight R. UCHIME improves sensitivity and speed of chimera detection. Bioinformatics. 2011;27(16):2194-2200.
23. Abarenkov K, Nilsson RH, Larsson KH, Alexander IJ, Eberhardt U, Erland S and Kõljalg U. The UNITE database for molecular identification of fungi - recent updates and future perspectives. New Phytologist. 2010;186(2):281-285. <https://doi.org/10.1111/j.1469-8137.2009.03160.x>
24. Selim KA, Nagia MM and Ghwas DE. Endophytic fungi are multifunctional biosynthesizers: Ecological role and chemical diversity. In: Evelyn Hughes (ed.). Endophytic Fungi: Diversity, Characterization and Biocontrol. Microbiology Research Advances. Nova Publishers, New York; 2017.
25. Tibpromma S, Hyde KD, Bhat JD, Mortimer PE, Xu J, Promputtha I, Doilom M, Yang J B., Tang AM and Karunarathna SC. Identification of endophytic fungi from leaves of Pandanaceae based on their morphotypes and DNA sequence data from Southern Thailand. MycoKeys. 2018;33:25-67.
26. Kembel SW and Mueller RC. Plant traits and taxonomy drive host associations in tropical phyllosphere fungal communities. Botany. 2014;92:303–311.
27. Darlison J, Mogren L, Rosberg AK, Grudén M, Minet A, Liné C, Mieli M, Bengtsson T, Håkansson A, Uhlig E, Becher PG, Karlsson M and Alsanius BW. Leaf mineral content govern microbial community structure in the phyllosphere of spinach (*Spinacia oleracea*) and rocket (*Diplotaxis tenuifolia*). Science of the Total Environment. 2019;675:501–512.
28. Sandoval-Denis M, Gene N, Sutton D, Wiederhold N, Cano-Lira J and Guarro J. New species of *Cladosporium* associated with human and animal infections. PubMed. 2016;36:281-298.
29. Luo Y, Li J, Zhang X, and Gao W. Characterization of potential pathogenic *Cladosporium* exposure risks from heating, ventilation and air conditioning (HVAC) in two cities, China. Medical Mycology: Open Access. 2016;16(2):18-25.
30. Zhao Q, Wu J, Zhang L, Xu L, Yan C and Gong Z. Identification and characterization of Cucurbita gummy stem blight fungi in Northeast China. Journal of Phytopathology.
31. Wipornpan N, Worawoot A, Nakarin S and Jaturong K. First Report of Gummy Stem Blight on *Stagonosporosis cucurbitacearum* on Cantaloupe in Thailand. Canadian Journal of Plant Pathology. 2018;40(20):306-311.

32. Bhuiyan MA, Groom T, Nicholas ME and Taylor PW. Disease cycle of *Stagonosporopsis tanacetii* in pyrethrum plants. Australian Plant Pathology. 2017;46(1):83-90.
33. Vaghefi N, Pethybridge SJ, Ford R, Nicolas ME, Crous PW and Taylor PW. *Stagonosporopsis* spp. associated with ray blight disease of Asteraceae. Australasian Plant Pathology. 2012;41:675-686.
34. Kumar S, Singh R, Gond DK and Saini DC. Two new species of *Corynespora* from Uttar Pradesh, India. Mycosphere. 2012;3:864-869.
35. Kumar S, Singh A, Singh R and Dubey NK. *Corynespora bombacina* causing foliar disease on *Bombax ceiba* from Sonebhadra Forest of Uttar Pradesh, India. Canadian Journal of Plant Protection. 2013;1(2):76-77.
36. Singh A, Kumar S, Singh R, Dubey NK. *Corynespora clerodendrigena* sp. nov. causing foliar disease on *Clerodendrum viscosum* from Sonebhadra forest of Uttar Pradesh, India. Plant Pathology and Quarantine. 2013;3(1):15-17.
37. Singh A, Kumar S, Singh R, Dubey NK. A new species of *Corynespora* from Sonebhadra forest of Uttar Pradesh, India. Current Research in Environmental and Applied Mycology. 2014;4(1):149-151.
38. Garibaldi AG, Gilardi GO and Gullino ML. First report of leaf spot of lettuce (*Lactuca sativa* L.) caused by *Phoma tropica* in Italy. Plant Disease. 2012;96(9):1380-1380.
39. Pintore I, Gilardi G, Garibaldi A and Gullino M. Efficacy of different fungicides against the leaf spot of lettuce caused by *Allophoma tropica*. Journal of Plant Diseases and Protection. 2018;125(3):297-309.
40. Parisa R and Doustmorad Z. Characterization of fungi causing lesion blight on *Papaver dubium* in Iran. Antonie van Leeuwenhoek. 2018;111(3):437-455.
41. Razaghi P and Zafar D. Characterization of fungi causing lesion blight on *Papaver dubium* in Iran. Antonie van Leeuwenhoek. 2017;111(6):1-20. doi: 10.1007/s10482-017-0966-8
42. Marin-Felix Y, Hernández-Restrepo M, Iturrieta- González I, Gené GD, Groenewald J, Ca L, Chen Q, Quaedvlie W, Schumacher R, Taylor P, Ambers C, Bonthond G, Edwards J, Krueger-Hadfield S, Luangsaard J Morton L, Moslemi A, Sandoval-Denis, M, Tan Y, Thangavel R, Vaghefi N, Cheewangkoon R and Crous P. Genera of phytopathogenic fungi: GOPHY 3. Studies in Mycology. 2019;94:1-124.
43. Kamarudin NA and Zakaria L. Characterization of two xerophilic *Aspergillus* spp. from peanuts (*Arachis hypogaea*). Malaysian Journal of Microbiology. 2018;14(1):41-48.
44. Chukwu MO, Ibiam OFA and Okoi A. Studies on the fungi and phytochemical and proximate composition of dry and fresh tiger nuts (*Cyperus esculentus* L.). International Research Journal of Biotechnology. 2013;4(1):11-14.
45. Nura M, Abubakar A, Auyo MI, Sunday E and Kutama AS. Isolation and identification of fungi associated with tigernut milk drink (kunum aya) in Dutse, Jigawa State. Global Advanced Research Journal of Agricultural Science. 2016;57:302-308.

46. Amaike S and Nancy P. *Aspergillus flavus*. Annual Review of Phytopathology. 2011;49:107-133.
47. Wang M, Liu F, Crous P and Cai L. Phylogenetic reassessment of *Nigrospora*: ubiquitous endophytes, plant and human pathogens. Persoonia. 2017;39:118-142.
48. Thalavaipandian A, Ramesh V, Bagyalakshmi A, Muthuramkumar S and Rajendran A. Diversity of fungal endophytes in medicinal plants of Courtallam Hills, Western Ghats, India. Mycosphere. 2011;2(5):575-582.
49. Uzor PF, Ebrahim W and Osadebe PO. Metabolites from *Combretum dolichopetalum* and its associated endophytic fungus *Nigrospora oryzae* – evidence for a metabolic partnership. Fitoterapia. 2015;105:147-150.
50. Zhai L, Liu J, Zhang M and Wang L. First report of leaf spots in aloe vera caused by *Nigrospora oryzae* in China. Plant Disease. 2013;9(7):1256.
51. Sharma P, Meena PD and Chauhan JS. First report of *Nigrospora oryzae* (Berk. and Broome) Petch causing stem blight on *Brassica juncea* in India. Journal of Phytopathology. 2013;161:439-441.
52. Liu YJ, Tang Q and Fang L. First report of *Nigrospora sphaerica* causing leaf blight on *Camellia sinensis* in China. Plant Disease. 2015;100:221.
53. Alam MW, Rehman A, Gleason ML, Riaz K Saira M, Aslam S, Rosli H and Muhammad S. First report of *Nigrospora sphaerica* causing leaf spot of kinnow mandarin in Pakistan. Journal of Plant Pathology. 2017;99(1):295.
54. Ananya TS, Kindo AJ, Subramanian A and Suresh K. *Nigrospora sphaerica* causing corneal ulcer in an immunocompetent woman: A case report. International Journal of Case Reports and Images. 2014;10:675-679.
55. Abass MH, Hameedm MA and Ahmedm A N. First report of *Nigrospora sphaerica* (Sacc.) Mason as a potential pathogen of date palm (*Phoenix dactylifera* L.). Canadian Journal of Plant Pathology. 2013;35(1):75-80.
56. Abass MZ and Mohammed NJ. Morphological, molecular and pathological study on *Nigrospora oryzae* and *Nigrospora sphaerica*, the leaf spot fungi of date palm. Basra Journal for Date Palm Researches. 2014;13(1-2):26-38.