

Original Research Article

A COMPARATIVE STUDY OF LOW VERSUS STANDARD INTRAPERITONEAL PRESSURES IN GYNAECOLOGICAL LAPAROSCOPIC SURGERY

ABSTRACT

BACKGROND -:Laparoscopic surgery has several advantages when compared to open surgery, including faster postoperative recovery and lower pain scores. However, Increased intraabdominal pressure may also have negative implications on cardiovascular, pulmonary, and intraabdominal organ functioning. To overcome these negative consequences, several trials have been performed comparing low- versus standard-pressure pneumoperitoneum.

AIM - The aim of present study was to evaluate the effectivity of low intraperitoneal pressures in comparison with standard intraperitoneal pressure during laparoscopic hysterectomies.

STUDY DESIGN- Experimental study

MATERIAL AND METHOD– 40 cases with uncomplicated symptomatic benign uterine pathologies who were posted for laparoscopic hysterectomy were selected out of which 20-20 cases were randomized into low and standard pneumoperitoneum groups.

RESULT– in patients in whom low pressure pneumoperitoneum is employed are better recovered in terms of pain than standard pressure pneumoperitoneum. This means hospital stay can be shortened in low pressure pneumoperitoneum groups which will be more economical and comfortable for patients.

CONCLUSION–

Laparoscopic hysterectomy can be done at 10 mmhg with the benefits of :

- Optimum visualization with low pressure
- Reduction in post operative pain helping the patient for early ambulation so that patient will get back to routine work and normal life earlier, it is the main purpose of minimal invasive surgery.

KEY WORDSlaparoscopy, pneumoperitoneum, hysterectomy.

INTRODUCTION

The optimal intraperitoneal pressure during laparoscopy is not known. Recent literature found benefits of using lower pressures, but the effectivity of doing abdominal surgery with low peritoneal pressures needs to be assessed. This study compares low with standard pneumoperitoneum during gynecological laparoscopy.

Intraperitoneal pressures at or above 12 mm Hg are usually used for intra-abdominal laparoscopy.^{1,2} Some authors have postulated that reducing abdominal distention might be able to decrease postoperative pain and the risk of laparoscopy-related complications such as air embolism, pneumothorax, pneumomediastinum, arrhythmia, and ventilation issues.³⁻⁸ There is a concern of a poorer visualization of the operative field⁹ and therefore an increase in the occurrence of complications. The optimal pneumoperitoneum pressure would allow proper visualization while having the fewest intra and postoperative complications.¹ Furthermore, given the differences in patient positioning (Trendelenburg vs Fowler) and the nature of the surgeries in gynecologic laparoscopy.¹⁰⁻¹²

However, little data exist about the possibility of abdominal endoscopic surgery with peritoneal pressures below the standard value of 12 mm Hg.^{13,14} Hence the purpose of this study to compare the gynaecologiclaparoscopies done with low and standard Intraperitoneal pressure.

AIMS AND OBJECTIVES

- The aim of present study was to evaluate the effectivity of low intraperitoneal pressures in comparison with standard intraperitoneal pressure during laparoscopic hysterectomies.

OBJECTIVES

-To compare the effectivity of low intraperitoneal pressure with standard intraperitoneal pressure in terms of

1. duration of operating time related to adequate exposure
2. Intraoperative haemodynamic changes

Postoperative pain

METHOD.

- **Study type** – Experimental study
- **Sample size** – out of total 40 cases, in which 20 were randomised in to low pneumoperitoneum pressure group and 20 patients were randomized in to standard pressure group.

INCLUSION CRITERIA

- All consecutive patients with uncomplicated symptomatic benign uterine pathologies posted for laparoscopic hysterectomies who gave consent for study

EXCLUSION CRITERIA

- Patients with complicated uterine pathologies, with large size masses (more than 20 weeks size), and with cancer.
 - Cases with medical and surgical problems.
 - Cases with acute abdomen
- Patients were randomized into two groups. One group with patients undergoing laparoscopic hysterectomy with standard pressure pneumoperitoneum at 14 mm Hg while the other group with patients undergoing laparoscopic hysterectomy with low pressure pneumoperitoneum at 10 mm Hg
 - A standard laparoscopic hysterectomy was performed with the insertion of four ports at the start of surgery.

- At admission, the patient's blood pressure and heart rate was noted.
- Intraoperative blood pressure and heart rate was noted. The difference between the readings at admission and those taken intraoperatively was calculated.
- Postoperative analgesia was administered in the form of diclofenac 12 hourly with additional doses where necessary. Postoperative pain was assessed at 6, 12 and 24 hours using a visual analogue scale.
- Need for additional analgesia over and above the 12 hourly diclofenac and incidence of shoulder tip pain was also noted.

RESULTS

Table no 1 : Time required while using Standard pressure v/s Low pressure pneumoperitoneum

| | Standard pressure | Low pressure |
|-----------------------|-------------------|--------------|
| Average time required | 139.2 | 147.3 |
| Minimum time required | 105 | 120 |

| | | |
|-----------------------|-----|-----|
| Maximum time required | 195 | 180 |
|-----------------------|-----|-----|

- Laparoscopic hysterectomy with standard pressure pneumoperitoneum took an average of 139.2 ± 6.9 minutes with a minimum of 105 minutes and a maximum of 195 minutes.
- Laparoscopic hysterectomy with low pressure pneumoperitoneum took an average of 147.3 ± 5.7 minutes with a minimum of 120 minutes and a maximum of 180 minutes.
- Low pressure laparoscopic hysterectomy took on average eight minutes more than standard pressure laparoscopic hysterectomy but this difference was not statistically significant ($p = 0.1$).

Table no 2 : Changes in blood Pressure while using Standard pressure v/s Low pressure pneumoperitoneum

| Average change in Blood Pressure | Standard pressure | Low pressure |
|----------------------------------|---------------------|-----------------------|
| Systolic Bp | 0.8 ± 8.9 mm Hg | 0.96 ± 6.27 mm Hg |
| Diastolic Bp | 2.8 ± 4.2 mm Hg | 1.8 ± 5.2 mm Hg |

- The average change in systolic BP in patients who underwent low pressure laparoscopic hysterectomy was an increase of 0.96 ± 6.27 mm Hg with a maximum rise of 13 mm Hg and a maximum fall of 10 mm Hg.
- The average change in systolic BP in patients who underwent standard pressure laparoscopic hysterectomy was an increase of 0.8 ± 8.9 mm Hg with a maximum rise of 18 mmHg and a maximum fall of 16mm Hg. This difference was not statistically significant.
- Average change in diastolic blood pressure in patients who underwent low pressure laparoscopic hysterectomy was increase of 1.8 ± 5.2 mm Hg with a maximum rise of 13 mm Hg and a maximum fall of 7 mm Hg.
- The average change in diastolic BP in patients who underwent standard pressure laparoscopic hysterectomy was an increase of 2.8 ± 4.2 mm Hg with a maximum rise of 10 mm Hg and a maximum fall of 7 mm Hg. This difference was not statistically significant.

Table 3 : Changes in heart rate while using Standard pressure v/s Low pressure pneumoperitoneum

| | Standard pressure | Low pressure |
|------------------------------|---------------------------------|---------------------------------|
| Average change in heart rate | 1.5 ± 6.02 beats per minute | 0.5 ± 5.28 beats per minute |

- The average change in heart rate in patients who underwent low pressure laparoscopic hysterectomy was a decrease of 0.5 ± 5.28 beats per minute.
- Average change in heart rate in patients who underwent standard pressure laparoscopic hysterectomy was an increase of 1.5 ± 6.02 beats per minute.

This difference was not statistically significant.

Table 4 : Pain Score noted while using Standard pressure v/s Low pressure pneumoperitoneum

| | Standard pressure | Low pressure |
|-------------------------------------|-------------------|--------------|
| Average pain score at time interval | | |
| 6 hours | 59.1 | 62.2 |
| 12 hours | 62.2 | 54.2 |
| 24 hours | 5.2 | 4.6 |

- The average pain score at 6 hours for patients who underwent low pressure laparoscopic hysterectomy was 62.2 ± 11.7 with a minimum of 36 and a maximum of 82. The pain score at 6

hours for standard pressure laparoscopic hysterectomy was 59.1 ± 18.0 with a minimum of 35 and a maximum of 100. This difference was not statistically significant ($p = 0.4$).

- The average pain score at 12 hours for patients who underwent low pressure laparoscopic hysterectomy was 54.2 ± 8.5 with a minimum of 38 and a maximum of 69. The average pain score at 12 hours for patients who underwent standard pressure laparoscopic hysterectomy was 62.2 ± 12.0 with a minimum of 35 and maximum of 100. This difference was statistically significant. ($p = 0.04$)
- The average pain score at 24 hours for patients who underwent standard pressure laparoscopic hysterectomy was $5.2 \pm .8$ with a minimum of 3.5 and a maximum of 10. Average pain score at 24 hours for patients who underwent low pressure laparoscopic hysterectomy was $4.60 \pm .81$ with a minimum of 3.6 and a maximum of 8.2. This difference was not statistically significant.

Table 5: Shoulder Tip Pain noted while using Standard pressure v/s Low pressure pneumoperitoneum

| | Standard pressure | Low pressure |
|---------------------------|-------------------|--------------|
| No. of cases | 18 | 17 |
| Shoulder tip pain present | 2(11.1%) | 1(5.8%) |

| | | |
|--------------------------|-----------|-----------|
| Shoulder tip pain absent | 16(88.8%) | 16(94.2%) |
|--------------------------|-----------|-----------|

- One (5.8%) of the 17 patients who underwent low pressure laparoscopic hysterectomy and two (11.11%) of the 18 patients who underwent standard pressure laparoscopic hysterectomy had post operative pain referred to the tip of the right shoulder. This difference was not statistically significant ($p = 1.0$).

DISCUSSION

CO₂ is the insufflation gas of choice in laparoscopy. It is preferred over air insufflation, which affects the systemic and peritoneal response to a larger degree than CO₂.¹⁵ The usage of CO₂ has some important advantages. It is transparent, non-inflammable, and well dissolvable in blood. There are, however, some disadvantages associated with its usage. The increased intra-abdominal pressure increases the absorption of CO₂, causing hypercapnia and acidosis, which has to be avoided by hyperventilation.¹⁶ It also pushes the diaphragm upwards, decreasing the pulmonary compliance,^{16,17} and increases the peak airway pressure.^{17,18} Pneumoperitoneum also increases systemic vascular resistance^{18,19} and pulmonary vascular resistance.¹⁸ CO₂ pneumoperitoneum also predisposes the patient to cardiac arrhythmias.²⁰ During the early phase of pneumoperitoneum, there is a reduction in the cardiac output^{17,18} by decreasing the venous return.²¹ Although these cardiorespiratory changes may be tolerated by healthy adults with adequate cardiopulmonary reserve, people with cardiopulmonary disease may not be able to tolerate these cardiopulmonary changes. Abdominal wall lift, using a special device, e.g., Laparolift²⁰ or Laparo-tensor,¹⁷ introduced through a port in the abdominal wall has been applied to decrease the cardiopulmonary changes.²²

Randomised clinical trials have shown that using a lower pressure of pneumoperitoneum decreases the cardiac changes,²³ the number of people complaining of shoulder-tip pain,²⁴ the intensity of pain,²⁵ and the analgesic requirement.²⁴⁻²⁷ Low intra-abdominal pressure may prevent mortality due to CO₂ embolism.²⁸ Moreover, Schwarte *et al.*²⁹ found that increasing intra-abdominal pressure decreased gastric mucosal O₂ saturation. The European Association for Endoscopic Surgery (EAES) guidelines²¹ also recommend use of the lowest intra-abdominal pressure rather than a routine pressure (14 mmHg) to allow adequate exposure of the operative field.

Conclusion

- Both groups are equally comparable in almost all above parameters. There is no significant differences in both groups, this means low pressure pneumoperitoneum can be used as effectively as standard pressure pneumoperitoneum.
- In patients in whom low pressure pneumoperitoneum is employed are better recovered in terms of pain than standard pressure pneumoperitoneum. This means hospital stay can be shortened in low pressure pneumoperitoneum groups which will be more economical and comfortable for patients.
- Impact of low pressure pneumoperitoneum on intra-operative hemodynamics is not significant. This needs to be examined through a more complex set up and probably a larger sample size that includes a significant numbers of patients with cardiovascular comorbid conditions.

REFERENCES

1. Neudecker J, Sauerland S, Neugebauer E, Bergamaschi R, Bonjer HJ, Cuschieri A, et al. The European Association for Endoscopic Surgery clinical practice guideline on the pneumoperitoneum for laparoscopic surgery. *Surg Endosc.* 2002;16(7): 1121–1143. doi:10.1007/s00464-001-9166-7.
2. laChapelle CF, Bemelman WA, Rademaker BM, van Barneveld TA, Jansen FW. A multidisciplinary evidence-based guideline for minimally invasive surgery: Part 1: entry techniques and the pneumoperitoneum. *Gynecol Surg.* 2012;9(3): 271–282.
3. Galizia G, Prizio G, Lieto E, Castellano P, Pelosio L, Imperatore V, et al. Hemodynamic and pulmonary changes during open, carbon dioxide pneumoperitoneum and abdominal walllifting cholecystectomy. A prospective, randomized study. *SurgEndosc.* 2001;15(5):477–483. doi:10.1007/s004640000343.
4. Leonard IE, Cunningham AJ. Anaesthetic considerations for laparoscopic cholecystectomy. *Best Pract Res Clin Anaesthesiol.* 2002;16(1):1–20.
5. Henny CP, Hofland J. Laparoscopic surgery: pitfalls due to anesthesia, positioning, and pneumoperitoneum. *Surg Endosc.* 2005;19(9):1163–1171. doi:10.1007/s00464-004-2250-z.
6. Wallace DH, Serpell MG, Baxter JN, O'Dwyer PJ. Randomized trial of different insufflation pressures for laparoscopic cholecystectomy. *Br J Surg.* 1997;84(4):455–458.

7. Sarli L, Costi R, Sansebastiano G, Trivelli M, Roncoroni L. Prospective randomized trial of low-pressure pneumoperitoneum for reduction of shoulder-tip pain following laparoscopy. *Br J Surg*. 2000;87(9):1161–1165. doi:10.1046/j.1365–2168. 2000.01507.x.
8. Chok KS, Yuen WK, Lau H, Fan ST. Prospective randomized trial on low-pressure versus standard pressure pneumoperitoneum in outpatient laparoscopic cholecystectomy. *Surg LaparoscEndosc Percutan Tech*. 2006;16(6):383–386. doi:10.1097/01.sle.0000213748.00525.1e.
9. Angioli R, Terranova C, Plotti F, Cafa EV, Gennari P, Ricciardi R, et al. Influence of pneumoperitoneum pressure on surgical field during robotic and laparoscopic surgery: a comparative study. *Arch Gynecol Obstet*. 2015;291(4):865– 868. doi:10.1007/ s00404–014-3494-z.
10. Hirvonen EA, Nuutinen LS, Kauko M. Hemodynamic changes due to Trendelenburg positioning and pneumoperitoneum during laparoscopic hysterectomy. *Acta AnaesthesiologicaScandinavica*. 1995;39(7):949 –955.
11. Hirvonen EA, Poikolainen EO, Paakkonen ME, Nuutinen LS. The adverse hemodynamic effects of anesthesia, head-up tilt, and carbon dioxide pneumoperitoneum during laparoscopic cholecystectomy. *Surg Endosc*. 2000;14(3):272–277.
12. Rist M, Hemmerling TM, Rauh R, Siebzehnruhl E, Jacobi KE. Influence of pneumoperitoneum and patient positioning on preload and splanchnic blood volume in laparoscopic surgery of the lower abdomen. *J Clin Anesth*. 2001;13(4):244 –249.
13. Gurusamy KS, Vaughan J, Davidson BR. Low pressure versus standard pressure pneumoperitoneum in laparoscopic cholecystectomy. *Cochrane Database Syst Rev*. 2014;3:CD006930.doi:10.1002/14651858.CD006930.pub3.
14. Ozdemir-van Brunschot DM, van Laarhoven KC, Scheffer GJ, Pouwels S, Wever KE, Warle MC. What is the evidence for the use of low-pressure pneumoperitoneum? A systematic review. *SurgEndosc*. 2015. doi:10.1007/s00464–015-4454 –9.
15. Watson RW, Redmond HP, McCarthy J, Burke PE, Bouchier-Hayes D. Exposure of the peritoneal cavity to air regulates early inflammatory responses to surgery in a murine model. *Br J Surg* 1995;82:1060-5.
16. Henny CP, Hofland J. Laparoscopic surgery: Pitfalls due to anesthesia, positioning, and pneumoperitoneum. *Surg Endosc* 2005;19:1163-71.
17. Alijani A, Hanna GB, Cuschieri A. Abdominal wall lift versus positive pressure capnoperitoneum for laparoscopic cholecystectomy: Randomized controlled trial. *Ann Surg* 2004;239:388-94.

18. Galizia G, Prizio G, Lieto E, Castellano P, Pelosio L, Imperatore V, *et al.* Hemodynamic and pulmonary changes during open, carbon dioxide pneumoperitoneum and abdominal wall-lifting cholecystectomy. A prospective, randomized study. *Surg Endosc* 2001;15:477-83.
19. Mertens zur Borg IR, Lim A, Verbrugge SJ, IJzermans JN, Klein J. Effect of intraabdominal pressure elevation and positioning on hemodynamic responses during carbon dioxide pneumoperitoneum for laparoscopic donor nephrectomy: A prospective controlled clinical study. *Surg Endosc* 2004;18:919-23.
20. Egawa H, Morita M, Yamaguchi S, Nagao M, Iwasaki T, Hamaguchi S, *et al.* Comparison between intraperitoneal CO₂ insufflation and abdominal wall lift on QT dispersion and rate-corrected QT dispersion during laparoscopic cholecystectomy. *Surg Laparosc Endosc Percutan Tech* 2006;16:78-81.
21. Neudecker J, Sauerland S, Neugebauer E, Bergamaschi R, Bonjer HJ, Cuschieri A, *et al.* The European Association for Endoscopic Surgery clinical practice guideline on the pneumoperitoneum for laparoscopic surgery. *Surg Endosc* 2002;16:1121-43.
22. Gurusamy KS, Samraj K, Davidson BR. Abdominal lift for laparoscopic cholecystectomy. *Cochrane Database Syst Rev* 2008;CD006574.
23. Dexter SP, Vucevic M, Gibson J, McMahon MJ. Hemodynamic consequences of high- and low-pressure capnoperitoneum during laparoscopic cholecystectomy. *Surg Endosc* 1999;13:376-81.
25. Sarli L, Costi R, Sansebastiano G, Trivelli M, Roncoroni L. Prospective randomized trial of low-pressure pneumoperitoneum for reduction of shoulder-tip pain following laparoscopy. *Br J Surg* 2000;87:1161-5.
26. Wallace DH, Serpell MG, Baxter JN, O'Dwyer PJ. Randomized trial of different insufflation pressure for laparoscopic cholecystectomy. *Br J Surg* 1997;84:455-8.
27. Sandhu T, Yamada S, Ariyakachon V, Chakrabandhu T, Chongruksut W, Ko-iam W. Low-pressure pneumoperitoneum versus standard pneumoperitoneum in laparoscopic cholecystectomy, a prospective randomized trial. *Surg Endosc* 2009;23:1044-7.
28. Beebe DS, Zhu S, Kumar MV, Komanduri V, Reichert JA, Belani KG. The effect of insufflation pressure on CO₂ pneumoperitoneum and embolism in piglets. *Anesth Analg* 2002;94:1182-7.
29. Schwarte LA, Scheeren TW, Lorenz C, De Bruyne F, Fournell A. Moderate increase intraabdominal pressure attenuates gastric mucosal oxygen saturation in patients undergoing laparoscopy. *Anesthesiology* 2004;100:1081-7.