

## Original Research Article

### Free-Radical Scavenging Assay and Quantitative Estimation of Flavonoids from *Corchorus depressus*

#### ABSTRACT

**Background:** The investigation of chemical constituents (Qualitative & Quantitative) and the evaluation of antioxidant activities of *Corchorus depressus* is the major aim of this study. As observed from ancient literatures & folkloric claims the Plant *Corchorus depressus* worshipped by the married women of Odisha, in the rituals called as “Jama Jutia”, possesses different biological activities including antioxidant property.

**Methods:** The DPPH (Diphenyl picryl hydrazine), hydroxyl radical (OH) and Nitric oxide (NO) radical scavenging assay was performed for measurement of the antioxidant activity at different extracts. The flavonoid rich fraction and phenolic content of the extracts were determined by using Aluminium chloride and Folin-Ciocalteu's Reagent (FCR) methods respectively.

**Results:** The results for estimation of total phenolic content (mg/ 100g) in Gallic acid Equivalent (GAE) & total flavonoid (mg/ 100g) in weight of Quercetin equivalent (QE) was highest in methanolic extract 78.46 & 21.2 respectively, followed by 18.18 mg/100g in GAE and 1.80mg/100g in QE for aqueous extract.

**Conclusion:** The methanolic extract of *C. depressus* (MECD) at 100µg/ml showed highest DPPH, OH & NO radical scavenging activity and this activity may be attributed due to the presence of saponins and flavonoids as detected in the extract.

*Key words:* Antioxidant, Flavonoids, *Corchorus depressus*, Hydroxyl radical

## 1. INTRODUCTION

A majority of the present day disease are reported to be due to the shift in the balance of the pro-oxidant and the anti-oxidant homeostatic phenomenon in the body. Pro-oxidant conditions dominate either due to the increased generation of free radicals caused by excessive oxidative stress of the present day life or due to the poor scavenging or quenching effect in the body caused by depletion of the dietary anti oxidant [1-2].

A lot of synthetic antioxidants like BHT, BHA and gallates are reported to produce cancer and hence their use have been restricted, as such there is an increased demand for the search of natural products that can be utilized as an antioxidant [3].

Plants are considered as wide source of antioxidants. The polyphenolic compounds like flavonoids present in them are excellent antioxidants. The proton donating ability of polyphenols empowers it to interrupt the oxidation mechanism so there is prevention of oxidative damage [4-5].

The strong biological activity of Plant phenolics and flavonoids outlines their necessity for quantitative determination.

The selected plant *Corchorus depressus* Linn. (Tiliaceae) is treated as religious and worshipped by the married women of Odisha, India, in the rituals called as “Jama Jutia”. The women make a daylong fasting, prepare different sweets, cakes and worship the plant in the evening. An old man of the society will sit at a distance and act as “Yamaraj” (The Lord of Death). The women sweep the road with the plants up to the old man and offer the cakes. The worshipped plants were taken up by the women and softly swept over on the body of their family members and it is believed that by doing so the family members will be free from attack of any disease and have a long life (**Figure 1**).

**Figure 1.** The Plant *C.depressus* worshipped in the festival “Jamajutia”, in Odisha, India.



In the Indigenous system of medicine the above plant used as a cooling medicine in fevers, as tonic; plants' mucilage is prescribed in gonorrhoea. On stone, the roots are rubbed and smeared over forehead to get relief in migraine; to cure leucorrhoea dried fruits are powdered and taken orally with milk for 2 to 3 days. The plant is crushed with tender twigs of *Prosopis cineraria*, mixed with whey and sugar and taken as a drink to treat body ache, protrusion of uterus, urinal inflammation and to avoid abortion. Leaves are made into paste and mixed with curd or whey and given orally to cure diarrhoea in children for 2-3 days [6].

It is also used to increase the viscosity of seminal fluid, to set-up menstrual disorder [7]. An extract of plant is used for its anti-diabetic activity and applied as a paste in healing of wounds (The wealth of India) [8]. Ikram *et al* studied the hexane and chloroform soluble whole plant extract of *C. depressus* which exhibited prominent antipyretic activity in rabbits receiving subcutaneous yeast injections and it did not show any toxic or adverse effect up to an oral dose of 1.6g/kg [9].

Pareek *et al* studied the invitro hepatoprotective activity of *Corchorus depressus* L. against CCl<sub>4</sub> induced toxicity in HepG<sub>2</sub> cell line. It was observed that the ethanolic extract alleviated the changes induced by CCl<sub>4</sub> in a concentration dependent manner [10].

Kataria *et al* through their research demonstrated the *invitro* and *invivo* aphrodisiac properties of *Corchorus depressus* Linn. on rabbit corpus cavernosum smooth muscle relaxation and sexual behavior of normal male rats [11].

A survey of the published literatures revealed that the antioxidant activity of this plant has not yet been subjected for scientific investigation. The aim of present research therefore is the quantative analysis of the Phenolics and flavonoids content in different extracts of *Corchorus depressus* and study their antioxidant property by using different protocols so as to authenticate the folkloric information about the utilization of the plant.

## **2. MATERIALS AND METHODS**

### **2.1 Collection of Plant Material**

*Corchorus depressus* Linn. (Tiliaceae) was authenticated by Botanical Sur. of India (BSI), Central National Herbarium, Kolkata, INDIA (Authentication No- CNH/I-I/28/2009/Tech.II/93) and (Figure-3&4) a voucher specimen (SJCPS-T) was preserved in the herbarium, SJCPS, Naharkanta, Bhubaneswar, Odisha, INDIA after its collection from local area of Salipur, present in state Odisha, India since August 2009.

### **2.2 Chemicals**

Standard quercetin, Gallic acid, Luteolin was obtained from Sigma Chemicals. All other chemicals and reagent used were used from E-Merck and of analytical grade.

### **2.3 Experimental**

#### **2.3.1 Preparation of Plant material**

The plants were washed thoroughly under running tap water, chopped; air dried for a week at 35 to 40 °C. The leaves, stem and the root were separated to prepare the powders separately. The respective plant materials were pulverized in electric grinder separately to moderately fine

powders (355/180) [All particles pass through a No. 355 sieve and not more than 40% through a No. 180 sieve] [12].

### **2.3.2 Extraction Process**

The moderately fine powder of the plant material (1 kg) was initially defatted with petroleum ether. The defatting process concerning mainly with the elimination of chlorophyll (colouring matter, wax in leaves or fixed oils and proteins) which may interfere with isolation of other phyto-constituents from the extract. The defatted material was processed to successive extraction employing solvents of different polarity in ascending order. The powder and the solvent was presented to maceration by stirring at each 4hrs interval for 48hrs at room temperature ( $25\pm5^{\circ}\text{C}$ ) and then filtered with what man filter paper of 2 to 3  $\mu\text{m}$  pore size [13]. The procedure adopted was for shortening the process of extraction, and to minimize the contact time of plant sample with solvent [14].

### **2.3.3 Qualitative Phytochemical analysis**

In the present work, the Qualitative Phytochemical analysis was carried out in the different extracts obtained from the selected plant. All the extracts were subjected to various chemical tests as described in the different literatures (Wager, 1984; Odebiyi, 1978; Trease and Evans, 1987) for preliminary identification of various phytoconstitutents [15-17].

### **2.3.4 Total Flavonoid Content:**

Aluminium chloride colorimetric method with a little modification was used to determine the flavonoid content as per Chang *et.al.* 2002 [18]. Standard Quercetin 10 mg dissolved in 96% ethanol and then it was used for preparation of calibration curve with different concentrations like 2,4,6,8,10 & 12  $\mu\text{g/ml}$ . 1 ml of standard & 1ml of extract sample were mixed with 3 ml of 96% ethanol, 0.2ml of aluminium chloride 10%, 0.2 ml of Potassium

acetate 1M and 5.6 ml distilled water. The mixture was incubated at room temperature for 10 mins with occasional shaking. The absorbance was measured at 376 nm against a blank without aluminium chloride using UV-Visible spectrophotometer.

### **2.3.5 Determination of Total Phenolic Content**

Folin-Ciocalteu's Reagent (FCR) method with little modification was used for quantification of total Phenolic compounds present in the extract of *C. depressus*. In a 10 ml volumetric flask, 0.2ml of plant extract and 0.5ml of Folin-Ciocalteu reagent (2N) was added. After 3 minutes, 1ml of saturated sodium carbonate (20% in distilled water) was added in the same volumetric flask. Final volume was made up to 10 ml with the distilled water. At 725nm the absorbance of the blue coloured formed was measured after 1 hr against a distilled water (blank) using UV-Visible spectrophotometer. A standard calibration curve with gallic acid was plotted using different concentrations (Standard, 100-600 µg/ ml) [19, 20].

### **2.3.6 Evaluation of *In-vitro* Antioxidant Activity**

The evaluation of the *in-vitro* antioxidant activity of *C. depressus* was carried out by using the following methods [21];

#### **2.3.6.1 DPPH Radical scavenging activity**

The DPPH assay measured hydrogen atom (or one electron) donating activity and hence provided an evaluation of anti-oxidant activity due to free radical scavenging. DPPH, a stable and coloured (Purple) free radical which was reduced into the yellow-coloured Diphenyl-picryl hydrazine. 0.1mM solution of DPPH solution in methanol was prepared. 1 ml of this solution was mixed with 3ml of sample solutions in water at different concentrations. Finally after 30 minutes of incubation period at room temperature, the absorbance was measured at 517 nm against the blank.

DPPH radical scavenging activity was calculated according to the following equation;

$$\% \text{ inhibition} = (A_0 - A_1) / A_0 \times 100$$

$A_0 \longrightarrow$  Absorbance of the Control (without extract)

$A_1 \longrightarrow$  Absorbance of the Sample.

### 2.3.6.2 *In-vitro* Hydroxyl radical scavenging activity

The formation of hydroxyl radical (OH) from Fenton reagent was quantified using 2-deoxyribose oxidative degradation. The principle of the assay is the quantification of the 2-deoxyribose degradation product, malonaldehyde, by its condensation with thiobarbituric acid (TBA). The reaction mixture contained deoxyribose (2.8 mM),  $\text{FeCl}_3$  (100mM),  $\text{KH}_2\text{PO}_4$ -KOH buffer (20mM,  $\text{p}^{\text{H}}$  7.4), EDTA (100mM),  $\text{H}_2\text{O}_2$  (1mM), ascorbic acid (100mM) and various concentrations of the extracts in a final volume of 1 ml. Ferric chloride and EDTA (when added) were remixed just before the addition to the reaction mixture. Reaction mixture was incubated at  $37^\circ\text{C}$  for 30 mins. After incubation at  $37^\circ\text{C}$  for 30 mins, 1 ml of 2.8% trichloro acetic acid and 1 ml of 1 % aqueous solution of TBA were added to the sample and the test tubes were heated at  $100^\circ\text{C}$  for 20 min to develop the colour. After cooling, TBARS formation was measured spectrophotometrically (Perkin-Elmer) at 532 nm against an appropriate blank. The hydroxyl radical scavenging activity was determined by comparing absorbance of the control with that of test compounds. The hydroxyl radicals scavenging activity was calculated according to the following equation;

$$\% \text{ inhibition} = (A_0 - A_1) / A_0 \times 100$$

$A_0 \longrightarrow$  Absorbance of the Control (without extract)

$A_1 \longrightarrow$  Absorbance of the Sample.

### 2.3.6.3 *In-vitro* Nitric oxide radical scavenging activity:

Nitric oxide was generated from sodium nitroprusside, which at physiological P<sup>H</sup> liberates nitric acid. This nitric acid gets converted to nitrous acid and further forms nitrite ions which diazotize with sulphanilic acid and couple with naphthylenediamine (Griess reagent), producing pink colour which can be measured at 546 nm. Sodium nitroprusside (10 mM, 2 ml) in phosphate buffer saline (P<sup>H</sup> 7.4) was incubated with 0.5 ml of test compounds in different concentrations at room temperature for 30 minutes. After 30 minutes, 0.5 ml of the incubated solution was added with 1 ml of Griess reagent and the absorbance was measured at 546 nm. The nitric oxide radical scavenging activity was calculated according to the following formula;

$$\% \text{ inhibition} = (A_0 - A_1) / A_0 \times 100$$

A<sub>0</sub> → Absorbance of the Control (without extract)

A<sub>1</sub> → Absorbance of the Sample.

The same experiment was repeated with ascorbic acid, as taken in different concentrations for methanolic and aqueous leaf extracts of *C. depressus*.


## 3. RESULTS AND DISCUSSION

Different extracts of *C. depressus*, were subjected to qualitative phytochemical analysis where the methanolic and the aqueous leaf extracts displayed variety of Phytochemicals present in the extract. The maximum number of phytochemicals like the alkaloids, glycosides, steroids, flavonoids, Saponins, tannins, and phenolic compounds were detected from the methanolic leaf extract, followed by the detection of Saponins, flavonoids, tannins and phenolic compounds from the aqueous extract.




With the measured absorbance data obtained spectrophotometrically, a standard calibration curve was plotted for determination of total flavonoid content which was expressed as weight of Quercetin equivalent (QE) at 100g extract (**Figure 2**).

**Figure 2: Calibration curve of Standard Quercetin.**



The results obtained by FCR method, the total phenolic content of the extract was determined as Gallic acid equivalents (GAE) from the plotted standard calibration curve of Gallic acid (**Figure 3**).

**Figure 3: Calibration curve of Standard Gallic acid.**



The results of total phenolic and flavonoid content of *Corchorus depressus* leaf extracts were presented in (**Table 1**).

**Table 1. Total Phenolic and Flavonoid content of *C. depressus* leaf extracts expressed as (mg/100g).**


<b>Leaf Extracts of <i>C. depressus</i></b>	<b>Total Phenolics (mg/100g) in GAE</b>	<b>Total Flavonoids (mg/100g) in QE</b>
Aqueous	18.18	1.80
Methanolic	78.46	21.2
Pet Ether	0.95	0.11
Chloroform	4.14	0.54
Ethyl acetate	7.2	0.99

#### **The DPPH scavenging activity**

Anti-oxidant reacts with DPPH, which is a nitrogen-centered radical with a characteristic absorption at 517 nm and convert it to 1, 1-diphenyl-2- picryl hydrazine, due to its hydrogen donating ability at a very rapid rate. The degree of discoloration indicates the scavenging potential of the anti-oxidant [22-23]. The extracts and the standard ascorbic acid were tested from lower to higher concentration (20-100 µg/ml). All the extracts & the ascorbic acid exhibited DPPH radical scavenging activity in a concentration dependant manner. Among the extracts, the methanolic leaf extract of *C. depressus* (MECD) showed a mean percentage inhibition of (71.6%) at the concentration of 100µg/ml; whereas the mean percentage inhibition for the aqueous leaf extract was found to be of 31.1% at 100 µg/ml. The results were well compared to standard drug ascorbic acid, which showed the highest mean percentage of

inhibition (84.09%) at the concentration of 100 $\mu$ g/ml. A graph between the mean percentage inhibitions of the DPPH radical scavenging activity of ascorbic acid, MECD & AECD were presented as (Figure 4).

**Figure 4. DPPH radical Scavenging Activity of Ascorbic acid, MECD & AECD.**




### Hydroxyl radical scavenging activity

The test extracts MECD & AECD along with the standard ascorbic acid, suppressed hydroxyl radical mediated deoxyribose degradation in a concentration dependant manner. The hydroxyl radical is a highly potent oxidant that reacts with almost all biomolecules found in the living cells [24]. When it reacts with poly unsaturated fatty acid moieties of cell membrane phospholipids, lipid hydroperoxide is produced. Lipid hydroperoxide can be decomposed to alkoxyl and peroxy radical and numerous carbonyl products such as malondialdehyde (MDA). The carbonyl products are responsible for DNA damage, generation of cancer, respiratory and aging related diseases [25, 26]. The hydroxyl radical scavenging activity of the

methanolic leaf extract of *C. depressus* (MECD) showed 56.31% at 100 $\mu$ g/ml, which is higher in comparison to the aqueous extract (AECD) which showed 22.5 % at 100 $\mu$ g/ml. However the ascorbic acid showed the highest mean percentage of hydroxyl radical scavenging activity of 71.04% at 100 $\mu$ g/ml. A graph showing the mean percentage inhibition of hydroxyl radical scavenging activity of the standard Ascorbic acid, the MECD & AECD was depicted in (Figure 5).

**Figure 5. Evaluation of Hydroxyl radical scavenging activity of Ascorbic acid, MECD and AECD.**




#### **Nitric oxide scavenging assay:**

Nitric oxide (NO) is a diffusible free radical which plays many roles as an effector molecule in diverse biological systems, including neuronal messenger, vasodilation, anti-microbial and anti-tumor activities [27, 28]. Studies in animal models have suggested a role for NO in the pathogenesis of inflammation and pain. NOS inhibitors have been shown to have beneficial effects on some aspects of inflammation and tissue changes seen in models of inflammatory bowel disease [29]. The mean percentage inhibition of the nitric oxide scavenging activity of methanolic leaf extract of *C. depressus* was 58.44 % at the concentration of 100 $\mu$ g/ml. This

value was higher in comparison to the mean percentage inhibition of aqueous leaf extract (AECD) of the same plant which showed 27.99 % at 100 $\mu$ g/ml. However the standard ascorbic acid showed highest mean percentage inhibition (74.07 %) at the concentration of 100  $\mu$ g/ml (Figure 6).

**Figure 6. Nitric oxide radical scavenging activity of Ascorbic acid, MECD and AECD.**



#### 4. CONCLUSION

Normally free radicals of different forms are generated at a low level in cells to help in modulation of several physiological functions and are quenched by an integrated anti-oxidant system in the body. However if the free radicals are produced in excess amount, they can be destructive, leading to generation of various types of diseases including diabetes, causing delay in the healing of wound and its repair. Therefore in the current study, different extracts of *C. depressus* were studied for their possible anti-oxidant activity by DPPH radical, Hydroxyl & Nitrous oxide scavenging activity at different concentrations. Among all, the methanolic leaf

extract of *C. depressus* (MECD) showed better anti-oxidant activity in comparison to the aqueous extract which may be due to the presence of saponins and flavonoids in the methanolic extract. Thus the methanolic leaf extract of the plant have the potency for treating oxidative stress during different diseases and may be responsible for different biological activity of the plant.

## CONSENT

All the authors have given their consent for submission of the manuscript to the esteemed Journal.

## ETHICAL APPROVAL

It is not applicable for this article.

## COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## REFERENCES:

1. Dringen R. Glutathione metabolism and oxidative stress in neurodegeneration. European J. Bio Chem. 2000; 267(16):4903.
2. Schulz JB, Lindnau J, Seyfried J, Dichgans J. Glutathione oxidative stress and neurodegeneration. European J. of Biochemistry. 2000; 267(16):4904-4911.

3. De Silva MC, Paiva SR. Antioxidant activity and flavonoid content of *Clusia fluminensis* Planch. & Triana. Ann. Braz. Academy of Sciences. 2012; 84(3):609-616.
4. Pavithra GM, Saba S, Naik AS, Kekuda PTR, Vinayaka KS. Anti oxidant and antimicrobial activity of flowers of *Wendlandia thyroidea*, *Olea dioica*, *Lagerstroemia speciosa* and *Bombax malabaricum*. Journal of Applied Ph. Sciences. 2013; 3(6):114-120.
5. Glucin I, Topal F, Sarikaya SBO, Bursal E, Bilsel G, Goren AC. Polyphenol contents and antioxidant properties of Medlar (*Mespilus germanica* L.). Records of Natural Products. 2011; 5(3):158-175.
6. Upadhyay B, Parveen AK, Dharker AK. Ethnomedicinal and ethnopharmaco-statistical studies of Eastern Rajasthan, India. Journal of Ethnopharmacology. 2010; 129(1):64-86. PMID: 20214972
7. Jain A, Katewa SS, Chaudhary BL, Galav P. Folk herbal medicines used in birth control and sexual diseases by tribals of southern Rajasthan, India. Journal of Ethnopharmacology. 2004; 90(1):171-177. PMID: 14698527
8. The Wealth of India, A Dictionary of Indian Raw Materials & Industrial Products, Raw Materials, Vol. 10, New Delhi: (National Institute of Science Communication and Information Resources (CSIR); 1993.
9. Ikram M, Khattak SG, Gilani SN. Antipyretic studies on some indigenous Pakistani Medicinal Plants II, Journal of Ethnopharmacology. 1987; 19(2):185-192. PMID: 3497307
10. Pareek A, Godavarthi A, Nagori BP. Invitro hepatoprotective activity of *Corchorus depressus* L. against CCl<sub>4</sub> induced toxicity in HepG<sub>2</sub> cell line. Pharmacognosy Journal. 2013; 5(4):191-195.

11. Kataria S, Kar D, Rao KS, Khajuria RK. *Invitro* and *invivo* aphrodisiac properties of *Corchorus depressus* Linn. on rabbit *Corpus cavernosum* smooth muscle relaxation and sexual behavior of normal male rats. Journal of Ethnopharmacology. 2013;148(1):210-217. PMID: 23612424
12. Anonymous, World Health Organization, Geneva, Quality control methods for Medicinal plant materials, 1998.
13. Anonymous. The Indian Pharmacopoeia. New Delhi: Govt. of India publication; 1966.
14. Cujic N, Savikin K, Jankovic T, Pljevljakusic D, Zdunic G, Ibric S. Optimization of polyphenols extraction from dried chokeberry using maceration as traditional technique. Food Chemistry, 2016; 194:135-142. PMID: 26471536
15. Wager H, Bladt S, Zgainski EM. Plant Drug Analysis. Berlin, Germany: Springer- Verlag, 1984.
16. Odebiyi OO, Sofowora EA. Phytochemical screening of Nigerian Medicinal plants II. Lloydia, 1978; 41(3):234-246. PMID: 672462
17. Trease GE, Evans WC. A Text Book of Pharmacognosy: ELSB Baillere Tindal, Oxford, 1987.
18. Chang CC, Yang MH, Chen JC. Estimation of total flavonoid content in Propolis by two complementary colorimetric methods. J Food Drug Anal. 2002; 10(3): 178-182.
19. Siger A, Nogala-Kalucka M, Lampart SE. The content and antioxidant activity of phenolic compounds in cold-pressed plant oils. J Food Lipids. 2008; 15 (2):137-149.
20. Junaid S, Rakesh KN, Dileep N, Poornima G, Kekuda TRP, Mukunda S. Total phenolic content and antioxidant activity of seed extract of *Lagerstroemia speciosa* L. Chemical Science Transactions. 2013;2(1):75-80.



21. Ganu GP, Jadhav SS, Deshpande AD. Anti-oxidant and antihyperglycemic potential of methanolic extract of bark of *Mimusops elengi* in mice. *Int. J. Phytomedicine*. 2010; 2:116-123.
22. Kitima S, Souksanh N, Yupaporn S. Paper-based DPPH Assay for Antioxidant Activity. *Anal Sci*. 2018; 34(7):795-800. PMID: 29998961.
23. Huang D, Boxin O, Ronald LP. The Chemistry behind antioxidant capacity assay. *J Agric Food Chem*. 2005; 53(6): 1841-1856. PMID: 15769103
24. Fejer J, Ivan K, Grulova D, Eliasova A. Seasonal variability of *Juniperus communis* L. Berry Ethanol Extracts: 1. *Invitro* Hydroxyl Radical Scavenging Activity. *Molecules*. 2020; 25(18):4114. PMID: 32916817
25. Gutowski M, Kowal Czyk S. A study of free radical chemistry: Their role and pathophysiological significance. *Acta Biochem. Polonica*. 2013;60:1-16. PMID: 23513192.
26. Li Q, Sun X, Gu G, Guo Z. Novel water soluble chitosan Derivatives with 1,2,3-Triazolium and their Free radical-scavenging Activity. *Mar Drugs*. 2018; 16(4):107. PMID: PMC5923394.
27. Khan FZ, Dervan E, Bhattacharyya DD, McAuliffe JD, Miranda KM, Glynn SA. The role of Nitric Oxide in cancer: Master Regulator or not?. *Int. J Mol Sci*. 2020; 21(24):9393. PMID: PMC7763974
28. Oleson BJ, Corbett JA. Dual role of Nitric oxide in regulating response of  $\beta$ -cells to DNA damage. *Antioxidant Redox Signal*. 2018; 29(14):1432-1445. PMID: PMC6166691.

29. Vishwakarma A, Wany A, Pandey S, Bulle M, Kumari A, Reddy K, Igamberdiev AU, Mur LAJ, Gupta KJ. Current approaches to measure nitric oxide in Plants. J Exp Bot. 2019; 70(17):4333-4343. PMID: PMC6736158.

UNDR PEER REVIEW