

ANTICARIOGENIC ACTIVITY OF SELENIUM NANOPARTICLES WITH *Pterocarpus Santa*.

RUNNING TITLE : Anticariogenic activity of selenium nanoparticles with *Pterocarpus santa*.

ABSTRACT:

Introduction: Nanoparticles (NPs) are used to minimise toxicity, increase bioactivity, improve targeting, and monitor the release profile of the encapsulated moiety in a variety of ways. Inorganic NPs such as Ag, Au, Ce, Fe, Se, Ti, and Zn hold a special position among NPs due to its versatile bioactivities in nanoforms. Selenium (Se) is an essential trace mineral. SeNPs have been studied in the treatment of oxidative stress and inflammation-related diseases such as arthritis, cancer, diabetes, and nephropathy. Medicinal plants have long been a valuable source of natural active constituents that have been used in products for preserving human health. *Pterocarpus santalinus* (Fabaceae) has been used as a folk medicine and has been shown to have anti-inflammatory, anti-ulcer, and anti-cancer properties. The aim of the present work is to evaluate synergistic Anticariogenic properties of *Pterocarpus santalinus* (red sandalwood).

Materials and methods Anticariogenic activity of respective nanoparticles against the strain staphylococcus aureus, Candida albicans, and Enterococcus faecalis and s.mutans . MHA agar was utilized for this activity to determine the zone of inhibition. Muller Hinton agar was prepared and sterilized for 45 minutes at 120lbs. Media poured into the sterilised plates and let them stabilize for solidification. The wells were cut using the well cutter and the test organisms were swabbed. The nanoparticles with different concentrations were loaded and the plates were incubated for 24 hours at 37 ° C. After the incubation time the zone of inhibition was measured.

Results: Agar well diffusion demonstrated that *Pterocarpus santa* significantly inhibited the growth of *S.Aureus*, *S.mutans*, *E. faecalis* and *C.albicans* . The increase in concentration increases the zone of inhibition. Data were implied as mean \pm SEM with the level of statistical significance at $p < 0.05$.

Conclusion : Selenium nanoparticles were synthesised from *Pterocarpus santa* and they are characterised .Their special study showed the potent anticariogenic activity against pathogens.

Key words: Anticariogenic activity; selenium nanoparticles; *Pterocarpus selenium* ;.

INTRODUCTION:

Nanotechnology is the study and control of matter with dimensions of 1–100 nanometers. The use of precisely designed materials at this length scale to establish novel therapeutic and diagnostic modalities is known as nanomedicine, which is the application of nanotechnology to medicine . Nanomaterials have unique physicochemical properties that distinguish them from bulk materials of the same composition, such as ultra-small scale, large surface area to mass ratio, and high reactivity. These features can be used to get around some of the disadvantages of conventional therapeutic and diagnostic agents (1).The promise of breakthrough developments in medicine, electronics, genomics, and robotics fuelled the growth of nanotechnology (2).Current diagnostic and treatment modalities for a variety of diseases, especially cancer, have significant flaws such as low sensitivity or precision, as well as drug toxicity. On the basis of nanoparticles, new and improved cancer detection methods are being developed (3). They're used as contrast agents, fluorescent products, molecular testing instruments, and medications that have antibodies that target them.Magnetic nanoparticles, quantum dots, nanoshells, and nanosomes are only a few examples of nanoparticles used in diagnostics (4). With the help of nanotechnology, medicines with a high toxic potential, such as cancer chemotherapeutic drugs, can be provided with a better safety profile (5).

Nanoparticles (NPs) are used to minimise toxicity, increase bioactivity, improve targeting, and monitor the release profile of the encapsulated moiety in a variety of ways. Inorganic NPs such as Ag, Au, Ce, Fe, Se, Ti, and Zn hold a special position among NPs due to its versatile bioactivities in nanoforms (6).Particulate systems, such as nanoparticles, have been used to modify and enhance the pharmacokinetic and pharmacodynamic properties of different drug molecules utilising a physical method (7). (Se) is an essential trace mineral(8). As

selenocysteine, it is integrated into selenoproteins and serves as the most essential component of the active core of their enzymatic activities. Many selenoproteins have oxidoreductase activity and thus control the redox balance in the body (8). Se has a small therapeutic window and fragile toxicity margins, whereas Se nanoparticles (SeNPs) have significantly reduced toxicity (6,8). SeNPs have been studied in the treatment of oxidative stress and inflammation-related diseases such as arthritis, cancer, diabetes, and nephropathy (9). Selenium is commonly used as a dietary supplement, owing to its links to immunity and cancer prevention. Heavy metals and selenium have a well-known relationship (10). SeNPs are an appealing carrier vehicle for transporting different drugs to the site of operation (11). Seleno-compounds can be used as chemopreventive and chemotherapeutic agents. SeNPs play a key role in bacterial growth inhibition at very low protein concentrations, while a large amount of protein is needed to inhibit bacterial growth individually (12).

Medicinal plants have long been a valuable source of natural active constituents that have been used in products for preserving human health and treating a wide range of diseases. *Pterocarpus santalinus* wood is in high demand and earns a lot of money in foreign exchange. The species is native to Andhra Pradesh, restricted to a small region in the districts of Cuddapah, Kurnool, and Chittoor. The tract's ecological features include Algonkian slate and quartzite geological terrain, as well as a rainfall regime that is halfway between the tropical form of the interior Deccan and the Coromandel-coast (13). Scientific name: *P. santalinus* Linn. f. belongs to fabaceae family. *Pterocarpus santalinus* (Fabaceae) has been used in Korea as a folk medicine and has been shown to have anti-inflammatory, anti-ulcer, and anti-cancer properties (14). Inflammations, emotional aberrations, ulcers, and cancer have all been treated with this herb as a folk remedy. Isoflavonoids, terpenoids, aurone glycosides, and lignans were discovered in previous phytochemical studies on the *Pterocarpus santalinus* (14). we had previously done research on these topics for the past 5 year (red sandalwood) (15–23). Our team has extensive knowledge and research experience that has translate into high quality publications(24–28),(29),(30),(31),(32),(33),(34),(26,35,36),(37–41),(42),(43). The aim of the present work is to evaluate synergistic Anticariogenic properties of *Pterocarpus santalinus*.

MATERIALS AND METHODS:

Synthesis of selenium nanoparticles

Selenium nanoparticles were made by measuring 0.5g of red sandal selenium and adding 50 ml of distilled water to it. The extract is now boiled for 10 minutes at 55 degrees Celsius. The extract was then purified until sodium selenite was added.

Antibacterial Activity

Anticariogenic activity of respective nanoparticles against the strain *Staphylococcus aureus*, *Candida albicans*, and *Enterococcus faecalis* and *S.mutans*. MHA agar was utilized for this activity to determine the zone of inhibition. Muller Hinton agar was prepared and sterilized for 45 minutes at 120lbs. Media poured into the sterilised plates and let it stable for solidification. The wells were cut using the well cutter and the test organisms were swabbed. The nanoparticles with different concentration were loaded and the plates were incubated for 24 hours at 37 ° C. After the incubation time the zone of inhibition was measured. Data were implied as mean \pm SEM with the level of statistical significance at $p < 0.05$.

RESULTS :




FIGURE 1: Green synthesis of *pterocarpus santa* mediated selenium nanoparticles .




FIGURE - 2 : Image showing the Zone of Inhibition of *P.santa* mediated selenium nanoparticle




FIGURE 3 : The bar graph depicts the Anticariogenic activity of selenium nanoparticles synthesised from the extract of *Pterocarpus santalinus*. x axis represents the concentration and y axis represents the zone of inhibition in mm, data implies as mean \pm SEM

The zone of inhibition for *E. faecalis* at 25µl is 22, the zone of inhibition of 50 µl is 21, the zone of inhibition of 100µl is 20, the zone of inhibition of standard is 40 .The zone of inhibition for *S. mutans* at 25µl is 20, the zone of inhibition of 50 µl is 21, the zone of inhibition of 100µl is 22,

the zone of inhibition of standard is 30. The zone of inhibition for *S. aureus* at 25 µl is 20, the zone of inhibition of 50 µl is 21, the zone of inhibition of 100 µl is 25, the zone of inhibition of standard is 27. The zone of inhibition for *C. albicans* at 25 µl is 20, the zone of inhibition of 50 µl is 22, the zone of inhibition of 100 µl is 25, the zone of inhibition of standard is 15. Among all the bacteria and fungi the zone of inhibition of *C. albicans* is good. As the concentration increases the zone of inhibition also increases.

DISCUSSION:

Pterocarpus santa along with selenium nanoparticles has a positive outcome in the present study. It has also shown that it shows a great effect on anticariogenic activity. Previous studies have been done using *P. santalinus* L. has long been used to treat diabetes and the heartwood extract has been shown to have antioxidant, anti-inflammatory, anti-cancer, gastro-protective and wound-healing properties. The antioxidant potential of methanol extracts from bark, and wood of *P. santalinus* L. was very evident. The powder made from the wood of *P. santalinus* L. could speed wound healing without causing any negative side effects (44). Previous article stated that the yields of *Pterocarpus macrocarpus* and *Pterocarpus soyauxii* were particularly high among the ethanol extracts, with 28.39 percent and 28.59 percent, respectively (45,46). At 2 g/L, 4 g/L, and 2 g/L, respectively, *Pterocarpus angolensis*, *Pterocarpus macrocarpus*, and *Pterocarpus soyauxii* achieved grade I corrosion resistance (46).

Selenium nanoparticles can be made utilizing a variety of processes, including biological, chemical, and physical processes (47). The manufacture of selenium nanoparticles from biological sources such as bacteria, fungus, yeasts, and plants has been reported in a number of works (47). Chemicals have also been employed to convert sodium selenite into selenium nanoparticles as a reducing agent. TEM, SEM, FTIR, XRD, and UV-vis are used to analyze synthesized selenium nanoparticles (47). Nanotechnology advancements have enabled the creation of customized metal/metalloid nanoparticles with physicochemical characteristics that

inhibit bacteria. Existing drug resistance mechanisms, such as slow drug uptake and accelerated efflux, biofilm formation, and intracellular bacterial parasitism, have been shown to be overcome by these nanoparticles (48). SeNPs have antibacterial, antiviral, and antioxidant activities, indicating that they might be used to treat infectious illnesses as therapeutic possibilities (48). Nanostructured particles, in particular, may be produced utilizing bacterial and fungal cells as biological catalysts, resulting in a non-toxic and ecologically friendly method of producing nanoparticles, such as SeNPs (48). From the previous article, Lampis described SeNPs as having a typical spherical form and a diameter of 50–400 nm. Furthermore, biogenic SeNPs have been shown to exhibit anti-biofilm action against clinical isolates of bacterial pathogens in recent investigations (48)

The small margin between therapeutic and hazardous levels of selenium, on the other hand, limits its use in dietary supplements(45). The therapeutic and harmful effects of selenium are strongly tied to its chemical forms, according to a large body of evidence. The redox process has been shown to produce orange-red and zero-valent selenium nanoparticles (SeNPs) with better properties(45). When compared to other chemical species of selenium, orange-red and zero-valent selenium nanoparticles (SeNPs) generated through the redox process exhibit superior bioavailability but lower toxicity. SeNPs, on the other hand, are often unstable and easily aggregate into grey and black elemental selenium(45). Polysaccharides have been employed as templates to increase the stability of SeNPs in previous studies. Fungi have long been known to be rich in bioactive polysaccharides. Fungal polysaccharides have a wide range of biological functions, including gut microbiota control, immunomodulatory, and anti-cancer effects. The limitations of this study was that the zone of inhibition was only tested on a few microbes. Future studies will be done on *Pterocarpus santa* and selenium nanoparticles to implement it in the drugs formulation for clinical trials (49) (50) (51) (52) (53) (54) (55) (56) (57) (58) (59) (60) (61) (62).

CONCLUSION :

Selenium nanoparticles were synthesised from *Pterocarpus santa* and they are characterised. Their special study showed the potent anticariogenic activity against pathogens(63-72). As a

result, SeNPs capped with polysaccharides are predicted to improve the benefits of selenium and may have synergistic activity.

REFERENCE:

1. Zhang L, Gu FX, Chan JM, Wang AZ, Langer RS, Farokhzad OC. Nanoparticles in Medicine: Therapeutic Applications and Developments [Internet]. Vol. 83, Clinical Pharmacology & Therapeutics. 2008. p. 761–9. Available from: <http://dx.doi.org/10.1038/sj.clpt.6100400>
2. Nie S. Nanotechnology for personalized and predictive medicine [Internet]. Vol. 2, Nanomedicine: Nanotechnology, Biology and Medicine. 2006. p. 305. Available from: <http://dx.doi.org/10.1016/j.nano.2006.10.115>
3. Rauta PR, Mohanta YK, Nayak D. Nanotechnology in Biology and Medicine: Research Advancements & Future Perspectives. CRC Press; 2019. 273 p.
4. Bharali DJ, Pradhan V, Elkin G, Qi W, Hutson A, Mousa SA, et al. Novel nanoparticles for the delivery of recombinant hepatitis B vaccine [Internet]. Vol. 4, Nanomedicine: Nanotechnology, Biology and Medicine. 2008. p. 311–7. Available from: <http://dx.doi.org/10.1016/j.nano.2008.05.006>
5. Bawa R. NanoBiotech 2008: Exploring global advances in nanomedicine [Internet]. Vol. 5, Nanomedicine: Nanotechnology, Biology and Medicine. 2009. p. 5–7. Available from: <http://dx.doi.org/10.1016/j.nano.2009.01.004>
6. A.t. K, Keshta AT, Emam M, Attia YA. Effect of Selenium nanoparticles in wound healing [Internet]. Vol. 16, Biochemistry Letters. 2020. p. 160–8. Available from: <http://dx.doi.org/10.21608/blj.2020.146617>
7. Mohanraj VJ, Chen Y. Nanoparticles - A review [Internet]. Vol. 5, Tropical Journal of Pharmaceutical Research. 2007. Available from: <http://dx.doi.org/10.4314/tjpr.v5i1.14634>
8. Vrček IV. Selenium Nanoparticles: Biomedical Applications [Internet]. Molecular and Integrative Toxicology. 2018. p. 393–412. Available from: http://dx.doi.org/10.1007/978-3-319-95390-8_21
9. Kontoghiorghes GJ. Chemical, pharmacological, toxicological and therapeutic advances of orally active iron and aluminium chelators. deieriprone (11) [Internet]. Vol. 78, Toxicology Letters. 1995. p. 7. Available from: [http://dx.doi.org/10.1016/0378-4274\(95\)94626-r](http://dx.doi.org/10.1016/0378-4274(95)94626-r)
10. (efsa) EFSA, European Food Safety Authority (EFSA). L-selenomethionine as a source of selenium added for nutritional purposes to food supplements [Internet]. Vol. 7, EFSA Journal. 2009. p. 1082. Available from: <http://dx.doi.org/10.2903/j.efsa.2009.1082>
11. Xiao Y, Huang Q, Zheng Z, Ma H. Selenium release kinetics and mechanism from

Cordyceps sinensis exopolysaccharide-selenium composite nanoparticles in simulated gastrointestinal conditions. *Food Chem.* 2021 Feb 6;350:129223.

12. Rangrazi A, Bagheri H, Ghazvini K, Boruziniat A, Darroudi M. Synthesis and antibacterial activity of colloidal selenium nanoparticles in chitosan solution: a new antibacterial agent [Internet]. Vol. 6, *Materials Research Express*. 2020. p. 1250h3. Available from: <http://dx.doi.org/10.1088/2053-1591/ab6a56>
13. Kukrety S, Dwivedi P, Jose S, Alavalapati JRR. Stakeholders' perceptions on developing sustainable Red Sanders (*Pterocarpus santalinus* L.) wood trade in Andhra Pradesh, India [Internet]. Vol. 26, *Forest Policy and Economics*. 2013. p. 43–53. Available from: <http://dx.doi.org/10.1016/j.forpol.2012.08.014>
14. Kwon HJ, Hong YK, Kim KH, Han CH, Cho SH, Choi JS, et al. Methanolic extract of *Pterocarpus santalinus* induces apoptosis in HeLa cells [Internet]. Vol. 105, *Journal of Ethnopharmacology*. 2006. p. 229–34. Available from: <http://dx.doi.org/10.1016/j.jep.2005.10.025>
15. Shankar SB, Barani Shankar S, Arivarasu L, Rajeshkumar S. Biosynthesis of Hydroxy Citric Acid Mediated Zinc Nanoparticles and Its Antioxidant and Cytotoxic Activity [Internet]. *Journal of Pharmaceutical Research International*. 2020. p. 108–12. Available from: <http://dx.doi.org/10.9734/jpri/2020/v32i2630845>
16. Karthik V, Arivarasu L, Rajeshkumar S. Hyaluronic Acid Mediated Zinc Nanoparticles against Oral Pathogens and Its Cytotoxic Potential [Internet]. *Journal of Pharmaceutical Research International*. 2020. p. 113–7. Available from: <http://dx.doi.org/10.9734/jpri/2020/v32i1930716>
17. J A, Aparna J, Rajeshkumar S. Cytotoxic and Antioxidant Activity of Zinc Oxide Nanoparticles Synthesised Using *Maranta Arundinacea* Root Extract [Internet]. Vol. 11, *International Journal of Research in Pharmaceutical Sciences*. 2020. p. 4372–7. Available from: <http://dx.doi.org/10.26452/ijrps.v11i3.2655>
18. Jaisankar AI, Arivarasu L. Free Radical Scavenging and Anti-Inflammatory Activity of Chlorogenic Acid Mediated Silver Nanoparticle [Internet]. *Journal of Pharmaceutical Research International*. 2020. p. 106–12. Available from: <http://dx.doi.org/10.9734/jpri/2020/v32i1930715>
19. Shree MK, Kavya Shree M, Arivarasu L, Rajeshkumar S. Cytotoxicity and Antimicrobial Activity of Chromium Picolinate Mediated Zinc Oxide Nanoparticle [Internet]. *Journal of Pharmaceutical Research International*. 2020. p. 28–32. Available from: <http://dx.doi.org/10.9734/jpri/2020/v32i2030726>
20. Aathira CM, Arivarasu L, Rajeshkumar S. Antioxidant and Anti-Inflammatory Potential of Chromium Picolinate Mediated Zinc Oxide Nanoparticle [Internet]. *Journal of Pharmaceutical Research International*. 2020. p. 118–21. Available from: <http://dx.doi.org/10.9734/jpri/2020/v32i1930717>

21. Akash N, Arivarasu L, Rajeshkumar S. Anti-inflammatory and Antioxidant Potential of Hyaluronic Acid Mediated Zinc Nanoparticles [Internet]. *Journal of Pharmaceutical Research International*. 2020. p. 33–7. Available from: <http://dx.doi.org/10.9734/jpri/2020/v32i2030727>
22. Niveditha AS, Sankari Niveditha A, Geetha RV, Arivarasu L. Will Alternative Medicine Help Us to Fight Against COVID-19 [Internet]. *International Journal of Current Research and Review*. 2020. p. 112–6. Available from: <http://dx.doi.org/10.31782/ijcrr.2020.sp47>
23. Devaraj E, Roy A, Veeraragavan GR, Magesh A, Sreeba AV, Arivarasu L, et al. β -Sitosterol attenuates carbon tetrachloride-induced oxidative stress and chronic liver injury in rats [Internet]. Vol. 393, *Naunyn-Schmiedeberg's Archives of Pharmacology*. 2020. p. 1067–75. Available from: <http://dx.doi.org/10.1007/s00210-020-01810-8>
24. Rajeshkumar S, Kumar SV, Ramaiah A, Agarwal H, Lakshmi T, Roopan SM. Biosynthesis of zinc oxide nanoparticles using *Mangifera indica* leaves and evaluation of their antioxidant and cytotoxic properties in lung cancer (A549) cells. *Enzyme Microb Technol*. 2018 Oct;117:91–5.
25. Nandhini NT, Rajeshkumar S, Mythili S. The possible mechanism of eco-friendly synthesized nanoparticles on hazardous dyes degradation. *Biocatal Agric Biotechnol*. 2019 May 1;19:101138.
26. Vairavel M, Devaraj E, Shanmugam R. An eco-friendly synthesis of *Enterococcus* sp.–mediated gold nanoparticle induces cytotoxicity in human colorectal cancer cells. *Environ Sci Pollut Res*. 2020 Mar 1;27(8):8166–75.
27. Gomathi M, Prakasam A, Rajkumar PV, Rajeshkumar S, Chandrasekaran R, Anbarasan PM. Green synthesis of silver nanoparticles using *Gymnema sylvestre* leaf extract and evaluation of its antibacterial activity [Internet]. Vol. 32, *South African Journal of Chemical Engineering*. 2020. p. 1–4. Available from: <http://dx.doi.org/10.1016/j.sajce.2019.11.005>
28. Rajasekaran S, Damodharan D, Gopal K, Rajesh Kumar B, De Poures MV. Collective influence of 1-decanol addition, injection pressure and EGR on diesel engine characteristics fueled with diesel/LDPE oil blends. *Fuel*. 2020 Oct 1;277:118166.
29. Santhoshkumar J, Sowmya B, Venkat Kumar S, Rajeshkumar S. Toxicology evaluation and antidermatophytic activity of silver nanoparticles synthesized using leaf extract of *Passiflora caerulea*. *S Afr J Chem Eng*. 2019 Jul;29:17–23.
30. Raj R K, D E, S R. β -Sitosterol-assisted silver nanoparticles activates Nrf2 and triggers mitochondrial apoptosis via oxidative stress in human hepatocellular cancer cell line. *J Biomed Mater Res A*. 2020 Sep;108(9):1899–908.
31. Saravanan M, Arokiyaraj S, Lakshmi T, Pugazhendhi A. Synthesis of silver nanoparticles from *Phanerochaete chrysosporium* (MTCC-787) and their antibacterial activity against human pathogenic bacteria. *Microb Pathog*. 2018 Apr;117:68–72.

32. Gheena S, Ezhilarasan D. Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells. *Hum Exp Toxicol*. 2019 Jun 1;38(6):694–702.
33. Ezhilarasan D, Sokal E, Najimi M. Hepatic fibrosis: It is time to go with hepatic stellate cell-specific therapeutic targets. *Hepatobiliary Pancreat Dis Int*. 2018 Jun;17(3):192–7.
34. Ezhilarasan D. Oxidative stress is bane in chronic liver diseases: Clinical and experimental perspective. *Arab J Gastroenterol*. 2018 Jun;19(2):56–64.
35. Gomathi AC, Xavier Rajarathinam SR, Mohammed Sadiq A, Rajeshkumar S. Anticancer activity of silver nanoparticles synthesized using aqueous fruit shell extract of *Tamarindus indica* on MCF-7 human breast cancer cell line. *J Drug Deliv Sci Technol*. 2020 Feb 1;55:101376.
36. Dua K, Wadhwa R, Singhvi G, Rapalli V, Shukla SD, Shastri MD, et al. The potential of siRNA based drug delivery in respiratory disorders: Recent advances and progress. *Drug Dev Res*. 2019 Sep;80(6):714–30.
37. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. *J Periodontol*. 2018 Oct;89(10):1241–8.
38. Arumugam P, George R, Jayaseelan VP. Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma. *Arch Oral Biol*. 2021 Feb;122:105030.
39. Joseph B, Prasanth CS. Is photodynamic therapy a viable antiviral weapon against COVID-19 in dentistry? *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2021 Jul;132(1):118–9.
40. Ezhilarasan D, Apoorva VS, Ashok Vardhan N. *Syzygium cumini* extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. *J Oral Pathol Med*. 2019 Feb;48(2):115–21.
41. Duraisamy R, Krishnan CS, Ramasubramanian H, Sampathkumar J, Mariappan S, Navarasampatti Sivaprakasam A. Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant-Abutment Interface, With Original and Nonoriginal Abutments. *Implant Dent*. 2019 Jun;28(3):289–95.
42. Gnanavel V, Roopan SM, Rajeshkumar S. Aquaculture: An overview of chemical ecology of seaweeds (food species) in natural products. *Aquaculture*. 2019 May 30;507:1–6.
43. Markov A, Thangavelu L, Aravindhan S, Zekiy AO, Jarahian M, Chartrand MS, et al. Mesenchymal stem/stromal cells as a valuable source for the treatment of immune-mediated disorders. *Stem Cell Res Ther*. 2021 Mar 18;12(1):192.
44. Gao W, Lin P, Hwang E, Wang Y, Yan Z, Ngo HTT, et al. *Pterocarpus santalinus* L. Regulated Ultraviolet B Irradiation-induced Procollagen Reduction and Matrix Metalloproteinases Expression Through Activation of TGF- β /Smad and Inhibition of the

- MAPK/AP-1 Pathway in Normal Human Dermal Fibroblasts [Internet]. Vol. 94, Photochemistry and Photobiology. 2018. p. 139–49. Available from: <http://dx.doi.org/10.1111/php.12835>
45. Gao X, Li X, Mu J, Ho C-T, Su J, Zhang Y, et al. Preparation, physicochemical characterization, and anti-proliferation of selenium nanoparticles stabilized by Polyporus umbellatus polysaccharide [Internet]. Vol. 152, International Journal of Biological Macromolecules. 2020. p. 605–15. Available from: <http://dx.doi.org/10.1016/j.ijbiomac.2020.02.199>
 46. Cai M, Lv H, Cao C, Zhang L, Cao R, Xu B. Evaluation of antimicrobial activity of Pterocarpus extracts [Internet]. Vol. 140, Industrial Crops and Products. 2019. p. 111668. Available from: <http://dx.doi.org/10.1016/j.indcrop.2019.111668>
 47. Rajeshkumar S, Veena P, Santhiyaa RV. Synthesis and Characterization of Selenium Nanoparticles Using Natural Resources and Its Applications [Internet]. Exploring the Realms of Nature for Nanosynthesis. 2018. p. 63–79. Available from: http://dx.doi.org/10.1007/978-3-319-99570-0_4
 48. Cremonini E, Zonaro E, Donini M, Lampis S, Boaretti M, Dusi S, et al. Biogenic selenium nanoparticles: characterization, antimicrobial activity and effects on human dendritic cells and fibroblasts [Internet]. Vol. 9, Microbial Biotechnology. 2016. p. 758–71. Available from: <http://dx.doi.org/10.1111/1751-7915.12374>
 49. Pushpaanjali G, Geetha RV, Lakshmi T. Knowledge and Awareness about Antibiotic Usage and Emerging Drug Resistance Bacteria among Dental Students. Journal of Pharmaceutical Research International. 2020 Aug 24;34–42.
 50. Aathira CM, Geetha RV, Lakshmi T. Knowledge and Awareness about the Mode of Transmission of Vector Borne Diseases among General Public. Journal of Pharmaceutical Research International. 2020 Aug 24;87–96.
 51. Baskar K, Lakshmi T. Knowledge, Attitude and Practices Regarding HPV Vaccination among Undergraduate and Postgraduate Dental Students in Chennai. Journal of Pharmaceutical Research International. 2020 Aug 25;95–100.
 52. Manya Suresh LT. Wound Healing Properties of Aloe Barbadensis Miller-In Vitro Assay. Journal of Complementary Medicine Research. 2020;11(5):30–4.
 53. First Report on Marine Actinobacterial Diversity around Madras Atomic Power Station (MAPS), India [Internet]. [cited 2021 Aug 31]. Available from: <http://alinteridergisi.com/article/first-report-on-marine-actinobacterial-diversity-around-madras-atomic-power-station-maps-india/>
 54. Physicochemical Profile of Acacia Catechu Bark Extract – An in Vitro Stud - International Journal of Pharmaceutical and Phytopharmacological Research [Internet]. [cited 2021 Aug 31]. Available from: <https://eijppr.com/article/physicochemical-profile-of-acacia-catechu-bark-extract-an-in-vitro-stud>

55. Lakshmi T. Antifungal Activity of *Ficus racemosa* Ethanolic Extract against Dermatophytes-An in vitro Study. *Journal of Research in Medical and Dental Science*. 2021;9(2):191–3.
56. Awareness of Drug Abuse among Teenagers - *International Journal of Pharmaceutical and Phytopharmacological Research* [Internet]. [cited 2021 Aug 31]. Available from: <https://eijppr.com/article/awareness-of-drug-abuse-among-teenagers>
57. Mangal CSK, Anitha R, Lakshmi T. Inhibition of Nitric oxide Production and Nitric oxide Synthase Gene Expression in LPS Activated RAW 264 .7 Macrophages by Thyme oleoresin from *Thymus vulgaris*. *J Young Pharm*. 2018;10(4):481.
58. COX2 Inhibitory Activity of *Abutilon Indicum* - *Pharmaceutical Research and Allied Sciences* [Internet]. [cited 2021 Aug 31]. Available from: <https://ijpras.com/article/cox2-inhibitory-activity-of-abutilon-indicum>
59. Jibu RM, Geetha RV, Lakshmi T. Isolation, Detection and Molecular Characterization of *Staphylococcus aureus* from Postoperative Infections. *Journal of Pharmaceutical Research International*. 2020 Aug 24;63–7.
60. Sindhu PK, Thangavelu L, Geetha RV, Rajeshkumar S, Raghunandhakumar S, Roy A. Anorectic drugs: an experimental and clinical perspective ♦A Review. *Journal of Complementary Medicine Research*. 2020;11(5):106–12.
61. Nivethitha R, Thangavelu L, Geetha RV, Anitha R, RajeshKumar S, Raghunandhakumar S. In Vitro Anticancer Effect of *Sesamum Indicum* Extract -. *Journal of Complementary Medicine Research*. 2020;11(5):99–105.
62. Mariona P, Roy A, Lakshmi T. Survey on lifestyle and food habits of patients with PCOS and obesity. *Journal of Complementary Medicine Research*. 2020;11(5):93–8.
63. Rajendran R, Kunjusankaran RN, Sandhya R, Anilkumar A, Santhosh R, Patil SR. Comparative Evaluation of Remineralizing Potential of a Paste Containing Bioactive Glass and a Topical Cream Containing Casein Phosphopeptide-Amorphous Calcium Phosphate: An in Vitro Study. *Pesqui Bras Odontopediatria Clin Integr*. 2019 Mar 12;19(0):4668.
64. Ashok BS, Ajith TA, Sivanesan S. Hypoxia-inducible factors as neuroprotective agent in Alzheimer's disease. *Clin Exp Pharmacol Physiol* [Internet]. 2017 Mar [cited 2021 Sep 15];44(3). Available from: <https://pubmed.ncbi.nlm.nih.gov/28004401/>
65. Malli SN, Selvarasu K, Jk V, Nandakumar M, Selvam D. Concentrated Growth Factors as an Ingenious Biomaterial in Regeneration of Bony Defects after Periapical Surgery: A Report of Two Cases. *Case Rep Dent* [Internet]. 2019 Jan 22 [cited 2021 Sep 15];2019. Available from: <https://pubmed.ncbi.nlm.nih.gov/30805222/>
66. Mohan M, Jagannathan N. Oral field cancerization: an update on current concepts. *Oncol Rev* [Internet]. 2014 Jun 30 [cited 2021 Sep 15];8(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/25992232/>

67. Menon S, Ks SD, R S, S R, Vk S. Selenium nanoparticles: A potent chemotherapeutic agent and an elucidation of its mechanism. *Colloids Surf B Biointerfaces* [Internet]. 2018 Oct 1 [cited 2021 Sep 15];170. Available from: <https://pubmed.ncbi.nlm.nih.gov/29936381/>
68. Samuel SR, Acharya S, Rao JC. School Interventions-based Prevention of Early-Childhood Caries among 3-5-year-old children from very low socioeconomic status: Two-year randomized trial. *J Public Health Dent* [Internet]. 2020 Jan [cited 2021 Sep 15];80(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/31710096/>
69. Praveen K, Narayanan V, Muthusekhar MR, Baig MF. Hypotensive anaesthesia and blood loss in orthognathic surgery: a clinical study. *Br J Oral Maxillofac Surg* [Internet]. 2001 Apr [cited 2021 Sep 15];39(2). Available from: <https://pubmed.ncbi.nlm.nih.gov/11286449/>
70. Neelakantan P, Subbarao C, Subbarao CV, De-Deus G, Zehnder M. The impact of root dentine conditioning on sealing ability and push-out bond strength of an epoxy resin root canal sealer. *Int Endod J* [Internet]. 2011 Jun [cited 2021 Sep 15];44(6). Available from: <https://pubmed.ncbi.nlm.nih.gov/21255047/>
71. Oligonucleotide therapy: An emerging focus area for drug delivery in chronic inflammatory respiratory diseases. *Chem Biol Interact*. 2019 Aug 1;308:206–15.
72. Kumar MS, Vamsi G, Sripriya R, Sehgal PK. Expression of matrix metalloproteinases (MMP-8 and -9) in chronic periodontitis patients with and without diabetes mellitus. *J Periodontol*. 2006 Nov;77(11):1803–8.