

## EVALUATION OF PROAPOPTOTIC POTENTIAL OF *Saraca asoca* FLOWER EXTRACT ON SKIN CANCER CELL LINE

**Running Title:** Assessment of proapoptotic potential of *Saraca asoca* flower extract on skin cancer cell line

### Abstract

**Background:** Cancer persists as a major health issue globally due to its high rate of morbidity and mortality. Skin cancer is the most common cancer and accounts for at least 40% of cancer cases worldwide. Search for tumour-selective and novel anticancer compounds with lesser side effects remains a major focus of cancer research. *Saraca asoca* is a traditional Indian medicinal plant, known to have anti-cancer, anti-menorrhagic, anti-oxidant, anti-oxytocic and anti-microbial activities. Though phytoconstituents of the *Saraca asoca* leaves, bark, and flowers have been reported in few studies, the cytotoxic potential of *Saraca asoca* flowers has not been evaluated.

**Aim:** To evaluate the proapoptotic potential of *Saraca asoca* flower extract on skin cancer cell line.

**Materials and methods:** In this present study, the cytotoxic potential of *Saraca asoca* flower extract (10 to 60 µg/ml) was evaluated by MTT assays in B16-F10 skin cancer cells. According to the MTT assay, we determined the optimal doses (IC-50: 30 µg/ml) which were used for further analyses. Analysis of changes in cell morphology is examined by a phase-contrast microscope. The impacts of *Saraca asoca* in B16-F10 cell death were also determined by AO/EtBr dual staining under a fluorescence microscope.

**Results:** In our study, the cell viability assay results showed that 50% of growth inhibition was observed at 30 µg/ml concentration of *Saraca asoca* flower extract treated B16-F10 cells, which has been taken as an inhibitory concentration (IC-50) dose value and fixed for further experiments. The morphological changes in B16-F10 skin cancer cell line with the treatment of *Saraca asoca* at 30 µg/mL for 24hrs has significantly altered the morphology of B16-F10 cell lines. AO/EtBr dual staining results showed the early apoptotic cells having bright orange areas

of condensed or fragmented chromatin in the nucleus after *Saraca asoca* flower extract treated skin cancer cells.

**Conclusion:** The results of this present study showed that the flower extracts of *Saraca asoca* were cytotoxic and induced apoptosis to the cancer cells at a concentration of 30µg/ml at the 24th-time point.

**Keywords:** *Saraca asoca*, Flower extract, Skin cancer, Cytotoxicity, Apoptosis

## Introduction

Cancer persists as a major health issue globally due to its high rate of morbidity and mortality (1). Skin cancer is the most common cancer and accounts for at least 40% of cancer cases worldwide (2). Skin cancers are cancers that emerge from the skin due to the development of abnormal or mutated cells which can invade or spread to other parts of the body. The main types of skin cancers include basal-cell skin cancer (BCC), squamous-cell skin cancer (SCC) and melanoma (3). The basal-cell skin cancer (BCC), squamous-cell skin cancer (SCC) along with a number of less common skin cancers, are recognized as nonmelanoma skin cancer (NMSC) (4). Non-melanoma skin cancer, which occurs in at least 2–3 million people per year, is the most common type of skin cancer (5). An alarming rate of incidence of skin cancer in India has been seen in recent times (6). The most important reason for the occurrence of skin cancer in India, as well as other countries, is attributed to increased urbanization, increased pollution due to the vehicles, smoke emitted from different types of industries causing the depletion of the ozone layer. These changes result in the passage of ultraviolet rays from the sun into the earth and cause genetic mutations in humans, leading to skin cancer (7).

Developing novel strategies for the management of skin cancer represents a desirable goal due to the increasing rise in the incidence of skin cancer patients throughout the world. Search for tumour-selective and novel anticancer compounds with lesser side effects remains a major focus of cancer research (8). Moreover, the side effects linked with synthetic drugs severely influence the quality of life of patients (9). Also, drugs obtained from natural sources have attained considerable attention worldwide. *Saraca asoca* belongs to the family Caesalpiniaceae is one of the most ancient trees seen in India, and is frequently known as an “Ashoka ” (10). *Saraca asoca*

is a traditional Indian medicinal plant, known to have anti-cancer, anti-menorrhagic, anti-oxidant, anti-oxytotic and anti-microbial activities (11). Researchers have reported that almost every part of the plant is known to have a medicinal property and has a role in ayurvedic medicine for centuries (12). Flowers of *Saraca asoca* have been widely used in the Ayurvedic system of medicine for years extremely owing to its wound healing property. Studies documented the phytochemical constituents of *Saraca asoca* flowers containing oleic, linoleic, palmitic, and stearic acids, quercetin-3-O-P-D-glucoside, P-sitosterol, gallic acid, kaempferol-3-O-P-D-glucoside, p and y sitosterols (13) quercetin, apigenin-7-O-p-D-glucoside, pelargonidin-3,5-diglucoside, cyanidin-3,5-diglucoside, and leucocyanidin (14). The cytotoxic activity of *Saraca asoca* bark and its possible application in cancer prevention has been recently reported. Though phytoconstituents of the *Saraca asoca* leaves, bark, flowers have been reported in few studies the cytotoxic potential of *Saraca asoca* flowers has not been evaluated.

Apoptosis means the ability of a cell to cause self-destruction by the activation of intrinsic cellular suicidal programs when the cells are no longer needed, genetically mutated or when they are extremely damaged (15). Induction of apoptosis in tumour cells is the most established anticancer mechanism and employed in many cancer therapies (16). Human cancer cell lines have been the most commonly used as experimental models in research as they can retain the characteristic features of cancer cells, purity, easily assessed and can be manipulated genetically to contribute reproducible findings (17). Outcomes obtained with cell lines are often used to understand human tumours in vivo. Our team has extensive knowledge and research experience that has translated into high quality publications (18–38). With this background, our study aims in evaluating the proapoptotic potential of *Saraca asoca* flower extract on skin cancer cell lines (39) (40) (41) (42).

## **Materials and methods**

### **Reagents used:**

DMEM (Dulbecco's Modified Eagle Medium), Phosphate Buffered Saline (PBS), Trypsin-EDTA, Fetal bovine serum (FBS), were purchased from Gibco, Canada. Acridine orange (AO), ethidium bromide (EtBr), Dimethyl sulfoxide (DMSO), [3-(4,5-dimethylthiazol-2-yl) 2,5-

diphenyl tetrazolium bromide (MTT), AO/EtBr were purchased from Sigma Chemical Pvt Ltd, USA. All other chemicals used were extra pure of molecular grade and were obtained from SRL, India.

#### **Cell line maintenance:**

Skin cancer cell lines B16-F10 were acquired from the NCCS, Pune. After that, the cells were grown in T25 culture flasks containing DMEM supplemented with 10% FBS and 1% antibiotics.

#### **Preparation of the Herbal Extract:**

*Saraca asoca* flower powder acquired from IMPCOPS (Chennai, India) was used for this current study. 50g of the *Saraca asoca* flower powder was soaked in 500ml of 95% ethanol and was maintained at room temperature for 3 days under static condition.

#### **Cell viability (MTT) assay:**

The cytotoxic potential of *Saraca asoca* flower extract treated with B16-F10 skin cancer cells was evaluated by using MTT assays. The principle of this assay is based on the reduction of soluble yellow tetrazolium salt to insoluble purple formazan crystals by metabolically active cells. The concentration of  $2 \times 10^4$  cells/well was plated in 48 well plates and incubated for 24 hours and further starved by incubating it in a serum-free medium for 3 hours at a temperature of 37°C. Later then the cells were treated with *Saraca asoca* flower extract at different concentrations (10 to 60µg/ml) for 24 hours. The MTT containing medium was then discarded and the cells were washed with 1x PBS. The crystals were then dissolved by adding 200µl of solubilization solution and this was mixed appropriately by pipetting up and down. Then the formazan crystals formed were dissolved in dimethyl sulfoxide (200µl) and incubated in dark for an hour. The intensity of the colour developed was assayed with the help of a Micro ELISA plate reader at 570nm. The number of viable cells was expressed as the percentage of control cells cultured in a serum-free medium. The cell viability is calculated using the following formula: % cell viability =  $[A_{570\text{nm}} \text{ of treated cells} / A_{570 \text{ nm}} \text{ of control cells}] \times 100$ .

#### **Morphology study:**

According to the MTT assay, we determined the optimal doses (IC-50: 30µg/ml) which were used for further analyses. Analysis of changes in cell morphology is examined by a phase-contrast microscope.  $3 \times 10^4$  cells were seeded in 6 well plates and treated with *Saraca asoca* (30µg/ml for B16-F10 cells) for an interval of 24h. After the incubation period, the medium was removed and the cells were washed once by phosphate buffer saline (PBS pH 7.4). The obtained plates were observed under a phase-contrast microscope.

#### **Determination of mode of cell death by acridine orange (AO)/ethidium bromide (EtBr) dual staining:**

The impacts of *Saraca asoca* in B16-F10 cell death were also determined by AO/EtBr dual staining. The cells were treated with *Saraca asoca* flower extract for a duration of 24 h and further, the cells were harvested. The pellets were resuspended in acridine orange and ethidium bromide dual staining and the cell death was examined under a fluorescence microscope.

#### **Statistical analysis**

Statistical analyses were executed using one-way ANOVA followed by Student–Newman–Keuls (SNK) tests for comparison between treatment values and control values. Data were implied as mean  $\pm$  SEM with the level of statistical significance at  $p < 0.05$ .

#### **Results**

##### **The cytotoxic potential of *Saraca asoca* flower extract on the B16 - F10 skin cancer cell line was determined by using an MTT assay.**


The cells were treated with different concentrations (10, 20, 30, 40, 50 and 60µg/ml) over a time duration of 24hrs. Our study observed, *Saraca asoca* flower extract treatment significantly decreased the viability of B16 - F10 skin cancer cells in comparison to the control at 24 hrs time point (Figure-1). The percentage of cell viability decreased gradually with an increase in the concentration. At a concentration of 30 µg/ml, we observed a 50% of growth inhibition which has been taken as inhibitory concentration (IC-50) dose value and was considered for further experiments.

**The cell morphological analysis of *Saraca asoca* flower extract treated skin cancer cells were observed through a phase-contrast microscope.**

The B16 - F10 skin cancer cell line was treated with *Saraca asoca* flower extract (30 µg/ml) for a duration of 24 hrs and compared with the untreated cells, the treated cells demonstrated significant morphological changes, such as cell shrinkage and reduced cell density which are characteristic of apoptotic cells were observed in the *Saraca asoca* flower extract-treated cells (Figure-2). In addition to it, cells undergoing apoptosis also exhibited other types of morphological changes such as rounded up cells that shrink and lose contact with neighbouring cells. Few sensitive cells were also detached from the surface of the plates.

**Acridine orange/Ethidium bromide (AO/EtBr) dual staining were used to confirm the induction of apoptosis in *Saraca asoca* flower extract-treated skin cancer cells.**


AO/EtBr dual staining is used in assessing the nuclear morphology of apoptotic cells for that the cells were treated with *Saraca asoca* flower extract (30 µg/ml) for 24h. After treatment, the cells were stained with both AO/EtBr stain for 20 mins and examined under fluorescence microscopy. The obtained result showed that AO stained both live as well as dead cells, whereas EtBr stains only that have lost their membrane integrity. Cells stained green represent viable cells, whereas yellow staining represents early apoptotic cells, and orange staining represents late apoptotic cells. In the present study, control cells expressed a uniform green colour and in *Saraca asoca* flower extract-treated cells showed a yellow, orange and red signal (Figure-3). From these results obtained, it is confirmed that *Saraca asoca* flower extract induces apoptosis in skin cancer cells.



*Figure 1: The cytotoxic potential of Saraca asoca on the B16 - F10 cell line was determined by MTT assay. The cells were treated with different concentrations (10, 20, 30, 40, 50 and 60 µg/ml) for 24hrs. The 50% of inhibition observed in a concentration of 30 µg/ml, (p value: 0.0086) which has been taken as inhibitory concentration (IC-50) dose value and fixed for further experiments. \* represents statistical significance between control versus treatment groups at  $p < 0.05$  level using Student's-Newman-Keuls test.*



*Figure 2: Represents the morphological changes in B16 - F10 skin cancer cell line upon and without and with the treatment of Saraca asoca at 30  $\mu\text{g/mL}$  for 24hrs by phase-contrast microscope at 20x magnification. The number of cells decreased after treatment and the cells exhibited cell shrinkage and cytoplasmic membrane blebbing.*



*Figure 3: Represents the acridine orange and ethidium bromide dual staining of treated B16 - F10 skin cancer cell line upon without and with the treatment of Saraca asoca at 30  $\mu\text{g/mL}$  for 24hrs, viewed under fluorescence microscope at 20x magnification. The viable cells are possessing a uniform bright green nucleus, early apoptotic cells having bright orange areas of condensed or fragmented chromatin in the nucleus and late apoptotic cells having a uniform bright red nucleus.*



## Discussion

Plants are notable for their therapeutic value and are considered to be a common source of medicine globally (43). Plants have been used for treatments in several countries over the years. *Saraca asoca* is an important ancient medicinal plant seen in India, Sri Lanka, Burma and Malaysia. Studies have reported *S.asoca* extracts to have antitumor and cytotoxic effects. As the skin is the most widely exposed tissue to environmental carcinogens, skin cancer accounts for the most common type of cancer (14). With an increasing prevalence of skin cancers, there is an urgent requirement for the development of various treatment options. Apoptosis or programmed cell death is represented by cell shrinkage, condensation of chromatin, DNA fragmentation and the activation of specific enzymes known as caspases (44). The process of apoptosis is being arrested during cancer progression. Induction of apoptosis in tumour cells is the most established anticancer mechanism and employed in many cancer therapies (45).

In our present study, the cytotoxicity potential of flower extracts of *Saraca asoca* is observed at a concentration of 30 µg/mL for 24hrs. Previous studies evaluated the flavonoid fraction from *Saraca asoca* as a potent chemopreventive property against DMBA-induced skin carcinogenesis and observed that the apoptosis-inducing biochemical markers were significantly restored to near-normal levels (46). The inhibitory concentration (IC-50) dose value found in our study from the flower extracts of *Saraca asoca* was 30 µg/ml. Similar study evaluated the cytotoxicity of crude methanolic extract of *Saraca indica* bark extract on the cervical cancer cell lines and the IC50 value was determined to be 14.63µg/ml (47). In our current study, the morphological changes in B16 - F10 skin cancer cell line upon treatment of *Saraca asoca* at 30 µg/mL for 24hrs has been observed, and the number of cells decreased after treatment, also the cells exhibited cell shrinkage and cytoplasmic membrane blebbing.

Induction of apoptosis in cancer cells is observed in our study after the acridine orange and ethidium bromide dual staining of treated B16 - F10 skin cancer cell line with the treatment of *Saraca asoca* at 30 µg/mL for 24hrs, when viewed under a fluorescence microscope. The viable cells were possessing a uniform bright green nucleus, early apoptotic cells having bright orange

areas of condensed or fragmented chromatin in the nucleus and late apoptotic cells having a uniform bright red nucleus. Studies reported that the presence of compounds such as quercetin and gallic acid in *S.asoca* flowers shows chemopreventive properties. The same study also documented a significant reduction in the expression of ornithine decarboxylase, which is a key enzyme in the promotion stage of 2-stage skin cancer, was observed in the *S.asoca* treated group (48). Another study assessed the anticancer activity of endophytic fungi associated with the medicinal plant *Saraca asoca* and provided promising lead molecules for the development of novel anti-cancer agents (49).

From this study, it can be concluded that the flower extracts of *Saraca asoca* possess potent cytotoxic properties against skin carcinogenesis. Also, further studies are required to understand and ascertain the component that plays a role in proapoptotic potential and their various mechanisms regulating the cytotoxic action of the *S.asoca* flower. The use of the flowers of the *Saraca asoca* can be suggested due to its edible nature, easy availability, and cost effectiveness(13) (50) (51) (52) (53) (54) (55) (56) (57) (58) (59) (60) (61) (62) (63). This work determines significance since the current trend worldwide is to identify therapeutics from natural sources particularly because most of the plants and their products are extensively free from adverse effects and also to interpret traditional knowledge to a scientific platform.

## Conclusion

Overall, the results of the current study revealed that the flower extracts of *Saraca asoca* were cytotoxic and induced apoptosis to the skin cancer cells at a concentration of 30µg/ml at the 24th-time point. However, further research is required to understand the mechanisms of anti-cancer effect of this *Saraca asoca* flower extract.

## References

1. Li C-C, Yu F-S, Fan M-J, Chen Y-Y, Lien J-C, Chou Y-C, et al. Anticancer effects of cantharidin in A431 human skin cancer (Epidermoid carcinoma) cells in vitro and in vivo. *Environ Toxicol* [Internet]. 2017 Mar;32(3):723–38. Available from: <http://dx.doi.org/10.1002/tox.22273>

2. PDQ Adult Treatment Editorial Board. Skin Cancer Treatment (PDQ®): Health Professional Version. In: PDQ Cancer Information Summaries [Internet]. Bethesda (MD): National Cancer Institute (US); 2020. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26389366>
3. Reddy JM, Anitha R, Rajeshkumar S, Lakshmi T. Characterisation of cumin oil mediated silver nanoparticles using UV-visible spectrophotometer and TEM. Research Journal of Pharmacy and Technology [Internet]. 2019;12(10):4931–3. Available from: <https://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=12&issue=10&article=065>
4. Zhang N, Cai Y-X, Wang Y-Y, Tian Y-T, Wang X-L, Badami B. Skin cancer diagnosis based on optimized convolutional neural network. Artif Intell Med [Internet]. 2020 Jan;102:101756. Available from: <http://dx.doi.org/10.1016/j.artmed.2019.101756>
5. Cingi C. Nonmelanoma Skin Cancer of the Head and Neck [Internet]. Vol. 20, Facial Plastic Surgery Clinics of North America. 2012. p. xv. Available from: <http://dx.doi.org/10.1016/j.fsc.2012.08.006>
6. Gomathi AC, Xavier Rajarathinam SR, Mohammed Sadiq A, Rajeshkumar S. Anticancer activity of silver nanoparticles synthesized using aqueous fruit shell extract of Tamarindus indica on MCF-7 human breast cancer cell line [Internet]. Vol. 55, Journal of Drug Delivery Science and Technology. 2020. p. 101376. Available from: <http://dx.doi.org/10.1016/j.jddst.2019.101376>
7. Zeng N, Hongbo T, Xu Y, Wu M, Wu Y. [Retracted] Anticancer activity of caffeic acid n-butyl ester against A431 skin carcinoma cell line occurs via induction of apoptosis and inhibition of the mTOR/PI3K/AKT signaling pathway. Mol Med Rep [Internet]. 2021 May;23(5). Available from: <http://dx.doi.org/10.3892/mmr.2021.12011>
8. Sarojini K, Arivarasu L, Rajeshkumar S, Lakshmi T. GREEN SYNTHESIS OF Solanum trilobatum MEDIATED SELENIUM NANOPARTICLES AND ITS ANTI-INFLAMMATORY AND ANTI-MICROBIAL ACTIVITY. PLANT CELL BIOTECHNOLOGY AND MOLECULAR BIOLOGY. 2020;75–81.
9. Rajeshkumar S. Anticancer activity of eco-friendly gold nanoparticles against lung and liver cancer cells. J Genet Eng Biotechnol [Internet]. 2016 Jun;14(1):195–202. Available from: <http://dx.doi.org/10.1016/j.jgeb.2016.05.007>
10. Dr S, Sherin DR. Flavanoids from Saraca asoca- Ideal Medication for Breast Cancer: A Molecular Simulation Approach [Internet]. Vol. 1, Biomedical Journal of Scientific & Technical Research. 2017. Available from: <http://dx.doi.org/10.26717/bjstr.2017.01.000533>
11. Anitha R, Prathoshni S, Lakshmi T. The effect of capsicum oleoresin on nitric oxide production and nitric oxide synthase gene expression in macrophage cell line [Internet]. Vol. 10, Pharmacognosy Research. 2018. p. 343. Available from: [http://dx.doi.org/10.4103/pr.pr\\_46\\_18](http://dx.doi.org/10.4103/pr.pr_46_18)

12. Cytotoxic Effect and Antibacterial Activity of Coriander Oleoresin Mediated Zinc Oxide Nanoparticles [Internet]. Vol. 12, International Journal of Pharmaceutical Research. 2020. Available from: <http://dx.doi.org/10.31838/ijpr/2020.sp1.411>
13. Ezhilarasan D, Apoorva VS, Ashok Vardhan N. Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. J Oral Pathol Med [Internet]. 2019 Feb;48(2):115–21. Available from: <http://dx.doi.org/10.1111/jop.12806>
14. Saha J, Mukherjee S, Gupta K, Gupta B. High-performance thin-layer chromatographic analysis of antioxidants present in different parts of Saraca asoca (Roxb.) de Wilde [Internet]. Vol. 7, Journal of Pharmacy Research. 2013. p. 798–803. Available from: <http://dx.doi.org/10.1016/j.jopr.2013.10.004>
15. Nandhini JT, Ezhilarasan D, Rajeshkumar S. An ecofriendly synthesized gold nanoparticles induces cytotoxicity via apoptosis in HepG2 cells. Environ Toxicol [Internet]. 2020 Aug 14; Available from: <http://dx.doi.org/10.1002/tox.23007>
16. Suhasini SJ, Roy A, Sosa G, Lakshmi T. The cytotoxic effect of Caralluma fimbriata on KB cell lines. Research Journal of Pharmacy and Technology [Internet]. 2019;12(10):4995–8. Available from: <https://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=12&issue=10&article=075>
17. Mohansrinivasan V, C. SD, Deori M, Biswas A, Jemimah NS. Exploring the Anticancer Activity of Grape Seed Extract on Skin Cancer Cell Lines A431 [Internet]. Vol. 58, Brazilian Archives of Biology and Technology. 2015. p. 540–6. Available from: <http://dx.doi.org/10.1590/s1516-8913201500076>
18. Rajeshkumar S, Kumar SV, Ramaiah A, Agarwal H, Lakshmi T, Roopan SM. Biosynthesis of zinc oxide nanoparticles using Mangifera indica leaves and evaluation of their antioxidant and cytotoxic properties in lung cancer (A549) cells. Enzyme Microb Technol [Internet]. 2018 Oct;117:91–5. Available from: <http://dx.doi.org/10.1016/j.enzmictec.2018.06.009>
19. Nandhini NT, Rajeshkumar S, Mythili S. The possible mechanism of eco-friendly synthesized nanoparticles on hazardous dyes degradation. Biocatal Agric Biotechnol [Internet]. 2019 May 1;19:101138. Available from: <https://www.sciencedirect.com/science/article/pii/S1878818118308235>
20. Vairavel M, Devaraj E, Shanmugam R. An eco-friendly synthesis of Enterococcus sp.–mediated gold nanoparticle induces cytotoxicity in human colorectal cancer cells. Environ Sci Pollut Res [Internet]. 2020 Mar 1;27(8):8166–75. Available from: <https://doi.org/10.1007/s11356-019-07511-x>
21. Gomathi M, Prakasam A, Rajkumar PV, Rajeshkumar S, Chandrasekaran R, Anbarasan PM. Green synthesis of silver nanoparticles using Gymnema sylvestre leaf extract and evaluation of its antibacterial activity. S Afr J Chem Eng [Internet]. 2020 Apr 1;32:1–4. Available from: <https://www.sciencedirect.com/science/article/pii/S1026918519300800>

22. Rajasekaran S, Damodharan D, Gopal K, Rajesh Kumar B, De Pours MV. Collective influence of 1-decanol addition, injection pressure and EGR on diesel engine characteristics fueled with diesel/LDPE oil blends. *Fuel* [Internet]. 2020 Oct 1;277:118166. Available from: <https://www.sciencedirect.com/science/article/pii/S0016236120311625>
23. Santhoshkumar J, Sowmya B, Venkat Kumar S, Rajeshkumar S. Toxicology evaluation and antidermatophytic activity of silver nanoparticles synthesized using leaf extract of *Passiflora caerulea*. *S Afr J Chem Eng* [Internet]. 2019 Jul 1;29:17–23. Available from: <https://www.sciencedirect.com/science/article/pii/S1026918519300253>
24. Raj R K.  $\beta$ -Sitosterol-assisted silver nanoparticles activates Nrf2 and triggers mitochondrial apoptosis via oxidative stress in human hepatocellular cancer cell line. *J Biomed Mater Res A* [Internet]. 2020; Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/jbm.a.36953>
25. Saravanan M, Arokiyaraj S, Lakshmi T, Pugazhendhi A. Synthesis of silver nanoparticles from *Phenerochaete chrysosporium* (MTCC-787) and their antibacterial activity against human pathogenic bacteria. *Microb Pathog* [Internet]. 2018 Apr;117:68–72. Available from: <http://dx.doi.org/10.1016/j.micpath.2018.02.008>
26. Gheena S, Ezhilarasan D. Syringic acid triggers reactive oxygen species-mediated cytotoxicity in HepG2 cells. *Hum Exp Toxicol* [Internet]. 2019; Available from: <https://journals.sagepub.com/doi/abs/10.1177/0960327119839173>
27. Ezhilarasan D, Sokal E, Najimi M. Hepatic fibrosis: It is time to go with hepatic stellate cell-specific therapeutic targets. *Hepatobiliary Pancreat Dis Int* [Internet]. 2018 Jun;17(3):192–7. Available from: <http://dx.doi.org/10.1016/j.hbpd.2018.04.003>
28. Ezhilarasan D. Oxidative stress is bane in chronic liver diseases: Clinical and experimental perspective. *Arab J Gastroenterol* [Internet]. 2018 Jun;19(2):56–64. Available from: <http://dx.doi.org/10.1016/j.ajg.2018.03.002>
29. Gomathi AC, Xavier Rajarathinam SR, Mohammed Sadiq A, Rajeshkumar S. Anticancer activity of silver nanoparticles synthesized using aqueous fruit shell extract of *Tamarindus indica* on MCF-7 human breast cancer cell line. *J Drug Deliv Sci Technol* [Internet]. 2020 Feb 1;55:101376. Available from: <https://www.sciencedirect.com/science/article/pii/S1773224719313693>
30. Dua K, Wadhwa R, Singhvi G, Rapalli V, Shukla SD, Shastri MD, et al. The potential of siRNA based drug delivery in respiratory disorders: Recent advances and progress. *Drug Dev Res* [Internet]. 2019 Sep;80(6):714–30. Available from: <http://dx.doi.org/10.1002/ddr.21571>
31. Ramesh A, Varghese S, Jayakumar ND. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients—A case-control study. *Journal of* [Internet]. 2018; Available from: <https://aap.onlinelibrary.wiley.com/doi/abs/10.1002/JPER.17-0445>

32. Arumugam P, George R, Jayaseelan VP. Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma. Arch Oral Biol [Internet]. 2021 Feb;122:105030. Available from: <http://dx.doi.org/10.1016/j.archoralbio.2020.105030>
33. Joseph B, Prasanth CS. Is photodynamic therapy a viable antiviral weapon against COVID-19 in dentistry? Oral Surg Oral Med Oral Pathol Oral Radiol [Internet]. 2021 Jul;132(1):118–9. Available from: <http://dx.doi.org/10.1016/j.oooo.2021.01.025>
34. Ezhilarasan D, Apoorva VS, Ashok Vardhan N. Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells. J Oral Pathol Med [Internet]. 2019 Feb;48(2):115–21. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/jop.12806>
35. Duraisamy R, Krishnan CS. Compatibility of Nonoriginal Abutments With Implants: Evaluation of Microgap at the Implant–Abutment Interface, With Original and Nonoriginal Abutments. Implantologist [Internet]. 2019; Available from: [https://journals.lww.com/implantdent/Fulltext/2019/06000/Compatibility\\_of\\_Nonoriginal\\_Abutments\\_With.11.aspx](https://journals.lww.com/implantdent/Fulltext/2019/06000/Compatibility_of_Nonoriginal_Abutments_With.11.aspx)
36. Gnanavel V, Roopan SM, Rajeshkumar S. Aquaculture: An overview of chemical ecology of seaweeds (food species) in natural products. Aquaculture [Internet]. 2019 May 30;507:1–6. Available from: <https://www.sciencedirect.com/science/article/pii/S0044848618328072>
37. Markov A, Thangavelu L, Aravindhan S, Zekiy AO, Jarahian M, Chartrand MS, et al. Mesenchymal stem/stromal cells as a valuable source for the treatment of immune-mediated disorders. Stem Cell Res Ther [Internet]. 2021 Mar 18;12(1):192. Available from: <http://dx.doi.org/10.1186/s13287-021-02265-1>
38. Veerasamy R, Roy A, Karunakaran R, Rajak H. Structure-Activity Relationship Analysis of Benzimidazoles as Emerging Anti-Inflammatory Agents: An Overview. Pharmaceuticals [Internet]. 2021 Jul 11;14(7). Available from: <http://dx.doi.org/10.3390/ph14070663>
39. An experimental analysis on the influence of fuel borne additives on the single cylinder diesel engine powered by Cymbopogon flexuosus biofuel. J Energy Inst [Internet]. 2017 Aug 1 [cited 2021 Sep 16];90(4):634–45. Available from: <http://dx.doi.org/10.1016/j.joei.2016.04.010>
40. Campeau PM, Kasperaviciute D, Lu JT, Burrage LC, Kim C, Hori M, et al. The genetic basis of DOORS syndrome: an exome-sequencing study. Lancet Neurol [Internet]. 2014 Jan;13(1):44–58. Available from: [http://dx.doi.org/10.1016/S1474-4422\(13\)70265-5](http://dx.doi.org/10.1016/S1474-4422(13)70265-5)
41. Sathish T, Karthick S. Wear behaviour analysis on aluminium alloy 7050 with reinforced SiC through taguchi approach [Internet]. Vol. 9, Journal of Materials Research and Technology. 2020. p. 3481–7. Available from: <http://dx.doi.org/10.1016/j.jmrt.2020.01.085>
42. Krishnaswamy H, Muthukrishnan S, Thanikodi S, Arockiaraj G, Venkatraman V. Investigation of air conditioning temperature variation by modifying the structure of

- passenger car using computational fluid dynamics [Internet]. Vol. 24, Thermal Science. 2020. p. 495–8. Available from: <http://dx.doi.org/10.2298/tsci190409397k>
43. K S, Sarojini K, Arivarasu L, Smiline GAS. Herbal formulation: Review of efficacy, safety, and regulations [Internet]. Vol. 11, International Journal of Research in Pharmaceutical Sciences. 2020. p. 1506–10. Available from: <http://dx.doi.org/10.26452/ijrps.v11ispl3.3467>
  44. Ashwini S, Ezhilarasan D, Anitha R. Cytotoxic Effect of *Caralluma fimbriata* Against Human Colon Cancer Cells [Internet]. Vol. 9, Pharmacognosy Journal. 2017. p. 204–7. Available from: <http://dx.doi.org/10.5530/pj.2017.2.34>
  45. Teoh PL, Liao M, Cheong BE. *Phyllanthus nodiflorus* L. Extracts Induce Apoptosis and Cell Cycle Arrest in Human Breast Cancer Cell Line, MCF-7. *Nutr Cancer* [Internet]. 2019 Jan 19;71(4):668–75. Available from: <http://dx.doi.org/10.1080/01635581.2018.1559942>
  46. Cibir TR, Gayathri Devi D, Abraham A. Chemoprevention of skin cancer by the flavonoid fraction of *Saraca asoka* [Internet]. Vol. 24, Phytotherapy Research. 2010. p. 666–72. Available from: <http://dx.doi.org/10.1002/ptr.2950>
  47. Asokan A, Department of Microbiology, SreeNarayana Guru College, K G Chavady P, Coimbatore, Thangavel M. In vitro Cytotoxic Studies of crude methanolic extract of *Saraca indica* bark extract [Internet]. Vol. 9, IOSR Journal of Pharmacy and Biological Sciences. 2014. p. 26–30. Available from: <http://dx.doi.org/10.9790/3008-09412630>
  48. Cibir TR, Devi DG, Abraham A. Chemoprevention of two-stage skin cancer in vivo by *Saraca asoka*. *Integr Cancer Ther* [Internet]. 2012 Sep;11(3):279–86. Available from: <http://dx.doi.org/10.1177/1534735411413264>
  49. Jinu MV. Diversity and anticancer activity of endophytic fungi associated with the medicinal plant *Saraca asoka* [Internet]. Vol. 5, Current Research in Environmental & Applied Mycology. 2015. p. 169–79. Available from: <http://dx.doi.org/10.5943/cream/5/3/2>
  50. Danda AK, Krishna TM, Narayanan V, Siddareddi A. Influence of primary and secondary closure of surgical wound after impacted mandibular third molar removal on postoperative pain and swelling--a comparative and split mouth study. *J Oral Maxillofac Surg* [Internet]. 2010 Feb [cited 2021 Sep 15];68(2). Available from: <https://pubmed.ncbi.nlm.nih.gov/20116700/>
  51. Ramadurai N, Gurunathan D, Samuel AV, Subramanian E, Rodrigues SJL. Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial. *Clin Oral Investig* [Internet]. 2019 Sep [cited 2021 Sep 15];23(9). Available from: <https://pubmed.ncbi.nlm.nih.gov/30552590/>
  52. Sathivel A, Raghavendran HR, Srinivasan P, Devaki T. Anti-peroxidative and anti-hyperlipidemic nature of *Ulva lactuca* crude polysaccharide on D-galactosamine induced hepatitis in rats. *Food Chem Toxicol* [Internet]. 2008 Oct [cited 2021 Sep 15];46(10). Available from: <https://pubmed.ncbi.nlm.nih.gov/18706469/>

53. Panda S, Doraiswamy J, Malaiappan S, Varghese SS, Del Fabbro M. Additive effect of autologous platelet concentrates in treatment of intrabony defects: a systematic review and meta-analysis. *J Investig Clin Dent* [Internet]. 2016 Feb [cited 2021 Sep 15];7(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/25048153/>
54. Neelakantan P, Varughese AA, Sharma S, Subbarao CV, Zehnder M, De-Deus G. Continuous chelation irrigation improves the adhesion of epoxy resin-based root canal sealer to root dentine. *Int Endod J* [Internet]. 2012 Dec [cited 2021 Sep 15];45(12). Available from: <https://pubmed.ncbi.nlm.nih.gov/22612994/>
55. Govindaraju L, Neelakantan P, Gutmann JL. Effect of root canal irrigating solutions on the compressive strength of tricalcium silicate cements. *Clin Oral Investig* [Internet]. 2017 Mar [cited 2021 Sep 15];21(2). Available from: <https://pubmed.ncbi.nlm.nih.gov/27469101/>
56. Sekhar CH, Narayanan V, Baig MF. Role of antimicrobials in third molar surgery: prospective, double blind, randomized, placebo-controlled clinical study. *Br J Oral Maxillofac Surg* [Internet]. 2001 Apr [cited 2021 Sep 15];39(2). Available from: <https://pubmed.ncbi.nlm.nih.gov/11286448/>
57. DeSouza SI, Rashmi MR, Vasanthi AP, Joseph SM, Rodrigues R. Mobile phones: the next step towards healthcare delivery in rural India? *PLoS One* [Internet]. 2014 Aug 18 [cited 2021 Sep 15];9(8). Available from: <https://pubmed.ncbi.nlm.nih.gov/25133610/>
58. Nasim I, Neelakantan P, Sujeer R, Subbarao CV. Color stability of microfilled, microhybrid and nanocomposite resins--an in vitro study. *J Dent* [Internet]. 2010 [cited 2021 Sep 15];38 Suppl 2. Available from: <https://pubmed.ncbi.nlm.nih.gov/20553993/>
59. Danda AK, Muthusekhar MR, Narayanan V, Baig MF, Siddareddi A. Open versus closed treatment of unilateral subcondylar and condylar neck fractures: a prospective, randomized clinical study. *J Oral Maxillofac Surg* [Internet]. 2010 Jun [cited 2021 Sep 15];68(6). Available from: <https://pubmed.ncbi.nlm.nih.gov/20303209/>
60. Molecular structure and vibrational spectra of 2,6-bis(benzylidene)cyclohexanone: A density functional theoretical study. *Spectrochim Acta A Mol Biomol Spectrosc* [Internet]. 2011 Jan 1 [cited 2021 Sep 15];78(1):113–21. Available from: <http://dx.doi.org/10.1016/j.saa.2010.09.007>
61. Putchala MC, Ramani P, Herald J, Sherlin, Premkumar P, Natesan A. Ascorbic acid and its pro-oxidant activity as a therapy for tumours of oral cavity – A systematic review [Internet]. Vol. 58, *Archives of Oral Biology*. 2013. p. 563–74. Available from: <http://dx.doi.org/10.1016/j.archoralbio.2013.01.016>
62. Neelakantan P, Grotra D, Sharma S. Retreatability of 2 mineral trioxide aggregate-based root canal sealers: a cone-beam computed tomography analysis. *J Endod* [Internet]. 2013 Jul;39(7):893–6. Available from: <http://dx.doi.org/10.1016/j.joen.2013.04.022>
63. Suresh P, Marimuthu K, Ranganathan S, Rajmohan T. Optimization of machining parameters in turning of Al-SiC-Gr hybrid metal matrix composites using grey-fuzzy



algorithm [Internet]. Vol. 24, Transactions of Nonferrous Metals Society of China. 2014. p. 2805–14. Available from: [http://dx.doi.org/10.1016/s1003-6326\(14\)63412-9](http://dx.doi.org/10.1016/s1003-6326(14)63412-9)

UNDER PEER REVIEW