

Original Research Article

Demographic Profiling of Patients with Leukoplakia Visiting a Private Dental Hospital in Chennai - An Institutional Study

Running Title: Demographic profiling of Leukoplakia patients in a Private dental hospital in Chennai

Abstract

Introduction: Oral potentially malignant disorders (OPMDs) are a group of conditions that predispose to the development of oral cancers. A few of the commonly recurring OPMDs include leukoplakia, oral submucous fibrosis, Smokers' palate, and oral lichen planus. Leukoplakia is more common in specific populations, attributed to their habits and environmental circumstances. Hence, early diagnosis with biopsies on suspecting these lesions during regular oral health check-ups ensure timely referral and addressal of the disorder. The present study aims to demographically profile patients with Leukoplakia visiting a private dental hospital in Chennai.

Aim: To demographically profile patients diagnosed with Leukoplakia, visiting a private dental hospital in Chennai

Materials and Methods: From the patients visiting the private dental hospital, those clinically diagnosed with Leukoplakia were considered in the inclusion criteria. Exclusion criteria involved those with poor maintenance of records and lack of follow-up of uncooperative patients. The sample size was n=100. The gender, age and locality of residence were considered and tabulated, followed by their statistical analysis using SPSS software version 22.0.

Results: From the results of the current study, it was observed that there was a greater male predilection to the occurrence of Leukoplakia (99%) compared to females (1%). A greater number of cases were seen in the age group of 41-50 years (34%), while the least number of cases were seen in patients of 21-30 years (7%). Among the cases reported, most patients were from Thiruvallur (40%), Chennai district (30%) followed by Kancheepuram district (17%) in Tamilnadu .

Conclusion: From the study, it can be concluded that Leukoplakia has a male preponderance, occurring predominantly in the age group of 41-50 years in the districts of Thiruvallur, followed by Chennai district, and Kancheepuram.

Keywords: Leukoplakia, demographic profiling, Private dental hospital, Oral potentially malignant disorders, Novel analysis, Innovative method

Introduction

Oral potentially malignant disorders (OPMDs) are a group of conditions that predispose to the development of oral cancers. The World Health Organization (WHO) earlier termed these as precancerous lesions (1) which were later accepted to be potentially malignant given the fact that they may or may not proceed into malignant forms (2). A few of the commonly recurring OPMDs include leukoplakia, oral submucous fibrosis, Smokers' palate, oral lichen planus, lichenoid reactions in the oral mucosa, graft-versus-host disease, lupus erythematosus, dyskeratosis congenita, epidermolysis bullosa and actinic cheilitis (3). These disorders are further clinically typed to aid in predicting the condition's prognosis (4). Further, early diagnosis with biopsies on suspecting these lesions during regular oral health check-ups ensure timely referral and addressal of the disorder (5). This is of prime concern as most of these conditions are asymptomatic on onset, and cannot be identified unless suspected by professional intervention (6). Besides, particular OPMDs are more common in specific populations, attributed to their habits and environmental circumstances, for example, certain studies show a greater prevalence of Oral submucous fibrosis in Asian populations (7).

The progressive transformation of these disorders are implacable as they often end up in malignancy (8). This feature gave these types of disorders, the previous name 'precancerous lesions'. However, with advancements in the medical field, it has been proved repeatedly that certain morbidities indicate a pattern in their occurrence and can be traced in the population (9). This supports both diagnosis and treatment, allowing the progression of these disorders to be ceased, provided they are identified early. Hence, they have the potential to turn malignant without timely intervention.(10). Thus, the WHO Collaborating Center adopted the latter term recently (11).

A very common OPMD is oral leukoplakia defined as a white mucosal lesion that cannot be characterized pathologically or clinically as any other lesion (12). It is closely related to the habit of smoking or exposure to tobacco (13). Some synergistic risk factors include consuming alcohol, acquiring bacterial and fungal infections, chronic irritation of the oral mucosa due to sharp teeth or ill-fitted dentures or due to syphilis, a sexually transmitted disease (14).

Sometimes even exposure to ultraviolet rays and oral galvanism caused by old metallic restorations may also lead to its development (15).

The histological presentation of oral leukoplakias begin with hyperkeratosis and progress into the different stages of dysplasia, i.e. mild, moderate and severe dysplasia, finally ending in carcinoma in-situ (16). These encompass transformations at the cellular level, in the cytological aspects of the cells with hyperkeratosis and acanthosis, parakeratinization, followed by widening and elongation of rete pegs, increase in the mitotic figures, and formation of keratin pearls, while finally reaching the stage of carcinoma in-situ, where there is no breach in the basement membrane or invasion into the connective tissue (17). Meanwhile, pleomorphic nuclei, increased nuclear-cytoplasmic ratio, reversal of basal cell polarity, prominent nucleoli and hyperchromasia are some cytological changes observed (18). Further, based on their clinical presentation, they can be classified as verrucous, speckled, granular or homogenous leukoplakia (19). For its diagnosis, biopsy from the site of the oral mucosal lesion is the standard procedure (20). It requires a trained health professional to do the biopsy, as it is an invasive procedure. Large lesions require an incisional biopsy inclusive of surrounding normal tissue while excisional biopsy of the entire lesion is indicated in case of small lesions (21). The treatment for leukoplakia generally involves surgical excision of the lesion or laser mediated surgery, if the dysplasia is identified between moderate and severe, especially with tongue, floor of the oral cavity, oropharynx or soft palate involvement (22).

The need for documenting these incidences at a hospital level lies in not only enhancing awareness of the same, but also for eliciting their patterns of occurrence, identifying more prone populations, predisposing behaviours, and recognising their prevalence in various socio-economic situations (23). These documentations further would aid the evidence based segmentation analysis of patients, in the context of culture, health needs and health status of the entire population involved. Thereby, identifying that group of the population predisposed to developing Leukoplakia can be identified and the progression of these conditions into malignant forms can be curbed with timely prophylactic management. Our team has extensive knowledge and research experience that has translated into high quality publications in the same field (24),(25),(26),(27),(28),(29),(30),(31),(32),(33),(34),(35),(36),(37),(38),(39),(40),(41),(42),(43). Thus, the present study aims to assess the demographic details of patients clinically diagnosed with Leukoplakia.

Materials And Methods

The present study was a retrospective study, performed under a University setting, among patients who reported to a Private Dental College and Hospital in Chennai. The data was collected from a total of 5,35,941 patients who reported to the dental college from June 2019 to February 2021. The ethical approval was obtained from the Institutional Ethical Committee

(ethical approval number:SDC/SIHEC/2020/DIASDATA/0619-0320). The population size was n=100, based on diagnosis of leukoplakia, and complete record maintenance. The data was cross verified using photographic record and was compiled under the categories of age, gender, and locality of residence. A descriptive statistical analysis was performed using the SPSS software (version 22.0). Any sampling bias was minimised with data collected from within the University and by using simple random sampling methods. This ensured a high internal validity, with, however, a low external validity. Any improper, incomplete or repeated data was ruled out. Pearson's Chi Square test was used to compare the groups where $p < 0.05$ was considered significant, and the results were interpreted.

Results

In the present study, the total number of patients with leukoplakia was n=100, out of which 7% belonged to 20-30 years, 20% in 31-40 years, 34% in 41-50 years, 29% in 51-60 years and 10% in 61-70 years (Fig. 1). Among these, most patients belonged to the Thiruvallur district (40%) followed by the Chennai district (30%), Kancheepuram district (17%), Chengalpattu (4%), Vellore (3%), Dharmapuri (2%), and 1% each in Sivagangai, Kallakurichi, Pudukottai and Tiruvannamalai districts (Fig. 2). Gender distribution of the patients with leukoplakia showed 99% of the patients were males while 1% of the population was female (Fig. 3). Association between age groups and locality of residence (Fig. 4) showed patients predominantly belonged to the Thiruvallur district from all age groups, including 20-30 years (4%), 31-40 years (8%), 41-50 years (10%), 51-60 years (14%), 61-70 years (4%), with $p=0.840$ ($p > 0.05$) indicating no significant association between them. Association between gender and locality of residence (Fig. 5) showed that male patients predominantly belonged to Thiruvallur district (40%), and females to Kancheepuram district (1%), with $p=0.854$ ($p > 0.05$) indicating no significant association between them.




Figure 1: The graph represents the frequency distribution of age groups of patients visiting a private dental college and hospital in Chennai, diagnosed with Leukoplakia. The X-axis is representative of different age groups while the Y-axis is representative of the percentage of patients with leukoplakia of scale 0-40. According to the graph, 7% belonged to 20-30 years (Dark blue), 20% in 31-40 years (Light blue), 34% in 41-50 years (Yellow), 29% in 51-60 years (Green) and 10% in 61-70 years (Red). Hence, leukoplakia was more prevalent in the age group of 41-50 years (34%).




Figure 2: The graph represents the frequency distribution of the locality of residence of patients diagnosed with leukoplakia, visiting a private dental college and hospital in Chennai. The X-axis is representative of different districts in Tamil Nadu that the patients belonged to, while the Y-axis is representative of the percentage of patients with leukoplakia with a scale of 0-40. According to the graph, 40% patients belonged to the Thiruvallur district (Dark Green) followed by 30% in the Chennai district (Pink), 17% in Kancheepuram district (Light green), 4% in Chengalpattu (Orange), 3% in Vellore (Dark blue), 2% in Dharmapuri (Purple), and 1% each in Sivagangai (Yellow), Kallakurichi (Red), Pudukottai (Light blue) and Thiruvannamalai (Grey). Hence, it can be inferred that leukoplakia was more prevalent in patients from the Thiruvallur district (40%).




Figure 3: The graph represents the frequency distribution of the Gender of patients diagnosed with leukoplakia, visiting a private dental college in Chennai. The X-axis is representative of gender while the Y-axis is representative of the percentage of patients with leukoplakia from a scale of 0-100. According to the graph, 99% of patients were males (Blue) with 1% female patients (Red). Hence it can be inferred that leukoplakia was more prevalent among males than females.




Figure 4: The graph represents the correlation between age groups and area-wise distribution of the patients with leukoplakia. The X-axis is representative of age groups while the Y-axis is representative of the percentage of patients with leukoplakia belonging to different localities within Tamil Nadu. Pink represents Chennai district, yellow represents Sivagangai district, dark green represents Thiruvallur district, light green represents Kancheepuram, orange represents Chengalpattu, dark blue represents Vellore, light blue represents Pudukottai, purple represents Dharmapuri, red represents Kallakurichi and grey represents Thiruvannamalai. According to the graph, from the age group of 41-50 years in which leukoplakia was more prevalent, patients predominantly belonged to the Chennai district (11%). From 51-60 years, more patients were from Thiruvallur district (14%), from 31-40 years 8% patients were from Thiruvallur district. From 20-30 years and 61-70 years each, 4% patients were from the Thiruvallur district. Pearson's chi-square test showed $p=0.840$; indicating that results were statistically not significant ($p>0.05$). It can thus be inferred that from the age group of 41-50 years in which leukoplakia was more prevalent, patients mostly belonged to the Chennai district (11%). However, in all other age groups, Thiruvallur district was more predominant.




Figure 5: The graph represents the correlation between the gender of patients diagnosed with leukoplakia and their locality of residence. The X-axis is representative of gender while the Y-axis is representative of the area-wise percentage of patients with leukoplakia with a scale of 0-40. According to the graph, pink represents Chennai district, yellow represents Sivagangai district, dark green represents Thiruvallur district, light green represents Kancheepuram, orange represents Chengalpattu, dark blue represents Vellore, light blue represents Pudukottai, purple represents Dharmapuri, red represents Kallakuruchi and grey represents Thiruvannamalai. It is evident from the graph that 40% of the male patients predominantly belonged to Thiruvallur district, while the female patient (1%) was from Kancheepuram district. Pearson's chi-square test gave $p=0.854$; indicating that results were statistically not significant ($p>0.05$), yet it can be inferred that predominantly, male patients were from the Thiruvallur district (40%).

Discussion

The requirement of profiling cases categorically based on the areas patients hail from is essential in identifying patterns of occurrence of the concerned health conditions. Previous studies (44,45) have indicated a greater occurrence of leukoplakia in the South-Asian population, providing evidence to practitioners on what can be expected in patients hailing from these demographic regions. Further, other studies (46) have also illustrated the necessity of profiling patients on a socio-economic scale, relating their economic background and their associated habits, to their exposure to risk factors, and thus accounting the same to be causative of leukoplakias (47). Besides, the polymorphisms in the gene pool of a population or a locality may also be traced based on similar diseases seen in them (27). Moreover, demographic profiling enables us to elicit and associate risk factors to the malignant transformation of the oral mucosa. It also enables us to prophylactically manage and halt the progression of leukoplakia into cancers, when predisposed individuals approach the practitioner (48). Thus, the informed practitioner will be able to identify mucosal abnormalities in the oral cavity aptly, and provide timely intervention and assistance (49).

In the present retrospective study, of the 100 patients assessed, area wise profiling of patients was done to identify which areas in the neighbourhood of the city, can be plausible hotspots for the occurrence of leukoplakia. In the current study, patients were categorized into 5 age groups (Fig 1) in the present study. More cases of leukoplakia were seen in the age group of 41-50 years (34%), while the least number of cases were seen in patients of 21-30 years (7%). This finding is concordant with a previous study by It has been recorded earlier by Pires et al, (50) that a higher number of patients belonging to the middle ages were affected with leukoplakia due to the long duration of constant exposure to the mucosal irritants that are causative of malignant transformation of the oral mucosa. However this is discrepant with another study by Balsaraf et al (51) who discussed the current trends of increased use of tobacco in the adolescent ages itself, to be causative of leukoplakia by the age of 30.

Further, from the results of the present study (Fig 2) it was observed that there was a male predilection to the occurrence of leukoplakia (99%) compared to females (1%). This was concordant with a previous study by Mello FW et al (52), who indicated a male preponderance (95%) to the occurrence of leukoplakia, probably attributed to the increased risk factors that males are prone to, a major one being the use of tobacco (53). Other similar studies (54) also indicated a similar pattern in its occurrence among male patients, coherent with the results in the current study.

Considering that the study was conducted in patients visiting a private dental hospital in the city Chennai, Tamil Nadu, the patients who reported here were from districts within and neighbouring Chennai, while few were from distant places (Fig 3). Patients hailed from the districts of Chennai, Chengalpattu, Dharmapuri, Kancheepuram, Thiruvallur, Thiruvannamalai, Pudukottai, Sivagangai and Vellore. Among these, most patients were from Thiruvallur (40%), Chennai district (30%), Kancheepuram (17%). This could indicate these regions to be populous in relation to leukoplakia patients. However, one cannot overlook the preference of patients to particularly seek the present dental hospital in discussion, due to its accessibility and convenient location close to their localities.

Our study shows that among all the age groups, more patients were from the Thiruvallur district over the other localities (Figure 4). Patients predominantly belonged to the Thiruvallur district, from all age groups, 20-30 years (4%), 31-40 years (8%), 41-50 years (10%), 51-60 years (14%), 61-70 years (4%). Pearson's chi-square value $p=0.840$; it was statistically not significant ($p>0.05$). The study also highlighted the correlation between gender and locality of residence of patients with leukoplakia. Male patients predominantly belonged to Thiruvallur district (40%), followed by Chennai district (30%) and Kancheepuram district (16%). There was only 1% of female patients who were also from Kancheepuram district. Pearson's chi-square test showed $p=0.854$; which was statistically not significant ($p>0.05$) (Figure 5). The insignificant p-values could be attributed to the small sample size or due to the location related bias of the patients turning up in the hospital on account of proximity. However, demographic profiling provides data indicative of what the practitioner may be able to find or expect in patients, who give details of living in the same localities that provide evidence of being leukoplakia hotspots.

Thus, the present study, with its limitations of the small sample size, and restricted demographic area taken in consideration, has attempted to categorize and demographically profile patients to provide evidence on what practitioners may find on examining the oral cavity of patients hailing from particular localities.

Conclusion

From the study, it can be concluded that leukoplakia has a male preponderance, occur more often in the age group of 41-50 years and are reported more in the districts of Thiruvallur, followed by Chennai district, and Kancheepuram. However, the study is done within the limits of the small sample size and bias of locality of residence of the patient which may be close to the hospital's location. Yet, the prevalence of leukoplakia based on different gender and age groups within a specific population has been assessed. The demographic distribution of the disorder has also been highlighted. However, to expand on the study, it is necessary to encompass a larger population, so as to avoid any bias related to the ease of approach of the patient to the hospital considered here, due to the patient's proximity to the locality.

References

1. Warnakulasuriya S, Johnson NW, van der Waal I. Nomenclature and classification of potentially malignant disorders of the oral mucosa. *J Oral Pathol Med*. 2007 Nov;36(10):575–80.
2. Mello FW, Melo G, Guerra ENS, Warnakulasuriya S, Garnis C, Rivero ERC. Oral potentially malignant disorders: A scoping review of prognostic biomarkers. *Crit Rev Oncol Hematol*. 2020 Sep;153:102986.
3. Dilhari A, Weerasekera MM, Siriwardhana A, Maheshika O, Gunasekara C, Karunathilaka S, et al. Candida infection in oral leukoplakia: an unperceived public health problem. *Acta Odontol Scand*. 2016 Oct;74(7):565–9.
4. Suvarna K, Abilasha R, Gheena S. Analysis of Prevalence of Oral Squamous Cell Carcinoma in Patients with History of Chronic Irritation of Oral Tissues-A Retrospective Study. *Indian Journal of [Internet]*. 2020; Available from: <http://search.ebscohost.com/login.aspx?direct=true&profile=ehost&scope=site&authtype=crawler&jrnl=09739122&AN=148410023&h=XYlwRUZ0p54Y6mZHESI7LVl228pkq%2FRhoPO4Jt4231oGPh6KuqA4bM6e9ZslMs9HCBZY6ZF8tl2TIR3cgyExXQ%3D%3D&crl=c>
5. Ramani P, Gheena S, Karunakaran M, Hannah R. Clear-cell variant of oral squamous cell carcinoma: A rare entity. *J Oral Maxillofac Pathol*. 2021 Mar 1;25(4):22.
6. Ramasubramanian A, Ramani P, Sherlin HJ, Premkumar P, Natesan A, Thiruvengadam C. Immunohistochemical evaluation of oral epithelial dysplasia using cyclin-D1, p27 and p63 expression as predictors of malignant transformation. *J Nat Sci Biol Med*. 2013 Jul;4(2):349–58.
7. Cheng YI, Gan YC, Liu D, Davies MPA, Li WM, Field JK. Potential genetic modifiers for somatic EGFR mutation in lung cancer: a meta-analysis and literature review. *BMC Cancer*. 2019 Nov 8;19(1):1068.
8. E A, Aswani E, Gheena S, Pratibha R, Abilasha R, Hannah R, et al. Overexpression of HNRNPA2B1 is Associated with Poor Prognosis in Head and Neck Squamous Cell Carcinoma [Internet]. *International Journal of Current Research and Review*. 2020. p. 15–8. Available from: <http://dx.doi.org/10.31782/ijcrr.2020.122502>
9. Yamunadevi A, Pratibha R, Rajmohan M, Ganapathy N, Porkodisudha J, Pavithrah D, et al. Molecular insight into odontogenesis in hyperglycemic environment: A systematic review. *J Pharm Bioallied Sci*. 2020 Aug 1;12(5):49.
10. Ramani P, Krishnan RP, Karunakaran M, Muthusekhar MR. Odontogenic sarcoma: First report after new who nomenclature with systematic review. *J Oral Maxillofac Pathol*. 2020 Jan;24(1):157–63.
11. Warnakulasuriya S. Clinical features and presentation of oral potentially malignant disorders. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2018 Jun;125(6):582–90.
12. Parlatescu I, Gheorghe C, Coculescu E, Tovar S. Oral leukoplakia - an update. *Maedica*. 2014 Mar;9(1):88–93.
13. Liu W, Wang Y-F, Zhou H-W, Shi P, Zhou Z-T, Tang G-Y. Malignant transformation of oral leukoplakia: a retrospective cohort study of 218 Chinese patients. *BMC Cancer*. 2010 Dec 16;10:685.

14. Pentenero M, Giaretti W, Navone R, Rostan I, Gassino L, Broccoletti R, et al. Evidence for a possible anatomical subsite-mediated effect of tobacco in oral potentially malignant disorders and carcinoma. *J Oral Pathol Med*. 2011 Mar;40(3):214–7.
15. Napier SS, Speight PM. Natural history of potentially malignant oral lesions and conditions: an overview of the literature. *J Oral Pathol Med*. 2008 Jan;37(1):1–10.
16. van der Waal I. Potentially malignant disorders of the oral and oropharyngeal mucosa; terminology, classification and present concepts of management. *Oral Oncol*. 2009 Apr;45(4-5):317–23.
17. Nasser W, Flechtenmacher C, Holzinger D, Hofele C, Bosch FX. Aberrant expression of p53, p16INK4a and Ki-67 as basic biomarker for malignant progression of oral leukoplakias. *J Oral Pathol Med*. 2011 Sep;40(8):629–35.
18. Smitha T, Sharada P, Girish H. Morphometry of the basal cell layer of oral leukoplakia and oral squamous cell carcinoma using computer-aided image analysis. *J Oral Maxillofac Pathol*. 2011 Jan;15(1):26–33.
19. Bagan JV, Jimenez Y, Murillo J, Gavalda C, Poveda R, Scully C, et al. Lack of association between proliferative verrucous leukoplakia and human papillomavirus infection. *J Oral Maxillofac Surg*. 2007 Jan;65(1):46–9.
20. Lingen MW, Kalmar JR, Karrison T, Speight PM. Critical evaluation of diagnostic aids for the detection of oral cancer. *Oral Oncol*. 2008 Jan;44(1):10–22.
21. Sankaranarayanan R, Ramadas K, Thomas G, Muwonge R, Thara S, Mathew B, et al. Effect of screening on oral cancer mortality in Kerala, India: a cluster-randomised controlled trial [Internet]. Vol. 365, *The Lancet*. 2005. p. 1927–33. Available from: [http://dx.doi.org/10.1016/s0140-6736\(05\)66658-5](http://dx.doi.org/10.1016/s0140-6736(05)66658-5)
22. Warnakulasuriya S. Histological grading of oral epithelial dysplasia: revisited. *J Pathol*. 2001 Jul;194(3):294–7.
23. Sukumaran G, Ramani P, Ramasubramanian A, Karunakaran M, Ravikumar H. Implantation Dermoid Cyst. *J evol med dent sci*. 2019 Dec 30;8(52):4023–5.
24. Princeton B, Santhakumar P, Prathap L. Awareness on Preventive Measures taken by Health Care Professionals Attending COVID-19 Patients among Dental Students. *Eur J Dent*. 2020 Dec;14(S 01):S105–9.
25. Mathew MG, Samuel SR, Soni AJ, Roopa KB. Evaluation of adhesion of *Streptococcus mutans*, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: randomized controlled trial. *Clin Oral Investig*. 2020 Sep;24(9):3275–80.
26. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma. *J Oral Pathol Med*. 2019 Apr;48(4):299–306.
27. R H, Ramani P, Ramanathan A, R JM, S G, Ramasubramanian A, et al. CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene. *Oral Surg Oral Med Oral Pathol Oral Radiol*. 2020 Sep;130(3):306–12.
28. Antony JVM, Ramani P, Ramasubramanian A, Sukumaran G. Particle size penetration rate and effects of smoke and smokeless tobacco products - An invitro analysis. *Heliyon*. 2021 Mar

29. Sarode SC, Gondivkar S, Sarode GS, Gadbail A, Yuwanati M. Hybrid oral potentially malignant disorder: A neglected fact in oral submucous fibrosis. *Oral Oncol.* 2021 Jun 16;105390.
30. Hannah R, Ramani P, WM Tilakaratne, Sukumaran G, Ramasubramanian A, Krishnan RP. Author response for “Critical appraisal of different triggering pathways for the pathobiology of pemphigus vulgaris—A review” [Internet]. Wiley; 2021. Available from: <https://publons.com/publon/47643844>
31. Chandrasekar R, Chandrasekhar S, Sundari KKS, Ravi P. Development and validation of a formula for objective assessment of cervical vertebral bone age. *Prog Orthod.* 2020 Oct 12;21(1):38.
32. Subramanyam D, Gurunathan D, Gaayathri R, Vishnu Priya V. Comparative evaluation of salivary malondialdehyde levels as a marker of lipid peroxidation in early childhood caries. *Eur J Dent.* 2018 Jan;12(1):67–70.
33. Jeevanandan G, Thomas E. Volumetric analysis of hand, reciprocating and rotary instrumentation techniques in primary molars using spiral computed tomography: An in vitro comparative study. *Eur J Dent.* 2018 Jan;12(1):21–6.
34. Ponnulakshmi R, Shyamaladevi B, Vijayalakshmi P, Selvaraj J. In silico and in vivo analysis to identify the antidiabetic activity of beta sitosterol in adipose tissue of high fat diet and sucrose induced type-2 diabetic experimental rats. *Toxicol Mech Methods.* 2019 May;29(4):276–90.
35. Sundaram R, Nandhakumar E, Haseena Banu H. Hesperidin, a citrus flavonoid ameliorates hyperglycemia by regulating key enzymes of carbohydrate metabolism in streptozotocin-induced diabetic rats. *Toxicol Mech Methods.* 2019 Nov;29(9):644–53.
36. Alsawalha M, Rao CV, Al-Subaie AM, Haque SKM, Veeraraghavan VP, Surapaneni KM. Novel mathematical modelling of Saudi Arabian natural diatomite clay. *Mater Res Express.* 2019 Sep 4;6(10):105531.
37. Yu J, Li M, Zhan D, Shi C, Fang L, Ban C, et al. Inhibitory effects of triterpenoid betulin on inflammatory mediators inducible nitric oxide synthase, cyclooxygenase-2, tumor necrosis factor- α , interleukin-6, and proliferating cell nuclear antigen in 1, 2-dimethylhydrazine-induced rat colon carcinogenesis. *Pharmacogn Mag.* 2020;16(72):836.
38. Shree KH, Hema Shree K, Ramani P, Herald Sherlin, Sukumaran G, Jeyaraj G, et al. Saliva as a Diagnostic Tool in Oral Squamous Cell Carcinoma – a Systematic Review with Meta Analysis [Internet]. Vol. 25, *Pathology & Oncology Research*. 2019. p. 447–53. Available from: <http://dx.doi.org/10.1007/s12253-019-00588-2>
39. Zafar A, Sherlin HJ, Jayaraj G, Ramani P, Don KR, Santhanam A. Diagnostic utility of touch imprint cytology for intraoperative assessment of surgical margins and sentinel lymph nodes in oral squamous cell carcinoma patients using four different cytological stains. *Diagn Cytopathol.* 2020 Feb;48(2):101–10.
40. Karunakaran M, Murali P, Palaniappan V, Sivapathasundharam B. Expression and distribution pattern of podoplanin in oral submucous fibrosis with varying degrees of dysplasia – an immunohistochemical study [Internet]. Vol. 42, *Journal of Histotechnology*. 2019. p. 80–6. Available from: <http://dx.doi.org/10.1080/01478885.2019.1594543>

41. Sarode SC, Gondivkar S, Gadbail A, Sarode GS, Yuwanati M. Oral submucous fibrosis and heterogeneity in outcome measures: a critical viewpoint. *Future Oncol.* 2021 Jun;17(17):2123–6.
42. Raj Preeth D, Saravanan S, Shairam M, Selvakumar N, Selestina Raja I, Dhanasekaran A, et al. Bioactive Zinc(II) complex incorporated PCL/gelatin electrospun nanofiber enhanced bone tissue regeneration. *Eur J Pharm Sci.* 2021 May 1;160:105768.
43. Prithiviraj N, Yang GE, Thangavelu L, Yan J. Anticancer Compounds From Starfish Regenerating Tissues and Their Antioxidant Properties on Human Oral Epidermoid Carcinoma KB Cells, In: PANCREAS. LIPPINCOTT WILLIAMS & WILKINS TWO COMMERCE SQ, 2001 MARKET ST, PHILADELPHIA ...; 2020. p. 155–6.
44. Peng Q, Li H, Chen J, Wang Y, Tang Z. Oral submucous fibrosis in Asian countries. *J Oral Pathol Med.* 2020 Apr;49(4):294–304.
45. Parakh MK, Ulaganambi S, Ashifa N, Premkumar R, Jain AL. Oral potentially malignant disorders: clinical diagnosis and current screening aids: a narrative review. *Eur J Cancer Prev.* 2020 Jan;29(1):65–72.
46. Conway DI, Petticrew M, Marlborough H, Berthiller J, Hashibe M, Macpherson LMD. Socioeconomic inequalities and oral cancer risk: a systematic review and meta-analysis of case-control studies. *Int J Cancer.* 2008 Jun 15;122(12):2811–9.
47. Thamilselvan S, Abilasha, Ramani P, Gheena, Hannah. Evaluation of accuracy between habit history and incidence of oral squamous cell carcinoma. *Int J Curr Res Rev.* 2020;30–5.
48. A study on the variability of drug responsiveness to anti inflammatory drugs - A pilot survey. *Int J Pharm Res [Internet].* 2020 Oct 2;12(02). Available from: <http://www.ijpronline.com/ViewArticleDetail.aspx?ID=17202>
49. Umashankar K, Abilasha, Hannah, Ramani P, Gheena. Knowledge and attitude about COVID-19 pathogenesis among oral pathologists in Chennai. *Int J Curr Res Rev.* 2020;12(19):143–51.
50. Pires F-R, Barreto M-E, Nunes J-G, Carneiro N-S, Azevedo A-B, Dos Santos T-C. Oral potentially malignant disorders: clinical-pathological study of 684 cases diagnosed in a Brazilian population. *Med Oral Patol Oral Cir Bucal.* 2020 Jan 1;25(1):e84–8.
51. Balsaraf S, Bhambal A, Chole R. Study of oral potentially malignant disorders related to various risk factors amongst the patients attending hospitals in Bhopal, India. *Med Pharm Rep.* 2019 Jan;92(1):66–71.
52. Mello FW, Miguel AFP, Dutra KL, Porporatti AL, Warnakulasuriya S, Guerra ENS, et al. Prevalence of oral potentially malignant disorders: A systematic review and meta-analysis. *J Oral Pathol Med.* 2018 Aug;47(7):633–40.
53. Ganesh D, Sreenivasan P, Öhman J, Wallström M, Braz-Silva PH, Giglio D, et al. Potentially Malignant Oral Disorders and Cancer Transformation. *Anticancer Res.* 2018 Jun;38(6):3223–9.
54. Gopinath D, Thannikunnath BV, Neermunda SF. Prevalence of Carcinomatous Foci in Oral Leukoplakia: A Clinicopathologic Study of 546 Indian Samples. *J Clin Diagn Res.* 2016 Aug;10(8):ZC78–83.

UNDER PEER REVIEW