

TITLE: EVALUATION AND VALIDATION OF CARVEDILOL IN BULK AND PHARMACEUTICAL DOSAGE FORMS.

ABSTRACT

Background: Quality may be described as the character, which determines the degree of excellence. A good quality medication is anything, which will match the specified product requirements, may be safely purchased and confidently used for the purpose for which it is designed. To obtain a high quality medication, the manufacturing for producing a drug should have quality integrated into it. Analytical chemistry is the discipline that seeks ever better methods of assessing the chemical composition of natural and manmade materials. **Aims and objective:** Evaluation and validation of carvedilol in bulk and pharmaceutical dosage forms. **Material and methods:** preparation of standard stock solution and preparation of sample solution as per standard protocol. **Result:** In RP-HPLC method, a wavelength of 242 nm was retention time was found to be 2.9 with optimized conditions. Carvedilol showed the linearity in the range of 15.62 -93.75µg/ml. Where the peak shape was symmetrical and a good correlation coefficient value was obtained. **Conclusion:** RP-HPLC, HPTLC, UV spectroscopy were found to be sensitive, precise, and accurate. However these three methods can be used for the routine analysis of carvedilol from formulation.

Keywords: validation, carvedilol, pharmaceutical, dosage

INTRODUCTION

Quality can be defined as the character, which defines the grade of excellence. A good quality drug is something, which will meet the established product specifications, can be safely bought and confidently used for the purpose for which it is intended¹. To get a good quality drug, the manufacturing for making a drug should have quality built into it. Analytical chemistry is the science that seeks ever improved means of measuring the

chemical composition of natural and artificial materials. Analytical chemistry is a sub-discipline of chemistry that has the broad mission of understanding the chemical composition of all matter and developing the tools to elucidate such compositionsⁱⁱ.

Spectrophotometry is generally preferred by industries as the cost of the equipment is less and the maintenance problems are minimal. The method of analysis based on measuring the absorption of a monochromatic light by colourless compounds in the near ultraviolet path of spectrum (200-380nm).ⁱⁱⁱ The photometric methods of analysis are based on the Bouguer- Lambert Beer's Law, which establishes that the absorbance of a solution is directly proportional to the concentration of the analyte. The fundamental principle of operation of spectrophotometer covering UV region consists in that light of definite interval of wavelength passes through a cell with solvent and falls on to photoelectric cell that transforms the radiant energy into electrical energy measured by galvanometer. Molecular absorption in the ultraviolet and visible region of the spectrum is dependent on the electronic structure of the molecule. Absorption of energy is quantized, resulting in the elevation of electrons from orbital in the ground state to higher energy orbital in the excited state.^{iv}

AIM AND OBJECTIVE

Evaluation and validation of carvedilol in bulk and pharmaceutical dosage forms.

MATERIAL AND METHODS

PREPARATION OF STANDARD STOCK SOLUTION:

Weigh accurately and transfer about 62.52 mg of carvedilol into a 100 ml volumetric flask. 50 ml methanol added and sonicated to dissolve, and then make up to the volume with methanol. Pipette out 10 ml of this solution into a 100 ml volumetric flask and diluted up to the mark with mobile phase. Filtered through 0.45 μ membrane filter.

PREPARATION OF SAMPLE SOLUTION:

Transfer the accurately weighed samples 15.62, 31.25, 46.88, 62.50, 78.13, 93.75 mg respectively into individual 100 ml flask. 50 ml methanol added and sonicated to dissolve,

then make up to the volume with methanol. Pipette out 10 ml of this solution into a 100 ml volumetric flask and diluted up to the mark with mobile phase. Filtered through 0.45 μ membrane filter. Inject 10 μ l of blank solution and each linearity level standard solutions into the chromatographic system and measure the peak area.

The linearity of carvedilol was performed in the range of 15.62 μ g/ml to 93.75 μ g/ml (25% - 150 % of working concentration). A graph was plotted with concentration in μ g/ml on x axis and peak area on y axis. Slope, y intercept, correlation coefficient (r value), were determined.

RESULT & OBSERVATION

Table 1 :RP-HPLC profile

Name of Drug	Retention Time	Area	Theoretical Plate	Tailing Factor
Carvedilol	2.96	4121905	5350.4	1.40

The analytical method meets the acceptance criteria for accuracy study. Hence the method is accurate for the determination of assay of carvedilol tablets. [Fig 1]

HPTLC profile [Fig 2 and 3]

Linearity:

Aliquots of 0.1-0.5 μ g/spot of standard solution of Carvedilol is applied on the plate with the help of micro liter syringe using an automatic sample applicator. The plates were developed, dried and scanned densitometrically at 254 nm. The drug peak-area was calculated for each concentration level and a graph was plotted of drug concentration against the peak area and shown in (Fig.3). Calibration parameters are given in table 2 : 5

Table 2 : Linearity of Carvedilol

SL.No	Standard concentration μ g/spot	Peak Area	Peak Height

1	0.1	756.35	35.8
2	0.2	1387.96	72.4
3	0.3	2081.94	123.9
4	0.4	2775.92	168.8
5	0.5	3469.9	203.4

UNDER PEER REVIEW

CALIBRATION CURVE OF CARVEDILOL




Table 3 : Calibration Parameters for Carvedilol

Parameters	Carvedilol
Linearity Range(µg/Spot)	0.1-0.5
Slope	6886
Intercept	23.76
Regression Co-Efficient	0.999

Accuracy:

Accuracy of the developed method was confirmed by doing a recovery study as per ICH guidelines at three different concentration levels (80%, 100% and 120%) by replicate analysis (n=3). Standard drug solutions were added to a preanalyzed sample solution, and then percentage of drug content was calculated. The results of the accuracy study are reported in Table 4. From the recovery study, it was clear that the method is very accurate for quantitative estimation of Carvedilol in tablet dosage form because all the statistical results were within the acceptance range (i.e., % RSD <2.0).

Table 4 :Recovery studies for Carvedilol (n=3)

Label claim (mg /tablet)	Recovery level (%)	Amount added (mg)	Amount recovered (mg) ± % RSD	% Recovery
--------------------------	--------------------	-------------------	-------------------------------	------------

25	80	20	19.96±0.42	99.82
25	100	25	24.65±0.85	98.67
25	120	30	30.45±0.59	101.5

Precision:

The precision of the method (system reproducibility) was assessed by spotting 0.3µl of drug solution six times on a TLC plate, followed by development of plate and recording the peak area for 6 spots. The % RSD for peak area values of carvedilol was found to be 0.58. The results were shown in Table-6.

The method reproducibility (intra-day precision) was determined by analyzing standard solution in the concentration range of 0.1 µg/spot to 0.5 µg/spot of drug for 3 times on the same day and inter-day precision was determined by analysing corresponding standards daily for 3 day over a period of one week. The intra-day and inter-day coefficients of variation (%RSD) are in range of 0.13 to .36 and 0.30 to 0.56, respectively. The results were shown in Table- 6a, 6b.

Table 5: Precision of Carvedilol

S.No	Concentration (µg/ spot)	Peak Area
1.	0.3	2081.94
2.	0.3	2111.40
3.	0.3	2090.52
4.	0.3	2075.36
5.	0.3	2087.25
6.	0.3	2092.17
Mean	-	2089.77
Percentage Relative Standard-		0.58

Deviation		
-----------	--	--

Table 6a: Intra-day Precision of Carvedilol

S.No	Concentration (µg / spot)	Area	Mean	Standard Deviation	% RSD
1.	0.1	756.35	757.3	3.37	0.44
2.	0.1	761.05			
3.	0.1	754.50			
1.	0.3	2080.50	2080.82	2.74	0.13
2.	0.3	2083.72			
3.	0.3	2078.25			
1.	0.5	3469.50	3470.32	5.13	0.14
2.	0.5	3465.64			
3.	0.5	3475.82			

Table 6b: Inter-day Precision of Carvedilol

S.No	Concentration (µg / spot)	Area	Mean	Standard Deviation	% RSD
1.	0.1	757.98	758.28	3.93	0.52
2.	0.1	754.50			
3.	0.1	762.35			
1.	0.3	2081.94	2083.56	6.28	0.30
2.	0.3	2090.50			
3.	0.3	2078.25			


UV Profile

The developed method was validated in terms of linearity, accuracy and stability studies.

Linearity

Carvedilol was found to be linear in a concentration range of 1-5 µg/ml. The absorbance of this solution were measured at 242 nm and a calibration graph was plotted using absorbances versus concentration. The correlation co-efficient value was found to be 0.998.

Fig 4 : CALIBRATION CURVE OF CARVEDILOL



Accuracy

The accuracy, specificity, suitability and validity of the present method were satisfied by conducting percentage recovery studies. A known quantity of the drug was added to the pre-analyzed sample formulation at 50% and 100% levels. The percentage recovery and standard deviation were calculated. (Table 3).

The % recovery was calculated by using the following formula

$$\% \text{ Recovery} = \frac{\text{Amount of drug found in sample before addition of standard drug} + \text{Amount of standard drug added}}{\text{Amount of drug found after addition of standard drug}} \times 100$$

Table 7 : RECOVERY STUDIES

Drug	Level	Amount found in µg	Actual amount Added in µg	%Recovery	% RSD
Carvedilol	50%	12.30	12.22	100.6	0.12
		12.28	12.22	100.4	
		12.30	12.22	100.6	
	100%	24.71	24.64	100.3	0.06
		24.69	24.64	100.2	
		24.70	24.64	100.2	

Stability

The drug solution was found to be stable for about three hours at room temperature.

Stability data reported.

Table 8 : STABILITY DATA

Concentration in µg/ml	Time (min)	Absorbance
1	0	0.113
	30	0.110
	60	0.109
	90	0.117
	120	0.112
	150	0.107
	180	0.108

Table 9 : REPEATABILITY STUDIES

Concentration in µg/ml	Absorbance	%RSD
	0.117	
	0.118	

1	0.119	0.93
	0.117	
	0.117	
	0.116	
5	0.516	0.47
	0.513	
	0.511	
	0.510	
	0.515	
	0.511	

Table 10 : INTER-DAY PRECISION

Concentration µg/ml)	Day	Absorbance	%RSD
1	1	0.116	1.02
	2	0.117	
	3	0.115	
	4	0.115	
	5	0.116	
	6	0.114	
5	1	0.512	0.40
	2	0.513	
	3	0.512	
	4	0.516	
	5	0.510	
	6	0.513	

Table 11: INTRA- DAY PRECISION

Concentration $\mu\text{g/ml}$	6 times in a day	Absorbance	%RSD
1	1	0.115	1.47
	2	0.113	
	3	0.111	
	4	0.114	
	5	0.115	
	6	0.112	
5	1	0.513	0.45
	2	0.515	
	3	0.510	
	4	0.512	
	5	0.515	
	6	0.516	

DISCUSSION

Validated analytical methods are aimed for the estimation of carvedilol in formulation. Simple, precise, rapid, accurate methods were developed for the estimation of carvedilol in formulation by following methods^v. In RP-HPLC method, a wavelength of 242 nm was selected and the mobile phase which consist potassium di hydrogen phosphate buffer : acetonitrile, in the ratio of (60:40). pH 3 adjusted with formic acid at a flow rate of 1ml/min were found to be optimum condition for analysis. The retention time was found to be 2.9 with optimized conditions. Carvedilol showed the linearity in the range of 15.62 -93.75 $\mu\text{g/ml}$. Where the peak shape was symmetrical and a good correlation coefficient value was obtained. The percentage label claim and recovery at three different levels,

80%, 100%, 120%, level was carried out. The suitability of the method was thus proved. Precision of the method was studied by making repeated injection of the same sample and standard deviation was determined. Inter day and intraday precision was also carried out and % RSD was calculated. In HPTLC during the stage of method development different mobile phase were tried and mobile phase comprising of ethyl chloroform: methanol : toluene in the proportion of 1.5: 3:3.5 v/v/v for carvedilol , were found to be better and produced the R_f value of 0.72 for carvedilol. The linearity of drug was determined by calibration curve and the linearity based on the area observed in the range of 0.1 – 0.5 $\mu\text{g/ml}$. The regression coefficient value for carvedilol is 0.999. Interday precision of the drugs was studied. No interference with the additives of the formulation was reported. Recovery studies were carried out for the accuracy parameter and were reported. The validated method was applied for the analysis of tablet containing 25 mg carvedilol drug as the label claim. The method developed was simple. It has showed a good peak and good R_f values. In case of UV-spectroscopic method solubility is the important parameter. Solubility parameter was studied and methanol was selected as the solvent, since it gave a maximum absorbance and a good spectral pattern when compared with other solvents^{vi}. The linearity was found to be in the range of 1 -5 $\mu\text{g/ml}$ at the maximum absorbance of 242 nm. The marketed formulation was extracted and diluted to get the concentration in the linearity range. The solution was scanned and measured at 242 nm. Percentage recovery, linearity, stability studies were also carried out. The above method gave a satisfactory recovery values and found to be stable, linear, hence it can be used for routine analysis of the drug formulation^{vii}.

These methods, RP-HPLC, HPTLC, UV spectroscopy were found to be sensitive, precise, and accurate. However these three methods can be used for the routine analysis of carvedilol from formulation.

CONCLUSION

RP-HPLC, HPTLC should spend most of the impact on the creation and optimization of a technique as this enhances the performance of the last method. It should be simple to verify a properly designed technique. In order to analyse preclinical samples, formulations and commercial samples, a technique should be established quickly. The observations of validation factors like as accuracy, accuracy, speciality and linearity have shown that the techniques devised may be used to routinely analyse carvedilol in bulk and tablets. The results of the validation parameters fulfilled the requirements of ICH and USP and complied with the legislation of BEER.

Fig 1: RP-HPLC




Fig 2: HPTLC profile











Fig 3: HPTLC Densitogram




Carvedilol Densitogram-0.1µg

3D display of Peak densitogram – All tracks (Carvedilol standards and sample) @ 254nm



Carvedilol densitogram -Sample

Fig 4: UV spectrum



REFERENCES

- ⁱ Merit.W, Dean.S, Instrumental method of analysis, 7th edition, 75-83.
- ⁱⁱ Sharma.B.K, Instrumental method of chemical analysis, 1994, 18th edition, 3-12.
- ⁱⁱⁱ Gaten.W.E, Instrumental method of chemical analysis, 5th edition, 33-52.
- ^{iv} Beckett.A.H, Stenlake J.B, Practical pharmaceutical chemistry, 4th edition, 249-255.
- ^v Gadape H, Parikh K. Quantitative determination and validation of Carvedilol in pharmaceuticals using quantitative nuclear magnetic resonance spectroscopy. Analytical methods. 2011;3(10):2341-7.
- ^{vi} Kim SH, Lee SH, Lee HJ. Rapid and sensitive carvedilol assay in human plasma using a high-performance liquid chromatography with mass/mass spectrometer detection employed for a bioequivalence study. American Journal of Analytical Chemistry. 2010 Nov 30;1(03):135.
- ^{vii} Kaplan K, Gürkan Polat T, Duman O, Tunç S. Development of a simple, rapid, accurate, and sensitive method for carvedilol analysis in human blood serum by reversed phase-high performance liquid chromatography with diode array detector. Journal of Liquid Chromatography & Related Technologies. 2018 May 28;41(9):532-43.