

Review Article

A REVIEW ON ANTIUROLITHIATIC(PASHANABEDA) EFFECTS OF HERBAL AND MARINE RESOURCES

ABSTRACT

Urolithiasis (urinary calculi) is stone development in bladder, urinary tract and kidney. The rise in frequency and occurrence of kidney stones is talking point all over the world. 12% of the global population suffer from urinary stone formation. The phenomenon effect of urolithiasis is lowered by using the pashanabeda elements. Pashanabeda (Pashan: stone; bheda: to break) is a Sanskrit locution which implies the breakage of stone. The adverse reactions caused due to man made drugs has prompted to rejoin with the natural safe medication. The evolution of plant-based therapeutics has shown terrific notice and demand as an modernistic drug entities all over the sphere. The present review aims the investigators to easily identify and develop plant and marine resources beneficial in management of urolithiasis.

Keywords: Urolithiasis, Antiurolithiatic , Pashanabeda, Herbal, Animal, Marine, Microbial , Mineral

1. INTRODUCTION

Humankind is known to be suffering from urinary stone disease since ancient times, and it was first found in tombs of Egyptian mummies dating back to 4000BC and in the graves of North American Indians from 1500 BC – 1000BC. (Mohammad Shamim Khan¹ et al 2016)

UROLITHIASIS OR NEPHROLITHIASIS:


Urolithiasis or nephrolithiasis is one of the oldest and endemic painful urological disorder. The genesis of stone in urinary system i.e., in the kidney, ureter and urinary bladder or in the

urethra is known as urolithiasis (ouron =urine, lithos = stone). (Mansoor, M., Jamil et al 2019)

World population of 3-20% have propensity to form single urinary stone in their lifespan of 70 years. (Ram Prakash et al, 2019). 12% of global population suffer from urinary stone formation. (shruthi gupta et al 2018)

The average lifetime risk of stone formation has been reported in the range of 5-10% population in which there is a predominance of men over women. A handful of lithiogenic factors that leads to formation of stones such as sexuality, age, profession, food habits, water intake, education level, national diffusion, socioeconomic status, metabolic and genetic disorders and diseases. (Mohammad Shazib Faridi et al. 2020; Victoriano Romero 2010; Yu Liu et al 2018)

Fig 1. FLOWCHART of MECHANISM OF URINARY STONE FORMATION



MECHANISM OF URINARY STONE FORMATION:

Stone genesis is a complicated process which results because of succeeding Physicochemical occurrences such as excellent saturation, growth, nucleation, retention and aggregation within the renal tubules.

The mechanism of stone formation comprises seedling of stone crystals (nuclei formation, nucleation), assemblage of nuclei to interact with inter renal structure (aggregation) and thus

maintain themselves in renal system further aggregate to form secondary nuclei which shape into clinical stone (crystal development). (Vishal N Rathkalar et al.; 2011)

Constituents of stone components may differ, but the severity it causes may vary based on its site, severity of stone formation and its action at site.

The mechanism of stone formation is depicted in flowchart format as indicated below

TYPES OF UROLITHIASIS:

The kidney stone size varies from 5mm to 7mm. different types of kidney stones are included in table format. The type of stone formed is named after its mineral composition. The most frequent stones are struvite (magnesium ammonium phosphate), calcium oxalate, urate, cysteine and silica. (Shruthi gupta et al. 2018)

TABLE:1 TYPES OF KIDNEY STONES (Shruthi gupta et al. 2018)

S.NO.	TYPES OF KIDNEY STONES	PERCENTAGE OF OCCURANCE	CAUSES FOR KIDNEY STONES	CHARACTERISTIC FEATURE OF STONES
1.	Calcium stones	-	Calcium combines with oxalate and phosphate to form calcium oxalate and calcium phosphate stones. The rise in oxalate and phosphate levels in dietary supplements leads to formation of stones	White, black or grey colored components, radio opaque
2.	Struvite stones	10-15%	Stones are formed due to bacterial infections	Large, glazed and laminated
3.	Uric acid stones	5-10%	People with malformation like gout syndrome, obesity and food source rich in purine, protein (meat & fish) results in stone formation	Yellow orange colored, smooth, round or square or diamond or rod shape, pleomorphic

4.	Protease related stones	4-12%	Formation of crystals are recognized in HIV positive patients using protease inhibitors (indinavir sulphate)	-
5.	Cysteine stones	-	Hereditary disorders induce formation of stones	Greenish yellow color, round, shiny, radiopaque
6.	Silica stones	-	Synthetic and herbal medicines induce formation of stones. These are crystal called as drug induced stones	-

PASHANABHEDA:

Pashanabeda is a Sanskrit phrase where the word Pashan means stone and bheda means to break, which literally means the breakage of stones. Several plants such as *Bergenia ligulate* syn., *Saxifraga ligulate* etc. are denominated as pashanabeda as they show properties similar to that of water pills and lithiotriptic activities which is included in classics of Ayurveda. (www.ccras.nic.in) (Poonam varma et al.; 2014)

ANTIROLITHIATIC ACTIVITY:

The components that reduce the stone shaping components in urine and reduce renal retention are known as antirolithiatic components. The activity is known as antirolithiatic activity

This review article mainly focuses on various plants and marine accredited with diuretic and antirolithiatic activities.

METHODOLOGY:

The components are classified into natural sources and synthetic sources. Natural sources are further classified into herbal source, animal source, marine source, microbial source and mineral source.

HERBAL SOURCES:

The plants that show antiurolithiatic activity and diuretic activity are classified based on their morphological characteristics and depicted examples in alphabetical order. It is further depicted in the form of table as shown below (table:2) (Salman ahmed et al., 2016 different articles)

TABLE:2 PLANTS WITH ANTIUROLITHIATIC AND DIURETIC ACTIVITY

S.NO.	CLASS (ROOT)	ANTIUROLITHIATIC PLANTS
1.	<i>Abutilon muticum</i> (Delile ex DC.)	<i>Hygrophila auriculata</i>
	<i>Achyranthus aspera</i> L.	<i>Hygrophila schulli</i>
	<i>Acorus calamus</i>	<i>Ichnocarpus frutescens</i> (L.) W.T.Aiton
	<i>Aerva javanica</i>	<i>Indigofera tinctoria</i>
	<i>Ageratum conyzoides</i>	<i>Juniperus chinensis</i> . <i>Juniperus polycarpus</i> K.Koch
	<i>Alcea fasciculiflora</i> Zohary	
	<i>Alhagi mannifera</i> Jaub. & Spach	<i>Lawsonia inermis</i> L.
	<i>Alhagi maurorum</i> Medik.	<i>Levisticum officinale</i>
	<i>Alisma orientale</i> (Sam.) Juz.	<i>Medicago sativa</i> L.
	<i>Angelica sinensis</i> polysaccharide	<i>Mimosa pudica</i> L.
	<i>Apium graveolens</i> L.	<i>Moringa oleifera</i> root-wood
	<i>Aquilegia fragrans</i> Benth.	<i>Musa balbisiana</i> Colla.
	<i>Arctium lappa</i>	<i>Musa × paradisiaca</i> L.
	<i>Arnebia euchroma</i>	<i>Nothosaerva brachiate</i> (L.) Wight.
	<i>Asclepias syriaca</i> L.	<i>Ononis spinosa</i>
	<i>Asparagus officinalis</i>	<i>Petroselinum crispum</i>
	<i>Asparagus racemosus</i>	<i>Petroselinum sativum</i>
	<i>Asparagopsis adscendens</i> Kunth. / <i>Asparagus officinalis</i>	<i>Phragmites australis</i> (Cav.) Trin. Ex Steud.
	<i>Anneslea fragrans</i> Wall. / <i>Aquilegia fragrans</i> Benth.	<i>Piper methysticum</i> G.Forst.
	<i>Barbaarea vulgaris</i> R.Br.	<i>Plantago major</i> L.
	<i>Barleria prionitis</i> L.	<i>Raphanus sativus</i> L.
	<i>Berberis vulgaris</i>	<i>Rheum emodi</i> Wall.
	<i>Bergenia ciliate</i>	<i>Ribes triste</i> Pall.
	<i>Beta vulgaris</i> L.	<i>Ricinus communis</i> L.
	<i>Biophytum abyssinicum</i> Steud. Ex A.	<i>Rotula aquatica</i> Lour.

Rich.	
<i>Biophytum reinwardtii</i>	<i>Rubia tinctorum</i>
<i>Biophytum sensitivum</i> L.	<i>Rubus caesius</i> L.
<i>Boerhavia diffusa</i>	<i>Rubus fruticosus</i> Lour.
<i>Caesalpinia nuga</i> (Aiton.)	<i>Rubus sanctus</i> Schreb.
<i>Celosia argentea</i>	<i>Rumex acetosella</i>
<i>Ceterach aureum</i> Buch.	<i>Rumex hastatus</i>
<i>Chondrodendron tomentosum</i> Ruiz & Pav.	<i>Saccharum officinarum</i>
<i>Cissampelos parerira</i> L.	<i>Saccharum spontaneum</i>
<i>Citrus limon</i> (L.) Osbeck	<i>Sageretia brandrethiana</i> Aitch.
<i>Clerodendrum serratum</i>	<i>Saponaria mesogitana</i> Boiss.
<i>Clitoria ternatea</i>	<i>Scoparia dulcis</i> L.
<i>Coccinia grandis</i>	<i>Sida rhombifolia</i>
<i>Coix lacryma-jobi</i>	<i>Smilax aspera</i>
<i>Costus arabicus</i>	<i>Solanum anguivi</i> Lam.
<i>Costus igneus</i>	<i>Solanum incanum</i> L.
<i>Crotalaria albida</i> Roth.	<i>Solanum surattense</i> Burm.f.
<i>Crotalaria pallida</i> Aiton.	<i>Solanum virginianum</i> L.
<i>Cucumis sativus</i> L.	<i>Sphaeranthus indicus</i> L.
<i>Cynara scolymus</i> L.	<i>Strychnos potatorum</i> L.f.
<i>Cynodont dactylon</i> (L.) Pers.	<i>Taraxacum officinale</i>
<i>Daucus carota</i> L.	<i>Tectona grandis</i>
<i>Ecbolium viride</i> (Forssk.) Alston	<i>Tephrosia purpurea</i>
<i>Echinops spinosus</i> L.	<i>Tinospora purpuea</i>
<i>Ensete superbum</i> (Roxb.) Cheesman.	<i>Traxacum pseudobrachyglossum</i> Soest.
<i>Eryngium creticum</i> Lam.	<i>Trianthema portulacastrum</i>
<i>Eupatorium purpureum</i> L.	<i>Tribulus terrestris</i>
<i>Filipendula vulgaris</i> Moench	<i>Tropaeolum tuberosum</i> Ruiz & Pav
<i>Glycyrrhiza glabra</i> L.	<i>Valeriana officinalis</i> L.
<i>Gypsophila struthium</i> Loefl.	<i>Valeriana wallichii</i> DC.
<i>Hamelia patens</i> Jacq.	<i>Vetiveria zizanioides</i>
<i>Hemidesmus indicus</i>	<i>Vitex negundo</i>
<i>Helianthus annuus</i>	<i>Xanthium strumarium</i>

	<i>Homonía riparia</i> Lour.	<i>Xenostegia tridentata</i> Webb & Berthel
	<i>Holarrhena antidysenterica</i>	<i>Zaleya pentandra</i> (L.) C.Jeffrey
	<i>Hydrangea arborescens</i> L.	<i>Ziziphus lotus</i> (L.) Lam.
2.	RHIZOME	
	<i>Agropyron repens</i> (L.) P. Beauv.	<i>Cynodon dactylon</i>
	<i>Asparagus racemosus</i> Willd	<i>Elymus repens</i>
	<i>Bergenia ligulate</i> Engl.	<i>Eupatorium purpureum</i>
	<i>Beta vulgaris</i> L.	<i>Hedychium coronarium</i> J. Koenig.
	<i>Colocasia esculenta</i> (L.) Schott.	<i>Smilax lanceifolia</i> Roxb.
	<i>Curculigo orchioides</i> Gaertn.	<i>Zingiber officinale</i> Roscoe
3	BULB	
	<i>Allium cepa</i> L.	<i>Drimia indica</i> (Roxb.) Jessop
	<i>Allium sativum</i> L.	<i>Scilla indica</i> Roxb.
4	AERIAL PARTS	
	<i>Achillea falcata</i> L.	<i>Gymnocarpus decandrum</i> Forssk.
	<i>Arnica montana</i> L.	<i>Geranium robertianum</i> L.
	<i>Cardamine uliginosa</i> M.Bieb.	<i>Helichrysum plicatum</i> DC.
	<i>Chimaphila maculate</i> (L.) Pursh.	<i>Herniaria glabra</i> L.
	<i>Chimaphila umbellate</i> (L.) Nutt.	<i>Hyparrhenia hirta</i> (L.) Stapf.
	<i>Cocculus hirsutus</i> (L.) W.Theob.	<i>Lolium perenne</i> L.
	<i>Crataegus monogyna</i>	<i>Melilotus officinalis</i> (L.) Pall.
	<i>Crataegus pentagyna</i>	<i>Orthosiphon aristatus</i> (Blume) Miq.
	<i>Cyperus longus</i>	<i>Prosopis farcta</i> (Banks & Sol.)J.F.Macbr.
	<i>Derris trifoliata</i> Lour.	<i>Salvia canariensis</i>
	<i>Equisetum ramosissimum</i>	<i>Verbena officinalis</i> L.
	<i>Equisetum telmateia</i> .	<i>Zilla spinosa</i> (L.) Prantl.
5	BARK	
	<i>Ammi visnaga</i> (L.) Lam.	<i>Juniperus communis</i>
	<i>Bauhinia purpurea</i> L.	<i>Laurus nobilis</i> L.
	<i>Betula pendula</i> Roth.	<i>Lawsonia inermis</i> L.
	<i>Berberis vulgaris</i> L.	<i>Macaranga peltate</i>
	<i>Cinnamonum aromaticum</i> Nees.	<i>Mimusops elengi</i> L.
	<i>Cinnamonum bejolghota</i> (Buch. – Ham.) Sweet	<i>Moringa pterygosperma</i> Gaertn

	<i>Cinnamomum verum</i> J. Presl.	<i>Neolamarckia cadamba</i> (Roxb.) Bosser.
	<i>Clitoria ternatea</i> L.	<i>Olea europaea</i> L.
	<i>Cedrus deodara</i> (Roxb. Ex D. Don) G.	<i>Picea mariana</i> (Mill.) Britton, Sterns & Poggenb
	<i>Crateva adansonii</i> subsp	<i>Raphanus sativus</i> L.
	<i>Crataeva magma</i> (Lour.) DC	<i>Saraca asoca</i> (Roxb.) Willd.
	<i>Crataeva nurvala</i>	<i>Sonchus oleraceus</i> (L.) L.
	<i>Holarrhena pubescens</i>	<i>Terminalia arjuna</i>
	<i>Hydrangea arborescens</i>	<i>Zizipus lotus</i>
6	STEM	
	<i>Achyranthus aspera</i> L.	<i>Hedychium aurantiacum</i> Roscoe.
	<i>Butea monosperma</i> (Lam.) Taub.	<i>Musa × paradisiaca</i>
	<i>Bryonia alba</i> L.	<i>Silybum marianum</i> (L.) Gaertn.
	<i>Eryngium campsetre</i> L.	<i>Ruscus aculeatus</i> L.
	<i>Equisetum arvense</i> L.	<i>Tinospora cordifolia</i> (Willd.) Miers.
7	BAMBOO SHOOTS	<i>Bambusa nutans</i> Wall.ex Munro
8	LEAVES	
	<i>Aaronsohnia pubescens</i> (Desf.) K.Bremer & Humphries.	<i>Hypericum hypericoides</i>
	<i>Adiantum capillus-veneris</i>	<i>Ilex aquifolium</i> L.
	<i>Allium odorum</i> L.	<i>Larrea tridentata</i>
	<i>Alternanthera brasiliana</i>	<i>Launaea procumbens</i> L.
	<i>Althaea officinalis</i> L.	<i>Lavandula stoechas</i> L.
	<i>Amaranthus blitum</i>	<i>Lepidium latifolium</i> L.
	<i>Amaranthus caudatus</i>	<i>Lithospermum officinale</i> L
	<i>Ammi visnaga</i> (L.) Lam.	<i>Melia azadirachta</i> L.
	<i>Anisotes trisulcus</i> (Forssk.) Nees	<i>Mentha arvensis</i> .
	<i>Anacardium occidentale</i> L.	<i>Mentha spicata</i> .
	<i>Anneslea fragrans</i> Wall. / <i>Aquilegia fragrans</i> Benth.	<i>Mimosa pudica</i> L.
	<i>Arctostaphylos pungens</i>	<i>Moringa oleifera</i> Lam.
	<i>Arctosyaphylos uva ursi</i>	<i>Musa × paradisiaca</i> L.
	<i>Argyreia nervosa</i> (Burm. f.) Bojer.	<i>Ocimum sanctum</i>
	<i>Asphodelus tenuifolius</i> Cav	<i>Ocimum tenuiflorum</i> L.

	<i>Asplenium ceterach</i> L.	<i>Oenothera biennis</i> L.
	<i>Asplenium hemionitis</i> L.	<i>Olea europaea</i>
	<i>Asplenium scolopendrium</i> L.	<i>Oxalis corniculata</i> L.
	<i>Arum rupicola</i> Boiss.	<i>Paronychia argentea</i> Lam.
	<i>Azadirachta indica</i> A. Juss.	<i>Paederia foetida</i> L.
	<i>Basella alba</i> L.	<i>Peperomia pellucida</i> (L.) Kunth
	<i>Barbarea vulgaris</i> R.Br.	<i>Phyllanthus niruri</i>
	<i>Bauhinia forficata</i> Link	<i>Phyllanthus lanceolatus</i> Poir.
	<i>Berberis integerrima</i> Bunge.	<i>Picea smithiana</i> (Wall.) Boiss.
	<i>Betula pendula</i> Roth.	<i>Piper aduncum</i> L.
	<i>Betula lenta</i> L	<i>Piper longum</i> L
	<i>Betula utilis</i> D Don.	<i>Plantago lanceolata</i> L.
	<i>Biophytum abyssinicum</i> Steud. Ex A. Rich.	<i>Plectranthus amboinicus</i> (Lour.) Spreng.
	<i>Boldoa purpurascens</i> cav	<i>Polygonum cognatum</i> Meisn.
	<i>Brassica napus</i> L.	<i>Populus alba</i> L.
	<i>Bryophyllum calycinum</i> Salisb.	<i>Portulaca oleracea</i>
	<i>Cassia auriculata</i> L.	<i>Prosopis farcta</i>
	<i>Calendula officinalis</i> L.	<i>Pulmonaria officinalis</i> L.
	<i>Carissa opaca</i> Stapf ex Haines.	<i>Raphanus sativus</i> L
	<i>Celastrus paniculatus</i>	<i>Rosa canina</i> L.
	<i>Celtis timorensis</i> .	<i>Rosmarinus</i>
	<i>Chenopodium album</i> L.	<i>Rubus caesius</i>
	<i>Chimaphila maculate</i> (L.) Pursh.	<i>Ruscus aculeatus</i> L.
	<i>Cichorium intybus</i> L.	<i>Ruta graveolens</i> L.
	<i>Cissus adnata</i> Roxb.	<i>Sapium sebiferum</i> (L.) Roxb.
	<i>Cissus gongylodes</i> (Baker)	<i>Saponaria mesogitana</i> Boiss.
	<i>Coleus amboinicus</i> Lour	<i>Smilax aspera</i> L.
	<i>Corallodiscus lanuginosus</i>	<i>Solidago virgaurea</i> L.
	<i>Coriandrum sativum</i> L.	<i>Spergularia rubra</i>
	<i>Crataegus aronia</i> (L.) Bosc ex DC.	<i>Spermacoce hispida</i> L
	<i>Cucumis sativus</i> L.	<i>Stachytarpheta indica</i> (L.) Vahl.
	<i>Cyclea peltata</i> (Lam.) Hook.f. & Thomson	<i>Tephrosia purpurea</i> (L.) Pers.

	<i>Didymocarpus pedicellatus</i> R.Br.	<i>Teucrium polium</i> L
	<i>Duranta erecta</i>	<i>Thunbergia alata</i> Bojer ex Sims.
	<i>Ecbolium viride</i> (Forssk.) Alston	<i>Thymus kotschyanus</i>
	<i>Eleusine indica</i> Linn. (poaceae)	<i>Thymus migricus</i>
	<i>Enicostema axillare</i> (Poir. ex Lam.) A. Raynal.	<i>Tinospora purpurea</i> (L.) Pers.
	<i>Eriobotrya japonica</i> (Thunb.) Lindl.	<i>Tinospora cordifolia</i> (Willd.) Miers.
	<i>Eupatorium birmanicum</i> DC.	<i>Tournefortia acuminata</i> A.DC.
	<i>Ficus carica</i> L.	<i>Trapogon buphtalmoides</i> (DC.) Boiss.
	<i>Fraxinus excelsior</i> L.	<i>Trianthema portulacastrum</i> L
	<i>Forsskaolea tenacissima</i> L.	<i>Tribulus terrestris</i> L.
	<i>Glechoma hederaceae</i> L.	<i>Triclisia gilletti</i>
	<i>Hedera helix</i> L.	<i>Vitis vinifera</i>
9	FLOWERS	
	<i>Alcea pallida</i> (Willd.) Waldst. & Kit.	<i>Inula oculus-christi</i> L
	<i>Anthemis nobilis</i> L.	<i>Matricaria chamomilla</i> L.
	<i>Borassus flabellifer</i> L.	<i>Moringa oleifera</i> Lam.
	<i>Cassia occidentalis</i> L.	<i>Musa × paradisiaca</i> L.
	<i>Calendula officinalis</i> L.	<i>Opuntia ficus-indica</i> (L.) Mill.
	<i>Cocus nucifera</i> L.	<i>Panicum miliaceum</i> L.
	<i>Cymbopogon schoenanthus</i> (L.) Spreng.	<i>Phlogacanthus thyrsoformis</i> Hardow (Mabb)
	<i>Echinops spinosus</i> L.	<i>Rosa canina</i> L. (rose hips)
	<i>Eryngium campestre</i> L.	<i>Rosa indica</i> L.
	<i>Helichrysum arenarium</i> (L.) Moench	<i>Rubus fruticosus</i> Lour.
	<i>Helichrysum plicatum</i> DC. Subsp. <i>plicatum</i>	<i>Solidago virgaurea</i> L.
	<i>Hibiscus sabdariffa</i> L.	<i>Taraxacum hybernum</i> Steven
		<i>Zea mays</i> L. (tea of corn silk)
10	FRUITS	
	<i>Ammi visnaga</i> (L.) Lam.	<i>Neolamarckia cadamba</i>
	<i>Aegle marmelos</i> (L.) Correa	<i>Nigella sativa</i>
	<i>Ananas comosus</i> (L.) Merr.	<i>Paeonia officinalis</i>
	<i>Artemisia abrotanum</i> L.	<i>Peganum harmala</i>

	<i>Averrhoa carambola</i> L.	<i>Pedaliium murex</i>
	<i>Ananas comusus</i> (L.) Meerr.	<i>Peucedanum grande</i> C.B.Clarke.
	<i>Benincasa hispida</i> (Thunb.) Cogn.	<i>Phillyrea latifolia</i> L.
	<i>Bombax ceiba</i> L.	<i>Phyllanthus emblica</i> L.
	<i>Brassica oleracea</i> L.	<i>Physalis alkekengi</i> L
	<i>Bunium persicum</i>	<i>Pinus brutia</i> Ten
	<i>Cannabis sativa</i> L	<i>Pinus eldarica</i> Medw.
	<i>Cassia fistula</i> L.	<i>Piper cubeba</i>
	<i>Cordia ecalyculata</i> Vell.	<i>Piper longum</i> .
	<i>Cordia grandis</i> Roxb.	<i>Piper nigrum</i>
	<i>Citrullus vulgaris</i>	<i>Pimpinella anisum</i>
	<i>Cucumis melo</i>	<i>Platanus orientalis</i>
	<i>Cucumis sativus</i>	<i>Prunus avium</i>
	<i>Elettaria cardamomum</i> (L.) Maton.	<i>Punica granatum</i>
	<i>Emblica officinalis</i> Gaertn	<i>Rosa canina</i>
	<i>Gmelina arborea</i> Roxb.	<i>Rhus succedanea</i> L.
	<i>Gossypium herbaceum</i> L.	<i>Ruscus hypoglossum</i> L.
	<i>Ficus carica</i> L.	<i>Rubus ellipticus</i>
	<i>Ficus palmata</i> Forssk.	<i>Rubus fruticosus</i>
	<i>Foeniculum vulgare</i> Mill.	<i>Serenoa repens</i> (W.Bartram) Small
	<i>Juniperus pseudosabina</i> Fisch. & C.A.Mey.	<i>Solanum xanthocarpum</i>
	<i>Kigelia pinnata</i> (Jacq.) DC.	<i>Spondias axillaris</i> Roxb.
	<i>Lagenaria siceraria</i> (Molina) Standl.	<i>Syzygium cumini</i>
	<i>Levisticum officinale</i>	<i>Tamarindus indica</i> L.
	<i>Lithospermum officinale</i>	<i>Tribulus terrestris</i> L.
	<i>Manilkara zapota</i>	<i>Trigonella foenum-graecum</i> L.
	<i>Momordica cochinchinensis</i> (Lour.) Spreng.	<i>Viburnum opulus</i> L.
	<i>Morinda citrifolia</i>	<i>Xenostegia tridentata</i>
11	SEEDS	
	<i>Alcea pallida</i> (Willd.) Waldst & Kit.	<i>Nigella sativa</i> L.
	<i>Celosia argentea</i> L.	<i>Phaseolus vulgaris</i> L.
	<i>Ensete superbum</i> (Roxb.) Cheesman	<i>Piper nigrum</i>

	<i>Holarrhena pubescens</i>	<i>Pongamia pinnata</i>
	<i>Lupinus varius</i> subsp.varius	<i>Raphanus sativus</i> L.
	<i>Macrotyloma uniflorum</i> (Lam.) Verdc.	<i>Trachyspermum ammi</i>
	<i>Manilkara zapota</i> (L.) P. Royen (kernel)	<i>Trionella foenum-graecum</i> L.
	<i>Momordica cochinchinensis</i> (Lour.) Spreng.	<i>Vigna unguiculata</i> (L.) Walp.
	<i>Myrtus communis</i> L.	
12	WHOLE PLANT	
	<i>Achillea millefolium</i> L.	<i>Gallium verum</i> L.
	<i>Achyranthes aspera</i>	<i>Haloxylon stocksii</i> (Boiss.) Benth. & Hook. f.
	<i>Acmella oleracea</i>	<i>Heliotropium crispum</i> Desf.
	<i>Actinodaphne angustifolia</i> Nees.	<i>Herniaria hirsute</i> L.
	<i>Aeonium canariense</i> (L.) Webb & Berthel	<i>Hypericum montbretii</i> Spach.
	<i>Aerva lanata</i>	<i>Kalimeris indica</i> (L.) Sch. Bip.
	<i>Ajuga chamaepitys</i> (L.) Schreb.	<i>Lamium album</i> L.
	<i>Ammi majus</i> L.	<i>Lemanea fluviatilis</i> L.
	<i>Alcea flavovirens</i> (Boiss.) Boiss	<i>Lindernia ruellioides</i> (Colsm.) Pennnell.
	<i>Alisma plantago-aquatica</i> L.	<i>Ludwigia perennis</i> L.
	<i>Alternanthera tenella</i> colla	<i>Malvella sherardiana</i> (L.) Jaub. & Spach.
	<i>Amaranthus spinosus</i> L.	<i>Meiogyne minuta</i> (G.Forst.)Less
	<i>Amaranthus viridis</i>	<i>Mentha pulegium</i> L.
	<i>Bonnaya brachiate</i>	<i>Merremia emarginata</i>
	<i>Capsella bursa-pastoris</i> Medik	<i>Micromeria biflora</i>
	<i>Cassia italica</i> (Mill.) Spreng.	<i>Mukia maderaspatana</i>
	<i>Centella asiatica</i>	<i>Myriogyne minuta</i> (G.Forst.) Less.
	<i>Cerasus avium</i>	<i>Ocimum basilicum</i>
	<i>Ceterach aureum</i> Buch.	<i>Ocimum tenuiflorum</i>
	<i>Coccinia indica</i> Wight & Arn.	<i>Oldenlandia herbacea</i>
	<i>Costus spicatus</i> (Jacq.) Sw.	<i>Origanum majorana</i> L.
	<i>Citrullus colocynthis</i> (L.) Shrad.	<i>Orthosiphon grandiflorus</i> Bold.
	<i>Dendrophthoe falcata</i> (L.f.) Ettingsh.	<i>Orthosiphon stamineus</i> Benth
	<i>Desmodium microphyllum</i> (Thumb.) DC	<i>Parmelia perlata</i>

<i>Desmodium styracifolium</i> (Osbeck) Merr.	<i>Pedaliium murex</i> L.
<i>Didymocarpus tomentosus</i> Wight	<i>Pergularia daemia</i>
<i>Duchesnea indica</i> (Jacks.) Focke.	<i>Phyllanthus fraternus</i> G.L. Webster
<i>Echinops echinattus</i> Roxb.	<i>Phyllanthus niruri</i>
<i>Ehydra fluctuans</i> Lour.	<i>Phyllanthus urinaria</i>
<i>Enhydra fluctuans</i> Lour.	<i>Pistacia lentiscus</i>
<i>Erigeron karvinskianus</i> DC.	<i>Plantago coronopus</i>
<i>Equisetum arvense</i>	<i>Polygonum aviculare</i>
<i>Equisetum bogotense</i>	<i>Pratia nummularia</i> (Lam.) A.Braun & Asch.
<i>Equisetum debile</i>	<i>Primula veris</i> L.
<i>Euphorbia hirta</i> .	<i>Rubia cordifolia</i> L.
<i>Euphorbia prostrata</i> Aiton	<i>Solanum nigrum</i> L.
<i>Euphorbia serpens</i> Kunth.	<i>Sphaeranthus indicus</i> L.
<i>Forsskaolea angustifolia</i> Retz.	<i>Teucrium chamaedrys</i> L.
<i>Fragaria nilgerrensis</i>	<i>Teucrium scordium</i> L.
<i>Fumaria officinalis</i> L.	<i>Withania somnifera</i> (L.) Dunal.

ANIMAL SOURCE:

The animal source with lithiolytic activity are listed in table:3 (Mohammad Shamim Khan et al., 2016)

TABLE:3 ANIMAL SOURCE WITH ANTIUROLITHIATIC AND DIURTIC ACTIVITY

S.NO.	NAME	PARTS USED
1.	Burnt Scorpion	Whole part
2.	Sparrow	Brain
3.	Purified Earth worm	Whole part
4.	Burnt rabbit	Whole part

MARINE SOURCES:

The marine source with antiurolithiatic activity and diuretic property are listed in table:4(Mohammad Shamim Khan et al., 2016)

TABLE:4 MARINE SOURCE WITH ANTIUROLITHIATIC AND DIURETIC ACTIVITY

S.NO.	CHEMICAL CONSTITUENT	TYPE OF ORGANISM	REFERENCE
1.	Chitosan (chitin) biopolymer	Marine and terrestrial organisms Macroalgae, microalgae	Moacir Fernandes Queiroz et al., 2014
2.	Fucoxanthine (active) carotenoid	Macroalgae, microalgae	RulinWang et al., 2020
3.	C-Phycocyanin	Spirulina plantensis(blue-green algae)	DéboraPez Jaeschke et al., 2021

MICROBIAL SOURCE:

Oxalate degrading bacteria such as Bifidobacterium species, bacillus species, oxalobacter formigenes, porphyromonas gingivalis, bacillus subtilis and Lactobacillus plantarum show antiurolithiatic activity by degrading oxalate crystals.(Rouhi Afkari et al., 2019)

MINERAL SOURCE:

The mineral source with diuretic and lithiolytic activity are listed in table:5(Mohammad Shamim Khan et al., 2016)

TABLE:5 MINERAL SOURCE WITH LITHIOLYTIC AND DIURETIC ACTIVITY:

S.NO.	NAME
1.	Alum
2.	Burnt crystal
3.	Fish stones
4.	Lapis Judaicus / Jews Stone
5.	Potassium carbonate
6.	Potassium nitrate

4. CONCLUSION

Urolithiasis is outstanding considerable malady of the urogenital system and also a great origin of morbidity. With its several aetiologies and high incidence of re occurrence, renal tract stone malady come up with a medical dispute.

Despite of many traditional curatives feasible for the urolithiasis, but re occurrence of stone formation still remains as a problem still to date there is no standard drug available and

foremost disadvantage in the maturing of a standard drug may be the multifactorial kind of the disease and several chemical varieties of renal stones.

Many curatives has been serviced during the ages to cure renal stones. In the classic system of medicine, utmost of the therapies was taken from plants and they were proven to be useful though the rationale at the rear of their used is not Triumphant through systematic pharmacological and clinical studies apart from some compounded herbaceous medicines and plants. Herbal medicines contain several phytonutrients and exert their beneficial outcomes by various mechanisms in treatment of urolithiasis.

CONSENT

Not applicable

ETHICAL APPROVAL

Not applicable

exist

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. Ajj A, Abdul W, Nasreen J, Alia Bilal, Syeda H, Efficacy of *Adiantum capillus veneris* Linn in chemically induced urolithiasis in rats, *Journal of Ethnopharmacology*, 2013 Volume 146, Issue 1, Pages 411-416. DOI: [10.1016/j.jep.2013.01.011](https://doi.org/10.1016/j.jep.2013.01.011)
2. Atul M, Vishavas R, Dishant Desai, Jaymin M, Vivek Parekh, Evaluation for the anti-urolithiatic activity of *Launaea procumbens* against ethylene glycol-induced renal calculi in rats, *Toxicology Reports*, 2014 Volume 1, 2014, Pages 46-52. DOI: [10.1016/j.toxrep.2014.03.006](https://doi.org/10.1016/j.toxrep.2014.03.006)
3. Kaleeswaran B, Ramadevi S, Murugesan R, Srigopalram.S, Suman.T, Balasubramanian.T, Evaluation of anti-urolithiatic potential of ethyl acetate extract of *Pedaliu murex* L. on struvite crystal (kidney stone), *Journal of Traditional and Complementary Medicine*, 2017 Volume 9, Issue 1, Pages 24-37. <https://dx.doi.org/10.1016%2Fj.jtcme.2017.08.003>

4. Kumar B.N, Abdul Wadud, Nasreen Jahan, Ghulamuddin Sofi, Humaira Bano, Shaikh Ajj Ahmed Makbul, Sadique Husain, Antilithiatic effect of *Peucedanum grande* C. B. Clarke in chemically induced urolithiasis in rats, *Journal of Ethnopharmacology*, 2016 Volume 194,2016, Pages 1122-1129.DOI: [10.1016/j.jep.2016.10.081](https://doi.org/10.1016/j.jep.2016.10.081)
5. Vyas BA, Vyas R.B, Joshi. S.V, Santani D.D, Antirolithiatic Activity of Whole-Plant Hydroalcoholic Extract of *Pergularia daemia* in Rats,*Journal of Young Pharmacists*, 2011 Volume 3, Issue 1,2011,Pages 36-40.doi: [10.4103/0975-1483.76417](https://doi.org/10.4103/0975-1483.76417)
6. Basavaraj M. Dinnimath, Sunil S. Jalalpure, Umesh K. Patil, Antirolithiatic activity of natural constituents isolated from *Aerva lanata*,*Journal of Ayurveda and Integrative Medicine*, 2017 Volume 8, Issue 4, Pages 226-232.DOI: [10.1016/j.jaim.2016.11.006](https://doi.org/10.1016/j.jaim.2016.11.006)
7. Bawari S, Sah AN, Tewari D. Anticalcifying effect of *Daucus carota* in experimental urolithiasis in Wistar rats. *J Ayurveda Integr Med*. 2020 Jul-Sep;11(3):308-315. DOI: [10.1016/j.jaim.2018.12.003](https://doi.org/10.1016/j.jaim.2018.12.003)
8. Adegbaaju O D, Otunola G.A, Afolayan A.J, Effects of growth stage and seasons on the phytochemical content and antioxidant activities of crude extracts of *Celosia argentea* L., *Heliyon*, 2020 Volume 6, Issue 6.DOI: [10.1016/j.heliyon.2020.e04086](https://doi.org/10.1016/j.heliyon.2020.e04086)
9. DéboraPez Jaeschke^aIngridRocha Teixeira^aLigiaDamasceno Ferreira Marczak^a GiovanaDomeneghini Mercali^b; Phycocyanin from *Spirulina*: A review of extraction methods and stability; *Food Research International*; 2021 Volume 143, 110314.DOI: [10.1016/j.foodres.2021.110314](https://doi.org/10.1016/j.foodres.2021.110314)
10. Deepti Sharma, Yadu Nandan Dey, Indu Sikarwar, Richa Sijoria, Manish M. Wanjari, Ankush D. Jadhav, In vitro study of aqueous leaf extract of *Chenopodium album* for inhibition of calcium oxalate and brushite crystallization, *Egyptian Journal of Basic and Applied Sciences*, 2016 Volume 3, Issue 2, 2 ,Pages 164-171.<https://doi.org/10.1016/j.ejbas.2016.02.001>
11. Dulce María González Mosquera, Yannarys Hernández Ortega, Pedro César Quero, Rafael Sosa Martínez, Luc Pieters, Antirolithiatic activity of *Boldoa purpurascens* aqueous extract: An in vitro and in vivo study, *Journal of Ethnopharmacology*, 2020 Volume 253, 112691.DOI: [10.1016/j.jep.2020.112691](https://doi.org/10.1016/j.jep.2020.112691)
12. Humaira Bano, Nasreen Jahan, Shaikh Ajj Ahmed Makbul, Kumar B.N, Sadique Husain, Atiya Sayed, Effect of *Piper cubeba* L. fruit on ethylene glycol and ammonium chloride induced urolithiasis in male Sprague Dawley rats, *Integrative Medicine Research*, 2018 Volume 7, Issue 4, Pages 358-365.DOI: [10.1016/j.imr.2018.06.005](https://doi.org/10.1016/j.imr.2018.06.005)

13. Iman S, Saleem U and Ahmad B, Preclinical assessment of antiurolithiatic activity of mangifera indica seeds on ethylene glycol induced urolithiasis in rat model. *Pak Vet J*, 2021 41(1): 92-96. DOI: [10.29261/pakvetj/2020.028](https://doi.org/10.29261/pakvetj/2020.028)
14. Jagannath N, Chikkannasetty SS, Govindadas D, Devasankaraiah G. Study of antiurolithiatic activity of Asparagus racemosus on albino rats. *Indian J Pharmacol*. 2012;44(5):576-579. doi: [10.4103/0253-7613.100378](https://doi.org/10.4103/0253-7613.100378)
15. K V S R G Prasad, R Abraham, K Bharathi & K K Srinivasan Evaluation of Homonia riparia Lour. for Antiurolithiatic Activity in Albino Rats, *International Journal of Pharmacognosy*, 1997 35:4, 278-283. DOI: [10.1076/phbi.35.4.278.13309](https://doi.org/10.1076/phbi.35.4.278.13309)
16. Ma. Therese H. Garcia* , Carelle Juliene A. Malaguit, Ashley Joy D. Verte, Renz P. Cometa, Maricel E. Dimayacyac, April Bernadette P. Taganap, Ma. Cielo Aeb F. Quintia, Sheryl C. Aguila, and Omar A. Villalobos, Antiurolithiatic activity of ethanolic extract of Eleusine indica Linn. (Poaceae) leaves in ethylene glycol-induced urolithiasis in male Sprague Dawley rats, *Asia Pacific Journal of Allied Health Sciences*, Vol. 1, 2018, ISSN 2704-3568.
17. Mansoor, M., Jamil, M., Latif, N., Muhammad, S., Gull, J., Shoab, M., & Khan, A. Review Importance of Herbal Plants in the Management of Urolithiasis. *Biological Sciences - PJSIR*, 2019 62(1), 61-66. <https://doi.org/10.52763/PJSIR.BIOL.SCI.62.1.2019.61.66>
18. Mert İlhan, Burçin Ergene, Ipek Süntar, Serkan Özbilgin, Gülçin Saltan Çitoğlu, M. Ayşe Demirel, Hikmet Keleş, Levent Altun, Esra Küpeli Akkol, "Preclinical Evaluation of Antiurolithiatic Activity of *Viburnum opulus* L. on Sodium Oxalate-Induced Urolithiasis Rat Model", *Evidence-Based Complementary and Alternative Medicine*, vol. 2014, ArticleID 578103, 10 pages, 2014. <https://doi.org/10.1155/2014/578103>
19. Mittal, S. Tandon, S.K. Singla, C. Tandon, *In vitro* studies reveal antiurolithiceffectof *Terminalia arjuna* using quantitative morphological information from computerized microscopy, *International Brazilian Journal of Urology*, Vol,41(5):935-944, september - october, 2015, doi: 10.1590/S1677-5538.IBJU.2014.0547.
20. Moacir Fernandes Queiroz¹, Karoline Rachel Teodosio Melo², Diego Araujo Sabry³, Guilherme Lanzi Sassaki⁴, Hugo Alexandre Oliveira Rocha Does the use of chitosan contribute to oxalate kidney stone formation?; *Marine Drugs*; 2014 13(1):141-58. DOI: [10.3390/md13010141](https://doi.org/10.3390/md13010141)
21. Mohammad Shamim Khan, Qamrul Hasan Lari² and Mahmood Ahmad Khan; ANTI- UROLITHIATIC UNANI DRUGS – A REVIEW; *World Journal of Pharmaceutical Research*; 2016 Volume 5, Issue 12, 279-294.

22. Shazib Faridi MD¹, Somarendra Singh K²; Preliminary study of prevalence of urolithiasis in North-Eastern city of India; J Family Med Prim Care; 2020 Dec 31;9(12):5939-5943. DOI: [10.4103/jfmprc.jfmprc_1522_20](https://doi.org/10.4103/jfmprc.jfmprc_1522_20)
23. Mosquera DMG, Ortega YH, Quero PC, Martínez RS, Pieters L. Antiuro lithiatic activity of *Boldoa purpurascens* aqueous extract: An in vitro and in vivo study. J Ethnopharmacol. 2020 May 10;253:112691. DOI: [10.1016/j.jep.2020.112691](https://doi.org/10.1016/j.jep.2020.112691)
24. Rathod N.R, Dipak Biswas, Chitme H.R, Sanjeev Ratna, I.S. Muchandi, Ramesh Chandra, Anti-urolithiatic effects of *Punica granatum* in male rats, Journal of Ethnopharmacology, 2012 Volume 140, Issue 2, Pages 234-238. <https://doi.org/10.1016/j.jep.2012.01.003>
25. Nishi Saxena, Ameeta Agrgal, Study of Antiuro lithiatic activity of a formulated herbal suspension, Herba Polonica-agro, 2015 Vol. 61 No. 2 . DOI: <https://doi.org/10.1515/hepo-2015-0014>
26. Kachchhi NR*, Parmar RK, Tirgar PR, Desai TR, Bhalodia PN, EVALUATION OF THE ANTIUROLITHIATIC ACTIVITY OF METHANOLIC EXTRACT OF CELOSIA ARGENTEA ROOTS IN RATS, Kachchhi NR. et al. / International Journal of Phytopharmacology. 2012 3(3), 249-255.
27. Olayeriju, O.S., Crown, O.O., Elekofehinti, O.O. et al. Effect of moonseed vine (*Triclisia gilletii* Staner) on ethane-1,2-diol-induced urolithiasis and its renotoxicity in Wistar albino rats. Afr J Urol 2020 **26**, 4 . <https://doi.org/10.1186/s12301-020-0018-x>
28. Patel P.K, Patel M.A, Saralai M.G, Gandhi T.R, Antiuro lithiatic Effects of *Solanum xanthocarpum* Fruit Extract on Ethylene-Glycol-Induced Nephrolithiasis in Rats, Journal of Young Pharmacists, 2012 Volume 4, Issue 3, Pages 164-170. DOI: [10.4103/0975-1483.100022](https://doi.org/10.4103/0975-1483.100022)
29. Paula P.O. Salem, Nátalie B. Vieira, Daniela A. Garcia, Karen J. Nicácio, Danielle F. Dias, Ana C.C. de Paula, Diego M. Assis, Ivo S. Caldas, Romulo D. Novaes, Maria V. Marinho, Iara M.L. Rosa, Marisi G. Soares, Daniela A. et al., Anti-urolithiatic and anti-inflammatory activities through a different mechanism of actions of *Cissus gongylodes* corroborated its ethnopharmacological historic, Journal of Ethnopharmacology, 2020 Volume 253, 112655. DOI: [10.1016/j.jep.2020.112655](https://doi.org/10.1016/j.jep.2020.112655)
30. Poonam Verma¹, Vinod Gauttam¹, Ajudhia N Kalia¹ Comparative pharmacognosy of Pashanbhed; Ayurveda Integr Med; 2014 Apr;5(2):104-8. DOI: [10.4103/0975-9476.131728](https://doi.org/10.4103/0975-9476.131728)
31. Poppy Das, Kiran Kumar, Arunai Nambiraj, Reshma Rajan, Rajendra Awasthi, Kamal Dua, et al., Potential therapeutic activity of *Phlogacanthus thyriformis*

- Hardow (Mabb) flower extract and its biofabricated silver nanoparticles against chemically induced urolithiasis in male Wistar rats, *International Journal of Biological Macromolecules*, 2017 Volume 103, Pages 621-629. DOI: [10.1016/j.ijbiomac.2017.05.096](https://doi.org/10.1016/j.ijbiomac.2017.05.096)
32. R Vargas S, R.M Perez G, S Perez G, M.A Zavala S, C Perez G, Antiuro lithiatic activity of *Raphanus sativus* aqueous extract on rats, *Journal of Ethnopharmacology*, 1999 ,Volume 68, Issues 1–3, Pages 335-338. [https://doi.org/10.1016/S0378-8741\(99\)00105-1](https://doi.org/10.1016/S0378-8741(99)00105-1)
 33. Ram Prakash¹, Arunachalam² , Narayanasamy³ ; Prevalence and socio-demographic status on kidney stone patients in Thanjavur district, Tamil Nadu, India; *Int J Community Med Public Health*; 2019 6(5):1943-1947. DOI : <http://dx.doi.org/10.18203/2394-6040.ijcmph20191614>
 34. Ravindra V. Karadi, Navneet B. Gadge, K.R. Alagawadi, Rudraprabhu V. Savadi, Effect of *Moringa oleifera* Lam. root-wood on ethylene glycol induced urolithiasis in rats, *Journal of Ethnopharmacology*, 2006 Volume 105, Issues 1–2, Pages 306-311. DOI: [10.1016/j.jep.2005.11.004](https://doi.org/10.1016/j.jep.2005.11.004)
 35. Rouhi Afkari¹, Mohammad Mehdi Feizabadi², Alireza Ansari-Moghadam³, Tahereh Safari⁴, Mohammad Bokaeian¹; Simultaneous use of oxalate-degrading bacteria and herbal extract to reduce the urinary oxalate in a rat model: A new strategy; *Int Braz J Urol*; 2019 45(6):1249-1259. DOI: [10.1590/S1677-5538.IBJU.2019.0167](https://doi.org/10.1590/S1677-5538.IBJU.2019.0167)
 36. RulinWang^a Elsayed M.Younis^b VishnuPriya Veeraraghavan^c ChenfeiTian^d; Antiuro lithiatic effect of Fucoxanthin on ethylene glycol-induced renal calculus in experimental rats; 2020 *Journal of King Saud University - Science*; Volume 32, Issue 3, Pages 1896-1901. <https://doi.org/10.1016/j.jksus.2020.01.027>
 37. Sachin B. Agawane, Vidya S. Gupta, Mahesh J. Kulkarni, Asish K. Bhattacharya, Santosh S. Koratkar, Vamkudoth Koteswara Rao, Patho-physiological evaluation of *Duranta erecta* for the treatment of urolithiasis, *Journal of Ayurveda and Integrative Medicine*, 2019 Volume 10, Issue 1, 2019, Pages 4-11. DOI: [10.1016/j.jaim.2017.08.001](https://doi.org/10.1016/j.jaim.2017.08.001)
 38. Sahithi Adepu, Raviraj Anand Devkar, Shilpee Chaudhary, Manganahalli Manjunath Setty, WITHDRAWN: Antiuro lithiatic activity of *Alternanthera tenella* Colla in ethylene glycol induced urolithiasis in rats, *Asian Journal of Pharmaceutical Sciences*, 2015, ISSN 1818-0876. <https://doi.org/10.1016/j.ajps.2015.11.003>

39. Salman Ahmed, Muhammad Mohtasheemul Hasan, Zafar Alam Mahmood;
Antirolithiatic plants in different countries and cultures; JPP ; 2016 5(1): 102-115.
40. Salman Ahmed, Muhammad Mohtasheemul Hasan, Zafar Alam Mahmood;
Antirolithiatic plants: Multidimensional pharmacology; JPP ; 2016 5(2): 04-24
41. Salman Ahmed¹ , Muhammad Mohtasheemul Hasan^{1*} and Zafar Alam Mahmood²;
Antirolithiatic plants: Formulations used in different countries and cultures; Pak. J. Pharm. Sci., 2016 Vol.29, No.6, pp.2129-2139.
42. Satish, S.; Mahesh, C.M.; Gowda,K.P.S; David Banji, Study on the Antirolithiatic activity of *Cynodon dactylon* root stalk extract in Albino rats; CAB Direct, Biomed 2009, vol. 4, No. 4 pp.384-391.
43. Shengbao Wang, Xiaoran Li, Junsheng Bao & Siyu Chen Protective potential of *Angelica sinensis* polysaccharide extract against ethylene glycol-induced calcium oxalate urolithiasis, Renal Failure, 2018 40:1, 618-627.
44. Shruti Gupta, Shamsheer SK; Kidney stones: Mechanism of formation, pathogenesis and possible treatments; J Biomol Biochem ; 2018 2(1):1-5.
45. Tanzeer Kaur, Rakesh K. Bijarnia, Surinder K. Singla, Chanderdeep Tandon, In vivo efficacy of *Trachyspermum ammi* anticalcifying protein in urolithiatic rat model, Journal of Ethnopharmacology, 2009 Volume 126, Issue 3, Pages 459-462.
46. Tushar Shelke, Sandesh Wayal, Shiv Gunjegaokar, Sachin Gaikwad, Amol Shirsath, Sunil Hadke; An Overview on Indian Medicinal Plants with Antirolithiatic Activity; Journal of Pharmaceutical Research & Clinical Practice, 2014 4(3):33-40.
47. Sasikala V, Ramu Radha S, Bavaniamma Vijayakumari, In vitro evaluation of *Rotula aquatica* Lour. for antirolithiatic activity, Journal of Pharmacy Research, 2013 Volume 6, Issue 3, 2013, Pages 378-382.
<https://doi.org/10.1016/j.jopr.2013.02.026>

48. Victoriano Romero , Haluk Akpınar, Dean G Assimios; Kidney stones: a global picture of prevalence, incidence, and associated risk factors; Rev Urol.; Spring 2010;12(2-3):e86-96. PMID: PMC2931286
49. Vishal N Ratkalkar, MD and Jack G Kleinman, MD; Mechanisms of Stone Formation; Clin Rev Bone Miner Metab. 2011 Dec; 9(3-4): 187–197. doi: [10.1007/s12018-011-9104-8](https://doi.org/10.1007/s12018-011-9104-8)
50. Yu Liu, Yuntian Chen, Banghua Liao, Deyi Luo, Kunjie Wang, Hong Li. Guohua Zeng; Epidemiology of urolithiasis in Asia; Asian J Urol. 2018 5(4): 205–214. <https://doi.org/10.1016/j.ajur.2018.08.007>