

Original Research Article

ANTIBACTERIAL EFFECT OF CINNAMON OIL AGAINST UROPATHOGENIC MULTIDRUG RESISTANT

ABSTRACT

Cinnamon is known for its antimicrobial activity and the aim of this study was to investigate the effect of its essential oil against ten of the multidrug-resistant uropathogenic by agar well diffusion assays. The diameters of the inhibition zone to the Cinnamon oil were 27 mm for *S. aureus*, 24 mm for *E. coli*, 20 mm for *P. aeruginosa*, 22 mm for *K. pneumoniae*, 23 mm for *E. aerogenes* & *P. mirabilis*, 24 mm for *E. faecalis*, and 27 mm for *N. gonorrhoeae*, *A. baumannii*, *S. epidermis*. In this study, the antimicrobial effects of the cinnamon essential oil are evaluated by determining the minimum inhibitory concentration (MIC), the inhibition zone, and minimum bactericidal concentration (MBC). Cinnamon was the most effective agent in inhibiting *A. baumannii*, *N. gonorrhoeae*, *S. epidermis*, *E. faecalis* and *E. coli* with the lowest MIC (0.0313%) while *S. aureus*, *E. aerogenes* and *P. aeruginosa* with MIC (0.125%) while *P. mirabilis* was 0.0625% in our study. The MBC was 0.25% for *A. baumannii*, 0.5 % for *N. gonorrhoeae*, *S. epidermis*, *E. faecalis*, and *K. pneumoniae*, while 1% *S. aureus*, *E. aerogenes*, *P. mirabilis*, *E. coli*, and *P. aeruginosa*.

KEY WORDS: Antimicrobial, Cinnamon Oil, Agar Well Diffusion, MIC & MBC, Uropathogenic.

1. INTRODUCTION

Cinnamon is a common spice that has been used for centuries by different cultures around the world. Cinnamon oil is extracted from the shredded cinnamon tree's dried inner bark. It is native to Sri Lanka and India's Malabar Coast.

Jamaica and Brazil are also home to this species (Bhattacharya, 2016)¹. There has been a constant increase in the search for alternative and efficient compounds from cinnamon oil for multidrug-resistant bacteria, aimed at partial or total replacement of antimicrobial chemical

additives where is Cinnamon oil has been shown to have numerous health benefits, especially as an anti-inflammatory. Its phytochemical elements, such as phenolic and volatile compounds, are primarily responsible for this (Muhammad & Dewettinck, 2017)². Various types of extraction methods are used to obtain cinnamon oil, which are solvent extraction, ultrasonic extraction, hydro-distillation, shaking, and stirring with organic solvents (Kamaliroosta et al., 2012)³. The chief significance of the above the study was therefore to test the antimicrobial activity of cinnamon oil against uropathogenic bacteria where is urinary tract infection (UTI) is the second most common infection next to respiratory tract in human body. The disease affects people of all ages and both gender, about 150 million people are diagnosed with UTI yearly. As well as *Enterobacteriaceae*, gram-negative facultative anaerobic bacilli cause UTIs. The most common of these bacteria is *Escherichia coli* which forms about 90% of all Urinary tract infections. The other one is *Klebsiella* and *Proteus*, additionally, *Pseudomonas* that cause a complicated infections, especially in women. *Staphylococci* may cause 5-10% of UTIs in many populations. *E. coli* usually causes a child's first infection. (Brkic, et al., 2010)⁴. *Staphylococcal*

infections, especially those due to *Staphylococcus saprophyticus* are common causes of urinary tract infection among female adolescent.

2. MATERIALS AND METHODS

2.1 Essential Oils Distillation

Cinnamon commercial essential oils (purity $\geq 98\%$) was purchased from Bluray Nutritional Products runs under the brand name Bluray Nutritional. Essential oils (EO) were obtained by water-steam distillation for 6 hours by Clevenger apparatus at 100°C for 6 hours.

2.2 Antimicrobial activity test

Agar Well Diffusion Method (AWD)

Bacterial suspension ($1-2 \times 10^6$ cells/mL.) was spreaded on Mueller Hinton agar plates using sterile cotton swab, then wells with a diameter of 6 mm were made on the surface and filled with (20µl), of cinnamon Standard Ciprofloxacin (20µl) were added to the 5mm well on agar plates. The treated plates with *E.coli*, *P.aeruginosa*, *K.pneumoniae*, *P.mirabilis*, *E.aerogenes*, *E.faecalis*, *A.baumannii*, *N.gonorrhoeae*, *S.aureus* and *S.epidermis* were incubated at 37°C for 24hrs. After incubation the treated plates were observed for zone of

inhibition around the wells were recorded in millimeters (Kordali *et al.*, 2005)⁵.

2.3 Determination of minimal inhibitory and bactericidal concentrations

To facilitate the dilution of the oil and the reading of the results, Brain-Heart Infusion broth supplemented with 0.15% agar was used (Chuah, *et al.*, 2014 and Owuama, 2017)^{6&7}. The micro dilution broth technique was performed as follows: double serial dilutions of selected EOs were prepared ranging from 1%, 0.5%, 0.25%, 0.125%, 0.0625%, 0.03125% & 0.015625% in 280µl tryptone broth. Standard - Ciprofloxacin (8µg): 8, 4, 2, 1, 0.5, 0.25 and 0.125 µg in 280 µl tryptone broth. Control-Tryptone broth was inoculated with respective cultures and without a test sample. Then, the plate was incubated at 35°C for 20–24 hr under aerobic conditions. After incubation O.D @ 590 nm was measured using plate reader Resazurin (0.015 %) was added to all wells (30 µL per well) and further incubated for 2–4 h for the observation of colour change. On completion of the incubation, columns with no colour change (blue Resazurin colour remained unchanged) were scored as above the MIC value. Metabolism of Resazurin by active bacterial cells leads to reduction of Resazurin (Purple-blue) to resorufin (pink-colorless) pink color.

Determine MIC as Minimum concentration of drug giving 50% inhibition of OD as compared with control. Turbidity indicates growth of the microorganism and the MIC is the lowest concentration where no cell metabolism is observed visually. To determine the MBC, the dilution representing the MIC and at least two-four of the more concentrated test product dilutions are plated (0.1mL) on SCDA plates and incubated at 37°C for 24hrs and enumerated to determine viability of cells CFU/mL. The dilution concentration that produces no growth is recorded as the MBC. The MBC is the lowest concentration that demonstrates a pre-determined reduction (such as <99.9%) in CFU/mL when compared to the MIC.

3. RESULTS AND DISCUSSION

Cinnamon essential oil was screened for their antibacterial activity by two techniques. The target cultures used were *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumonia*, *Proteus mirabilis*, *Enterobacter aerogenes*, *Enterococcus faecalis*, *Acinetobacter baumannii*, *Neisseria gonorrhoeae*, *Staphylococcus aureus* and *Staphylococcus epidermis*. Our research exhibited a strong variation in the antibacterial activity selected essential oil against ten bacterial species is summarized in Table 1 and Plate 1. The

results revealed that the selected essential oil showed antibacterial activity with varying magnitudes. Cinnamon has been employed for centuries as food preservatives and as medicinal plants due to their antioxidant and antimicrobial activities. Nowadays, many reports confirmed the antibacterial, antifungal, antiviral, and anticarcinogenic properties of spice plants (Shan, *et al.*, 2005)⁸. In the present study, the antibacterial effect of cinnamon oil against some important uropathogens studied and their potencies were qualitatively assessed by the presence or the absence of inhibition zone diameter. The results indicated that the essential oil of cinnamon had substantial antibacterial activity against *E.coli*, *P.aeruginosa*, *K.pneumoniae*, *P.mirabilis*, *E.aerogenes*, *E.faecalis*, *A.baumannii*, *N.gonorrhoeae*, *S.aureus*, and *S.epidermis* (≥ 20 mm inhibition zone diameter). (Table 2 and plat11). The results of the present study agree with those reported by (Braga *et al.*, 2007; Bayoub *et al.*, 2010)^{9&10} who stated that against Gram-positive and Gram-negative

bacteria. In our study was find the results of showed antimicrobial activity against all of the bacterial strains used in this study: *Staphylococcus aureus* with an inhibitory zone of 27 mm, *E.coli* with an inhibitory zone of 24 mm *P.aeruginosa* with an inhibitory zones of 20 mm, *K.pneumoniae* with an inhibitory zone of 22 mm, *P.mirabilis* with an inhibitory zone of 23 mm, *E.aerogenes* with an inhibitory zone of 23 mm, *E.faecalis* with an inhibitory zone of 24 mm, *A.baumannii* with an inhibitory zone of 27 mm, *N.gonorrhoeae* with an inhibitory zone of 27 mm, and *S.epidermis* with an inhibitory zone of 27 mm.

Table 1: Inhibitory activity of test Cinnamon Oil against uropathogens

Z	Test Compounds	Conc.per well	Zone of inhibition (mm)
<i>E.coli</i>	Ciprofloxacin (Std)(20µl)	2µg	21
	Cinnamon oil(20µl)	2mg	24
<i>P.aeruginosa</i>	Ciprofloxacin (Std)(20µl)	2µg	20
	Cinnamon oil (20µl)	2mg	20

<i>K.pneumoniae</i>	Ciprofloxacin (Std)(20µl)	2µg	19
	Cinnamon oil (20µl)	2mg	22
<i>P.mirabilis</i>	Ciprofloxacin (Std)(20µl)	2µg	22
	Cinnamon oil (20µl)	2mg	23
<i>E.aerogenes</i>	Ciprofloxacin (Std)(20µl)	2µg	19
	Cinnamon oil (20µl)	2mg	23
<i>E.faecalis</i>	Ciprofloxacin (Std)(20µl)	2µg	23
	Cinnamon oil (20µl)	2mg	24
<i>A.baumannii</i>	Ciprofloxacin (Std)(20µl)	2µg	19
	Cinnamon oil (20µl)	2mg	27
<i>N.gonorrhoeae</i>	Ciprofloxacin (Std)(20µl)	2µg	20
	Cinnamon oil (20µl)	2mg	29
<i>S.aureus</i>	Ciprofloxacin (Std)(20µl)	2µg	22
	Cinnamon oil (20µl)	2mg	27
<i>S.epidermis</i>	Ciprofloxacin (Std)(20µl)	2µg	21
	Cinnamon oil (20µl)	2mg	27

Plate 1: Inhibitory activity of Cinnamon Oil against test organisms S – Standard (Ciprofloxacin); C - Control (distilled water)




Plate 2(a) - E.coli




Plate 2 (b) - P.aeruginosa





Plate 2(c) - *K.pneumoniae*




Plate 2(d) - *P.mirabilis*




Plate 2 (e) - *E.aerogenes*




Plate 2(f) - *E.faecalis*




Plate 2(g) - *A.baumannii*




Plate 2(h) - *N.gonorrhoeae*




Plate 2(i) - *S.aureus*

Plate 2(j) - *S.epidermis*

Plate 2(a-j): Inhibitory activity of Cinnamon Oil against test organisms S – Standard (Ciprofloxacin); C - Control (distilled water).

The effect of Cinnamon against tested pathogenic bacteria using Minimum Inhibitory Concentration (MIC) as the lowest concentration of an antimicrobial to prevent the visible growth of bacteria and Minimum Bactericidal Concentration

(MBC) is the lowest concentration of an antibacterial agent required to kill a bacterium. The results of *in vitro* antibacterial activity by crude Cinnamon at five concentrations (1, 50, 25, 125, and .00625 conc. (%)) against *E. coli*, *P.*

aeruginosa, *K. pneumonia*, *P.mirabilis*, *E. aerogenes*, *E. faecalis*, *A. baumannii*, *Neisseria gonorrhoeae*, *S. aureus* and *S.epidermis* by broth dilution technique are summarized in Table 2

We carried out antibiotic susceptibility testing to determine the activity of cinnamon essential oil against growing uropathogens. As shown in Table 2, Cinnamon was the most effective agent in inhibiting *A. baumannii*, *N.gonorrhoeae*, *S. epidermis*, *E. faecalis* and *E. coli* with the lowest MIC (0.0313%) while *S. aureus*, *E.aerogenes* and *P. aeruginosa*

with MIC (0.125%) while *Proteus mirabilis* was 0.0625% in our study. The MBC was 0.25% for *A.baumannii*, 0.5 % for *N.gonorrhoeae*, *S. epidermis*, *E. faecalis*, and *K. pneumonia*, while 1% *S. aureus*, *E. aerogenes*, *Pr. mirabilis*, *E. coli*, and *P.aeruginosa*. Therefore the growth of uropathogenic was efficiently suppressed by Cinnamon oil at the concentration of 0.03%, 0.0625% and 0.125. Clinical drug Ciprofloxacin included as a control inhibited the growth of uropathogenic growing cells with MIC of 1 and 0.5 µg/mL (Table 3).

Table 2: Minimum inhibitory activity of Samples against uropathogenic bacteria

Test Organism	OD & %	Cinnamon oil concentration %								MIC %
		0.0000	0.0156	0.0313	0.0625	0.1250	0.2500	0.5000	1.0000	
E.coli	OD at 590nm	0.5335	0.3591	0.2601	0.0929	0.0297	0.0085	0.0021	0.00019	0.0313%
	% Inhibition	0.0000	32.6898	51.2390	82.5867	94.4330	98.4049	99.6109	99.9765	
P.aeruginosa	OD at 590nm	0.5835	0.4725	0.3816	0.3130	0.2446	0.1405	0.0098	0.0034	0.125%
	% Inhibition	0.0000	19.0315	34.6041	46.3625	58.0891	75.9195	98.3171	99.4173	
K.pneumoniae	OD at 590nm	0.7033	0.4809	0.2809	0.0858	0.0404	0.0158	0.0028	0.0006	0.0313%
	% Inhibition	0.0000	31.6224	60.0597	87.8004	94.2557	97.7534	99.6006	99.9111	
P.mirabilis	OD at 590nm	0.6434	0.4812	0.3441	0.2603	0.0690	0.0102	0.0102	0.0102	0.0625%
	% Inhibition	0.0000	25.2129	46.5185	59.5431	89.2757	98.4162	99.7760	99.9501	
E.aerogenes	OD at 590nm	0.6492	0.5207	0.4805	0.3715	0.2527	0.1527	0.0270	0.0001	0.1250%
	% Inhibition	0.0000	19.7936	25.9858	42.7757	61.0752	76.4787	95.8410	99.9809	
E.faecalis	OD at 590nm	0.6985	0.4352	0.2649	0.0904	0.0413	0.0069	0.0007	0.0005	0.0313%
	% Inhibition	0.0000	37.9652	62.0759	87.0580	94.0873	99.0150	99.8933	99.9329	
A.baumannii	OD at 590nm	0.6981	0.4060	0.3010	0.2060	0.0251	0.0030	0.0025	0.0010	0.0313%
	% Inhibition	0.0000	41.8421	56.8787	70.4913	96.4045	99.5770	99.6405	99.8575	
N.gonorrhoeae	OD at 590nm	0.6865	0.4408	0.3080	0.0899	0.0469	0.0079	0.0019	0.0001	0.0313%
	% Inhibition	0.000	35.7902	55.1347	86.9046	93.1682	98.8536	99.7170	99.9836	
S.aureus	OD at 590nm	0.6745	0.4079	0.2776	0.0710	0.0297	0.0071	0.0002	0.0001	0.031

	% Inhibition	0.0000	39.5256	58.8436	89.4737	95.5967	98.9474	99.9703	99.9703	3%
S.epidermis	OD at 590nm	0.6855	0.4189	0.2886	0.0820	0.0307	0.0082	0.0003	0.0002	0.031 3%
	% Inhibition	0.0000	38.8972	57.8993	88.0379	95.5251	98.8083	99.9621	99.9752	

Therefore, Cinnamon were highly effective against uropathogenic bacteria more Ciprofloxacin. The main reason for inhibiting the growth of bacteria is the active constituents in cinnamon oil, the variation in the concentration of the active compounds in each extract contribute to prevent the normal growth of the pathogenic bacteria (Skandamis and Nychas, 2001; Carson et al., 2002)^{11&12}. Essential oil are potential source of novel antimicrobial activity especially pathogenic bacteria, cinnamon essential oil contain cinnamaldehyde and eugenol as major component (Ducke, 1994)¹³. Cinnamon oil also contain other active compounds which participate in antimicrobial activity as alkaloids, terpens, Cumarine and flavones (Alsengry, 2006)¹⁴. The MIC results in this study agreed with (Shareef, 2001)¹⁵, who mentioned that cinnamon oil gave (MIC) against some pathogenic bacteria (such as *E. coli*, *K pneumonia*, *S. aureus*,

P aeruginosa, *Proteus spp* and *Brucellaspp*) and observed that gram positive bacteria was more sensitive than gram negative tocinnamon oil. The result of this study were corresponding with (Bowels et al., 1995; and Helander et al., 1998)^{16&17}, they reported that *S. aureus*, *E. coli* and salmonella typhimurium were inhibited by essential oil of cinnamon. (Friedman et al., 2002)¹⁸ found that essential oil of cinnamon was active against *E. coli* and *Campylobacter jejuni*. In another study (Houqe et al., 2008)¹⁹ recorded that essential oil of cinnamon showed high antibacterial activity against *S. aureus*. (Trajano et al. 2009)²⁰ analyzed antimicrobial activity of 11 essential oils, including cinnamon, mint, black pepper, rosemary and ginger, against 10 strains of Gram-positive and Gram-negative bacteria. They reported that among the oils tested, cinnamon showed greater inhibitory act.

Table 3: Minimum inhibitory activity of standard (Ciprofloxacin) against uropathogenic bacteria

Test Organism	OD & %	Std (Cipro) Conc. 8µg/well								MI C %
		0	0.125	0.25	0.5	1	2	4	8	
<i>E.coli</i>	OD at 590nm	0.533	21.326	34.360	52.445	63.767	81.446	90.582	97.657	0.5%
	% Inhibition	0.000	21.326	34.360	52.445	63.767	81.446	90.582	97.657	

<i>P.aeruginosa</i>	OD at 590nm	0.541	0.433	0.349	0.241	0.215	0.171	0.089	0.034	0.5%
	% Inhibition	0.000	20.054	35.582	55.445	60.342	68.482	83.582	93.715	
<i>K.pneumoniae</i>	OD at 590nm	0.685	0.586	0.422	0.350	0.291	0.202	0.079	0.039	1%
	% Inhibition	0.000	14.359	38.417	48.866	57.546	70.470	88.533	94.376	
<i>P.mirabilis</i>	OD at 590nm	0.642	0.463	0.374	0.296	0.196	0.089	0.045	0.013	0.5%
	% Inhibition	0.000	27.852	41.708	53.897	69.483	86.191	93.033	97.974	
<i>E.aerogenes</i>	OD at 590nm	0.667	0.631	0.496	0.295	0.254	0.157	0.051	0.020	0.5%
	% Inhibition	0.000	5.424	25.697	55.769	61.912	76.551	92.433	97.048	
<i>E.faecalis</i>	OD at 590nm	0.724	0.620	0.526	0.361	0.225	0.087	0.068	0.025	0.5%
	% Inhibition	0.000	14.432	27.413	50.146	68.955	87.985	90.637	96.534	
<i>A.baumannii</i>	OD at 590nm	0.695	0.657	0.612	0.505	0.335	0.211	0.109	0.023	1%
	% Inhibition	0.000	5.413	11.863	27.296	51.828	69.565	84.308	96.743	
<i>N.gonorrhoeae</i>	OD at 590nm	0.666	0.613	0.517	0.349	0.269	0.134	0.092	0.013	1%
	% Inhibition	0.000	7.944	22.471	47.658	59.637	79.886	86.205	97.989	
<i>S.aureus</i>	OD at 590nm	0.772	0.651	0.551	0.340	0.268	0.181	0.088	0.045	0.5%
	% Inhibition	0.000	15.693	28.642	55.950	65.311	76.602	88.580	94.160	
<i>S.epidermis</i>	OD at 590nm	0.686	0.560	0.464	0.337	0.173	0.088	0.053	0.016	0.5%
	% Inhibition	0.000	18.379	32.313	50.896	74.803	87.203	92.231	97.639	

According to (Burt, 2004)²¹ natural compounds such as essential oils can act on bacterial cells by disintegration of the cell membrane by destabilising the proton force, electron flow, active transport and coagulation of cell contents. However, considering that essential oils have many different groups of chemical compounds, the antibacterial activity may be related to its composition and the mechanism of action cannot be assigned to a specific mechanism. Besides, there may be other targets in the cell, not only the cytoplasmic membrane. The direction for proper use of essential oil may be closely related to its composition. High concentrations can denature proteins and low concentrations can interfere with the

activity of enzymes involved in the energy production of the cell (Tiwari et al., 2009)²². Silva et al. (2010)²³ analyzed the effect of essential oils on *E. coli* and *Salmonella* spp. isolated from humans and ATCC cultures. Cinnamon essential oil showed the lowest MIC values for the studied microorganisms thereby confirming its high inhibitory activity. The active compounds present in the essential oil of cinnamon such as eugenol and cinnamaldehyde are responsible for causing damage to the structure of bacterial cell wall and has the capacity to interfere with the synthesis of certain bacterial enzymes (Matan et al., 2006).²⁴

Table 4: Minimum Bactericidal Concentration (MBC) of Samples against test organisms.

Organisms	Cinnamon oil Test conc. (%)					MBC
	1.00	0.50	0.25	0.125	0.0625	
<i>A.baumannii</i>	-	-	-	+	+	0.25
<i>N.gonorrhoeae</i>	-	-	+	+	+	0.5
<i>S.aureus</i>	-	+	+	+	+	1
<i>S.epidermis</i>	-	-	+	+	+	0.5
<i>E.faecalis</i>	-	-	+	+	+	0.5
<i>E.aerogenes</i>	-	+	+	+	+	1
<i>P.mirabilis</i>	-	+	+	+	+	1
<i>K.pneumoniae</i>	-	-	+	+	+	0.5
<i>E.coli</i>	-	+	+	+	+	1
<i>P.aeruginosa</i>	-	+	+	+	+	1

aerogenes, P. mirabilis, E. coli, and P. aeruginosa.

4. CONCLUSION

The antimicrobial effects of the cinnamon essential oil are evaluated by determining the minimum inhibitory concentration (MIC), the inhibition zone, and minimum bactericidal concentration (MBC). Cinnamon was the most effective agent in inhibiting A.baumannii, N.gonorrhoeae, S. epidermis, E. faecalis and E. coli with the lowest MIC (0.0313%) while S.aureus, E. aerogenes and P. aeruginosa with MIC (0.125%) while P. mirabilis was 0.0625% in our study. The MBC was 0.25% for

MIC and MBC results of cinnamon oil against uropathogenic bacteria are tabulated in Table 2 and 4. From the result we can conclude that Cinnamon Oil showed better inhibitory activity.

COMPETING INTERESTS

DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the

A.baumannii, 0.5 % for N, gonorrhoeae, S. epidermis, E. faecalis, and K. pneumonia, while 1% S.aureus, E.

research was not funded by the producing company rather it was funded by personal efforts of the authors.

5. REFERENCES

1. Bhattacharya, S. Cultivation of essential oils. In *Essential oils in food preservation, flavor and safety*, Academic Press, 20016; 16-29
2. Muhammad, D. R. A., & Dewettinck, K. Cinnamon and its derivatives as potential ingredient in functional food—a review. *International journal of food properties*, 2017; 20(2):2237-2263.
3. Kamaliroosta, L., Gharachorloo, M., Kamaliroosta, Z., Ali mohammad Zadeh K. H. Extraction of cinnamon essential oil and identification of its chemical compounds. *Journal of Medicinal Plants Research*, 2012; 6: 609-614.
4. Brkic, S., S. Mustafic, S. Nuhbegovic, F. Ljucam and L. Gavran. Clinical and epidemiology characteristics of urinary tract infections in childhood. *Med. Arh.*, 2010 64: 135-138.
5. Kordali S., Kotan R., Mavi A., Caki A., Ala A and Yildirim A. Determination of the chemical composition and antioxidant activity of the essential oil of *Artemisia draculus* and of the antifungal and antibacterial activities of Turkish *Artemisia absinthium*, *Artemisia draculus*, *Artemisia santonicum* and *Artemisia spicigera* essential oil. *J. Agric. Food Chem.* 2005; (53): 9452-9458.
6. Chuah, E.L., Zakaria, Z.A., Suhaili, Z., Bakar, S.A. and Desa, M.N.M. Antimicrobial activities of plant extracts against methicillin-susceptible and methicillin-resistant *Staphylococcus aureus*. *Journal of Microbiology Research*, 2014; 4(1): 6-13.
7. Owuama, C.I. Determination of minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) using a novel dilution tube method. *African journal of microbiology research*, 2017; 11(23), pp.977-980.
8. Shan, B., Cai, Y. Z., Sun, M., & Corke, H. Antioxidant capacity of 26 spice extracts and characterization of their phenolic constituents. *Journal of Agricultural and Food Chemistry*, 2005; 53(20): 7749-7759.
9. Braga, P.C., Dal-Sasso, M.; Culici, M. and Alfieri, M. Eugenol and thymol; alone or in combination; induce morphological alterations in the envelope of *Candida albicans*. *Fitoterapia*, 2007; 78: 396-400.
10. Bayoub, K.; Baibai, T.; Mountassif, D.; Retmane, A. and Soukri, A. Antibacterial activities of the crude ethanol extracts of medicinal plants against *Listeria monocytogenes* and some other pathogenic strains. *African Journal of Biotechnology*, 2010; 9: 4251-4258.
11. Skandamis P. N., Nychas G.-J. E. (2001). Effect of oregano essential oil on microbiological and physico-chemical attributes of minced meat stored in air and modified atmospheres.
12. Carson, C. F., Mee, B. J., and Riley, T. V. Mechanism of action of *Melaleuca alternifolia* (tea tree) oil on *Staphylococcus aureus* determined by time-kill, lysis, leakage, and salt tolerance assays and electron

- microscopy. *Antimicrob. Agents Chemother.* 2002; 46: 1914–1920.
13. Ducke JA. Biologically active compounds in important species. In species, herbs and edible fungi. (G, charalambous, ed.), Elsevier. New York, 1994 ; 225-250.
 14. Alsengry Wessal H. Studies on aflatoxins formation in ground nut *Arachishypogaea* L. and its inhibition in semi synthetic medium by cinnamon extract. Master thesis, department of Biotechnology, College of science, University of Baghdad, 2006.
 15. Shareef, A. Ali. Evaluation of antibacterial activity of essential oils of *Cinnamomum* sp. and *Boswellia* sp. *Journal of Basrah Researches (Sciences)*, 2001; 37(5):60-71.
 16. Bowels B. L., Sackitey S. K. and Williams A. C. Inhibitory effects of flavor compounds on *Staphylococcus aureus* WRRCB 124. *J. Food Safety*, 1995; 15: 337-347.
 17. Helander I.M., Alakomi H.L., Latva-Kala K., Mattila-Sandholm T., Pol I., Smid E.J., Gorris L.G.M., Von Wright A. Characterization of the action of selected essential oil components on gram-negative bacteria. *Journal of Agricultural and Food Chemistry*, 1998; 46: 3590–3595.
 18. Friedman M., Henika P. R and Mandrill R. E. (Bactericidal activities of plant essential oils and some of their isolated constituents against *C. jejune*, *E. coli*, *L. monocytogenes* and *S. enterica*. *J. Food Protection*, 2002; 65: 1545-1560.
 19. Houqe Md. Mahfuzul, M. L. Barib, Vijay K. Junejac, and S. Kawamoto. (2008). *Antimicrobial activity of clove and cinnamon extract against food borne pathogens and spoilage bacteria, and inactivation of Listeria monocytogenes in ground chicken meat with their essential oils. Rep, Nat'l. Food Res. Inst*, 2008; 72: 9-21.
 20. Trajano, V. N., Lima, E. O., Souza, E. L. and Travassos, A. E. R. Propriedade antibacteriana de óleos essenciais de especiarias sobre bactérias contaminantes de alimentos (Antibacterial property of spice essential oils on food contaminating bacteria). *Food Science and Technology*, 2009; 29(3): 542–545.
 21. Burt, S. Essential oils: their antibacterial properties and potential applications in foods - a review. *International Journal of Food Microbiology*, 2004; 94(3): 223–253.
 22. Tiwari, B. K., Valdramidis, V. P., O'Donnell, C. P., Muthukumarappan, K., Bourke, P. and Cullen, P. J. Application of natural antimicrobials for food preservation. *Journal of Agricultural and Food Chemistry*, 2009; 57(14): 5987–6000.
 23. Silva, C. J., Barbosa, L. C. A., Demuner, A. J., Montanari, R. M., Pinheiro, A. L., Dias, I. and Andrade, N. J. Chemical composition and antibacterial activities from the essential oils of Myrtaceae species planted in Brazil. *Química Nova*. 2010; 33(1): 104–108.
 24. Matan, N., Rimkeeree, H., Mawson, A. J., Chompreeda, P., Haruthaithanasan, V. and Parker, M. 2006. Antimicrobial activity of cinnamon and clove oils under modified atmosphere conditions. *International Journal of Food Microbiology*, 2006; 107(2): 180–185.

UNDER PEER REVIEW