

# **Patients' Perceptions and Satisfaction Regarding Teleconsultations during the COVID-19 Pandemic in Jeddah, Saudi Arabia**

## **Abstract**

**Background:** The coronavirus disease 2019 (COVID-19) pandemic has resulted in the widespread implementation of teleconsultation. Teleconsultation provides remote communication and reduces the risk of virus transmission. However, patients' perceptions and satisfaction are relevant to ensuring the quality of the service. Thus, this study aimed to evaluate patients' perceptions and satisfaction regarding teleconsultations during the COVID-19 pandemic in Jeddah.

**Methods:** This was a cross-sectional study using an electronic questionnaire to assess patients' perceptions and satisfaction regarding teleconsultations during May 2020. A total of 347 participants were included.

**Results:** More than one third of the sample were above forty years, women (73.2%), and had a university education (70.8%). Patients' perceptions of the effectiveness of teleconsultation ranged from 71.43% to 88.77% and their satisfaction level ranged between 59.40% and 83.96%. Older patients who had favorable outcomes from teleconsultation strongly agreed about the effectiveness of teleconsultation and were more satisfied. Patients with higher education levels who used the service only once were more satisfied with the experience. Moreover, those patients with greater age and educational levels and those who called for emergencies and to request medication refills were more likely to use teleconsultation in the future beyond the COVID-19 pandemic. When asked about reasons for dissatisfaction, 70% of the participants were unhappy with the wait time for a teleconsultation.

**Conclusions:** Most patients were satisfied with the teleconsultations during the COVID-19 pandemic. However, a better understanding of patients' perceptions and reasons for dissatisfaction are fundamental in the planning and implementation of teleconsultations in Saudi Arabia.

**Keywords:** COVID-19, Patient satisfaction, Teleconsultations, Saudi Arabia

## 1. Background

Telemedicine has been defined by the World Health Organization as “the delivery of health care services by all health care professionals using technology for the exchange of valid information for the diagnosis, treatment, and prevention of disease and injuries.”[1] One of the main categories of telemedicine is teleconsultation, which is defined as interactive communication between patients and healthcare providers at a distance by using technology.[2] A recent meta-analysis has shown that teleconsultation provided a rapid alternative to face-to-face clinical visits.[3] It is considered as an effective triage method to evaluate patients' complaints, prevent unnecessary clinical visits, and reduce waiting time.[4] Although some government and private health institutes already established teleconsultation services years ago, the implementation of teleconsultation has not been popular in Saudi Arabia.[5,6]

On March 2, 2020, Saudi Arabia announced the first case of coronavirus disease (COVID-19).[7] Shortly thereafter, the Saudi government imposed a 24-hour curfew for highly affected cities, including Jeddah. At the same time, the Ministry of Health (MOH), as well as numerous private hospitals and clinics, have activated teleconsultation services to provide health care to citizens without leaving their homes. Teleconsultation is an attractive solution

to minimize the risk of virus transmission,[8] and it provides not only emergency medical services but also routine medical consultations and follow-up.

The COVID-19 pandemic has offered an unexpected opportunity to increase the use of teleconsultation services in various medical specialties. Although many countries have adopted the idea of teleconsultation during the COVID-19 pandemic,[9,10,11,12] there has been limited research in regard to patient acceptance and satisfaction.[13,14] To date, no Saudi-based study has assessed patient satisfaction with this type of service during the COVID-19 pandemic. Thus, the aim of this project was to evaluate the patients' satisfaction with teleconsultation during the COVID-19 pandemic in Jeddah, Saudi Arabia. Continuing to provide teleconsultation services beyond the COVID-19 pandemic could save time and minimize the economic burden of providing health services. However, ensuring patients' satisfaction and acceptance of teleconsultation is a crucial step to ensuring success in this endeavor in Saudi Arabia.[15]

## **2. Methods**

### **2.1 Study design**

This cross-sectional study was conducted in Jeddah in May 2020. An electronic questionnaire was distributed online through social media (WhatsApp and Twitter). All the participants included in the study were 18 years or older, lived in Jeddah, and used teleconsultation services at least once. At the beginning of the questionnaire, the participants were briefly informed of the aim of the study and the term “teleconsultation” was clearly defined as the telephonic communication between patient and healthcare providers. Further, participants were asked to complete the questionnaire if they agreed to share their answers. Any incomplete questionnaires were excluded. This study was approved by the Unit of

Biomedical Ethics Research Committee at the Faculty of Medicine, King Abdulaziz University (Reference No. 229-20), dated May 4, 2020.

## **2.2 Sample size**

The sample size was calculated based on MOH reports regarding the call service center (937) that showed the number of medical calls in Jeddah was 906 per week.[7] This number was used as a guide to calculate the sample size. Because the study duration was planned to be for a month, the expected calls was 3,624. Then, Raosoft sample size calculator (<http://www.raosoft.com/samplesize.html>) was used to calculate the sample size. Based on achieving a confidence level (*P*-value) of 5% and a confidence interval of 95%, the sample size was 347.

## **2.3 Questionnaire**

For this study, a validated electronic questionnaire was used. The validity and internal consistency of the questionnaire were tested during a pilot study on 30 subjects who were excluded from the main study. The pretesting of the questionnaire showed that the Cronbach's alpha scores for all items were above 0.7. The questionnaire was translated from English into Arabic using the backward-forward method and was double-checked by the authors and a translator.

The questionnaire consisted of 18 questions distributed in 4 domains. The first question was a screening question to ensure the inclusion criteria (adult, living in Jeddah, and used teleconsultation service during COVID-19 pandemic). The first domain comprised demographic characteristics, including age, sex, nationality, and education level. The second domain was related to the personal experience of the teleconsultation (the number of times teleconsultation services were used during the COVID-19 pandemic, the reason for the call, and the outcome). The third and fourth domains addressed patients' perspectives regarding the effectiveness of teleconsultation and their satisfaction regarding the teleconsultation.

Respondents were asked to answer from ‘strongly agree’ to ‘strongly disagree’ on a 5-point Likert scale. The maximum total score was 25 and the minimum total score was 5, with higher scores reflecting greater satisfaction. Based on the results of this questionnaire, the participants were classified as follows: satisfied (satisfaction score from 25 to 16), neutral (satisfaction score from 15 to 11), and unsatisfied (satisfaction score from 10 to 5). Moreover, the Likert scores were transformed into satisfaction percentages using the following formula: Satisfaction % = (strongly agree + agree)/all response × 100.

## **2.4 Statistical analysis**

Data cleaning and data analysis for this study were performed using the Statistical Package for Social Sciences (SPSS) v. 23.0. Descriptive statistics were used to characterize the study population. Demographic data were described with frequency and percentages, while satisfaction scores were described using mean (M) and standard deviation (SD). Comparisons between the two groups were represented by the Mann–Whitney U test. A *P*-value < .05 was considered statistically significant. Qualitative thematic analysis for open-ended questions was used.

## **3. Results**

A total of 347 responses (279 in Arabic and 68 in English) were received during the study period. Fifteen questionnaires were excluded from the analysis due to incomplete information. The completion rate was 96.7%

### **3.1 Demographic characteristics**

The results obtained for the demographic data are listed in Table 1. The participants’ age range was 18 to older than 60 years. Most (38.6%) of the participants were aged 40 to 59 years, 28.3% were 18 to 29, and 25.9% were 30 to 39. Some 73.2% were women. Most

(92.5%) participants were Saudi nationals. Most (70.8%) participants had a university education.

The prevalence of using teleconsultation during the COVID-19 pandemic showed that almost 70% used the service one time only, whereas 30% used this service more than once. The main reason for using the service was experiencing a new complaint (26%), next, for follow-up (24%), then for emergencies and for COVID-19-related medical issues at 18% each, and finally 14% for medication refills.

The most common outcome from the teleconsultation was prescribed drugs (39.2%), followed by reassurance and self-management (35.2%). Making an appointment (5.1%) and sending for an ambulance (3.6%) were the least common reasons for using the service.

**Table 1. Demographic characteristics.**

Characteristics		Frequency (N)	Percentage (%)
Age	18–29	94	28.3%
	30–39	86	25.9%
	40–59	128	38.6%
	>60	24	7.2%
Sex	Men	89	26.8%
	Women	243	73.2%
Nationality	Saudi	307	92.5%
	Non-Saudi	25	7.5%
Educational level	Elementary	2	0.6%
	Secondary	37	11.1%
	University	235	70.8%
	Higher education	58	17.5%
How many times	Once	232	69.9%
did you use the	More than one	100	30.1%
service?			

For what reason did you use the service?	New complaint	86	25.9%
	Follow-up	80	24.1%
	Emergency	60	18.1%
	Refill medication	46	13.9%
	COVID-19 related	60	18.1%
What was the outcome or treatment from your call consultation?	Reassurance and self-management	117	35.2%
	Prescribed drugs	130	39.2%
	Appointment	17	5.1%
	Ambulance sent	12	3.6%
	Other	56	16.9%

### 3.2 Participants' satisfaction scores for various components of the teleconsultation

Table 2 shows the participants' perceptions of the effectiveness of the telemedicine offered. They stated they mostly agreed to strongly agreed regarding their ability to explain their medical problems to the doctor on the phone, with a median score of 4.33 out of 5. In addition, they believed that doctors understood their problems via telephone and were confident that the doctor could provide an appropriate management plan via teleconsultation, with median scores of 4.22 and 4.04, respectively. They expressed satisfaction with how much the doctors told them about the treatment, with a median score of 4.03. The lowest levels of agreement related to understanding their illness after the teleconsultation, with a median score of 3.91. Overall satisfaction rates with the service exceeded 70%.

**Table 2. Satisfaction scores on perceptions of the effectiveness of telemedicine (N = 332).**

Components	Mean	SD	Satisfaction %
------------	------	----	----------------

<b>I easily explained my medical problem to the doctor on the phone.</b>	4.33	0.76	88.47
<b>I believe the doctor understood my medical problem on the phone.</b>	4.22	0.84	83.71
<b>I was confident that the doctor could provide an appropriate management plan via phone consultation.</b>	4.04	1.02	75.19
<b>The doctor told me everything about my treatment.</b>	4.03	0.95	74.44
<b>I understood my illness much better after the phone consultation.</b>	3.9	1.02	72.43
<b>Overall Medical Satisfaction</b>	4.1	0.82	

Table 3 shows the participants' satisfaction with the telemedicine service. They were the least satisfied with sharing private or personal information, with a median score of 3.7 out of 5. However, there was greater satisfaction regarding the time offered for the consultation and the 24/7 availability of consultations, with median scores of 3.99 and 4.04 out of 5, respectively. The participants were satisfied that good medical service was provided that not require transportation, and that they would likely use it again in the future after the COVID-19 pandemic, with median scores of 4.3 and 4.05, respectively. Moreover, satisfaction rates exceeded 70% for all aspects assessed, except for the item related to sharing private information with a doctor face-to-face versus via the telephone. Some 59% of participants felt that they were more comfortable sharing private data via the phone than in face-to-face meetings.

**Table 3. Participants' satisfaction with the telemedicine service (N = 332).**

<b>Components</b>	<b>Mean</b>	<b>SD</b>	<b>Satisfaction %</b>
<b>I was more comfortable telling the doctor about private or personal information than in a face-face appointment</b>	3.70	1.13	59.40
<b>I was satisfied with the consultation time.</b>	3.99	0.98	77.94
<b>Medical consultations were offered at my preferred time 24/7.</b>	4.04	1.02	77.19
<b>Teleconsultation provided good medical service that did not</b>	4.30	0.88	83.96

require transportation.			
I would like to use teleconsultation again in the future after the COVID-19 pandemic.	4.05	1.12	76.19
Overall satisfaction	4.02	0.8	

### 3.3 Factors affecting participants' perception of the effectiveness of teleconsultation

Table 4 shows the factors affecting participants' perception of the effectiveness of teleconsultation. A significant association was observed between satisfaction levels and age ( $P = .037$ ), whereby older people were more likely to report higher levels of satisfaction.

Furthermore, we observed a significant relationship between levels of satisfaction and outcomes from teleconsultation ( $P = .000$ ), in which the highest satisfaction was shown in cases with consultation for an ambulance, then for prescribed drugs, and then for reassurance and self-management. There was no significant relationship between satisfaction and any other variable.

[Table 4. Factors affecting participants' perception toward telemedicine satisfaction.]

### 3.4 Factors affecting participants' satisfaction with the telemedicine experience

Table 5 shows 3 factors significantly affecting participant satisfaction with the telemedicine services. Participants with higher educational levels were more likely to be satisfied with the service compared with those with a lower educational background ( $P = .001$ ). On the other hand, participants who used the service more than once were more likely to report lower satisfaction compared with those who used the service one time only ( $P = .02$ ). There was also a significant relationship between the level of satisfaction and the outcome of the call consultation ( $P = .000$ ), in which satisfaction was highest among cases with reassurance and self-management and prescribed drugs as the outcomes.

[Table 5. Factors affecting participants' satisfaction regarding telemedicine experience.]

### 3.5 Characteristics of those who would be likely to use teleconsultation again in the future after the COVID-19 pandemic

Table 6 shows the characteristics of those who indicated they would likely use the phone consultation services again in the future after the COVID-19 pandemic. A total of 5 factors were shown to have an effect on this behavior: age, educational level, times of use of phone consultation, reasons for using the phone consultation, and outcomes from the phone consultation ( $P < .05$ ).

As can be observed, there was a significant association with age ( $P = .04$ ) and with using the service once or multiple times ( $P = .03$ ). The association with the outcome of the consultation was even more significant ( $P = .02$ ) than that for age or the number of times using the service. The strongest association in terms of using the service again was the reasons for using the service ( $P = .01$ ) and educational level ( $P = .01$ ).

[Table 6. Characteristics of those likely to use teleconsultation again in the future after the COVID-19 pandemic.]

### 3.6 Qualitative analysis of the main problems that the participants encountered during the teleconsultation experience

At the end of the questionnaire, the participants were asked an open-ended question about any problems they encountered during their teleconsultation. After ruling out the positive and unrelated comments, only 41 (10%) negative comments were included for further analysis (Table 7). A significant majority (70%) of the complaints with the service were related to the long waiting time for a consultation.

Table 7. Main problems encountered the participants during the teleconsultation.

Theme	Comments	N (%)
-------	----------	-------

Response and waiting time	Delayed response.	29 (70%)
	Long waiting time.	
	No response.	
Unsolved medical issue	The doctor did not understand my medical issues.	9 (%)
	They gave me an appointment without explaining my problem.	
	My problem was not solved.	
Technical issues	The doctor's voice was not clear.	3 (%)

#### 4. Discussion

This study aimed to evaluate the patients' perceptions and satisfaction regarding teleconsultations during the COVID-19 pandemic in Jeddah, Saudi Arabia. Factors that could influence patient satisfaction with the service were also considered. To our knowledge, this is the first study to assess patient satisfaction regarding teleconsultation in Saudi Arabia during the COVID-19 pandemic.

The demographic data indicated that the majority of the participants in this study were women between the ages of 30 and 59 with a college degree. It is worth mentioning that the results of this study showed that 70% of the study population used teleconsultation only once. This could be explained, to some extent, by the reported outcomes from their calls. The outcomes of the majority of the teleconsultation services result in either the patients' reassurance or medication prescription to them. However, more studies are needed to explore this area in-depth.

Although several studies[16,17] have indicated that these characteristics generally correlate with lower patient satisfaction, we found that patient satisfaction with the teleconsultation services was high across all domains tested. In over 70% of the cases, the participants felt that the doctor provided them with an appropriate management plan, that they understood their condition better, and that they were able to receive and convey information over the phone.

Similarly, in terms of satisfaction with the experience, >80% of the participants were satisfied with the fact that they did not need a physical meeting with the doctor. As first reported by Newbould *et al.*,[18] patient dissatisfaction with teleconsultation seems to increase when the patient still needs to have a physical visit despite the teleconsultation. Hence, it can be argued that the high rates of satisfaction, in this case, were obtained in part because the vast majority of the participants did not need to visit the doctor's office following the teleconsultation.

Several demographic variables appeared to impact satisfaction levels. In some cases, these elements were found to concur with previous reports.[16,17,18] Such aspects include older individuals and their likelihood of being satisfied with the service. Newbould *et al.*[18] stated that younger people are quicker to find solutions to their medical issues and therefore tend to be more dissatisfied with the services provided than older individuals. Similarly, the outcome of the service was also found to influence patient satisfaction regardless of the nature of the outcome. Some divergent aspects were also noted. Contrary to previous findings, [18,19,20] this study found that patients with lower levels of education tended to be more dissatisfied with the service.

Finally, contrary to other reports, sex was not found to influence levels of satisfaction.

Hence, by corroborating the results obtained by this study with similar studies, it appears that patient satisfaction with teleconsultation services is largely dictated by the reasons for which the service is requested, the outcomes of the consultation, and by specific patient characteristics.

Despite experiencing higher levels of dissatisfaction after using the service more than once, participants in this study were also more likely to use the service again if they had used it before. The interpretation of these results could be limited by the circumstances created by the COVID-19 pandemic.[21] Repeated use of the service might occur for these patients, if they lack other alternatives. Those with higher educational levels were also more likely to use the service and experienced greater satisfaction with the service, contrary to other findings. Finally, the reasons for use and the outcomes of the service also had a significant impact on the likelihood of reuse.

The reasons for service dissatisfaction identified by this study have also been reported elsewhere.[18] Over 70% of participants were dissatisfied with the long waiting time for a consultation, and 9% were unhappy because, despite the consultation, they had to attend a physical appointment. Only 3% had experienced a technical difficulty, which might indicate that the quality of the service could be influenced by the infrastructure available to carry out the teleconsultation.

Despite showing an overall high satisfaction rate with the teleconsultation service, these results must be considered with caution. First, the patient sample presents various homogenous characteristics, especially in terms of sex and education. These aspects limit the generalizability of the study across Saudi Arabia. It is therefore recommended that future studies include more diverse participant samples when assessing patient satisfaction with teleconsultations. Response bias is another limitation that can affect the outcome. Furthermore, the results of this study cannot be generalized to all medical subspecialties. The applicability of teleconsultation in various subspecialties must be assessed individually. Also, assessing the physicians' perceptions toward teleconsultation was beyond the scope of this study. It is worth mentioning that the adoption of telemedicine by health professionals is limited in Saudi Arabia despite the efforts and support provided by the Saudi MOH.[2,6,22]

A recent systemic review highlighted the factors that influenced the adoption of telemedicine in Saudi Arabia, which included a lack of healthcare provider knowledge and limited publications in support of such services.[5] Thus, it is crucial to increase the knowledge of healthcare providers about teleconsultation services and to incorporate this topic in all training programs and university medical curricula.

## **5. Conclusions**

We conclude that patients were generally satisfied with the teleconsultation service implemented in Jeddah during the COVID-19 pandemic. More studies will be necessary to obtain an in-depth understanding of patients' perception and satisfaction with these services as well as the necessary logistics. However, the telephone service could be extended to other medical areas and appears to function well as a viable alternative to physical consultations. Given the current circumstances created by the COVID-19 pandemic, investigating these services in terms of patient satisfaction, service quality, and potential improvements is a global necessity. The results of this study could shape the future of practicing medicine in Saudi Arabia and will help with establishing guidelines for conducting teleconsultation services. However, more studies are required to provide evidence-based data for teleconsultation in various medical subspecialties and specific patient populations.

## **List of abbreviations**

MOH	Ministry of Health
SD	Standard deviation
SR	Swain-Eng RJ

## **Declarations**

### **Ethics approval and consent to participate**

This study was approved by the Unit of Biomedical Ethics Research Committee at the Faculty of Medicine, King Abdulaziz University (Reference No. 229-20), dated May 4, 2020.

A written informed consent was at the beginning of the electronic questionnaire. The participants were informed about the aim and process of the study. Further, they were asked to complete the questionnaire only if they agreed to share their answers.

### **Consent for publication**

Not applicable.

### **Availability of data and materials**

Data that support the findings in the current study are available from the corresponding author on reasonable request.

## 6. References

1. WHO: A health telematics policy in support of WHO's health for all strategy for global health development: report of the WHO group consultation on health telematics, 11-16 December, Geneva 1997. Geneva: World Health Organisation; 1998. Accessed 23 July 2020.
2. El-Mahalli AA, El-Khafif SH, Al-Qahtani MF. Successes and challenges in the implementation and application of telemedicine in the eastern province of Saudi Arabia. *Perspect Health Inf Manag.* 2012;9:1-27.
3. Downes MJ, Mervin MC, Byrnes JM, Scuffham PA. Telephone consultations for general practice: a systematic review. *Syst Rev.* 2017;6(1):128.
4. Wechsler LR, Tsao JW, Levine SR, Swain-Eng RJ, Adams RJ, Demaerschalk BM, Hess DC, Moro E, Schwamm LH, Steffensen S, et al. Teleneurology applications: Report of the Telemedicine Work Group of the American Academy of Neurology. *Neurology.* 2013;80(7):670-6.
5. Alshahrani A, Stewart D, MacLure K. A systematic review of the adoption and acceptance of eHealth in Saudi Arabia: Views of multiple stakeholders. *Int J Med Inform.* 2019;128:7-17.
6. Ullah S, Maghazil AM, Qureshi AZ, Tantawy S, Moukais IS, Aldajani AA. Knowledge and attitudes of rehabilitation professional toward telerehabilitation in Saudi Arabia: A cross-sectional survey. *Telemed J E Health.* 2020; doi: 10.1089/tmj.2020.0016.
7. COVID 19 Dashboard: Saudi Arabia [<https://covid19.moh.gov.sa>] Accessed 23 July 2020.
8. Portnoy J, Waller M, Elliott T. Telemedicine in the era of COVID-19. *J Allergy Clin Immunol Pract.* 2020;8(5):1489-91.
9. Calton B, Abedini N, Fratkin M. Telemedicine in the time of coronavirus. *J Pain Symptom Manage.* 2020;60(1):e13-e14.
10. Latifi R, Doarn CR. Perspective on COVID-19: finally, telemedicine at center stage. *Telemed J E Health.* 2020; doi: 10.1089/tmj.2020.0132.
11. Liu S, Yang L, Zhang C, Xiang YT, Liu Z, Hu S, Zhang B. Online mental health services in China during the COVID-19 outbreak. *Lancet Psychiatry.* 2020;7(4):e17-e18.
12. Vidal-Alaball J, Acosta-Roja R, Pastor Hernandez N, Sanchez Luque U, Morrison D, Narejos Perez S, Perez-Llano J, Salvador Verges A, Lopez Segui F. Telemedicine in the face of the COVID-19 pandemic. *Aten Primaria.* 2020;52(6):418-22.
13. Capozzo R, Zoccolella S, Musio M, Barone R, Accogli M, Logroscino G. Telemedicine is a useful tool to deliver care to patients with Amyotrophic Lateral Sclerosis during COVID-19 pandemic: results from Southern Italy. *Amyotroph Lateral Scler Frontotemporal Degener.* 2020:1-7.
14. Layfield E, Triantafillou V, Prasad A, Deng J, Shanti RM, Newman JG, Rajasekaran K. Telemedicine for head and neck ambulatory visits during COVID-19: Evaluating usability and patient satisfaction. *Head Neck.* 2020;42(7):1681-89.
15. Kruse CS, Krowski N, Rodriguez B, Tran L, Vela J, Brooks M. Telehealth and patient satisfaction: a systematic review and narrative analysis. *BMJ Open.* 2017;7(8):e016242.
16. Woods SE, Heidari Z. The influence of gender on patient satisfaction. *J Gend Specif Med.* 2003;6(4):30-35.

17. Lopez C, Valenzuela JI, Calderon JE, Velasco AF, Fajardo R. A telephone survey of patient satisfaction with realtime telemedicine in a rural community in Colombia. *J Telemed Telecare*. 2011;17(2):83-7.
18. Newbould J, Ball S, Abel G, Barclay M, Brown T, Corbett J, Doble B, Elliott M, Exley J, Knack A, et al. A 'telephone first' approach to demand management in English general practice: a multimethod evaluation. In: A 'telephone first' approach to demand management in English general practice: a multimethod evaluation. edn. Southampton (UK); 2019.
19. Tabekhan AK, Alkhaldi YM, Alghamdi AK. Patients satisfaction with consultation at primary health care centers in Abha City, Saudi Arabia. *J Family Med Prim Care*. 2018;7(4):658-63.
20. Prakash B. Patient satisfaction. *J Cutan Aesthet Surg*. 2010;3(3):151-5.
21. Wosik J, Fudim M, Cameron B, Gellad ZF, Cho A, Phinney D, Curtis S, Roman M, Poon EG, Ferranti J, et al. Telehealth transformation: COVID-19 and the rise of virtual care. *J Am Med Inform Assoc*. 2020;27(6):957-62.
22. Albarrak AI, Mohammed R, Almarshoud N, Almujaalli L, Aljaeed R, Altuwaijiri S, Albohairy T. Assessment of physician's knowledge, perception and willingness of telemedicine in Riyadh region, Saudi Arabia. *J Infect Public Health* 2019;S1876-0341(19)30139-X.

**Table 4. Factors affecting participants' perception toward telemedicine satisfaction.**

Characteristics		Satisfied (25–16)	Neutral (15–11)	Unsatisfied (10–5)	P-value
Age	18–29	75 (79.8%)	16 (17%)	3 (3.2%)	.037
	30–39	80 (93%)	6 (7%)	0 (0%)	
	40–59	114 (89.1%)	8 (6.3%)	6 (4.7%)	
	>60	22 (91.7%)	2 (8.3%)	0 (0%)	
Sex	Men	75 (84.3%)	13 (14.6%)	1 (1.1%)	.111
	Women	216 (88.9%)	19 (7.8%)	8 (3.3%)	
Nationality	Saudi	270 (87.9%)	28 (9.1%)	9 (2.9%)	.385
	Non-Saudi	21(84%)	4 (16%)	0 (0%)	
Educational level	Elementary	2 (100%)	0 (0%)	0 (0%)	.606
	Secondary	33 (89.2%)	3 (8.1%)	1 (2.7%)	
	University	201 (85.5%)	26 (11.1%)	8 (3.4%)	
	Higher	55 (94.8%)	3 (5.2%)	0 (0.0%)	

	education				
How many times	Once	202 (87.1%)	25 (10.8%)	5 (2.2%)	
did you use the	More than				.379
service?	once	89 (89%)	7 (7%)	4 (4%)	
For what reason	New complaint	75 (87.2%)	10 (11.6%)	1 (1.2%)	
did you use the	Follow-up	71 (88.8%)	4 (5%)	5 (6.3%)	
service?	Emergency	52 (86.7%)	7 (11.7%)	1 (1.7%)	
	Refill				.145
	medication	44 (95.7%)	2 (4.3%)	0 (0%)	
	COVID-19				
	related	49 (81.7%)	9 (15%)	2 (3.3%)	
What was the	Reassurance				
outcome or	and self-	107 (91.5%)	9 (7.7%)	1 (0.9%)	
treatment from	management				
your call	Prescribed				
consultation?	drugs	124 (95.4%)	6 (4.6%)	0 (0%)	.000
	Appointment	15 (88.2%)	1 (5.9%)	1 (5.9%)	
	Ambulance				
	sent	12 (100%)	0 (0%)	0 (0%)	
	Other	33 (58.9%)	16 (28.6%)	7 (12.5%)	

**Table 5. Factors affecting participants' satisfaction regarding telemedicine experience.**

Characteristics		Satisfied (25–16)	Neutral (15–11)	Unsatisfied (10–5)	P-value
Age	18–29	76 (80.9%)	15 (16%)	3 (3.2%)	
	30–39	79 (91.9%)	6 (7%)	1 (1.2%)	
	40–59	115 (89.8%)	7 (5.5%)	6 (4.7%)	.092
	>60	22 (91.7%)	2 (8.3%)	0 (0%)	
Sex	Men	79 (88.8%)	8 (9%)	2 (2.2%)	.884

	Women	213 (87.7%)	22 (9.1%)	8 (3.3%)	
Nationality	Saudi	268 (87.3%)	29 (9.4%)	10 (3.3%)	.412
	Non-Saudi	24 (96%)	1 (4%)	0 (0%)	
Educational level	Elementary	0 (0%)	2 (100%)	0 (0%)	.001
	Secondary	32 (86.5%)	3 (8.1%)	2 (5.4%)	
	University	205 (87.2%)	22 (9.4%)	8 (3.4%)	
	Higher education	55 (94.8%)	3 (5.2%)	0 (0%)	
How many times did you use the service?	Once	208 (89.7%)	21 (9.1%)	3 (1.3%)	.020
	More than once	84 (84%)	9 (9%)	7 (7%)	
For what reason did you use the service?	New complaint	76 (88.4%)	9 (10.5%)	1 (1.2%)	.519
	Follow-up	67 (83.8%)	8 (10%)	5 (6.3%)	
	Emergency	53 (88.3%)	5 (8.3%)	2 (3.3%)	
	Refill medication	44 (95.7%)	2 (4.3%)	0 (0%)	
	COVID-19 related	52 (86.7%)	6 (10%)	2 (3.3%)	
What was the outcome or treatment from your call consultation?	Reassurance and self-management	108 (92.3%)	8 (6.8%)	1 (0.9%)	.000
	Prescribed drugs	121 (93.1%)	8 (6.2%)	1 (0.8%)	
	Appointment	15 (88.2%)	2 (11.8%)	0 (0%)	
	Ambulance sent	10 (83.3%)	2 (16.7%)	0 (0%)	
	Other	38 (67.9%)	10 (17.9%)	8 (14.3%)	

**Table 6. Characteristics of those likely to use teleconsultation again in the future after the COVID-19 pandemic.**

Characteristics		Strongly agree/agree	Not sure	Strongly disagree/disagree	<i>P</i> -value
Age	18–29	65 (69.1%)	11 (11.7%)	18 (19.1%)	.044
	30–39	68 (79.1%)	12 (14.0%)	6 (7%)	
	40–59	104 (81.3%)	10 (7.8%)	14 (10.9%)	
	>60	20 (83.3%)	4 (16.7%)	0 (0 %)	
Sex	Men	66 (74.2%)	15 (16.9%)	8 (9%)	.113
	Women	191 (78.6%)	22 (9.1%)	30 (12.3%)	
Nationality	Saudi	235 (76.5%)	35 (11.4%)	37 (12.1%)	.379
	Non-Saudi	22 (88%)	2 (8%)	1 (4%)	
Educational level	Elementary	0 (0%)	0 (0%)	2 (100%)	.001
	Secondary	27 (73.0%)	7 (18.9%)	3 (8.1%)	
	University	178 (75.7%)	27 (11.5%)	30 (12.8%)	
	Higher education	52 (89.7%)	3 (5.2%)	3 (5.2%)	
How many times did you use the service?	Once	177 (76.3%)	32 (13.8%)	23 (9.9%)	0.038
	More than once	80 (80%)	5 (5%)	15 (15.0%)	
For what reason did	New complaint	64 (74.4%)	18 (20.9%)	4 (4.7%)	

you use the service?	Follow-up	59 (73.8%)	3 (3.8%)	18 (22.5%)	.001
	Emergency	49 (81.7%)	4 (6.7%)	7 (11.7%)	
	Refill medication	38 (82.6%)	3 (6.5%)	5 (10.9%)	
	COVID-19 related	47 (78.3%)	9 (15%)	4 (6.7%)	
What was the outcome or treatment from your call consultation?	Reassurance and self-management	97 (82.9%)	13 (11.1%)	7 (6%)	.02
	Prescribed drugs	104 (80%)	14 (10.8%)	12 (9.2%)	
	Appointment	13 (76.5%)	2 (11.8%)	2 (11.8%)	
	Ambulance sent	9 (75%)	1 (8.3%)	2 (16.7%)	
	Other	34 (60.7%)	7 (12.5%)	15 (26.8%)	