

Original Research Article

Chemical Characterization of Solutions from Cooked Maize and ~~T~~their Comparison with ~~S~~solutions from ~~C~~ooked ~~B~~beans: Their ~~P~~osition ~~B~~etween ~~F~~ood and Fertilizer

ABSTRACT

The ~~present~~ paper ~~aims to~~ highlight the chemical characteristics of ~~solutions~~ ~~broth~~ from cooked maize and ~~to~~ compare them with ~~that solutions~~ from cooked beans. Solutions from cooked maize were produced by cooking variety of *Zea mays L.* with water. After this stage, samples of those solutions and samples of the water used for the cooking process were collected for laboratory analysis. Solutions from cooked maize are rich in mineral salts, particularly major macro elements (N and K) and minor macro elements (Ca, S, Mg). Concerning the third major macro element, notably the phosphorous, it is present in low amounts. The advantage of this fluid consists in its low electric conductivity. This fluid has a pH of 6.15. It is made of about 92% of water. A deep parallelism can be drawn between the end-of cooking solutions of beans and solutions from cooked maize. In fact, these two fluids are rich in nitrogen and potassium, and mainly made of water. But, in detail, some particularities are present. The end-of cooking solutions of maize have higher amounts of sodium and chlorides, this joined with a higher electric conductivity. Concerning solutions from cooked beans, they have higher amounts of the different macro elements, and a lower electric conductivity. The recycling of end-of cooking solutions of maize must gainfully become for the nutritionists a favorable target for the future, and this due to the numerous nutrients contained. However, its agri use is directly topical according to the results obtained in the case of the present studies. Using it as manure requires an earthing-up directly after its application in order to avoid the ~~loose~~ ~~loss~~ of sulfur and nitrogen through gas emanation. Consuming solutions from cooked maize as herbal tea could be an excellent way to recycle the nutrients that have diffused from the seeds during the cooking process.

Keywords: *Used water; sanitation; recycling; fertilization; agriculture; nutrition.*

The language of the abstract is not clear. Please take a close look at the comments and do a summary of your work by briefly introducing the work, stating the methodology, some of the results obtained and then the conclusion.

1. INTRODUCTION

In ~~ACP~~ (Africa, Caribbean, and Pacific) ~~(ACP)~~ countries, the demographic explosion [1], the reduction of ~~tillable~~ lands, and the reducing of the fallow periods [2] represent today the most perceptible phenomenon. This part of the world is peopled essentially by resource-poor farmers, with soils as their principal source of earnings [3]. Hence, to insure their survival, they cultivate even the most sloping plots [4]. Various and risky farming techniques are

Comment [M11]: A Comparison of the Chemical Composition of cooked maize and beans ~~left-over-water~~: Potentials as food drinks and Fertilizers. You may consider the suggested topic. This is because I believe that what you meant by solution is the water that remained after cooking the maize and beans.

Comment [M12]: What quantity of maize, beans and water did you use. Did you use maize on the cob or just maize grains. What were the cooking period and temperature.

Comment [M13]: What laboratory analysis methods did you use

Comment [M14]: Change the word solution to ~~left-over-water~~ wherever it appears in this write up

Comment [M15]: State the quantities of the individual minerals in comparison with the two ~~left-over-water~~

Comment [M16]: Did you do conductivity tests

Comment [M17]: Meaning?

Comment [M18]: State the figures

Comment [M19]: Meaning?

Formatted: Font: Not Italic

noticeable here [4], the aim being the desire of the daily satisfaction of their needs in food. In consequence, the yields decrease years after years [5, 3], attesting then the permanent depletion of soils fertility. If a minority among these peoples can easily acquire synthetic fertilizers to solve short term problem of soils fertility depletion [6, 7, 8], the majority can't. But, even in the first group of these populations, the solution for soil fertility is not completely acquired because of the remaining of queries. In fact, a fraction of them up to today lacks knowledge about the use of these chemicals, since they could be pollutants when poorly used [9]. Moreover, those chemicals are sometimes rare, and their solubility is not often guaranteed [10]. Concerning the second group of these peoples, represented by resource-poor farmers, possess synthetic fertilizers is simply a fairytale. Their survival then appears today with acuteness. That's why day after day, researchers investigate different sectors since the beginning of the twentieth century in order to provide fertilizers at low costs, easily used, efficient, and available for the poorest people worldwide. The tendencies of the results obtained today are encouraging. So, in some parts of the world, many were taught in how to use rocks [11, 12], plants such as *Titonia diversifolia* [13], and mixtures of organic fluids [14, 15] among all as fertilizers. These approaches already constitute exits for that problem despite their local character; in fact, geological and floral diversity all over the world generates variations in their application. In front of these signs of potential victories, scientists have not given up; in that way, [14] recently showed that the mixture of human urine and solutions from cooked beans can pertinently enhance the growth of plants. To comfort the contribution of the solutions from cooked beans as potentials nutrients provider, [16] successfully tested them as substrate for some heterotrophic organisms, notably moulds. He justified the choice of moulds by their inability to manufacture by themselves their food, and then, condemned to be often looking for external incoming of organic matters, mineral salts and water, both necessary for them to realize their vital functions as proved by [17]. To make deeper the approach started with the potential recycling of the end-of cooking solutions of beans, maize dragged the attention. This was justified by the existence of a strong parallelism between the cooking processes of beans and maize. Therefore, it appear interesting to question the impact of the boiling process on maize seeds, and consequently on the water used for cooking the maize. This would help to determine the chemical characteristics of the end-of cooking solutions of maize on the one hand, and to compare those characteristics to those of the end-of cooking solutions of beans on the other hand. Finally, it could be determined if those two solutions could be recycling for the same purposes. Those motivations then encouraged the present study.

Comment [M110]: Please,recast to bring out the meaning

The introduction is supposed to give background information on the theme of the paper namely;the nutrient content and potentials of maize and beans Left-over-water and similar studies.Then highlight their peculiar or novel contributions.

2. MATERIALS AND METHODS

2.1 Materials

2.1.1 Beans seeds

Most of the Africa inhabitants practice agriculture [18, 19, 20], with beans as one of the main crops [21]. That plant is appreciated all over Cameroon because of its dietetically potential [22, 23, 24]. In that way, 100 g of that food is made of 9.06 g of proteins, 27.91 g of

carbohydrates, 0.49 g of fats, 5.3 g of fibers, 61.2 g of water, 0.257 mg of B1 vitamin (thiamine), 0.063 mg of B2 vitamin (riboflavin), 0.57 mg of B3 vitamin (niacin), 0.299 mg of B5 vitamin (pantothenic acid), 0.175 mg of B6 vitamin, 0.98 mg of E vitamin (tocopherole), 3.7 µg of K vitamin, 168 µg of total folate, 1337 mg of potassium, 225 mg of sulphur, 414 mg of phosphorus, 140 mg of magnesium, 113mg of calcium, 47 mg of chloride, 6.5 mg of iron, 2.5 mg of zinc, 2.3 mg of iron, 3.5 mg of sodium, 0.96 mg of zinc, 0.635 mg of copper, 0.548 mg of manganese, 0.430 mg of boron, 0.271 mg of copper, 0.183 mg of nickel, 0.02 mg of chrome, 0.095 mg of fluoride, 0.0052 mg of cobalt, 0.0018 mg of iodate, and 0.0014 mg of selenium among others [25,26,27,28]. The amount of the total nitrogen can be deducted from the rate of the proteins with the help of the Jones factor [29]; in fact, the amount of the proteins = total nitrogen*6.25 (with 6.25 as the Jones factor). For the present case, we have 1.4496 g of total nitrogen in bean seeds.

About 85% of their product is directly eaten in some of the countries where it is produced. The remaining 15% is sold [30]. Particularly in Cameroon, large quantities are produced every year, and are mainly intended for export [31]. Bean seeds represent an important source of vegetal proteins, their flatulent effect is universally known and has, undoubtedly, been a source of discomfort throughout history [32]. Integument of bean seeds is permeable or semi permeable, allowing then diffusion phenomenon to occur through it [33].

2.1.1 Maize seeds

There are several varieties of corn. One can name the yellow, purple [34], white, blue, and red [35] varieties. The violet and blue color are due to the presence of anthocyanin pigments [36] while the carotenoid pigments give yellow and red maize their respective colors [37]. The differentiation of maize varieties also uses the shape, strength and composition of the kernels. Thus, there are "popcorn" varieties, others "soft" for human consumption fresh or canned, or even "horny" used in semolina [38]. According to consumers, the postharvest date and variety affects the taste and texture of boiled corn [39]. The color of the husks (leaves that wrap the cob) would be an indicator of the freshness and taste of the corn. Thus, the fresher the corn husks (bright green color), the more intense the sweetness of the corn, indicating that the corn is freshly harvested. In contrast, the less green color of corn husks indicates a less sweet taste and a long post-harvest period [39]. For 100g of corn, there is an average of 73.40g of water, 2.4g of fiber, 18.60g of carbohydrates, 1.7g of lipids, 3.41g of protein, 3mg of calcium, 0.05mg of copper, 0, 45mg of iron, 0.001mg of iodine, 107.9mg of magnesium, 0.17mg of manganese, 299.6mg of phosphorus, 324.8mg of potassium, 59.2mg of sodium, 0.62mg of zinc, 0.066mg of provitamin A Beta carotene, 0.011mg of vitamin A, 0.09mg of vitamin B1, 0.06mg of vitamin B2, 1.68mg of vitamin B3, 0.79 mg of vitamin B5, 0.14mg of vitamin B6, 0.023mg of vitamin B9, 5, 5mg of vitamin C, 0.09mg of vitamin E, 0.0004mg of vitamin K1, 0.01mg of flavonoids, 0.01mg of anthocyanins, 214.56mg of phenolic acids, 2.41mg of hydroxybenzoic acids, 212, 15mg of hydroxycinnamic acids, 0.33mg of lignans [40].

2.1.3 Solutions from cooked beans

Solutions from cooked beans are heterogeneous mixtures, and particularly a proteic globular suspension, with considerable amounts of carbohydrates. It has a pH value of 6.4. At rest, that water divides itself into two superimposed domains: a flaky superficial domain and a liquid

lower domain. The flaky domain is the organic part and the liquid domain is the water and the mineral salts provider. The density of the flaky domain is 0.964 and that of the liquid domain is 1.011. The average speed of the growth of *Aspergillus L.* at the surface of the solutions from cooked beans is 3,17 cm² /H; they cover in five days a surface of 379.74 cm². These solutions are made of about 90% of water. Their electric conductivity is 0.668 mS/cm. Their pH is 6.31. The ionic composition of the solutions from cooked beans reveals the presence of: Ammonium (5.41g/l), Potassium (2.63 g/l), Phosphate (0.0333 g/l), Magnesium (0.538 g/l), Calcium (0.403g/l), Sulfate (0.29g/l), Sodium (0.07487g/l), Chlorides (0.029g/l) [41]. The physico-chemical characteristics of the flaky domain floating on the liquid domain make it an adequate area for the development of moulds (*Aspergillus L.*). Solutions from cooked beans contain all the nutrients required for an optimal fertilization of soils; this include water, organic matters and mineral salts among which nitrogen, potassium, phosphorus, sulfur and calcium can be named. It is then a complete liquid organic fertilizer. Those solutions position themselves also as a high grade activator for soils micro flora. It represents finally a way for a sustainable improvement of agriculture in developing countries and a way for a sustainable development of soils micro flora, required for achieve the food self-sufficiency [16].

In daily life, many pig farmers feed their animals with spent grain mixed with solutions from the end of cooking the beans; results are wonderful. Moreover, those solutions are used in the confection of some meals in Cameroon known as “pilé” in many tribes. This action is justified by the increasing of the nourishing potential of the food. In addition, when solutions from beans cooking are spread out, after ten of minutes, one can easily feel the emanation of a loud smell; that odor disappears within two or three day according to the quantity of solutions thrown [33].

2.1.3 Water used for cooking the maize seeds

The water for beans cooking has an electric conductivity of 0.029mS/cm and a pH of 5.1. This water is composed, in decreasing weight order, of 0.0154 g/l of sodium, 0.0095 g/l of potassium, 0.009 g/l of ammonium, 0.0045 g/l of calcium, and 0.0034 g/l of magnesium. Concerning anions, it has, in decreasing weight order, 0.0085 g/l of sulfate, 0.0016 g/l of chlorides, 0.00146 g/l of phosphate, and 0.0013 g/l of carbonate. Finally, the water used for the beans cooking process is chemically under the control of sodium as major cation and sulfate as major anion [41].

2.1.4 Area of the studies

The well for domestic use that provided the cooking water is located in the locality of Yaoundé, at mid-slope of an interfluvies, at about 5.5m from latrines, and at open air. The climate of the locality is of Guinean equatorial type, with four different seasons [42]. The average annual rainfall in Yaoundé is 1,600 mm, for an average temperature of 23°C. [43]. The geological substratum is made of embrechites with quartz, garnet, biotite, feldspar, pyroxenes, plagioclase, and hornblende as main minerals. Soils are of lateritic type [44]. On the hydro geological point of view, the crystalline substratum of the Yaoundé region is essentially constituted by two superimposed aquifers: an upper aquifer (within alterites), located between 5 and 20 m depth, and a lower aquifer (in substratum discontinuities), located at depths greater than 20 m [45].

2.2 Methods

Field and laboratory studies help to reach the targets.

2.2.1 Field work

Field work made it possible to obtain the seeds (photograph 1) of maize necessary for the production of the studied solutions, to acquire the opinions of the populations on facts related to the daily management of maize, and finally to practice the tasting of the corn tea.



Photograph 1. « Maize seeds »

2.2.2 Laboratory tests

2.2.2.1 Manipulation of maize seeds samples and production of solutions from their end-of cooking process

In the laboratory, maize samples were sorted out in order to remove undesirable particles, and weighted. For the production of the samples of solutions from cooked maize, 2kg of seeds were quickly washed to remove dust and other impurities. Maize were cooked into water without salt to avoid any out coming electrolytes. The resulting suspension (so called solutions from cooked maize) were collected, cooled, and stored in the fridge for future lab tests.

2.2.2.2 Laboratory analysis

Physic-chemical analysis dealt with the identification of Ca^{2+} , Mg^{2+} , K^+ , Na^+ , PO_4^{3-} , SO_4^{2-} , NH_4^+ , Cl^- , and CO_3^{2-} since they are directly or indirectly present in maize seeds. Moreover, their choice is motivated by the fact that they can be organized as major macro elements (N, P, K), minor macro elements (S, Ca, Mg, Na), and oligo elements (Fe, Se, Mn, Zn, Cu) for plant nutrition [46]. The process started by the chemical characterization of the water used for the cooking process during which the quantities of the same ions were measured. The lab tests ended with the evaluation of the amount of water in the solutions from cooked maize.

The IonPac CS12A and IonPac AS12A analytical columns were respectively used for the separation of ammonium (NH_4^+), calcium (Ca^{2+}), magnesium (Mg^{2+}), sodium (Na^+), and potassium (K^+) as far as cations were concerned, and chloride (Cl^-) and sulfate (SO_4^{2-}) in the case of anions. The flow rate was set to 1.0 mL/min for both cations and anions optimization.

The different cations were isocratically separated with a 20 mM Methanesulfonic acid solution within 15 minutes. Concerning the ions studied here, isocratic elution with 4mM and 20mM Sodium hydroxide solution were employed for their separation, respectively for chloride and sulfate, during 15 minutes. All eluents were degassed and pressurized under high purity nitrogen to prevent dissolution of carbon dioxide and subsequent production of carbonate.

Concerning bicarbonate ion (HCO_3^{2-}), its amount was measured by using titrimetric method. The pH was determined with a pH-meter having a glass electrode. Electric conductivity for its own was measured using a conductivity meter fitted with a calibrated measuring cell; the results were expressed in mS/cm.

To determine the amount of water, heating was used. In fact, a given quantity of the solution was collected and weighted (M1). After the weighting, it was poured in a metallic container with known mass (M2). After the complete evaporation, the container and the residues were weighted (M3). The proportion of water was obtained through the following calculations: $(M3-M2)*100/M1$. The percentages of the macro elements were obtained as follow: $\%N = \text{NH}_4^+ * 100 / (\text{NH}_4^+ + \text{HPO}_4^{2-} + \text{K}^+)$, $\%K = \text{K}^+ * 100 / (\text{NH}_4^+ + \text{HPO}_4^{2-} + \text{K}^+)$, $\%P = \text{HPO}_4^{2-} * 100 / (\text{NH}_4^+ + \text{HPO}_4^{2-} + \text{K}^+)$, $\%Ca = \text{Ca}^{2+} * 100 / (\text{Ca}^{2+} + \text{Mg}^{2+} + \text{SO}_4^{2-})$, $\%Mg = \text{Mg}^{2+} * 100 / (\text{Ca}^{2+} + \text{Mg}^{2+} + \text{SO}_4^{2-})$, $\%S = \text{SO}_4^{2-} * 100 / (\text{Ca}^{2+} + \text{Mg}^{2+} + \text{SO}_4^{2-})$.

[Materials and methods entail information on the samples such as: sample collection, description, identification, treatments or processing, parameters to be measured, techniques used in assessing the parameters and then the instrument for data analysis.](#)

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Water for maize cooking

[The water for maize cooking has an electric conductivity of 0.029mS/cm and a pH of 5.1.](#)

This water is composed, in decreasing weight order, of 15.4 mg/l of sodium, 9.5 mg/l of potassium, 9 mg/l of ammonium, 4.5 mg/l of calcium, and 3.4 mg/l of magnesium. Concerning anions, it has, in decreasing weight order, 8.5 mg/l of sulfate, 1.6 mg/l of chlorides, 1.46 mg/l of phosphate, and 1.3 mg/l of carbonate. Finally, the water used for the beans cooking process is chemically under the control of sodium as major cation and sulfate as major anion (Table 1).

3.1.2 Solutions from cooked maize

[While the corn is cooking, a pleasant aroma emerges from the pot. Also, the end of cooking liquid, drunk hot, is excellent. In this sense, of the 100 people selected at random to taste this solution whose identity was not revealed, 83 of them took it again, which corresponds to 83% of favorable cases. Interviews with many other people have revealed that they like to delight in the solution contained in the cob.](#)

Comment [MI11]: Clearly bring out these features in your materials and method. As a guide, you can take a look at some previous publications of this Journal.

Comment [MI12]: Meaning?

Comment [MI13]: This sounds like sensory evaluation. If so, is it necessary?

Solutions from cooked maize are made of about 92% of water. Its electric conductivity is 0.8158 mS/cm. Compared to the water used for the cooking process, a sensible increasing is noticed, about 0.7868mS/cm (Fig. 1 and Table 1).

Its pH is 6.15. Compared to the water used for the cooking process, a sensible increase of the present parameter is detected too, of about 1.05 (Fig. 2 and Table 1).

The ionic composition of the solutions from cooked beans reveals the presence of:

- **Ammonium (NH_4^+)** : Its amount is 4508.108 mg/l. Compared to the water used for the cooking process, a high increasing (4499.108 mg/l) of the amount of that ion is noticed (Fig. 3 and Table 1);
- **Potassium (K^+)** : Its concentration is 1163.35 mg/l. Compared to the water used for the cooking process, a high increasing (1153.85 mg/l) of the amount of this ion is detected (Fig. 3 and Table 1);




Fig. 1. Evolution of the electric conductivity from the water used for the maize cooking process (E1) to solutions from maize cooking (E2)

Comment [MI14]: What is the meaning of this?




Fig. 2. Evolution of the pH from the water used for the maize cooking process (E1) to the solutions from maize cooking (E2)

UNDER PEER REVIEW

Table 1. Chemical characteristics of solutions from cooked maize and the water used for the cooking process

	pH	Ca²⁺ (mg/l)	Mg²⁺ (mg/l)	K⁺ (mg/l)	Na⁺ (mg/l)	CO₃²⁻ (mg/l)	SO₄²⁻ (mg/l)	HPO₄²⁻ (mg/l)	NH₄⁺ (mg/l)	E.C. (mS/cm)
Water used for the cooking process (E1)	5.1	4.5	3.4	9.5	15.4	1.3	8.5	1.46	9	0.0290
Solutions from cooked maize (E2)	6.15	196.54	380.43	1163.35	205.3	152	264.23	57.96	4508.108	0.8158




Fig. 3. Evolution of ions from the water used for the maize cooking process (E1) to solutions from maize cooking (E2)

Comment [MI15]: There is no colour difference.

- **Phosphate (PO_4^{3-})**: Its concentration is about 57.96 mg/l. Compared to the water used for the cooking process, a high increasing (56.5 mg/l) of the amount of this ion is however detected (Fig. 3 and Table 1);
- **Magnesium (Mg^{2+})**: Its amount is 380.43 mg/l. Compared to the water used for the cooking process, a high increasing (377.03 mg/l) of the quantity of the present ions is detected (Fig. 3 and Table 1);
- **Calcium (Ca^{2+})**: Its amount is 196.54 mg/l. Compared to the water used for the cooking process, a high increasing (192.04 mg/l) of the quantity of the present ions is highlighted (Fig. 3 and Table 1);
- **Sodium (Na^+)**: Its amount is 205.3 mg/l. Compared to the water used for the cooking process, a high increasing (189.9 mg/l) of the quantity of the present ions is highlighted (Fig. 3 and Table 1);
- **Chloride (Cl^-)**: Its amount is 38.92 mg/l. Compared to the water used for the cooking process, a high increasing (37.32 mg/l) of the quantity of the present ions is highlighted (Fig. 3 and Table 1);
- **Sulphate (SO_4^{3-})**: Its amount is 264.23 mg/l. Compared to the water used for the cooking process, a high increasing (255.73 mg/l) of the quantity of the present ions is highlighted (Fig. 3 and Table 1);
- **Carbonates (HCO_3^-)**: Its amount is 152 mg/l. Compared to the water used for the cooking process, a high increasing (150.7 mg/l) of the quantity of the present ions is highlighted (Fig. 3 and Table 1).

Comment [MI16]: Why the bullets?

~~In short,~~ Chemically, all the parameters targeted in the context of this study increase when comparing the water used for cooking the maize and the end-of cooking solutions of maize (Figs. 1, 2,3 and Table 1). In detail, if the water used for the cooking process of maize is shyly under the control of sodium, solutions from cooked beans is largely dominated by ammonia, potassium, magnesium, sulphate, sodium, and calcium. Further on, then come respectively carbonate, phosphate, and below, chloride. On a completely different level, Fig. 4 reveals a positive correlation between the pH and the electrical conductivity of the solutions studied.




Fig. 4. Correlation between pH and electric conductivity (E.C.) of fluids

3.1.3 Solutions from cooked maize

3.2 Discussion

3.2.1 Water used for maize cooking

The amount of sodium is greater in the water used for the cooking process of maize compared to the amount of potassium. This can be due to the greater stability of K-feldspar and K-micas in one hand, in other hand by the absorption of K in minerals newly formed during the process of alteration as showed by [47], and further on by the atomic mass of K, greater than that of Na, making then the motion of Na easier than that of K. Those two chemical elements are suspected to be from rocks alteration [48]. This is in accordance with the mineral composition (Feldspars, Plagioclases, Hornblende, and Ferromagnesian and the main minerals) of the geological substratum of Yaoundé. Calcium for its own could have been released after the weathering of Plagioclases, Garnet, and Hornblende contained in the substratum (Schist, Gneiss, and Granite) of Yaoundé. The presence of calcium can also be associated to the process of the impoverishment of soils as the consequence of lixiviation process according to [49]. Concerning magnesium ions, they are thought to be from the alteration of ferromagnesian minerals of the substratum. Nitrates are the major forms through which nitrogen is present in water. That ion can be used as an indicator to testify the pollution of water according to [50]. But when oxygen lacks in an aquatic area, nitrogen mainly appears

Comment [MI18]: Always use scientific terms.

as Ammonium. This corroborates the observations made by [51]. The presence of chlorides can be a direct consequence of the nature of some of the minerals from the Yaoundé substratum. In fact, [52, 53] demonstrated that their presence is one of the prime characteristics of groundwater in basement region; it is then in accordance with the presence of Hornblende and Biotite, which are among the chloride bearers. Moreover, [54] demonstrated the possible substitution of Cl-OH type, responsible of the liberation of chloride in water. Further, the presence of chlorides in this water can be justified by atmospheric inputs following infiltration of rainwater [55], and by the alteration of silicate minerals [56]. Bicarbonates are present in the water used for the cooking process. This can be due to the dissolution of atmospheric and pedological carbon dioxide [57]. It can be also due to the anthropic factors as showed by [58, 59]; this is in accordance with the presence of latrines around the well from which the water used for the cooking process of the maize was collected. Regarding the presence of sulfates, it can be due to the alteration of sulfur bearing minerals from the substratum. This corroborates the presence of Pyrite in the gneissic rocks from the Yaoundé region as revealed by [54]. However, meteoric inputs [60] as same as anthropogenic pollution [81] could be considered as far as the presence of sulfates in the water used for the cooking process of maize is of concerned. The pH of the present water is 5.1; after [61], it is acid. The electric conductivity of the same water is 0.029mS/cm. Such a value certifies a low mineralization of the concerned water since it remains very low compared to the standard established by [62]. According to the abacus proposed by [61], the present water is nonconductive.

Finally, the results of analyzes of the water used for cooking the maize in the context of this study are consistent with the data obtained by [63, 64].

3.2.2 Solutions from cooked maize

Significant enrichment in the different chemical elements followed up in the present study is noticed in solutions from cooked maize. This can be explained by the leaching of those elements by maize seeds in contact with water during the cooking process. In fact, when the seeds come into contact with water during cooking, they absorb water and swell. Such phenomenon was previously described in the different stages of the germination of seeds by [65]. This observations corroborates the conclusions of [66] and [33], who consider biological membranes as permeable or semi-permeable, and then suggested to diffusion phenomenon, responsible here to the enrichment of the solutions from cooked maize in mineral salts and in protein derivatives as stated by [16]. This can then be used to explain the fact that in Cameroon, some pigs breeders often mix up the food of their animals with that fluids or simply give it to them as drink; on the field, the results of such practice are wonderful as highlight by [33]. In the same vein, according to [33, 67], and [65], this enrichment is facilitated by the temperature, materialized here by the boiling which leads to the cooked stage of the seeds. If such observations reminds the fertilizing characteristics of the solutions from cooked beans as showed by [33, 14], and [16], it also induces a real dietary concern with regard to the consumption of maize. In fact, solutions from cooked maize, considered as waste water in Cameroon by many populations, are most of the time threw away at the end of the cooking process. Much then remains to be done to reconcile these latter and this fluid which undoubtedly has an important nutritional value as shown by the sudden improvement in the contents of exchangeable basic cations, nitrogen, and phosphorus, among others. Many other facts contribute to certify the richness of the solutions from maize cooking. In fact, when those solutions are abandoned at open air, few hours after, they attract

many drosophila. Such facts are often noticed when slice of fruits are exposed at open air. For [68], those insects always followed matters rich in carbohydrates. Such facts are consistent with the observations made by [69].

In order of decreasing enrichment, we have nitrogen (4200 mg/l), potassium (1163.35 mg/l) and magnesium (380.43 mg/l). These are respectively two of the major macro elements, that is to say entering mainly in plant nutrition, and one of the minor macro elements [46]. According to the work of [40], potassium and phosphorus are among the most concentrated chemical elements in maize. Thus, these authors show that for 100 g of maize, we have 324.8 mg of potassium and 77mg (299.6mg) of phosphorus. By applying the Jones factor according to which proteins = $6.25 \times$ total nitrogen [29], it comes that in 100g of beans, we have 1295mg (1.295g) of nitrogen, which is therefore the most concentrated chemical element in these seeds. The abundance of nitrogen and potassium in the maize end-cooking solutions is therefore consistent with their concentration in the seeds; their degree of abundance in the maize end-cooking solutions is in the same order as in the seeds; the third major macro element, however, is weakly expressed. This is in agreement with the observations of [33]. Beside the major macro elements, minor macro elements are also well expressed, In fact, calcium (196.54mg/l) and sulfate (264.23 mg/l) represent the most concentrated chemical elements after the previous in solutions from cooked maize.

The electrical conductivity remains below 1mS / cm. According to [61], the maize end cooking solution is a non-saline fluid. However, the electrical conductivity goes from 0.029 to 0.8158mS / cm from the water used for cooking the maize to solutions at the end of cooking. Even if this parameter is very low according to [61], it is however multiplied by approximately 28 (28.13) in solutions at the end of cooking of the maize. This therefore attests to a mineralization of the maize end-of-cooking solutions, which concretely results in a sudden increase in the contents of the various elements targeted in the context of the present study when switching from the water used for cooking the maize to the end-of-cooking solutions of this food. At the same time, there is a significant increase in the pH value, which goes from 5.1 in the water used for cooking the beans to 6.15 in the maize end solutions, an increase of 1.05 units. This is in agreement with the mineralization of the maize end cooking solutions, certified by the increase in the content of the different elements and the increase in electrical conductivity. Such an observation refers us to the remark made by [62] within the framework of their investigation on the mineralization of underground water in the locality of Yaoundé. The pH value (6.15) obtained in the case of the present study is closer to that previously obtained by [16] when studying solutions from cooked beans.

To help the characterization of drinkable water, [63] defined the following thresholds: less than 0.15 g / l for sodium, 0.25 g / l for sulfates, 0.25 g / l for calcium, 0.012 g / l for potassium, 0.05 g / l for magnesium, 0.05 g / l for nitrates, 0.25 g / l for chlorides, and 0.25 g / l for carbonates. Taken individually the chemical elements contained in the end of cooking solutions of the maize in relation to these standards, it returns that in the end of cooking solutions of the maize, the contents of the main major macro elements (nitrogen, potassium, calcium, magnesium) are well beyond the prefixed thresholds. Further on, the phosphate ion contents, according to the observations of [70], reached thresholds above which the phenomenon of eutrophication is noted in the environment. It is then a highly nutritious solution for plants. This is another argument that attests to the high chemical content of the maize end-of-cooking solutions. Such observations imposes the remark according to which

during the process of the maize cooking, the leaching of mineral salts occurs from the seed towards the cooking solution. This is consistent with the observations made by [14, 15, 16, 33, 64] during their studies on solutions from cooked beans.

3.2.3 Comparison between solutions from cooked maize and solutions from cooked beans

As in bean, so proved by [41], nitrogen is predominantly expressed in the form of ammonium ion in the maize end-of-cooking solutions. One can explain such a fact by relying on the theory of loss of oxygen by water when its temperature increases as showed by [46].

In bean end-of cooking solutions, the ammonium content is 5410 mg / l according to [64]. In the maize end-of cooking solutions, it is 5200 mg / l (Table 2). Nitrogen represents 66.6% of all major macro elements [46] in beans end-of cooking solutions while it represents 67.37% of the same elements for the maize end-of cooking solutions. As shown in Fig. 5, nitrogen (ammonium) may be a little more concentrated in bean end-of cooking solutions than in the maize end-of cooking solutions. This is in accordance with the ratio between the ammonium content in the bean end-of cooking solutions and the content of the same element in the maize end-of cooking solutions: 1.29 (Table 2).

Phosphate content is 33.3 mg/l in end-of cooking solutions of beans. In the maize end-of-cooking solutions, it is 29.3 mg / l (Table 2). It represents 0.47% of all major macro elements [46] in the maize end-of cooking solutions and 0.4% in the end-of cooking solutions of beans. As showed in Fig. 5, phosphates are a little more concentrated in bean end-of cooking solutions than in the maize end-of cooking solutions as comforted by the ratio between the phosphate content in the bean-end-of-cooking solutions and the content of the same element in the maize end-cooking solutions: 1.14 (Table 2).

Potassium content is 2630 mg/l in end-of cooking solutions of beans. In the maize end-of cooking solutions, it is 1163.35 mg / l (Table 2). It represents 16.66% of all major macro elements [46] in end-of-cooking solutions of maize and 33% of major macro elements in the end-of-cooking solutions of beans. As showed in Fig. 5, potassium is more concentrated in end-of cooking solutions of beans than in the maize end-of cooking solutions. To deepen this observation, one can refer to the ratio between the potassium content in the end-of cooking solutions of beans and the content of the same element in the maize end-of cooking solutions: 2.26 (Table 2). This parameter shows that the concentration of potassium in the end-of cooking solutions of beans is more than twice the concentration of the same element in solutions from the end of cooking the maize.

Table 2. Comparison between the human urine and water from cooked beans

	Solutions from cooked beans	Solutions from cooked maize	Percentage of macro elements		Solutions from cooked beans parameters/ Solutions from cooked maize parameters
			Solutions from cooked maize	Solutions from cooked beans	
NH ₄ ⁺ (mg/l)	5410	4200	67.37	66.6	1.29
HPO ₄ ²⁻ (mg/l)	33.3	29.3	0.47	0.4	1.14
K ⁺ (mg/l)	2630	1163.35	18.66	33	2.26
SO ₄ ²⁻ (mg/l)	290	264,23	4.24	3.11	1.097
Ca ²⁺ (mg/l)	403	196.54	3.15	32.73	2.05
Mg ²⁺ (mg/l)	538	380.43	6.1	43.70	1.41
Na ⁺ (mg/l)	74.84	185,3			0.82
Cl ⁻ (mg/l)	29	38.92			0.74
pH	6.31	6.15			1.02
E.C. (mS/cm)	0.668	0.8158			0.82
Water (%)	90	92			0.98




Fig. 5. Chemical composition of solutions from cooked maize and solutions from cooked beans

Sulfate content is 290 mg/l in end-of cooking solutions of beans. In the maize end-of cooking solutions, it is 264.23 mg / l (Table 2). It represents 4.24% of all major macro elements [46] in the maize end-of-cooking solutions and 3.11% in the end-of cooking solutions of beans. As showed in Fig. 5, sulfates are a little more concentrated in bean end-of cooking solutions than in the maize end-of cooking solutions. This is in accordance with the ratio between the phosphate content in the bean-end-of cooking solutions and the content of the same element in the maize end-cooking solutions: 1.097 (Table 2).

Calcium content is 403 mg/l in end-of cooking solutions of beans. In the maize end-of cooking solutions, it is 196.54 mg / l (Table 2). It represents 3.15% of all major macro elements [46] in the maize end-of-cooking solutions and 32.73% in the end-of cooking solutions of beans. As showed in Fig. 5, calcium is more concentrated in bean end-of cooking solutions than in the maize end-of cooking solutions as proved by the ratio between the calcium content in the bean end-of cooking solutions and the content of the same element in the maize end-of cooking solutions: 2.05 (Table 2).

Magnesium content is 538 mg/l in end-of cooking solutions of beans. In the maize end-of cooking solutions, it is 380.43 mg / l (Table 2). It represents 6.1% of all major macro elements [46] in the maize end-of-cooking solutions and 43.70% in the end-of cooking solutions of beans. As showed in Fig. 5, calcium is more concentrated in bean-end-of cooking solutions than in the maize end-of cooking solutions. This is consistent with the ratio between the calcium content in the bean end-of cooking solutions and the content of the same element in the maize end-of cooking solutions: 1.41 (Table 2).

Sodium content is 74.84 mg/l in end-of cooking solutions of beans. In the maize end-of cooking solutions, it is 205.3 mg / l (Table 2). As showed in Fig. 5, sodium is more concentrated in maize end-of cooking solutions than in the beans end-of cooking solutions as revealed by the ratio between the sodium content in the bean end-of cooking solutions and the content of the same element in the maize end-cooking solutions: 0.82 (Table 2).

Chlorides content is 29 mg/l in end-of cooking solutions of beans. In the maize end-of cooking solutions, it is 38.92 mg / l (Table 2). As showed in Fig. 5, Chlorides are more concentrated in maize end-of cooking solutions than in the beans end-of cooking solutions. The ratio between the sodium content in the bean end-of cooking solutions and the content of the same element in the maize end-of cooking solutions, of 0.74 (Table 2), is in agreement with that fact.

Electric conductivity of the end-of cooking solutions of beans is 0.668mS/cm. That of the maize end-of cooking solutions is 0.8158mS/cm (Table 2). As showed in Fig. 5, despite the low level in general, it is slightly higher in maize end-of cooking solutions than in the beans end-of cooking solutions. The ratio between its level in the bean end-of cooking solutions and in the maize end-cooking solutions, reaching 0.82 (Table 2), confirms it.

The pH of the end-of cooking solutions of beans is 6.31. That of the maize end-of cooking solutions is 6.15 (Table 2). As showed in Fig. 5, this parameter is comparable in the two solutions. The ratio between its level in the bean end-of cooking solutions and in the maize end-of cooking solutions, of about 1 (Table 2), is in accordance with that observation.

According to [64], water constitutes 90% of the beans end-of cooking solutions. Concerning the maize end-of cooking solutions, it represents 92%. Its quantities in both solutions are comparable (Table 2) as showed by the ration between its level in the bean end-of cooking solutions and in the maize end-of cooking solutions: 0.98.

Generally speaking, bean and maize end-of cooking solutions are two fluids that are very similar in composition. Thus, as shown in Fig. 6 below, these are two fluids whose composition is largely dominated by nitrogen and potassium respectively. But, with regard to their respective origin, they will logically present particularities. Thus, bean end-of cooking solutions are richer in ammonium, potassium, calcium, magnesium, and phosphate, with lower electrical conductivity. The maize end-of cooking solutions for their own are richer in sodium and chloride. It comes that the end-of cooking solutions of beans are richer in the different macro elements than the end-of cooking solutions of maize do. According to [72] who recommends a dilution of human urine in the proportions 1: 4 (urine: water) before application in view of its too high concentration of sodium and chlorides, and, further, because of its high electrical conductivity, the same recommendation seems to be valid for the end-of cooking maize solutions. According to the observations of [14], the agri use of these two solutions requires hilling to prevent losses of nitrogen and sulfur, particularly in gaseous form. The strong parallelism that exists between their chemical compositions makes these two fluids excellent liquid organic fertilizers, with end-of cooking solutions of beans as the best among the two.

As it was done with the end-of cooking solutions of beans, it would be further important that the maize end-of-cooking solutions be reconsidered from a dietetic point of view with regard to its high nutritional potential. Indeed, in view of its chemical composition, this solution could be consumed as a soup in order to recycle the cohort of nutrients it contains.




Fig. 6. Position of the end-of cooking solutions of beans and maize beside human urine in the N-P-K diagram of [73]

3.2.3. Phenomenon accompanying the cooking process of maize

While the corn is cooking, a pleasant aroma emerges from the pot. Also, the end of cooking liquid, drunk hot, is excellent. In this sense, from the 100 people selected at random to taste this solution whose identity was not revealed, 83 of them took it again, which corresponds to 83% of favorable cases. Interviews with many other people revealed that they like to delight in solutions contained in the cob; this fluid is nothing more than trapped corn end-cooking solution. Such a declaration gives meaning to the behavior of the 83 people who took up the herbal tea. It is common that smelling sauces are also palatable according to [74]. This is in accordance with the leaching of nutrients from maize seeds during cooking processes as previously demonstrated by [75]. According to the composition of the corn end-of-cooking solution, consuming it as a herbal tea could be an excellent way to recycle the nutrients that have diffused from the seeds during the cooking process; this mobility agrees with the observation made by [76] concerning the permeability and semi-permeability of biological membranes.

Results and discussion should go hand in hand(state each result and discuss) until you are done with the results in each table.Also,restrict yourself to works done for the purpose of this article.For instance,sensory evaluation has no place in this article.

4. CONCLUSION

Solutions from cooked maize are rich in mineral salts, particularly major macro elements (N and K) and minor macro elements (Ca, S, Mg). Concerning the third major macro element, notably the phosphorous, it is present in low amounts. The advantage of this fluid consists in its low electric conductivity. This fluid has a pH of 6.15. It is made of about 92% of water. A deep parallelism can be drawn between the end-of cooking solutions of beans and solutions from cooked maize. In fact, these two fluids are rich in nitrogen and potassium, and mainly made of water. But, in the detail, some particularities are present. The end-of cooking solutions of maize have higher amounts of sodium and chlorides, this joined with a higher electric conductivity. Concerning solutions from cooked beans, they have higher amounts of the different macro elements, and a low electric conductivity. The recycling of end-of cooking solutions of maize must gainfully become for the nutritionists a favorable target for the future, and this due to the numerous nutrients contained. However, its agri use is directly topical according to the results obtained in the case of the present studies. To use it as manure requires an earthing-up directly after its application in order to avoid the loss of sulfur and nitrogen through gas emanation. Consuming solutions from cooked maize as herbal tea could be an excellent way to recycle the nutrients that have diffused from the seeds during the cooking process.

5. RECOMMENDATION

The use of solutions from cooked maize as fertilizers requires dilution; its farm application must be directly followed by earthing up in order to avoid the loss of nitrogen and sulfur through gas emanation. These solutions could be gainfully consumed as herbal tea.

Comment [MI19]: Please still consult publications in this journal for guidance.

Conclusion and recommendation should be done after due attention to the comments on results and discussion

Comment [MI20]: Rewrite the results and discussion in line with the comments. You will then be able to conclude your work and make recommendations.

REFERENCES

Comment [MI21]: Follow the journal's referencing format.

- [1] Isolina Boto. Population growth and its implications for ACP rural development. Resources on population growth in ACP countries1. No. 16. CTA Office Brussels. 2012;55.
- [2] FAO. World population growth and decrease of cultivated areas per inhabitants; 2003.
- [3] Shackleton CM, Pasquini MW, Drescher AW. African Indigenous Vegetables in Urban Agriculture. Earth scan Publishing for a sustainable future. London, Sterling, VA. 2009;339.
- [4] Tematio P, Tsafack EI, Kegni L. Effects of tillage, fallow and burning on selected properties and fertility status of andosols in the Mounts Bambouto, West Cameroon. Agricultural Science. 2011;2(3):334-340.
- [5] Tematio P, Olson KR. Impact of industrialized agriculture on land in Bafou, Cameroon. J. of Soil and Water Conservation. 1997;404-405.
- [6] Sanchez PA, Shepherd KD, Soule MJ, Place FM, Buresh RJ, Izac NA. Soil fertility replenishment in Africa: An investment in natural resource capital. In: Buresh RJ, Sanchez PA and Calhoun F eds. Replenishing soil fertility in Africa. SSSA Special Publ. 1997;51:1-46.
- [7] Sanchez, Soil fertility and hunger in Africa. Science. 2002;295:2019-2020.
- [8] Zhou JM, Huang PM, Kinetics and mechanisms of monoammonium phosphate-induced potassium release from selected potassium-bearing minerals. Can. J. Soil Sci. 2008;799-811.
- [9] Brady NC. The nature and properties of soils. 8th edition. Macmillan Publishing C., 14 INC. New York. Collier Macmillan Publisher, London. 1974;639.
- [10] Conway G. The doubly green revolution: Food for all in the 21st century. Penguin Books, London, UK. 1997;335.
- [11] Yusdar H, Hanafi M. Use of phosphate rock for perennial and annual crops cultivation in Malaysia: A review. In: Rajan SSS and Chien SH (eds.) Direct application of phosphate rock and related appropriate technology latest developments and practical experiences: Proceedings of an international meeting, IFDC, Muscle Shoals, Alabama USA, Special P Ublications. 2001;37:78-91.
- [12] Prevosto B, Dambrine E, Moares C, Curt T. Effect of volcanic ash chemistry and former agricultural use on the soils and vegetation of naturally regenerated woodlands in the Massif Central, France. Catena. 2004;56:239-261.
- [13] Kaho F, Yemefack M, Feujio-Teguefouet P, Tchantchouang JC. Combined effect of *Tithonia diversifolia* leaves and inorganic fertilizers on maize yields and ferralitic soil properties in Central Cameroon. Tropicultura. 2011;29(1):39-45.
- [14] Fopoussi Tuebue JC, Basga SD, Tematio P, Nguetnkam JP. Impact of the mixture of water from cooked bean and human urine on the growth of some common plants in Cameroon: Case study of *Talinum fruticosum* L. and *Ocimum gratissimum* L. 2019;5(1):1-15. Article no. AJSSPN.51071 [ISSN: 2456-9682]
- [15] Fopoussi Tuebue JC, Basga SD, Tematio P, Nguetnkam JP. Effect of trachyte powder, human urine and reserved water from cooked beans on Andosols Fertility in Cameroonain Western Highlands. Asian Journal of Soil Science and Plants Nutrition. 2018;3(2):1-17. Article no.AJSSPN.41999. [ISSN: 2456-9682]
- [16] Fopoussi Tuebue JC, Basga SD, Tematio P, Nguetnkam JP. Water from cooked beans as substrate for some heterotrophic organisms: Case Study of Moulds 2020; 5(3):1-15. Article no.AJSSPN.53588.ISSN: pp 2456-9682.

- [17] Restino C. Production of itaconic acid by *Aspergilli* strains by fermentation in a solid medium. Thes Doct. Univ. Reims, Champagne-Ardenne, INRA. 2012;174.
- [18] CIAT. Genetic diversity in the genus *Phaseolus*, Study notebook, Intended to be used as a complement to the Audio tutorial Unit dealing with the same subject; Scientific Content: Daniel D. Debouck; Production: Luz Maria Médina Cali, Colombie. 1989;52.
- [19] Morin S. The fundamental dissymmetries of the West Cameroon Highlands and their consequences for human occupation. Example of the Bambouto mountains. The man and the tropical mountain. SEPANRIT éd., Bordeaux. 1988;49-51.
- [20] Shackleton CM, Pasquini MW, Drescher AW. African indigenous vegetables in urban agriculture. Earth scan Publishing for a sustainable future. London, Sterling, VA. 2009;339.
- [21] Aufrey P. Associated crops and cropping systems in Bamiléké country (West Cameroon). Centre Univ. Dschang-CNEARC, First year thes, ESAT. 1985;82. annexes.
- [22] Lauchaux C. Dried beans to tomato; 2003. Available: <http://pascal.moron.free.fr/rec>
- [23] Andrianiaina Harivola P. Laboratory study of the behavior of some varieties of *Phaseolus vulgaris*. L, IL5 (50-1), IL5 (50-7), IL5 (56-1), IL5 (56-4), RJ2 et Lb vis-a-vis de *Acanthoscelides*. Univ. Antananarivo, Ecole Norm. Sup., Dept of Initial Scientific Training, Studies and Research Sciences Natural Sciences. Thes. 2006;99.
- [24] Marcia LP, Bykowski C, Fujiko F Duke, Danielle R. Reed, « Excretion and Perception of a Characteristic Odor in Urine after Asparagus Ingestion: a Psychophysical and Genetic Study », Chemical Senses. 2011;36(1):9-17.
- [25] Boissard P. Malagasy cuisine, Creole cuisine. Bookstore, Madagascar.1907; 144.
- [26] Press L. The bean book: Over seventy incredible recipes. Globe Pequot Press; 2002. ISBN 9781585744732.
- [27] Souci SW, Fachmann E, Kraut H. Food composition and nutrition tables. Medpharm Scientific Publishers Stuttgart/ Taylor & Francis; 2008. (ISBN 978-3-8047-5038-8).
- [28] Journal des Femmes. Haricot blanc; 2019. Available:<https://cuisine.journaldesfemmes.fr/encyclopedie-produits/1958658-haricots-blancs/>
- [29] Jones DB. Eds. Factors for converting percentages of nitrogen in foods and feeds into percentages of protein, US Department of Agriculture - circ. 1941;83.
- [30] Fanantenana JO. Varietal improvement of white bean type white ingot by hybridization for future export: 2002 - 2003 campaignà Nanisana Mémoire de C.A.P.E.N. - ENS - Université d'Antananarivo. 2003;52.
- [31] Azeufouet A, Tsague E, Medjou S, David O, Temple L, Parrot L. Quantification of cross-border flows from Cameroon. Hal Id : cirad-00948344. 2006;53.
- [32] Roger Fenwick R, Price KR, Johnson Ian T. Flatulence December 1987 Nutrition & Food Science. (Fenwick, Roger & Price KR, Johnson Ian. Flatulence. Nutrition & Food Science. 1987;87(3):2-4. DOI: 10.1108/eb059428.
- [33] Fopoussi Tuebue JC. Petrological study of andosols in the West Cameroon Highlands and fertilization test: Case of a toposequence developed on trachyte of the upper zone of the southern slope of the Bambouto Mountains (West Cameroon). Doctorate / PhD thesis, University of Ngaoundere, Faculty of Sciences, Department of Earth Sciences, Laboratory of Sciences, Water, and Environment. 2018;273.
- [34] Ramos-Escudero F, Muñoz AM, Alvarado-Ortíz C, Alvarado Á, Yáñez JA. Purple corn (*Zea mays* L.) phenolic compounds profile and its assessment as an agent against oxidative stress in isolated mouse organs. *J Med Food*. 2012;15(2):206-15.

- [35] Lopez-Martinez LX, Parkin KL, Garcia HS. Phase II-inducing, polyphenols content and antioxidant capacity of corn (*Zea mays* L.) from phenotypes of white, blue, red and purple colors processed into masa and tortillas. *Plant Foods Hum Nutr.* 2011;66(1):41-7.
- [36] Lao F, Sigurdson GT, Giusti MM. Health benefits of purple corn (*Zea mays* L.) phenolic compounds. *Comprehensive Reviews in Food Science and Food Safety.* 2017;16(2):234-46.
- [37] Siyuan S, Tong L, Liu R. Corn phytochemicals and their health benefits. *Food Science and Human Wellness.* 2018;7(3):185-95.
- [38] Chégut M, Hardy C, Lebarbier R, Marot MF, Martin E, Nicolai L / Agreste Nouvelle-Aquitaine. Grain corn sector [en ligne]. [Consulté le 03/05/2020]. Disponible à l'adresse : http://www.epsilon.insee.fr/jspui/bitstream/1/84615/3/agr_NA_filiere_ma%C3%AFs_2018_09.pdf
- [39] Hongbete F, Kindossi JM, Hounhouigan JD, Nago MC. Production and nutritional quality of fresh boiled corn cobs consumed in Benin. *International Journal of Biological and Chemical Sciences.* 2017;11(5):2378-92.
- [40] A. J. Semassa , S.W. Padonou, V.B. Anihouvi , N.H. Akissoé, D. Aly, A. Adjanohoun, L. Baba-Moussa, Diversité Variétale, Qualité Et Utilisation Du Maïs *Zea Mays*) En Afrique De l'Ouest : Revue Critique, European Scientific Journal, June 2016, Edition 18, Vol. 12, ISSN : 1857-7881 (Print) e-ISSN 1857-7431
- [41] Fopoussi Tuebue J.C., Tchinda I.N., Chemical evolution of solutions from beans soaking and cooking processes: case study of *Phaseolus vulgaris* L. *European Journal of Nutrition & Food Safety*, 13(2): 24-51, 2021; Article no.EJNFS.65701 ISSN: 2347-5641
- [42] Suchel JB, The climates of Cameroon. PhD Thes, Univ. Saint Etienne. 1988;188.
- [43] Olivry JC. River and rivers of Cameroon. Hydrology Monograph Collection. MESRES/ORSTOM, n°9. 1986;733.
- [44] Yongue-Fouateu R, Contribution to the petrographic study of the alteration and facies of ferruginous armouring of migmatitic gneisses in the Yaoundé region. Th. Doct. 3eme cycle .Univ.Yaoundé. 1986;214.
- [45] Djeuda Tchapgna HB, Tanawa E, Temgoua E, Siakeu J. Ngo Massana B, et al. Mode of circulation, recharging mechanisms and relative residence times of groundwater from alterites in the crystalline environment: case of the Anga'a watershed, Yaounde, Cameroun. Press. Univ. Yaoundé. 1989;117-126.
- [46] Space For Life. Horticultural and botanical notebook; 2019. Available:<https://m.espacepurlavie.ca/besoins-nutritifs>
- [47] Le Marechal A. Geology and geochemistry of thermomineral sources in Cameroon. Works of Doc., ORSTOM, 1976;176 .
- [48] Post VEA. Chemistry for modellers aqueous geochemistry in coastal area. 17 th Salt Water intrusion Meeting. Delft. The Netherlands. 2002;3-12.
- [49] Kamgang Kabeyene Beyala V, Ekodeck GE. Alteration and geochemical balances of biotites in gneiss Nkolbisson (NW of Yaoundé, Cameroon). *Geodynamic.* 1991;6(2): 191-199.
- [50] Antonakos N, et Lambrakis A. The hydrogeochemistry of the lake Wako drainage basin, Texas. *Environmental Geology.* 2000;45(1):106-114.
- [51] Grambow B, Michel N. Solubility of solids in water: surface property or solid? »The 6th Scientific Days of Marcoule. 15 – 19 mai. Subatech, Nantes. 2006;1:53.
- [52] Oga YMS, Lazm T, Koffi Yao T, Soro N, Bachir Saley M, Dongo D, Gnanba F. Chemical characterization of water from fractured aquifers: Case of the Tiassalé Region in Cote d'Ivoire. Tiassalé in Ivory Coast. *European Journal of Scientific Research.* 2009;31(1):72-87.

- [53] Ngounou Ngatcha B., Murdry J., and Sarrot R.J., Groundwater recharge from rainfall in the southern border of Lake Chad in Cameroon. *World Appl Sci.* 2009; 2:125–131.
- [54] Nzenti JP. Neoproterozoic alkaline meta-igneous rock from the Pan-African North Equatorial Fold Belt (Yaoundé, Cameroon): Biotites and magnetite rich pyroxenites. *Journal of African Earth Sciences.* 1998;26(1):37-47.
- [55] Sigha-Nkamdjou , Galy-Lacaux C, Pont V, Richard S, Sighomnou D. Lacaux JP, et al. Rainwater chemistry and wet deposition over the Equatorial forested ecosystem of Zoétélé (Cameroon). *Journal of Atmospheric Chemistry.* 2003;173-198.
- [56] Nguengang P, Parrot L, Lejoly J. Joiris V, et al. Development of lowlands in Yaoundé: Production system, traditional know-how and potential for developing urban and peri-urban agriculture. In Parrot L. (coord.) *Agriculture and urban development in sub-Saharan Africa, environment and health issues.* L'Harmattan, Paris. 2008;97-108.
- [57] Kortatsi K, Tay K, Anomu G, Hayford E, Dartey A. Hydrogeochemical evaluation of groundwater in the lower Offin basin, Ghana. *Environ Geol* 2: Springer-Verlag, 2007;12.
- [58] Vicat JP, Mvondo H, Willems L, Pouclet A. Fossil and current karst phenomena within silicoaluminous metamorphic formations of the Pan-African water table of Yaoundé (Sud-Cameroun). *CR Geosci.* 2002;334: 545–550.
- [59] Michard G. *Chemical equilibrium in natural waters.* Ed, Publisud 2002;357. ISBN. 2- 86600-446-9.
- [60] Kuitcha D, Fouéfé Takounjou AL, Ndjama J, Takem Eneke G, Tita Awah M. Kamgang Kabeyene BV, Chemical and isotopic signal of precipitation in Yaounde-Cameroon. *Archives of Applied Science Research.* 2012;4(6):2591-2597.
- [61] El Oumlouki K, Moussadek R, Zouahri A, Dakak H, Chati M, EL Amrani M. Study of physico-chemical quality of water and soil in the region Souss Massa (Case perimetr Issen), Morocco. *J. Environ. Sci. EDE4, CODEN: JMESCN.* 2014;5(S2):12-24, 2365-2374. ISSN :2028-2508.
- [62] 5902OMS, Guideline for drinking water. Hygiene criteria and supporting document. World Health Organization. 2 édition. 2000; 1050.
- [63] Kuitcha D, Fouéfé Takounjou AL, Ndjama J. Contribution of hydrochemistry and the isotope of the environment to the knowledge of groundwater resources in Yaoundé, Cameroon, *Journal of Applied Biosciences.* 2013;67:5194–5208. ISSN 1997.
- [64] Fopoussi Tuebue J.C., Tchinda I.N., Djiotsa D., Physico-chemical characterization of solutions from cooked beans and their comparison with human urine: fertilizer and food? *European Journal of Nutrition & Food Safety*, 12(12): 9-27, 2020c; Article no.EJNFS.63712 ISSN: 2347-5641
- [65] Gimeno-Gilles C. Cellular and molecular study of germination in *Medicago truncatula*. PhD Thes. Biol. Cel. Et Mol. Veg. Ecole Doctorale Végétale. Environnement- Nutrition-Agro-Alimentaire -Mer. Univ. Angers. No d'ordre. 2009;973: 175.
- [66] Rachidi W. Biological membranes: Structures and functions. Health studies. Joseph Fourier University of Grenoble, First Year Medicine Course, UE2 Cell Biology. 2012;75.
- [67] Plots of INRA in Versailles. Management and soils studies. 2002;9(3):177-186.
- [68] Available: <https://www.hausinfo.ch> › home › assurance-securite › animaux-vermines
- [69] Fopoussi Tuebue J.C. Basga D.B., Tematio P., and Nguetnkam J.P. Water from Cooked Beans as Substrate for Some Heterotrophic Organisms: Case Study of Moulds 5(3): 1-15, 2020a; Article no.AJSSPN.53588 ISSN: 2456-9682
- [70] Camus G. Eutrophisation, Planet Vie; 2014. Available: <https://planet-vie.ens.fr>
- [71] CRE Laurentides. Dissolved oxygen; 2009. Available: <https://Crelaurentides.org>

- [72] Richert A, Gensch R, Håkan J, Stenström TA, et Dagerskog L. Practical Advice for Using Urine in Agricultural Production. EcoSaRes ser. 2011;3:SEI.73
- [73] Larsen TA, Udert KM, Lienert J. Source separation and decentralization for wastewater management (IWA Publ); 2013.
- [74] Charles D. G., Kitchen explained, Edition BPI, 2009, 306p.
- [75] Palacios-Pola G., Perales Rivera H., Figueroa-Cardenas J.D., Zorba J., Estrada H., Changes in the physical, chemical, and sensory properties from three native corn landraces from Chiapas using two nixtamalization times, International Journal of Gastronomy and Food Science, Elsevier, Vol. 25, 2021, 2021100373
- [76] Rachidi W. Biological membranes: Structures and functions. Health studies. Joseph Fourier University of Grenoble, First Year Medicine Course, UE2 Cell Biology. 2012;75.