

# PROXIMATE AND PHYSICOCHEMICAL PROPERTIES OF JELLY PRODUCED FROM BLENDS OF BEETROOT AND PINEAPPLE JUICE

Comment [RD1]: Quality

## Abstract

Jelly was produced from beetroot and pineapple and evaluated as a value added product for addressing beetroot and pineapple glut during their peak season. The fruits were cleaned, juiced and juices made into different formulations in the following ratio beetroot: pineapple, 50:50, 60:40, 70:30, 80:20 and 90:10. The formulations were identified with the letters; I, R, O, H, and A, respectively, with a commercial jelly sample, K made from beetroot as control. Proximate composition showed that the blends had better nutritional profile than the commercial sample except for energy value (262.72 KJ) and carbohydrate content (63.74 %) of the commercial jelly which was significantly ( $p < 0.05$ ) different. Protein levels within the blends were expectedly low, ranging from 1.52 % in sample I (50 % beetroot and 50 % pineapple juice) to 1.76 % in sample A. The same trend was observed in fat, ash, and moisture content except for the carbohydrate content where the sample with the least amount of beetroot, sample I (50 % beetroot and 50 % pineapple juice) had the highest value of 62.69 % within the blends thus provided the highest amount of energy, 261.30 KJ. Physicochemical properties showed that blends had less sugar contents. Both commercial sample and blends have comparable acidity which is an indication that the formulated product will be shelf stable. Total soluble solids, total sugars, reducing and non-reducing sugars increased as the proportion of pineapple increased. Titratable acidity and pH showed that an inverse relationship existed between them. The beetroot and pineapple jelly blends produced compared favourably with proximate composition and physicochemical properties of commercial jelly.

Comment [RD2]: Jelly was produced from beetroot and pineapple fruits as a value-added product for addressing their gluts during their peak season.

Comment [RD3]: and their juices were made

Comment [RD4]: , and

Comment [RD5]: a better

Comment [RD6]: except for the

Comment [RD7]: Fruit Jelly was produced from beetroot and pineapple fruits as a value-added product for addressing their gluts during its peak season. The fruits were cleaned and their juices were blended using the following beetroot: pineapple ratio; 50:50, 60:40, 70:30, 80:20, and 90:10. The fruit juice blends were then labeled I, R, O, H, and A, respectively, while the commercial jelly sample K, and beetroot jelly were used only as control. Proximate composition showed that the fruit jellies had a higher nutritional profile than the commercial jelly. The result revealed variation in moisture content (35.11 to 36.17% ), Ash content (0.12 to 0.26 %), Carbohydrate content (61.22% to 63.74%), Energy value (257.26 KJ in to 262.72 KJ) and protein content (0.88 to 1.77 %). The energy value (262.72 KJ) and carbohydrate content (63.74 %) of the commercial jelly were however significantly ( $p < 0.05$ ) higher than in the beetroot - pineapple fruits jelly. The protein content of the jelly ranged from 1.52 to 1.76 %. Overall, increasing the blending ratio increased the fat, ash, protein, and moisture contents of the jellies. Besides, the jellies with the least amount of beetroot I (50 % beetroot and 50 % pineapple juice) had the highest carbohydrate content (62.69 %) thus providing the highest amount of energy (261.30 KJ).

Comment [RD8]: Recast. Please use values of obtained to briefly summarized this section. Avoid lengthy sentences for clarity.

Comment [RD9]: carbohydrates

Comment [RD10]: , and

Comment [RD11]: vegetables

Comment [RD12]: , and

Comment [RD13]: , and

## Introduction

Fruits and vegetables have similar composition, having high water content (70-85%), relatively high carbohydrate but low in fat and protein and usually contain useful vitamins (Nzelu, 2010). The carbohydrate portion can be further broken down into digestible and indigestible parts which are sugars and starches versus pectin and cellulose material (Potter and Hotchkiss, 2017). Vegetables differ from fruits in chemical composition. Most vegetables contain more starch than sugar as contrasted with fruits which are higher in sugar than starch especially when ripe. Vegetables are edible parts of the plant which are usually cooked and salted before consumption with other foods. These may include leaves, stems, roots, flowers, seeds, fruits, bulbs, tubers, and fungi. (Nzelu, 2010).

Beetroot (*Beta vulgaris L.*) is one of the important root vegetable that belong to the *Chenopodiaceae* family and is originally from temperate climate regions of Europe and North Africa. It is a dark red vegetable whose taste is described as sweet, earthy and tender to eat (Nottingham, 2004). Beetroot is delicious if eaten raw but is more typically cooked or pickled (Partha et al., 2014). It is grown in the ground and is related to turnips, swedes and sugar beet. Beetroots are notable for their sweetness; they have the highest sugar content of vegetables but

they are low in calories. However, fresh beetroots are exposed to post-harvest spoilage due to their high nitrate content. Pineapple (*Ananas comosus*) is a tropical fruit, a member of the *Bromeliaceae* family (TFNET, 2016). Pineapples are rich in vitamins, enzymes and antioxidants. They help boost the immune system, build strong bones, and aid digestion. Despite their sweetness, pineapples are low in calories. They are members of the bromeliad family and the only member that produces edible fruit (Arshad et al., 2014).

Different food products like jam, jelly, fruit bar and marmalade are prepared from raw edible fruits. Fruit jams, jellies and marmalades are made by cooking fruits (pieces, pulps and or juice) with sugars, gelling agents (usually pectin) and edible acids, and concentrating the mixture until a characteristic and suitable consistency is obtained. The minimum amount of fruits in the final product may vary from about 35–45% (Fugel et al., 2015). Jellies are defined by Codex Alimentarius Commission, (CAC) section 2-2 as a product brought to semi solid gelled consistency and made from the juice and or aqueous extracts of one or more fruits or vegetables, mixed with foodstuffs with sweetening properties with or without the addition of water (CODEX STAN 2009).

Consumers find the flavour of beetroot repulsive and so avoid it, which calls for the need to make the vegetable available in more palatable forms. On the other hand, beetroots are rich in nitrates. This high concentration of nitrates makes it prone to microbial attacks responsible for the early deterioration of fresh raw beetroots (Nottingham, 2004). In essence, processing beetroots into jelly will solve the problem of high rate of spoilage of fresh beets, improve the usage of these nutritious root on the table of households and also enrich health with the numerous nutritional and health benefits it brings. More so, blending beetroot juice with pineapple juice will improve the flavour and provide the important nutrient, vitamin C which beetroot lacks in a nutritionally significant level, making it more acceptable and more nutritious. The objective is to produce and evaluate jelly from beetroot and pineapple juice as a value added product to curb peak season post –harvest losses of the two produce.

## Materials and Methods

Fresh beetroot (*Beta vulgaris L*) were purchased from Jos, Plateau state, Nigeria. Pineapples (*Ananas comosus*), golden penny granulated sugar and pfizer citric acid were purchased from Oigige main market in Nsukka, Enugu state Nigeria. Sure jell rapid set Pectin was ordered through Alibaba.com (online market) and was supplied through DHL courier service.

### Preparation of beetroot and pineapple juice

Fresh beetroots and pineapples were washed separately under running tap water to remove dirt and dusts. They were further peeled, sliced, blended and juiced separately before formulating into the different required blends.

### Making beetroot and pineapple blend jelly

Beetroot juice 2,500 ml and 2,000 ml of pineapple juice were extracted. The different formulations were mixed in the following ratio beetroot: pineapple, 50:50, 60:40, 70:30, 80:20 and 90:10 and 400 ml of each blend was weighed out identified with the letters; I, R, O, H, and A, respectively. Sugar 500 g was used for every 400 ml of juice. The sugar was measured and divided into three equal parts (166.67 g) each. First part was added into the juice and boiled at

**Comment [RD14]:** fruit bars,

**Comment [RD15]:** , and

**Comment [RD16]:** gelling agents (pectin), and edible acids (CODEX STAN 2009; Fugel et al., 2015). These mixtures are then concentrated until a characteristic and suitable consistency is obtained.

**Comment [RD17]:** semi-solid

**Comment [RD18]:** too lengthy and complex sentence. Use short and avoid combining phrases that will make the information wordy.

**Comment [RD19]:** Recast clumping so many information in a sentence. This result in too lengthy and complex sentences.

**Comment [RD20]:** This study aimed to determined the physicochemical and proximate composition jelly from beetroot and pineapple fruits blends. The production of jelly is a value addition that will curb peak season post –harvest fruit losses in Nigeria..

**Comment [RD21]:** What quantity of the fruits were purchased, state all these?

**Comment [RD22]:** Fruits were

**Comment [RD23]:** State

**Comment [RD24]:** , and Pfizer

**Comment [RD25]:** purchased

**Comment [RD26]:** dust

**Comment [RD27]:** Lumping so many important procedures. Scientist will adopt the procedure in future, please provide details. How was the juice extracted? What aseptic procedures were adopted?


**Comment [RD28]:** How?


**Comment [RD29]:** , and

130°C. The second part was mixed with pectin (6 g), the heat was reduced to 100 °C and the pectin and sugar mixture was heated until a gel is formed. The gel was added to the juice and was stirred continuously till it was totally dissolved, before the third part was added. Upon reaching 60 degree Brix, citric acid was added to get a pH of 3.4 for gel enhancement. Heating was discontinued at 65 degree brix and the jelly was allowed to cool to 80 °C before they were filled into already pre-sterilized glass jars leaving about ¼ inch head space for vacuum formation.

**Comment [RD30]: Making beetroot and pineapple blend jelly**

The beetroot and pineapple fruit juices obtained were then blended according to the following ratio: 50:50, 60:40, 70:30, 80:20, and 90:10. The resultant blends were then labeled as I, R, O, H, and A, respectively. Afterward, 400 ml of each blend was weighed as the working volume. A 500 g of sugar was weighed and divided into three equal portions (166.67 g) each. The first part was added to the juice and boiled at 130 oC. The second part was mixed with pectin (6 g), the heat was reduced to 100 oC and the pectin and sugar mixture was heated until a gel is formed. The gel was added to the juice and was stirred continuously till it was dissolved before the third part was added. Upon reaching 60 degrees Brix, citric acid was added to get a pH of 3.4 for gel enhancement. Heating was discontinued at 65-degree Brix and the jelly was allowed to cool to 80 OC before they were filled into already pre-sterilized glass jars leaving about ¼ inch of headspace for vacuum formation.





**Comment [RD31]:** Why sterilization? The detail of this procedure is not explain anywhere in this manuscript.

**Figure 1:** Flow diagram for beetroot and pineapple blend jelly production.  
Source: *Gava et al. (2008)*.

**Comment [RD32]:** Delete full stop on title

**Table 1:** Proportions of beetroot and pineapple juice in the jelly blends

Sample	Beetroot Juice (%)	Pineapple Juice (%)
K	--	--
I	50	50
R	60	40
O	70	30
H	80	20
A	90	10

**Keys:** Sample K= Commercial fruit jelly.  
 Sample I=50% beetroot juice and 50% pineapple juice.  
 Sample R= 60% beetroot juice and 40% pineapple juice.  
 Sample O=70% beetroot juice and 30% pineapple juice.  
 Sample H=80% beetroot juice and 20% pineapple juice.

Sample A=90% beetroot juice and 10% pineapple juice.

**Comment [RD34R33]:** This table is not a result and not necessary. You are to provide details in the procedures above, please delete

### Proximate Analyses

The proximate composition of jellies was determined according to the method of Association of Official Analytical Chemist (AOAC) (2010).

**Comment [RD35]:** the Association

**Comment [RD36]:** Chemists

### Determination of moisture content

Moisture content was determined by the hot air oven method. Stainless steel moisture dishes were cleaned and dried in the oven (Mettler Model UN30), at 100 °C for one hour to achieve constant weights. The moisture dishes were cooled in a desiccator and then weighed. Two grams of each sample was weighed into respectively labeled moisture dish and dried at 100 °C, for 1 hour, removed from the oven and placed in a desiccator to cool to room temperature before weighing. The moisture dishes were put back into the oven, dried and weighed intermittently until a constant weight was attained. The loss in weight from the original sample weight was calculated as the moisture content using the expression

**Comment [RD37]:** oven drying method to constant weight

**Comment [RD38]:** A stainless steel moisture dish was cleaned and pre dried in the oven (Mettler Model UN30), at 100 °C for 1 h until constant weights was attained and cooled in a desiccator .

$$\% \text{ moisture content} = \left( \frac{W_2 - W_3}{W_2 - W_1} \right) \times \frac{100}{1}$$

**Comment [RD39]:** Two grams of each sample was weighed into respectively labeled moisture dish and dried at 100 °C, for 1 h , This was followed by cooling in a desiccator . The moisture dishes were re dried and weighed intermittently until a constant weight was attained. The loss in weight from the original sample was calculated as the moisture content using the formula

Where:  $W_1$  = weight of empty moisture dish,  $W_2$  = weight of moisture dish + sample before drying,  $W_3$  = weight of moisture dish + sample after drying.

### Determination of ash content

Two grams of each sample was weighed into crucibles that had been previously washed, dried and weighed. The crucibles were placed in a muffle furnace (Vecstar, Model Lf3, U.S.A) and ignited at  $550 \pm 2$  °C for four hours, cooled and weighed. The ash content was calculated with the expression

**Comment [RD40]:** put

**Comment [RD41]:** clean crucibles previously weighed.

$$\% \text{ Ash content} = \left( \frac{W_2 - W_3}{W_2 - W_1} \right) \times \frac{100}{1}$$

**Comment [RD42]:** Two grams of each sample was weighed into clean crucibles previously weighed. It was then placed in a muffle furnace (Vecstar, Model Lf3, U.S.A) and ignited at  $550 \pm 2$  °C for 4 h. afterward. The muffle furnace was cooled and crucibles reweighed. The ash content was calculated as:

Where:  $W_1$  = weight of empty crucible,  $W_2$  = weight of crucible and sample before ashing,  $W_3$  = weight of crucible and sample after ashing.

### Determination of fat content

A Soxhlet extractor with a reflux condenser and a previously dried cooled and weighed 250 ml round bottom flask was assembled. Two grams of each sample was weighed into labeled thimble and petroleum ether (150 ml, boiling point 60 – 80 °C) filled into the round bottom flask. The extraction thimble was plugged with cotton wool. The Soxhlet apparatus after assembling was allowed to reflux for 6 hours. The thimble was removed with care and the petroleum ether was recovered for reuse. The round bottom flask containing the petroleum ether was dried at 70 °C for 1 hour in an oven (Mettler UN30, Germany), cooled in desiccator and weighed.

**Comment [RD43]:** a labeled

**Comment [RD44]:** was filled

Calculation:

$$\% \text{ Fat} = \frac{\text{Weight of fat}}{\text{Weight of sample}} \times \frac{100}{1}$$

**Comment [RD45]:** a desiccator, and weighed

### Determination of fiber content

Two grams of each sample was weighed and defatted using petroleum ether (Boiling point of 40 to 60°C). The defatted sample was boiled for 30 minutes in 200 ml of 1.25%  $\text{H}_2\text{SO}_4$  and the

**Comment [RD46]:** were

solution filtered through a funnel fitted with muslin cloth. It was washed with boiling water until it was free of acid. The residue was boiled for another 30 minutes with 100 ml of 0.02 M NaOH. It was further washed with boiling water then with 1 % hydrochloric acid and finally with boiling water to ensure that it was free of acid. The final residue was transferred into a crucible and dried in the oven for 1 hour. The crucible with its content was cooled in a desiccator and weighed. The residue was transferred into crucible and dried at 100 °C to a constant weight. Incineration to ash was done at 600 °C for 30 minutes, cooled in a desiccator and weighed. The difference in weight between oven dry weight and weight after incineration was taken as the fiber content of the sample. This was expressed as a percentage of the original weight of sample taken for analysis.

$$\text{Crude fiber (\%)} = \frac{\text{Weight of dried sample} - \text{weight of sample after incineration}}{\text{Initial weight of sample}} \times \frac{100}{1}$$

### Determination of crude protein content

Protein was determined using the Kjeldahl method. Two grams of each sample was weighed into a Kjeldahl flask and added anhydrous sodium sulphate (5g of Kjeldahl catalyst), concentrated H<sub>2</sub>SO<sub>4</sub> (25 ml) and few boiling chips. The samples were digested in the fume chamber to clear sample solution. Each sample digest was allowed to cool and then transferred into a 250 ml volumetric flask made up to volume with distilled water. Five milliliters of 2% boric acid solution with few drops of methyl red indicator was introduced into a distillate collector (100 ml conical flask) and placed under the condenser in a distillation unit. Five milliliters of each sample digest was pipetted into the distillation unit, washed down with distilled water followed by addition of five milliliters of 60% sodium hydroxide solution to the digest. The sample was heated until 50 ml of the distillate was collected in the receiving flask. The distillate was titrated against 0.01N HCl to a pink colored end point. For the blank diluted digest from filter paper was also distilled and the distillate titrated against 0.01N HCl Total nitrogen (%) was estimated using the expression:

$$\% \text{ Nitrogen} = \frac{\text{Titre value} - \text{Blank} \times \text{Normality of acid}}{\text{Weight of sample}} \times \frac{100}{1}$$

$$\text{Crude protein} = \% \text{N} \times 6.25$$

Where T= Titre

B= Blank

N= Normality

W= weight of acid

### Determination of carbohydrate content

Carbohydrate content of each sample was calculated by difference. The difference between 100 and the sum of percentages of moisture, protein, fat, fiber and ash of each sample was calculated and the result expressed as

$$\% \text{ carbohydrate} = 100 - (\% \text{ moisture} + \% \text{ crude protein} + \% \text{ crude fiber} + \% \text{ ash} + \% \text{ fat})$$

### Determination of energy value

The values obtained for protein, fat, and carbohydrate was used to calculate the energy value of the samples, using the At-water factor described by AOAC (2010).

$$\text{Calorific value (Kcal/100g)} = P \times 4.0 + F \times 9.0 + C \times 3.75$$

Where P, F and C are (%) protein, fat, and carbohydrate in the samples.

**Comment [RD47]:** was filtered through a funnel fitted with a muslin cloth.

**Comment [RD48]:** the crucible

**Comment [RD49]:** oven-dry

**Comment [RD50]:** the sample

**Comment [RD51]:** were

**Comment [RD52]:** , and a few boiling chips.

**Comment [RD53]:** to a clear solution

**Comment [RD54]:** a few drops

**Comment [RD55]:** were

**Comment [RD56]:** unit

**Comment [RD57]:** by the addition of 5 ml

**Comment [RD58]:** a pink-colored endpoint.. The

**Comment [RD59]:** was titrated

**Comment [RD60]:** The carbohydrate content

**Comment [RD61]:** , and

**Comment [RD62]:** Respectively.



## Physicochemical analyses

### Determination of Total soluble solid (TSS)

The TSS content of pulp was determined in accordance with AOAC [2010], using Erma hand refractometer at 20 °C and by reference tables expressed as % sucrose by weight (°Brix). The prism of refractometer was washed with water and wiped to dry after each reading.

Comment [RD63]: determined by

Comment [RD64]: the refractometer

### Determination of total sugars

Total sugar was determined using the method described by AOAC (2010). Five grams of sample was put in a 500 ml beaker, 100 ml warm water was added and it was neutralized with 10% NaOH. Two milliliters of lead acetate solution was added it was allowed to stand for 10 minutes. The necessary amount of sodium oxalate was added to remove excess lead in the solution. The volume was made up to 250 ml mark with distilled water and filtered. Fifty milliliters of the clarified and delead solution was transferred into a 250 ml flask. Ten milliliters of 1N HCL was added into the flask. This solution was boiled for two minutes. After cooling, 3 drops of phenolphthalein was added and the content was neutralized with NaOH. The solution was filtered and the volume, made up to 250 ml with distilled water. Ten milliliters of a mixed fehling's solution was pipetted into a conical flask. A burette was filled with the clarified sample solution and running the whole volume required to reduce the fehling's solution so that, 0.5-1.0 ml was still required to complete the titration. The content of the flask was mixed and heated to boiling for 2 minutes. Three drops of methylene blue indicator were added then the titration continued until color completely disappeared. The percentage total sugar was calculated as follows:

Comment [RD65]: of warm

Comment [RD66]: were

Comment [RD67]: the 250 ml

Comment [RD68]: to

Comment [RD69]: were

Comment [RD70]: filtered

Comment [RD71]: Fehling's

Comment [RD72]: Fehling's

Comment [RD73]: the color

Comment [RD74]: of total sugar

$$\text{Mg total sugar per 100 ml} = \frac{\text{Factor} \times 100}{\text{Titre}}$$

$$\text{Percentage total sugar} = \frac{\text{mg per 100ml} \times \text{Dilution}}{\text{weight of the sample} \times 1000} \times 100$$

Comment [RD75]: of total

### Determination of reducing sugars

Reducing sugars was determined using the method described by AOAC (2010). Five grams of sample was put in a 500 milliliters beaker, 100 milliliters warm water was added and it was neutralized with 10 % NaOH. Two milliliters of lead acetate solution was added it was allowed to stand for 10 minutes. The necessary amount of sodium oxalate was added to remove excess lead in the solution. The volume was made up to 250 milliliter mark with distilled water and filtered. Ten milliliters of mixed fehling's solution was pipetted into a conical flask. A burette was filled with the clarified sample solution and running the whole volume required to reduce the fehling's solution so that, 0.5-1.0 ml was still required to complete the titration. The content of the flask was mixed and heated to boiling for 2 minutes. Three drops of methylene blue indicator were added then the titration continued until color completely disappeared. The percentage reducing sugar was calculated as follows:

Comment [RD76]: were

Comment [RD77]: 500 ml

Comment [RD78]: of warm water was then

Comment [RD79]: delete

Comment [RD80]: and

$$\text{Mg reducing sugar per 100 ml} = \frac{\text{Factor} \times 100}{\text{Titre}}$$

$$\text{Percentage reducing sugar} = \frac{\text{mg per 100ml} \times \text{Dilution}}{\text{weight of the sample} \times 1000} \times 100$$

### Determination of non-reducing sugar

Non-reducing sugars was determined using the method described by AOAC (2010). Percentage reducing sugar was subtracted from total sugar and the result obtained were values for the non-reducing sugar.

### Determination of pH

pH was determined using a standard pH (model 20 pH) conductivity meter (Denver instrument united national inventory database) as described by AOAC (2010). Two grams of sample was homogenized in twenty millilitres of distilled water in a beaker. The pH meter was first standardized using buffer solutions of pH 4 and 9. The electrode was rinsed with distilled water and then dipped into the homogenate, allowing sufficient time for stabilization before taking the reading.

### Determination of titratable acidity

Titratable acidity was determined by the methods described by AOAC (2010). Ten grams of the sample was diluted into 250 ml of boiled distilled water and titrated just before the end point using 0.1 N NaOH using 0.3 ml phenolphthalein as an indicator. Two grams of sample was diluted in 20 ml of distilled water. With this, the colour was easily noticed and then it was titrated with NaOH solution until a pink colour was obtained that persisted for 30 seconds. The titration was done in duplicates and the end point was taken. The acidity was calculated using the formula:

$$\text{Titratable acidity (\%)} = \frac{0.1N(\text{NaOH}) \times T \times 75}{V \times 1000} \times 100$$

Where T = titre value

V = volume of sample

### Experimental design

The experiment was laid on a Completely Randomized Design (CRD). The mean and standard deviation were calculated by one-way analysis of variance using statistical package for service solution (SPSS) software version 22. Means were separated by Duncan's new multiple range test. Significant difference was at  $p < 0.05$  according to Steel and Torrie, (1980).

### Result and Discussion

#### Proximate composition of beetroot and pineapple blend jelly

Protein content of the samples ranged from 0.88 in sample K to 1.77 in sample H. No significant ( $p > 0.05$ ) difference was observed between samples A, H and O. Sample K varied significantly ( $p < 0.05$ ) from the rest of the samples and no significant ( $p > 0.05$ ) difference was observed between samples I and R. Sample K had the least protein content followed by I and R. Sample A had the highest. The protein content is generally about 1% of fresh fruits and 2% in most vegetables (Nzelu, 2010). It was observed that the protein content of the jellies increased with increase in beetroot. This is because beetroots have more protein than pineapples. The same trend was observed by Ajenifujah and Aina (2011), in the study of physicochemical properties

Comment [RD81]: delete

Comment [RD82]: 20 ml

Comment [RD83]: and

Comment [RD84]: A pink colour was obtained that persisted for 30 sec. The titration was done in duplicates

Comment [RD85]: Data Analysis

Comment [RD86]: Insert the Title: Result and Discussion section

Comment [RD87]: The protein content of the beetroot and pineapple blend jelly (Table 1) ranged from 0.88 to 1.77 %. The result showed no significant ( $p > 0.05$ ) difference between samples A, H and O. Sample K varied significantly ( $p < 0.05$ ) from the rest of the samples while no variation ( $p > 0.05$ ) was observed between samples I and R. Comparatively, sample K had the least protein content while sample A had the highest.

Comment [RD88]: In this study, the protein content of the jellies were observed to increased with an increase in the beetroot ratio.



and sensory evaluation of jam made from black-plum fruit and beetroot where the protein content was increasing with increase in beetroot.

Fat content of the samples ranged from 0.46 in sample K to 0.60 in sample A. No significant ( $p>0.05$ ) differences existed between samples K, I, R, and O but there was a significant ( $p<0.05$ ) difference between them and samples H and A. The amount of lipids found in fruits and vegetables are usually small, comprising less than 1% of the bulk with avocado as an exception (Nzulu, 2010) (They are usually associated with the cell membranes). The fat content in the samples showed that beetroots contains, even though they are in trace levels, higher amount of fat than pineapples as seen.

**Table 2: Proximate composition of beetroot and pineapple blend jelly**

Samples	Protein (%)	Fat (%)	Moisture (%)	Ash (%)	Fiber (%)	Carbohydrate (%)	Energy value (kj)
K	0.88 <sup>a</sup> ±0.02	0.46 <sup>a</sup> ±0.04	34.80 <sup>a</sup> ±0.00	0.12 <sup>a</sup> ±0.02	ND	63.74 <sup>1</sup> ±0.04	262.72 <sup>e</sup> ±0.09
I	1.52 <sup>b</sup> ±0.02	0.50 <sup>a</sup> ±0.00	35.11 <sup>b</sup> ±0.00	0.19 <sup>b</sup> ±0.01	ND	62.69 <sup>c</sup> ±0.01	261.30 <sup>d</sup> ±0.06
R	1.54 <sup>b</sup> ±0.01	0.53 <sup>ab</sup> ±0.04	35.14 <sup>b</sup> ±0.01	0.21 <sup>b</sup> ±0.01	ND	62.59 <sup>d</sup> ±0.03	261.25 <sup>d</sup> ±0.15
O	1.75 <sup>c</sup> ±0.00	0.53 <sup>ab</sup> ±0.04	35.40 <sup>c</sup> ±0.01	0.20 <sup>b</sup> ±0.00	ND	62.13 <sup>c</sup> ±0.02	260.23 <sup>c</sup> ±0.23
H	1.77 <sup>c</sup> ±0.02	0.58 <sup>bc</sup> ±0.04	35.75 <sup>d</sup> ±0.03	0.25 <sup>c</sup> ±0.01	ND	61.66 <sup>b</sup> ±0.03	258.88 <sup>b</sup> ±0.12
A	1.76 <sup>c</sup> ±0.01	0.60 <sup>c</sup> ±0.00	36.17 <sup>e</sup> ±0.02	0.26 <sup>c</sup> ±0.01	ND	61.22 <sup>a</sup> ±0.00	257.26 <sup>a</sup> ±0.03

Values are means ± standard deviation of 2 replications. Values with the same superscripts in a column are not significantly different ( $p>0.05$ ).

Key: Sample K=commercial mixed fruit jelly, I=50:50% beetroot and pineapple blend jelly, R=60:40% beetroot and pineapple blend jelly, O=70:30% beetroot and pineapple blend jelly, H=80:20% beetroot and pineapple blend jelly, A=90:10% beetroot and pineapple blend jelly.

Moisture content of the samples ranged from 34.80 in sample K to 36.17% in sample A. This is similar to but does not exactly conform to the 32 to 35% moisture levels for jellies (USDA 2015). No significant ( $p>0.05$ ) difference was found in sample I and R but the rest of the samples varied significantly ( $p<0.05$ ) from each other. Sample K had the least amount of moisture and sample A, the highest. Usually, fruits and vegetables have very high levels of moisture and as a result, they have low energy value and short shelf life (Nzulu, 2010). However, the heat treatment that was given to the fruit and vegetable raw materials resulted to about 60 % reduction in the moisture content, to give higher energy value and shelf stable products. The moisture content was observed to be increasing as the beetroot increased probably because beetroots, having 87.5% moisture (Gajanan et al, 2014) has about 1.5% more moisture than pineapples, with 86% (Black, 2012). Olugbenga et al. (2018) observed a similar trend in the moisture content of banana, watermelon and pineapple blend jam where the sample with the highest amount of pineapple had the least moisture level.

Ash content of the samples varied from 0.12 % in sample K to 0.26 % in sample A. Sample K, was significantly ( $p<0.05$ ) different from the rest of the samples, no significant ( $p>0.05$ ) difference was observed between samples I, R and O but they were different, significantly ( $p<0.05$ ) from sample H and A. Sample K had the least amount of ash. Within the blends, sample I had the least amount of ash and sample A the highest. The ash content was seen to be increasing with incorporation of beetroot. This confirms the report of Bakkali et al. (2009) that

**Comment [RD89]:** The same trend was observed by Ajenifujah and Aina (2011), in a study on the physicochemical properties and sensory evaluation of jam made from black-plum and beetroot. fruits. Attributably, the beetroots have higher protein content than pineapples that could contribute to the increase contents of the fruit blend.

**Comment [RD90]:** The fat content of the jellies ranged from 0.46 in sample K to 0.60 % in sample A. No significant ( $p>0.05$ ) differences existed between samples K, I, R, and O but there was a significant ( $p<0.05$ ) difference between them and samples H and A.

**Comment [RD91]:** Table 1

**Comment [RD92]:** The moisture content of the beetroot and pineapple blend jelly ranged from 35.11 to 36.17%.

**Comment [RD93]:** The finding of this study conform to the 32 to 35% moisture levels reported for jellies (USDA 2015).

**Comment [RD94]:** In this

**Comment [RD95]:** Usually, fruits are high moisture foods (Nzulu, 2010). However, with heat treatment of the fruit juice about 60 % reduction in the moisture content, could be attained. In this study, moisture content was observed to increase with increased beetroot proportion. This could be attributed to the higher moisture content (87.5% ) of beetroots than the pineapples fruit (Gajanan et al, 2014). In a study, about 1.5% more moisture content was reported in beetroots than pineapples fruits (Black, 2012). Olugbenga et al. (2018) observed a similar trend in the moisture content of banana, watermelon and pineapple blend jam where the sample with the highest amount of pineapple had the least moisture level.

**Comment [RD96]:** Recast, avoid excessive presentations and use of wordy sentences. This style make reading very dreary.

the ash content of most vegetables are generally higher than that of fruits. Since ash content of a food material is a measure of the inorganic compounds present, it is not wrong to assume that the samples with higher ash content are rich in inorganic compounds.

Crude fibre was tested for but was not detected in any of the samples because the raw materials were juiced and the fibrous materials were removed before further processing.

Carbohydrate content of the jellies ranged from 61.22% in sample A to 63.74% in sample K. All the samples varied significantly ( $p < 0.05$ ) from each other. Sample A had the least amount of carbohydrate and sample I, the highest within the blends. Sample I with the least amount beetroot and the highest amount of pineapple recorded the highest amount of carbohydrate. Similar trend was observed by Anuradha et al. (2017) in pineapple and papaya blend jam where the samples with more pineapple had more carbohydrate and also Olugbenga et al. (2018) observed same trend in banana, watermelon and pineapple blend where the sample with the most pineapple had the highest carbohydrate content. Increase in pineapple resulted in increase in the carbohydrate content. In general, fleshy fruits possess high sugar content than vegetables, glucose, fructose, sucrose and starch constitute the available carbohydrate of fruits and vegetables. Starches in fruits usually disappears on ripening (Nzulu, 2010) therefore sugars are the predominant carbohydrate in the product and are very necessary for proper functioning of the brain.

The energy value of the products ranged from 257.26 KJ in sample A to 262.72 KJ in sample K. No significant ( $p > 0.05$ ) difference was observed between sample I and R but the rest of the samples differed significantly from each other. Within the blends, sample I had the highest energy level of 261.30 KJ. The energy value of a food indicates its value to the body as fuel (Schmidt, 2015). Energy requirement can be thought of as the amount needed to maintain the basic processes of life at rest, that is, basal metabolism, plus the amount needed for physical activity under a variety of circumstances. Body weight is an important factor in determining how much energy we need, since more energy will be needed to sustain and move a greater body mass (Schmidt, 2015). USDA (2015) recommends that the daily energy intake of the average man should be about 10,500kj and 40 to 65% of this should come from dietary carbohydrate which is about 6300kj from carbohydrates. The carbohydrate in the product provides about 244.88kj which is roughly 4.0% of the daily requirements for carbohydrates

**Comment [RD97]:** In this study, the carbohydrate content of the jellies ranged from 61.22% to 63.74%. All the samples varied significantly ( $p < 0.05$ ) from each other in terms of carbohydrate content. The finding of this study revealed that jelly A had the least amount of carbohydrate. Sample I with the least proportion beetroot to pineapple fruits however, had the highest amount of carbohydrate. Similar trend was observed by Anuradha et al. (2017) in a study on pineapple and papaya blend jam where increased pineapple fruit proportion resulted to higher carbohydrate content. Olugbenga et al. (2018) had similarly observed that in banana, watermelon and pineapple blend, sample with increased pineapple fruits had the highest carbohydrate content. In general, fleshy fruits possess high sugar content, glucose, fructose, sucrose and starch constitute than vegetables. Besides, starches in the fruits usually could also disappears on ripening (Nzulu, 2010).

### Physicochemical properties

The total soluble solids (TSS) content of the jellies in the present study were in the range of 63.84 % in sample A to 65.20 % in sample K, with the commercial sample K, having the highest percentage. The final Total Soluble Solids (TSS) content of a jam (jelly) (also known as "Degrees Brix") shall in all cases be between 60 to 65 % (CODEX STAN, 2009). The TSS is a measure of the amount of material that is soluble in water. It is expressed as a percentage. A product with 100 % soluble solids, has no water and one with 0 % soluble solids is all water. The correct sugar content is critical for proper gel formation and for preservation of jelly. If the final TSS is lower than 60-65 % the shelf life will be reduced. The product will have a runny consistency and bacteria and moulds will be able to grow in the product. If the TSS is higher than 65%, the jelly will be very stiff and the sugar might start forming crystals in the product

**Comment [RD98]:** The total soluble solids (TSS) content of the jellies in the study range between 63.84 to 65.20 %. (Table 3), the result showed comparatively lower total soluble solids (TSS) content of the jelly with the commercial sample.

**Comment [RD99]:** The total soluble solids (TSS) content of the jelly observed is in conformity with standard of 60 to 65 % stipulated by CODEX STAN, (2009).

(CODEX STAN 2009). However, according to CODEX STAN (2009), all the products fell within this range therefore the products can be said to be of adequate quality.

**Comment [RD100]:** Recast, the paragraph is not logically presented, wordy with so many uncompleted statements. Important points are not backed up by citations/references.

**Table 3: Total soluble solids, total sugar, reducing and non-reducing sugar composition**

**Comment [RD101]:** Table 2

Samples	Total soluble solids (°Brix)	Total sugar (%)	Reducing sugars (%)	Non-reducing sugars (%)
K	65.20 <sup>e</sup> ±0.00	44.15 <sup>e</sup> ±0.02	28.90 <sup>e</sup> ±0.00	15.25 <sup>c</sup> ±0.02
I	64.89 <sup>d</sup> ±0.00	43.94 <sup>d</sup> ±0.00	28.77 <sup>d</sup> ±0.04	15.18 <sup>bc</sup> ±0.04
R	64.86 <sup>d</sup> ±0.01	43.92 <sup>d</sup> ±0.03	28.76 <sup>d</sup> ±0.00	15.16 <sup>bc</sup> ±0.03
O	64.60 <sup>c</sup> ±0.01	43.75 <sup>c</sup> ±0.05	28.60 <sup>c</sup> ±0.01	15.15 <sup>b</sup> ±0.06
H	64.25 <sup>b</sup> ±0.03	43.51 <sup>b</sup> ±0.00	28.49 <sup>b</sup> ±0.00	15.02 <sup>a</sup> ±0.00
A	63.84 <sup>a</sup> ±0.02	43.23 <sup>a</sup> ±0.02	28.30 <sup>a</sup> ±0.02	14.94 <sup>a</sup> ±0.04

Values are means ± standard deviation of 2 replications. Values with the same superscripts in a column are not significantly ( $p>0.05$ ) different.

Key: Sample K=commercial mixed fruit jelly, I=50:50% beetroot and pineapple blend jelly, R=60:40% beetroot and pineapple blend jelly, O=70:30% beetroot and pineapple blend jelly, H=80:20% beetroot and pineapple blend jelly, A=90:10% beetroot and pineapple blend jelly.

The values for the total sugars in the samples ranged from 43.23 in sample A to 44.15% in sample K. All the samples varied significantly ( $p<0.05$ ) from each other except for sample I and R that had no significant ( $p>0.05$ ) difference. As seen on the table, the total sugar level was decreasing as beetroot increased showing that beetroot contained less sugar than pineapples.

Reducing sugars ranged from 28.30 in sample A to 28.90 % in sample K. No significant ( $p>0.05$ ) difference was seen between sample I and sample R, every other sample differed significantly ( $p<0.05$ ) from each other. The result obtained was in accordance with that obtained by Lokonuzzaman et al. (2015) in quantitative estimation of the amount of sugar in fruit jams available in Bangladesh where the reducing sugars for pineapple jam was 28.00 %.

The values for the non-reducing sugars followed the same trend as that of reducing sugar where sample I and R had no significant ( $p>0.05$ ) difference between them. Sample A had the least value of 14.94 % while sample K had the highest value of 15.25 %. According to Lokonuzzama et al (2015), the values for non-reducing sugar was 10.00 % which corresponds to the values from the samples.

The pH of the products ranged from 3.41 in sample K to 3.46 in sample H. No significant ( $P>0.05$ ) difference existed between sample R, H, and A, but they differed ( $p<0.05$ ) significantly from sample K, I and O. Therefore it can be said that pineapple contributed significantly to the acid concentration of the product. The result was also in line with that from Olugbenga et al. (2018). The pH of jams and jellies are very essential as low pH is maintained in the product to prevent microbial growth and enhance the colour and flavour of the product. On the other hand, low pH is necessary for the setting of jams or jellies.

The acidity in the products ranged from 0.28 % in sample H to 0.32 % in sample K. Sample K varied significantly ( $p < 0.05$ ) from sample H and no significant ( $p > 0.05$ ) difference was seen between the rest of the samples. Acidity reduced with increase in beetroot. The level of acidity in jams and jellies are inversely related to the pH as seen from Table 4. High acidity is required for preservative actions, colour and flavour improvement and setting of the jelly. A moderate level of acid is critical as too much acid will cause the jelly to weep and too little acid, may deter the product from setting (Tomas *et al.* 2007).

**Comment [RD102]:** In complete sentence

**Comment [RD103]:** Discuss the effects of this finding?

**Table 4:** Physicochemical composition of beetroot and pineapple blend jelly

**Comment [RD104]:** Table 3

Sample	pH	Titrateable acidity(%)
K	3.41 <sup>a</sup> ±0.01	0.32 <sup>b</sup> ±0.03
I	3.42 <sup>a</sup> ±0.00	0.31 <sup>ab</sup> ±0.01
R	3.45 <sup>b</sup> ±0.00	0.30 <sup>ab</sup> ±0.00
O	3.42 <sup>a</sup> ±0.02	0.30 <sup>ab</sup> ±0.01
H	3.46 <sup>b</sup> ±0.01	0.28 <sup>a</sup> ±0.01
A	3.45 <sup>b</sup> ±0.01	0.29 <sup>ab</sup> ±0.01

Values are means ± standard deviation of 2 replications. Values with the same superscripts in a column are not significantly ( $p > 0.05$ ) different.

Keys: Sample K=commercial mixed fruit jelly, I=50:50% beetroot and pineapple blend jelly, R=60:40% beetroot and pineapple blend jelly, O=70:30% beetroot and pineapple blend jelly, H=80:20% beetroot and pineapple blend jelly, A=90:10% beetroot and pineapple blend jelly.

## Conclusion

This study has shown that beetroot utilization in Nigeria can be improved and postharvest losses reduced by processing them into value added products such as jellies. Proximate composition showed that the blends had better nutritional profile than the commercial sample except for the energy value and carbohydrate content where the commercial jelly was significantly ( $p < 0.05$ ) higher. Physicochemical properties showed that blends had less sugar contents and thus will be preferred by most adults. Both commercial sample and blends have comparable acidity which is an indication that the formulated product will be shelf stable.

**Comment [RD105]:** Recast this section to reflect conclusions that are directly based on the objectives of this study.

## COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

**Comment [RD106]:** Too wordy, please recast

## References

- Ajenifujah, S. S. and Ania, J. O. (2011). Physicochemical properties and sensory evaluation of jam made from black-plum fruit and beetroot. *African journal of food, agriculture, nutrition and development*, 11(3): 4772-4784.
- Anuradha, D., Arya, A., Jaishree, G., Bhalerao, A. and Rupali S. S. (2017) Development of value added papaya and pineapple jams. *Food science research journal*, 8(1) 20-21.
- AOAC, (2010). Association of official analytical chemists. Official methods of analysis, 18th edition, Washington D. C., USA.
- Arshad, Z., Amid, A., Yusof, F., Jaswir, I Ahmad, K., Loke S. P. (2014). Bromelain: An overview of industrial application and purification strategies. *Journal of applied microbiology and biotechnology*, 89(17): 7283-97.
- Bakkali, K., Martos, N. R., Souhail, B. and Ballesteros, E. (2009) Characterization of trace metals in vegetables by graphite furnace atomic absorption spectrometry after closed vessel microwave digestion. *Food Chemistry*, 116(2):590–594.
- Black, R. F. (2012). Pineapple growth and nutrition over a plant crop cycle in south eastern Queensland. 1. Root development and general growth features. *Journal of Agricultural Science*, 4(19):435-451.
- CODEX STAN (2009), FAO/WHO Food standards. Codex standard for jams, jellies and marmalades. Codex Stan 296–319. <http://codexalimentarius.net>. Accessed 27th August, 2019.
- Fugel, R., Carle R. and Schieber A. (2015). Quality and authenticity control of fruit purées, fruit preparations and jams – a review. *Trends in Food Science and Technology*, 3(16): 433–441.
- Gajanan, P., Naveen, J. and Rohit, S. (2014). Application of beetroot as natural colouring pigment and functional ingredient in dairy and food products. *International Journal of Current Microbiological Applied science*, 7(12):2010-2016.
- Gava, A. J., Silva, C. A. and Frias, J. R. (2008). Food technology principles and applications. *Journal of food science*, 14(71):100-106.
- Lokonuzzaman, A., Nazrul, I. M. and Saidul, I. (2015). A quantitative estimation of the amount of sugar in fruit jams available in Bangladesh. *Science journal of analytical chemistry*, 3(5) 52-55.
- Nancy, C. F. and Schlenker, C. D. (2010). Making Jam, Jelly, and Sweet Spreads at Home. <http://aces.nmsu.edu/county>) or online at [http://nchfp.uga.edu/publications/publications\\_usda.html](http://nchfp.uga.edu/publications/publications_usda.html). Accessed 23rd February, 2019.
- Nottingham, S. (2004). Beetroot by Stephen Nottingham. Zed books publishers, London pp 1-98.

**Comment [RD107]:** The references are obsolescent. The most current is 2017. This manuscript has not discussed the current trends in the production of jellies from beetroot and pineapple fruits blend. I strongly recommend that it is very necessary to update knowledge for the 5 years gap provided by this manuscript.

- Nzelu, I. C. (2010). Identification composition and processing of tropical food commodities. 2nd edition. Eaglet publishers, Enugu state, Nigeria. 126-139.
- Olugbenga, O. A., Grace, O. O., Modupe E. O., Modupe, E. O., and Fumilayo, G. O. (2018). Functional jam production from blends of banana, pineapple and watermelon pulp. *International Journal of Food Science and Biotechnology*, 3(1) 9-10.
- Partha, P. B., Narzary, K., and Rakhi, B. (2014). Antioxidant activity of red beet juices obtained after microwave and thermal pretreatment. *Journal of Food Science*. 16(31): 139-147.
- Pomeranz, Y., Benton-Jones, J., and Martha T. (2012). Developments in the measurement of trace metal constituents in foods. In *Analysis of Food Contaminants* (J. Gilbert, edition.), Elsevier Applied Science Publishers, London. 157–206.
- Potter, N. N., and Hotchkiss, J. H. (2017). Review of non-enzymatic browning and antioxidant capacity in processed foods. *Trends in Food Science and Technology*, 7(11): 340–346.
- Schmidt, R. K., (2015). Why Combustions are always exothermic, yielding about 418kj per mole of O<sub>2</sub>. *Journal of chemistry education*, 9(2): 2094-2099.
- Steel, R.G.D. and Torrie, J.H. (1980). *Principles and Procedures of Statistics*. Second edition. McGraw-Hill Publishers, New-York pp 176-183
- TFNET (2016). International Tropical Fruits Network news compilation. Major fruits, tropical fruits information. Retrieved May 12 2021 from <https://www.itfnet.org>
- Tomas, L. F., Garcia, V. C., Ferreres, F and Tomas, B. F. (2007). Phenolic compounds analysis in determining fruit jam genuineness. *Journal of Agriculture, food and chemistry*, 40(6): 1800-1804.
- United States Department of Agriculture. (2015). National nutrient database for standard reference, agricultural research service. National agricultural library, USA.
- Vanitha, T. and Khan, M.(2019). Role of pectin in food processing and food packaging. Peer reviewed chapter, advance online publication DOI:10.5772/intechopen.83677.