

# Case report SPONTANEOUS RUPTURE OF URINARY BLADDER: A Case Report

Comment [gl1]: not well written, it is SPONTANEOUS

## ABSTRACT

### ABSTRACT:

**INTRODUCTION:** Bladder rupture is most frequently seen after some form of trauma. However, Spontaneous bladder rupture (SBR) may also occur in rare cases and often presents with non-specific clinical features delaying the diagnosis and management.

**CASE PRESENTATION:** A 51-year-old male presented with abdominal distension, pain, vomiting, constipated and urinary retention. The patient was diagnosed with bladder rupture. Immediately was surgical explorer finding the bladder wall perforation; therefore, it closed by primary suturing. Patient was discharged 10 days after the procedure with active follow up for 2 months and experienced no further complaints.

**DISCUSSION:** In most of the cases of bladder rupture, there is a history of trauma; however, SBR can occur without history of trauma. It could present as a case of pseudo renal failure with acute abdomen. The suspect is infrequent; however, it must be considered. Almost always, there is an underlying abnormality.

**CONCLUSION:** Urinary Bladder rupture is an emergency condition. If it left untreated or undiagnosed for long it will be fatal. Therefore, high index of clinical suspicion is necessary for early diagnosis and proper treatment of bladder rupture.

*Keywords: Bladder rupture, Cystogram, Pseudorenal failure, Diagnostic challenge.*

## 1. INTRODUCTION

Spontaneous bladder rupture (SBR) encompasses cases where there is no history of trauma. It is known that a pressure of more than 300 cm H<sub>2</sub>O is required to rupture a normal bladder most commonly over the dome, the thinnest and least supported part of the bladder. [1] and as a rule, it is the peritoneal segment. [2]

Almost ever, SBR occurs if there is an underlying pathology. The ethologist of can be categorized as direct or indirect. Direct are those producing direct cellular death, for example, chemical solutions, excessive heat, radiation, indwelling catheters, calculi, infection, prostatic electrocoagulation, and carcinoma. Indirect are those that interfere with blood supply, for example, the presence of a gravid uterus, urinary retention, trauma, pelvic thromboembolic events, or diabetes. Bladder necrosis is most often seen in patients with prolonged labor journeys, intentional ligation of the internal iliac

arteries to control bleeding, or urinary tract infections. [3] Patients normally present with one of these conditions and have a short history of severe lower abdominal pain. [4]

The clinical presentation is variable and apart from abdominal pain includes different urinary complaints being a challenge the final diagnosis, even with the aid of Computed Tomography (CT). If untreated, it can lead to severe complications such as sepsis, renal failure and hyperkalemia, and can eventually death. [5]

## 2. PATIENT PRESENTATION:

A 50-year-old male presented to emergency with following abdominal complaints, distension for about 2 days initially in the lower part of the abdomen and then all over the abdomen. Also vomiting, constipation and urinary retention without fever or history of trauma.

### 2.1 CLINICAL FINDINGS:

The patient had a pulse rate of 96 beats per minute, blood pressure – 116/78 mm Hg, respiratory rate of 16 times per minute and temperature of 98.8 Fahrenheit. On examination abdomen was tense and distended with generalized guarding present all over the abdomen. After Foleys catheter insertion the distension and tenderness were relieved significantly. Patient was asymptomatic with no significant clinical findings for the entire duration when the Foley catheter was in situ. However, after removing urinary catheter, the patient developed similar clinical manifestation than before.

### 2.2 INVESTIGATIONS:

The patient was vitally stable. Blood was sent for standard workup including hemoglobin, total white blood count, platelet count, Prothrombin time and International Normalized Ratio (INR). The patient was sent for erect X ray chest and abdomen and kidney, ureters and bladder, followed by ultrasonography of the abdomen. After stabilization of the patient, the patient underwent a CT scan. Cystogram was done. Micturating Cystourethrogram was attempted after that which was unsuccessful as patient was not able to pass urine. Ascitic fluid tapping was done and sent for total white blood cells and levels of creatinine and urea.

### 2.3 THERAPEUTIC INTERVENTION :

Patient had tachycardia and guarding with distended abdomen on examination. However, he was first thought as a case of abdomen distension due to medical causes and managed clinically. When he was unable to urinate voluntarily another urinary catheter was inserted obtaining 4 liters of urine with a clear clinical relief. Ascitic tapping was done and it was suggestive of elevated WBC counts in the ascitic fluid. A **CECT** of abdomen and pelvis was done with catheter in situ and which was not suggestive of any abnormality so Micturating Cystourethrogram and Cystogram were attempted to rule out urethral and bladder pathology as patient was unable to pass urine voluntarily.




Figure 1:

- A- Cystogram immediately after injecting the radiopaque contrast.
- B- Cystogram after 30 minutes of injecting the radiopaque contrast.

Cystogram revealed leakage of dye into the peritoneum. Diagnostic aspiration of the ascitic fluid was done and sent for biochemical examination and it revealed elevated levels of creatinine indicating a leak from the urinary system.



Figure 2: X-ray abdomen erect taken 1 hour after Cystogram




Figure 3: Intraoperative finding of perforation in bladder. (Upward direction points to the caudal end of the patient where two Langenbeck retractors are visible)

Emergency exploratory laparotomy was done and patient had a bladder perforation which was closed in two layers. A pre-vesical and pelvic drains were left with a urinary catheter.

## 2.5 FOLLOW-UP AND OUTCOME:

The pelvic drain was removed on the 5<sup>th</sup> post-operative day. The pre-vesical drain was removed on 8<sup>th</sup> postoperative day. Patient was discharged on 10th day after removal of Foley catheter. Histopathological findings demonstrated a congestion of the thin-walled vascular capillaries in the deep muscular layer.

At follow up every month, the patient was without complain and at 3 months a new cystogram demonstrated no urinary leak.

## 3. RESULTS AND DISCUSSION

SBR is a rare entity and is often the result of an underlying pathology. [6] Some important factors in the etiology are weaknesses of the bladder wall such as enterocystoplasty; radiation injury; and surgical scars of cystolithotomy or diverticulum. These factors are compounded by decreased compliance, overdistension, increased bladder outlet resistance and severe cystitis. Because the weakening factor may not be modifiable, it is pertinent to meticulously manage the compounding factors to prevent rare event. [1] **The patients more predisposed are those with neurogenic bladder, a history of enterocystoplasty, after pelvic radiotherapy or malignant bladder tumors. [2] (it is repeated yet)**

Clinically, most patient present with lower abdominal pain with associated symptoms of dysuria, anuria, and hematuria. In most cases the symptoms of urinary tract infection were the initial complaints; these were later accompanied by peritonism. [7]

If a rupture exists for an extended period of time (>24 hours), urine tests may show microscopic hematuria. Furthermore, an elevated serum creatinine due to reuptake of urine creatinine through the peritoneum is commonly seen. [8]

The immediate relief of abdominal pain after catheterization could be explained by the significantly reduced intraperitoneal extravasation of urine, intraperitoneal pressure, and peritoneal irritation after the relief of urinary retention. [9]

The imaging test of choice is cystography, which shows intraperitoneal contrast extravasation. Accuracy is close to 100%. [3]

Computed tomography may show free intraperitoneal fluid, even though this finding alone does not warrant a definitive diagnosis. [3]

If intra-peritoneal rupture has occurred, patients present with peritonism and blood tests consistent with acute renal failure due to the intra-peritoneal resorption of urine. [4] Retroperitoneal rupture may be treated conservatively, but otherwise surgery is often the only modality of treatment. [4]

Conservative management should also be considered for temporization in severely septic patients with intra peritoneal bladder rupture if a delay in surgery is contemplated. Accurate diagnosis is mandatory before conservative management. [1]

Awareness of this serious condition, which has a high mortality rate (47%), as well as early surgical intervention is prerequisite to prevent the undesirable progression to abdominal sepsis and death. [2]

## 4. Conclusion

Spontaneous intraperitoneal bladder rupture is a rare event whose course can be severe and even lethal if early diagnosis and treatment are not achieved, **and it must be always suspected in high risk patients.** [7] Almost all the cases have some underlying pathology which makes their bladders prone to rupture. Early diagnosis can prevent significant mortality

associated with this condition as it prevents severe complications. Patients may recover after intervention for the bladder rupture, but it's prudent to find the underlying cause and treat that as well so to prevent any future of the bladder.

## CONSENT (WHERE EVER APPLICABLE)

Informed consent was obtained from the patient for treatment and management on admission. Also the patient consent was obtained for publication of article postoperatively.

## ETHICAL APPROVAL (WHERE EVER APPLICABLE)

None

## REFERENCES

1. Agarwal M, Singh S, Naja V, Mavuduru R, Mandal A. Spontaneous Bladder Rupture: A Diagnostic and Treatment Dilemma Case Studies and Literature Review. *UIJ* 2009 Apr;2(2). doi:10.3834/uij.1944-5784.2009.04.04.
2. Basavaraj DR, Zachariah KK, Feggetter JGW. Acute abdomen - remember spontaneous perforation of the urinary bladder. *J R Coll Surg Edinb*. 2001; 46(5):316-7.
3. Gomes CA, de Figueiredo AA, Soares Júnior C, Bastos Netto JM, Tassi FR. Acute abdomen: spontaneous bladder rupture as an important differential diagnosis. *Rev Col Bras Cir* 2009; 36(4):364-5.
4. Mitchell et al.: Acute abdomen caused by bladder rupture attributable to neurogenic bladder dysfunction following a stroke: a case report. *Journal of Medical Case Reports* 2011 5:254.
5. Palthe S, Dijkstra GA, Steffens MG. A case of spontaneous urinary bladder rupture secondary to urinary retention due to a urethral stricture. *Urology Case Reports*. 2018; 17:85-87.
6. Schraur WH, Huffman J, Bagfley DH. Acute abdominal pain caused by spontaneous perforation of the urinary bladder. *Surg Gynecol Obstet*. 1983; 156:589-592.
7. Ahmed J, Mallick IH, Ahmad SM. Rupture of urinary bladder: a case report and review of literature. *Cases J*. 2009; 2:7004.
8. Dubey I.B., Mohanty D., Jain B.K. Diverse presentation of spontaneous rupture of urinary bladder: review of two cases and literature. *Am J Emerg Med*. 2012; 30(5):832. E1-832.e3. [PubMed] [Google Scholar].
9. Zhang X, Zhang G, Zhang L, et al. Spontaneous rupture of the urinary bladder caused by eosinophilic cystitis in a male after binge drinking: a case report. *Medicine (Baltimore)* 2017; 96:e9170.

## YOU CAN MENTION THIS ARTICLE

[Spontaneous rupture of urinary bladder: a case report and review.](#)

Albino G, Bilardi F, Gattulli D, Maggi P, Corvasce A, Marucco EC. *Arch Ital Urol Androl*. 2012 Dec;84(4):224-6. PMID: 23427749 Review.

In most cases, the **spontaneous rupture** of the **bladder** takes place in presence of a urothelial neoplasm or after radiation therapy of the pelvic organs. The etiology of **spontaneous rupture** of the **bladder** in our case does not relate to a ...