

# Assesment of Tree tree vigour vigor models models in University of Port Harcourt Park

## ABSTRACT

Trees are the unsung heroes of the environment that requires care and maintenance so as to continue its provision of quantifiable benefits. Hence a better understanding of the physical aspect of tree mechanics would help to provide basis for the proper management of trees. The ~~studystudy: was to assess~~ ~~the~~ ~~assess the~~ vigor of ~~selected trees within-in~~ University of Port Harcourt Park. Data were collected from four ~~Purposively~~ selected tree species based on their abundance in the study area. Tree growth parameters were measured and tree vigor estimated using live crown ratio. Different regression functions were used to model tree vigor. ~~The selected tree species used were Terminalia mantaly, Azadirachta indica, Mangifera indica and Terminalia cattapa with frequency of 214 (32.8%), 123 (18.89%), 82 (12.60%) and 79 (12.14%) relatively.~~ Tree growth parameters such as height, Diameter at breast height, crown variable measured showed ~~significate-significant~~ correlation with tree ~~vigour- vigor~~ (live crown ratio). The prediction models revealed plausible regression coefficient of determination ( $R^2$ ) value ranging from 0.247 - 0.953 and standard Error of Estimate (SEE) ranging from 0.0003 – 6.209. From the validation test, it was observed that all the models were good in prediction. However, the double log function was selected as the best model with  $R^2$  value of 0.953 and SEE value Of 0.0003. Conclusively, the trees within the University of Port Harcourt Park are stable hence are not prone to wind throw.

**Comment [G1]:** Please take these materials under materials and methodes

**Key|words:** ~~Models~~, Stability, Tree, Urban, ~~Vigour~~ Vigor

## 1. INTRODUCTION

Trees are the unsung heroes of the environment that provide quantifiable benefits [1] including basic and naturally regenerating material for timber and pulp industries, a living space for animals, insects, and plants, and recreation opportunities for people. As an important asset of the environment, trees require care and maintenance. Maintaining trees to function adequately requires an intact forest cover with mechanically stable trees. Forest management policy depends on societal value of trees and the evaluated mechanical stability of the forest [2]. Different tree management strategies affect tree vigour and protection levels differently. According to [3], tree ~~vigour-vigor~~ is the capacity of a tree to resist strain or capacity to survive under increasing stimulus that threatens survival.

The load experienced by a tree can be considered from the tree itself or climatic factors. loading caused by the tree itself which is triggered by poor maintenance and management system such as excessive pruning, poor compartment, deformed stem, less branching pattern, thin bole and age effect, affects growth parameters thus reducing tree vigour. Secondly, climatic factors such as heavy rainfall, wind etc. can cause load on trees. Hence, trees has to be care and maintained in order to be vigorous enough to withstand the mechanical stress imposed on them, which in turn will prevent tree hazard. A better understanding of the physical aspects of tree mechanics would help to provide better basis for the management of tress with respect to their mechanical vigour and protective function [4, 5]. Vigorous trees which is characterized by well-formed root system has the ability to develop forces or moments that restore the original condition when disturbed from a condition of equilibrium or steady motion compared to trees with less vigor that is susceptible to falling.

Trees with less vigour can be classified as unhealthy trees. Unhealthy trees partly contradict the benefits provided to urban landscapes. Destruction or damage caused by unhealthy trees incurs more cost to the manager. The hazard created by unhealthy trees can disrupt everyday life by removing power to homes, destroying property such as cars or a roof, and possibly even lead to loss of life. These risks prove the importance of maintaining the health of our trees. Hazard created by the presence of a less vigorous tree can have multiplying effect on the environment. According to [6], the impacts of one hazardous event of a tree or trees are often aggravated by interaction with other hazards.

Tree vigour reduces mortality response to short and long term environmental stress, it improves the mechanical resistance to pest and diseases and also improves tree stability thereby making trees not to be susceptible to abiotic factors such as wind throw. University of Port Harcourt as an academic environment is highly populated and these human population must be protected from negative impact of less vigour tree hence it is imperative to know the vigour status of the trees within the university environment in relations to its growth parameters.

Therefore, this study focus on assessment of modelling—tree vigour—vigor status, effect and management modeling in University of Port Harcourt Park with the aim of knowing the vigour status of the trees, factors that threatens tree vigour, how to improve tree vigour and negative effect of trees with less vigour.

## 2. METHODOLOGY

### 2.1 Study Area

This study was carried out in University Park of University of Port Harcourt, Choba in Obio-Akpor Local Government Area of Rivers State, Nigeria. The University of Port Harcourt is located at Latitude  $4^{\circ} 53' 14''$  N through  $4^{\circ} 54' 42''$  N and longitude  $6^{\circ} 54' 00''$  E through  $6^{\circ} 55' 50''$  E (Figure 1). The vegetation of University of Port Harcourt is tropical rain forest.




Figure 1: Map of University of Port Harcourt park  
Source: [7]

Table 1 Tree species selected for study based on their abundance

Species	Family
<i>Terminalia mantaly</i>	Combretaceae
<i>Azadirachta indica</i>	Meliaceae
<i>Mangifera indica</i>	Anacardiaceae
<i>Terminalia catappa</i>	Combretaceae

Formatted: Indent: First line: 0 cm

## 2.2 Sampling Techniques

Reconnaissance survey was carried out in order to identify and take inventory of the trees species within the University. Purposive sampling technique was used to select four major trees species for the study based on the abundance of the trees. Complete enumeration of the selected sample trees ~~was~~ were done.

## 2.3 Data Collection

Data ~~was~~ were collected from all the selected trees species within the study area. The following tree growth variables were measured for all selected trees species:

- Total height
- Clear Bole height
- Merchantable height
- Crown length
- Crown diameter
- Diameter outside bark at breast height (DBH, 1.3m above the ground)

Clinometer and distance tape was used for the height and crown diameter measurements respectively while diameter tape was used to measure diameter outside bark at breast height.

## 2.4 Data Analysis

The data collected from tree's characteristic measurement was processed into suitable form for statistical analysis. Data processed includes Basal area estimation and tree vigour estimation.

## 2.5 Computation of Model Variables

### 2.5.1 Basal Area Estimation

The basal area for each tree selected by purposive sampling techniques and measured was estimated using this formula.

$$BA = \frac{\pi D^2}{4}$$

Where;

BA = Basal Area (m<sup>2</sup>)

D = Diameter at Breast Height (DBH)

### 2.5.2 Tree Vigour Estimation

A useful measurement to indicate tree vigour is live crown ratio. This is the ratio of crown length to total tree height or the percentage of a tree total height that has foliage.

Tree vigour was estimated for all trees using this formula

$$LCR = \frac{CL}{TH}$$

Where

LCR = Live Crown Ratio

CL = Crown Length

TH = Total Tree Height

## 2.6 Statistical Analysis

Descriptive Statistics was used to estimate growth parameters attributes and tree vigour. Inferential statistics such as regression and correlation analysis were also used in this study. Regression analysis was used to develop tree vigour models for the selected trees species. Correlation analysis was used to evaluate the association between measureable tree characteristics and vigour of some trees within the study area.

## 2.7 Development of Tree Vigor Model

### 2.7.1 Model Description

Model is expressed as mathematical equation showing basic structure of the relationship among variables. Two-third (2/3) of the data collected from the observed trees was used in developing the model while the remaining one-third (1/3) was used in model testing and validation. Different equation functions were used in developing tree vigour prediction model.

### 2.7.2 Model Evaluation

The model that was formulated was evaluated with a view of selecting the best estimate for tree vigour prediction. The evaluation is based on the following criteria

- 1 Coefficient of determination ( $R^2$ )
- 2 Standard Error of Estimate (SEE)
- 3 Significance of the overall Regression Equation (F-Ratio)
- 4 Significance of Regression Coefficient.

### 2.7.3 Model Validation

For a model to be valid, the observed and predicted value should not be significantly different at 0.05 level of significance. The selected model was used to predict tree vigour for the trees species in the University. The values was compared with the observed value and the differences is expressed as residual (bias). Paired student t-test procedure was used to compare predicted values with observed values.

## 3. RESULTS AND DISCUSSION

### 3.1 Tree Species Inventory

The reconnaissance survey carried out within the University of Port Harcourt Park enumerated a total of six hundred and fifty-one (651) trees comprising of thirteen (13) identified species belonging to eleven (11) families (Table 42). The first four species ~~on the table~~ were used for the study ~~due to their because they are the dominant dominance tree species~~ in the University Park. The most abundant species presented is the *Terminalia mantaly* ~~which accounted~~ that accounted for 32.87% (214) of the total population.

**Table 42: Composition of Trees within the University of Port Harcourt Park**

Species	Family	Frequency	Percentage composition
<i>Terminalia mantaly</i>	Combretaceae	214	32.87
<i>Azadirachta indica</i>	Meliaceae	123	18.89
<i>Mangifera indica</i>	Anacardiaceae	82	12.60
<i>Terminalia catappa</i>	Combretaceae	79	12.14
<i>Delonix regia</i>	Fabaceae	33	5.07
<i>Psidium guajava</i>	Myrtaceae	22	3.38
<i>Persea americana</i>	Lauraceae	20	3.07
<i>Anacardium occidentale</i>	Anacardiaceae	19	2.92
<i>Lagerstroemia speciosa</i>	Lythraceae	19	2.92
<i>Hura crepitans</i>	Euphorbiaceae	19	2.92
<i>Gmelina arborea</i>	Lamiaceae	9	1.38
<i>Prunus domestica</i>	Rosaceae	8	1.23
<i>Milicia excelsa</i>	Moraceae	4	0.61

### 3.2 Tree Growth Variable Measurement

Results of the tree growth variables showed that *Terminalia mantaly* (0.43m) had the lowest mean DBH while *Mangifera indica* (0.69m) had the highest mean DBH. On average, *Azadirachta indica* recorded the lowest crown length, crown diameter, total height values of 12.18m, 12.85m and 15.08 respectively (Table 23).

**Table 23: Descriptive Analysis of Tree Growth Parameters**

Parameters	Species	Min.	Max.	Mean	Std. Error	Std. Deviation
DBH (m)	<i>T. mantaly</i>	0.24	0.70	0.43	0.01	0.09
	<i>A. indica</i>	0.38	0.89	0.59	0.01	0.10
	<i>M. indica</i>	0.29	1.24	0.69	0.03	0.22
	<i>T. catappa</i>	0.14	0.97	0.50	0.03	0.21
	<b>Total</b>	<b>0.14</b>	<b>1.24</b>	<b>0.55</b>	<b>0.01</b>	<b>0.19</b>
Crown Length (m)	<i>T. mantaly</i>	5.18	20.04	13.25	0.35	2.88
	<i>A. indica</i>	5.98	19.00	12.18	0.34	2.79
	<i>M. indica</i>	5.98	29.80	14.09	0.54	4.42
	<i>T. catappa</i>	1.19	27.14	13.65	0.64	5.19
	<b>Total</b>	<b>1.19</b>	<b>29.80</b>	<b>13.29</b>	<b>0.25</b>	<b>4.00</b>
Crown Diameter (m)	<i>T. mantaly</i>	6.26	22.10	14.48	0.36	2.93
	<i>A. indica</i>	8.90	18.05	12.85	0.23	1.87
	<i>M. indica</i>	5.55	21.65	13.79	0.46	3.76
	<i>T. catappa</i>	5.55	29.90	16.33	0.62	5.02
	<b>Total</b>	<b>5.55</b>	<b>29.90</b>	<b>14.36</b>	<b>0.23</b>	<b>3.79</b>
Clear Bole Height (m)	<i>T. mantaly</i>	1.68	6.65	3.11	0.13	1.08
	<i>A. indica</i>	1.52	5.40	2.90	0.08	0.62
	<i>M. indica</i>	1.46	4.36	2.72	0.09	0.69
	<i>T. catappa</i>	1.52	6.77	2.88	0.12	0.94
	<b>Total</b>	<b>1.46</b>	<b>6.77</b>	<b>2.90</b>	<b>0.05</b>	<b>0.86</b>
Total Height (m)	<i>T. mantaly</i>	7.08	23.73	16.35	0.38	3.10
	<i>A. indica</i>	9.55	22.02	15.08	0.32	2.60
	<i>M. indica</i>	9.15	32.64	16.81	0.55	4.43
	<i>T. catappa</i>	3.81	28.97	16.52	0.65	5.28
	<b>Total</b>	<b>3.81</b>	<b>32.64</b>	<b>16.19</b>	<b>0.25</b>	<b>4.03</b>
Merchantable Height (m)	<i>T. mantaly</i>	2.50	12.20	6.25	0.24	1.99
	<i>A. indica</i>	3.72	12.05	6.42	0.21	1.74
	<i>M. indica</i>	2.44	19.52	7.17	0.33	2.67
	<i>T. catappa</i>	1.52	14.52	6.98	0.38	3.12
	<b>Total</b>	<b>1.52</b>	<b>19.52</b>	<b>6.71</b>	<b>0.15</b>	<b>2.46</b>
Basal Area (m <sup>2</sup> )	<i>T. mantaly</i>	0.05	0.39	0.15	0.01	0.07
	<i>A. indica</i>	0.11	0.62	0.28	0.01	0.09
	<i>M. indica</i>	0.06	1.21	0.42	0.03	0.26
	<i>T. catappa</i>	0.01	0.74	0.23	0.02	0.17
	<b>Total</b>	<b>0.01</b>	<b>1.21</b>	<b>0.27</b>	<b>0.01</b>	<b>0.19</b>

### Key

### 3.3 Tree Vigour Estimation


Live crown ratio was used to estimate tree vigour. The result shows that *Azadirachta indica* (79.98) had the lowest averaged live crown ratio while *Mangifera indica* (82.80) had the highest averaged value for live crown ratio. *Terminalia mantaly* and *Terminalia catappa* had almost similar averaged value of 80.80 and 80.78 respectively (table 3).

**Table 34: Descriptive Analysis of Estimated Tree Vigour (Live Crown Ratio)**

Parameters	Species	Min.	Max.	Mean	Std. Error	Std. Deviation
Live Crown Ratio	<i>T. mantaly</i>	65.99	91.27	80.80	0.72	5.86
	<i>A. indica</i>	61.09	90.85	79.98	0.76	6.20
	<i>M. indica</i>	64.55	91.62	82.80	0.74	6.03
	<i>T. catappa</i>	31.20	93.68	80.78	1.14	9.23
	<b>Total</b>	<b>31.20</b>	<b>93.68</b>	<b>81.09</b>	<b>0.43</b>	<b>7.01</b>

### 3.4 Evaluation of Relationship between Vigour Measure and Tree Growth Parameter

There is significant relationship between LCR and all tree growth parameters (Total height, Merchantable height, Crown diameter, Clear bole height, Crown length, Diameter at breast height and basal area) as they yielded a p-value less than 0.05. Positive correlation was observed in LCR relationship with Total height ( $r = 0.57$ ), Merchantable height ( $r = 0.34$ ), Crown diameter ( $r = 0.43$ ), Basal area ( $r = 0.39$ ), Crown length ( $r = 0.71$ ) and DBH ( $r = 0.40$ ) while negative correlation was observed only in LCR relationship with Clear bole height ( $-0.62$ ) (Fig 2).



**Fig. 2: 2 x 4 Scatter plot grid of Live Crown Ratio and Tree Growth Parameters showing trend Line and Correlation Coefficient**

### 3.5 Tree Vigour Predictive Model

Table 54 indicates various simple linear models developed with various growth parameters as independent and live crown ratio as the dependent. All models were statistically significant as they yielded a probability value less than 0.001. Among the simple linear regression model, having clear bole height as the predictor was the best model as it explained 53.2% ( $R^2 = 0.532$ ) of the variability in the independent variable with SEE of 4.399 and the model with least representation is observed in equation involving merchantable height which produced an  $R^2$  value of 0.067 and SEE of 6.209.

**Table 45: Tree Vigour Predictive Model using Simple linear models**

S/N	Model	$R^2$	SEE	Prob	Coefficient in equation
1	$Y=b_0+b_1X_1$	0.247	5.578	<0.001	$Y = 67.574 + 0.825(X_1)$
2	$Y=b_0+b_1X_2$	0.532	4.399	<0.001	$Y = 95.685 - 5.030(X_2)$
3	$Y=b_0+b_1X_3$	0.067	6.209	<0.001	$Y = 76.327 + 0.671(X_3)$
4	$Y=b_0+b_1X_4$	0.464	4.704	<0.001	$Y = 65.802 + 1.147(X_4)$
5	$Y=b_0+b_1X_5$	0.152	5.918	<0.001	$Y = 71.269 + 0.671(X_5)$
6	$Y=b_0+b_1X_6$	0.129	5.997	<0.001	$Y = 74.008 + 12.337(X_6)$
7	$Y=b_0+b_1X_7$	0.134	5.983	<0.001	$Y = 77.385 + 12.875(X_7)$

Y = Live crown ratio;  $X_1$  = Total height;  $X_2$  = Clear bole height;  $X_3$  = Merchantable height;  $X_4$  = Crown length;  $X_5$  = Crown diameter;  $X_6$  = Diameter at breast height;  $X_7$  = Basal area;  $b_0$  = intercept;  $b_1, b_2 \dots b_7$  = slope for respective independent variable; Significant at  $p \leq 0.05$

Result from Table 65 shows that in developing models using multiple predictors (growth parameters) to predict live crown ratio (LCR), the models were statistically significant at  $p \leq 0.05$ . All levels of combination of total height ( $X_1$ ), clear bole height ( $X_2$ ) and crown length ( $X_4$ ) produced same coefficient of determination of 0.934 while the combination of all independent variables (0.941) yielded the best result of  $R^2$ .

**Table 56: Tree Vigour Predictive Model using Multiple linear models**

S/N	Model	$R^2$	SEE	Prob	Coefficient in equation
1	$Y=b_0+b_1X_1+b_2X_2$	0.934	1.654	<0.001	$Y = 80.820 + 1.070(X_1) - 5.810(X_2)$
2	$Y=b_0+b_1X_1+b_2X_4$	0.934	1.653	<0.001	$Y = 80.822 - 4.740(X_1) + 5.810(X_4)$
3	$Y=b_0+b_1X_2+b_2X_4$	0.934	1.654	<0.001	$Y = 80.820 - 4.740(X_2) + 1.070(X_4)$
4	$Y=b_0+b_1X_1+b_2X_2+b_3X_4$	0.934	1.658	<0.001	$Y = 80.822 - 4.362(X_1) - 0.377(X_2) + 5.432(X_4)$
5	$Y=b_0+b_1X_1+b_2X_2+b_3X_3+b_4X_4+b_5X_5+b_6X_6+b_7X_7$	0.941	1.585	<0.001	$Y = 78.120 - 6.575X_1 + 1.911X_2 - 0.214X_3 + 7.740X_4 + 0.042X_5 + 8.448X_6 - 9.659X_7$

Y = Live crown ratio;  $X_1$  = Total height;  $X_2$  = Clear bole height;  $X_3$  = Merchantable height;  $X_4$  = Crown length;  $X_5$  = Crown diameter;  $X_6$  = Diameter at breast height;  $X_7$  = Basal area;  $b_0$  = intercept;  $b_1, b_2 \dots b_7$  = slope for respective independent variable.

Amongst the models in Table 67, semi log models developed using single predictors produced the least coefficient of determination and largest SEE ( $X_2 = 0.535, 0.056$ ;  $X_4 = 0.443, 0.061$ ). Multiple combination of all the independent variables was the best ( $R^2 = 0.923, SEE = 0.023$ ).

**Table 67: Tree Vigour Predictive Model using Semi log models**

S/N	Model	$R^2$	SEE	Prob	Coefficient in equation
1	$\ln Y=b_0+b_1X_2$	0.535	0.056	<0.001	$\ln Y = 4.580 - 0.065(X_2)$
2	$\ln Y=b_0+b_1X_4$	0.443	0.061	<0.001	$\ln Y = 4.201 + 0.014(X_4)$
3	$\ln Y=b_0+b_1X_1+b_2X_2$	0.917	0.024	<0.001	$\ln Y = 4.394 + 0.013(X_1) - 0.074(X_2)$
4	$\ln Y=b_0+b_1X_1+b_2X_4$	0.917	0.024	<0.001	$\ln Y = 4.394 - 0.061(X_1) + 0.074(X_4)$

5	$\ln Y = b_0 + b_1 X_2 + b_2 X_4$	0.917	0.024	<0.001	$\ln Y = 4.394 - 0.061(X_2) + 0.013(X_4)$
6	$\ln Y = b_0 + b_1 X_1 + b_2 X_2 + b_3 X_4$	0.917	0.024	<0.001	$\ln Y = 4.394 - 0.072(X_1) + 0.011(X_2) + 0.085(X_4)$

Y = Live crown ratio;  $X_1$  = Total height;  $X_2$  = Clear bole height;  $X_3$  = Merchantable height;  $X_4$  = Crown length;  $X_5$  = Crown diameter;  $X_6$  = Diameter at breast height;  $X_7$  = Basal area;  $b_0$  = intercept;  $b_1, b_2 \dots b_7$  = slope for respective independent variable.

Also in Table 78, double log models produced with single independent variables produced the least coefficient of determination and largest SEE ( $X_2 = 0.511, 0.058$ ;  $X_4 = 0.459, 0.061$ ) respectively. Multiple combination of the independent variables ( $X_1$  and  $X_4$ ;  $X_1, X_2$  and  $X_4$ ; and  $X_1, \dots, X_7$ ) produced absolute value of 1.000 for coefficient of determination and SEE less than 0.000

**Table 78: Tree Vigour Predictive Model using Double log models**

S/N	Model	$R^2$	SEE	Prob	Coefficient in equation
1	$\ln Y = b_0 + b_1 \ln X_2$	0.511	0.058	<0.001	$\ln Y = 4.594 - 0.197(\ln X_2)$
2	$\ln Y = b_0 + b_1 \ln X_4$	0.459	0.061	<0.001	$\ln Y = 3.909 + 0.190(\ln X_4)$
3	$\ln Y = b_0 + b_1 \ln X_1 + b_2 \ln X_2$	0.929	0.022	<0.001	$\ln Y = 4.025 + 0.222(\ln X_1) - 0.237(\ln X_2)$
4	$\ln Y = b_0 + b_1 \ln X_1 + b_2 \ln X_4$	1.000	0.0003	<0.001	$\ln Y = 4.605 - 1.000(\ln X_1) + 0.999(\ln X_4)$
5	$\ln Y = b_0 + b_1 \ln X_2 + b_2 \ln X_4$	0.953	0.018	<0.001	$\ln Y = 4.119 - 0.194(\ln X_2) + 0.186(\ln X_4)$

Y = Live crown ratio;  $X_1$  = Total height;  $X_2$  = Clear bole height;  $X_3$  = Merchantable height;  $X_4$  = Crown length;  $X_5$  = Crown diameter;  $X_6$  = Diameter at breast height;  $X_7$  = Basal area;  $b_0$  = intercept;  $b_1, b_2 \dots b_7$  = slope for respective independent variable.

### 3.6 Model validation

The results of the validation test of the selected models are presented in Table 89. There were no Non significant differences between the observed value and the predicted values of the selected simple, multiple, semi logarithm and double logarithm model. Hence, all models are therefore suitable for current and future prediction but double log model gave the best result with highest  $R^2$  and lowest SEE.

**Table 8: Model validation**

Model	MOV	MPV	P-Value	Remark
<b>Simple linear models</b> $Y = b_0 + b_1 X_2$	81.797	81.795	0.999	NS
<b>Multiple linear models</b> $Y = b_0 + b_1 X_1 + b_2 X_2 + b_3 X_4$	81.797	82.503	0.548	NS
<b>Semi log models</b> $\ln Y = b_0 + b_1 X_1 + b_2 X_2$	4.397	4.410	0.452	NS
<b>Double log models</b> $\ln Y = b_0 + b_1 \ln X_1 + b_2 \ln X_4$	4.397	4.397	0.999	NS



Y = Live crown ratio;  $X_1$  = Total height;  $X_2$  = Clear bole height;  $X_4$  = Crown length;  $b_0$  = intercept;  $b_1, b_2 \dots b_7$  = slope for respective independent variable; MOV = Mean observed value; MPV = Mean predicted value; significant at  $p \leq 0.05$ ; NS = Not significant

### 3.7 Discussion

The University being a place of learning attracts numerous people together which make it not to differ from other urban environment. The inventory of tree species in University of Port Harcourt Park shows the presence of different tree species in which *Mangifera indica*, *Terminalia mantaly*, *Azadirachta indica* and *Terminalia catappa* are most dominant. Sociologist, psychologist and the masses supports the notion that the quality of urban life rely greatly on the quantity and quality of green spaces close to it or around it. The university environment is rich with diverse tree species because of the benefit of trees in the environment [8]. *Terminalia catappa* and *Azadirachta indica* are among the most dominant species in the University of Port Harcourt park. Researcher have also observed these species in various urban environment [9,10]. Tree vigour classes as modified by [11] are low vigour for LCR < 0.30, moderate vigour for LCR= 0.30 - 0.50 and high vigour for LCR > 0.50. it was observed that the studied trees species in the University park had high vigour (79.98-82.80) which implies high stability and low risk or susceptibility to wind throw and breakage.

In this study effort was directed towards obtaining tree vigour prediction models. Assessing tree vigour gives an understanding of the structural stability of tree roots which are inaccessible and also create a guideline on the maximum and stable height of a tree. Live crown ratio has been used as an indicator of tree vigour [12, 13] and a useful variable for assessing forest health condition. Before the models were developed, correlation analysis was carried out to give an insight of the association between tree vigour (live crown ratio) and the growth variables. Positive correlation was observed in LCR relationship with tree total height, merchantable height, crown diameter, Basal area, crown length and diameter at breast height while negative correlation was observed only in LCR relationship with clear bole height. Similar report of both positive and negative correlation between LCR and growth variables was observed in the findings of [14]

Model fitting and evaluation are important parts of model building. Fitting of tree vigour models were based on the total data set. A number of different models were examined for predicting tree vigour using simple linear equation, multiple linear equation, semi-logarithm and double logarithm functions. All the models showed strong fit to tree vigour though double logarithm gave the best result. One unique independent variable that features in all the models is clear bole height. This proves that clear bole height is one of the factors contributing to the tree vigour.

The result of the test of significance of tree vigour sampled showed no significant difference ( $P=0.05$ ) between the observed value and the predicted values, hence all model are therefore suitable for current and future prediction. This study is in agreement with the study of [15], in which all models were found to be significant at ( $P=0.05$ ) using slenderness coefficient to assess the risk of *Tectona grandis* plantation. The selected models can be useful for prediction of tree vigour among trees within the range of data used in model development.

## 4. CONCLUSION

Tree vigour model was developed in order to assess the stability of most abundant trees in University of Port Harcourt Park. Based on the parameters measured, the result of modelling

tree vigour with growth parameters showed that live crown ratio is a useful tool in tree vigour estimation. The observation from the study can be useful information on stability of trees to wind throw. This information can help forest managers know the trees which have good vigour in relationship with its growth parameters and the right silvicultural treatment in order to prevent damages by windthrow.

## REFERENCES

1. Oyebade, B. A., Popo-ola, F. S. and Itam, I. S., "Perception of urban dwellers about the benefits and management of urban trees in Uyo metropolis, Akwa Ibom State, Nigeria". *ARNP Journal of Science and Technology*, Vol 3, pp 767-773, 2013.
2. Brang P., Schönenberger W., Frehner M., Schwitter R., Thormann J., and Wasser B., "Management of protection forests in the European Alps: an overview". *Forest Snow and Landscape Research*, vol 80, pp 23-44, 2006.
3. Petrokas R. 2008, "Growth vigor in Wych Elm (*Ulmus glabra* Huds.)", *Baltic Forestry*, vol 14, pp 204-215, 2008.
4. Nicoll B.C., Gardiner B.A., Rayner B., Peace A.J., "Anchorage of coniferous trees in relation to species, soil type, and rooting depth", *Canadian Journal of Forest Research*, vol 36, pp 1871-1883, 2006.
5. Sellier D, Brunet Y, Fourcaud T., "A numerical model of tree aerodynamic response to a turbulent airflow", *Forestry*, vol 81, pp 279-297, 2008.
6. Marzocchi, W., Garcia-Aristizabal, A., Gasparini, P., Mastellone, M. L. and Di Ruocco A., "Basic principles of multi-risk assessment: a case study in Italy", *Nat. Hazards*, vol 62, pp 551 -573, 2012.
7. Eludoyin, O.S., Oladele, A.T. and Iyanda, O.M., "Mapping and Assessment of Ethno-Medicinal Trees in Built Up Areas - University of Port Harcourt, Nigeria", *South-east European forestry*, vol 6, 129-140, 2015.
8. Oladele, A.T., Eguakun, F.S., Ugbaja, U.C., "Amenity Trees Diversity in Selected Tertiary Institutions within Port Harcourt Metropolis, Rivers State, Nigeria", *J. Appl. Sci. Environ. Manage.* vol 24, pp 2175-2181, 2020.
9. Mulani R.M., and Khan. K. Z, "Study of Avenue Tree Diversity of Urban Area of Nanded District Maharashtra State India" *International Journal of Science and Research (IJSR)*, vol 6, 2154-2156, 2015.
10. Ogwu, M.C., Osawaru, M.E. and Obayuwana, O.K. "Diversity and abundance of tree species in the University of Benin City, Nigeria" *Applied Tropical Agriculture*, vol 21, pp 46-54, 2016.
11. Schultz, P.W, "The Structure of Environmental Concern: Concern for Self, Other People and the Biosphere" *Journal of Environmental Psychology*, vol 21, pp 327-339, 2001.
12. Hasenauer H. and Monserved R. A., "A crown ratio model for Australia forest", *Forest ecology and management*, vol 84, pp 49-60, 1996.
13. Popoola, F.S. and Adesoye, P.O, "Crown Ratio Models for *Tectona grandis* (Linn. f) Stands in Osho Forest Reserve, Oyo State, Nigeria", *Journal of Forest Science*, vol 28, 63-67, 2012.

14. Chukwu, O., Ezenwenyi, J.U., “Models for estimating crown projection area from stump diameter for *Tectona grandis* Linn. f. in the tropical rainforests of Nigeria”, *Forestist* 8 July 2020. 10.5152/ forestist.2020.

15. Eguakun, F. S., Oyebade, B. A. and Olayiwola, V. A, “Simulation study on risk assessment of *Tectona grandis* plantation in osho forest reserve, Oyo State, Nigeria”, *J. of Trop. For. Res.*, vol 30, pp 119-126, 2014.

**Comment [G2]:**

- ✓Make sure that the references in the text should be exists in the references
- ✓Please follow the guide line and instructions of Asian Journal of Environment & Ecology

UNDER PEER REVIEW