

Original Research Article

Increasing Growth, EPA, and HSI in Biofloc Catfish with the Addition of Soybean Oil

ABSTRACT

This study aims to determine the concentration of soybean oil added to feed to increase levels of omega-3 fatty acids EPA, *Hepatosomatic Index* (HSI), *Specific Growth Rates* (SGR), and *Average Daily Growth* (ADG). This research method uses a Complete Randomized Design (CRD) consisting of 4 treatments and 4 tests. The treatment applied is the difference in the concentration of soybean oil in the feed, namely treatments A (0%); B (2%); C (4%); and D (6%). The test fish used was catfish (*Clarias sp*) from Bandung Regency, West Java. Feed as much as 2.5% of the biomass weight is given 2 times a day, namely at 07 a.m and 16 p.m for 30 days in a biofloc pond. The results showed that the addition of soybean oil of as much as 4% to the feed was able to have the best influence on the increase in Omega-3 eicosapentaenoic acid/EPA fatty acids by 25 mg/100g with the highest Hepatosomatic Index (HSI) value of 1.52 ± 0.11 , the best Specific Growth rate (SGR) value of 1.1% and Average Daily Growth (ADG) value of 0.27%.

Comment [s1]: The research conclusions in the abstract have not been written yet

Keywords: Soybean Oil, Catfish, Eicosapentaenoic Acid, Specific Growth Rates, Average Daily Growth

1. INTRODUCTION

Soybeans are a type of legume plant that has high nutrition. Every 100 g of soy dry matter contains 35 g of protein, 35 g of carbohydrates, 18-20 g of fat as well as other nutritional content. Soybeans are usually processed as the basic ingredients of various products, one of which is soybean oil. The virtue of soybean oil is that it contains high protein levels compared to other types of nuts and contains high levels of unsaturated fatty acids and contains natural antioxidants [1]. Most of the fatty acid content of soy consists of essential fatty acids that are needed by the body. The main components contained in soybean oil are unsaturated fatty acids (85%) and saturated fatty acids (15%) [2].

Comment [s2]: Write down the complete nutritional content of soybeans along with the percentage value

Catfish is one of the most popular types of fish commodities in Indonesia that can be cultivated on limited land and water sources with high stocking density, the cultivation technology used is relatively easy to master by the community, with low business capital, easy marketing, and short cultivation time [3]. Another advantage of catfish is that it contains omega-3 fatty acids which are relatively higher compared to other types of freshwater fish but lower when compared to seawater fish [4].

Comment [s3]: Write down the protein content in soybean oil and other content levels along with the percentage value.

Omega-3 belongs to a type of polyunsaturated fatty acid that has multiple bonds. The first bond lies in the third carbon atom of the methyl omega group, the next bond lies in the third carbon atomic number of the previous double bond. Natural fatty acids include omega-3 fatty acids, one of which is *eicosapentaenoic acid* (C20: 5) or EPA [5].

Comment [s4]: Explain the type of catfish, because in the title catfish biofloc

EPA fatty acids are abundant in fish and foods that contain high fatty acids such as soybean oil. This fatty acid belongs to the LCPUFA (*Long Chain Polyunsaturated Fatty Acid*) group which has an important role in visual function, and brain development helps in the

formation of blood cells and the heart, nourishes the blood, and also helps good blood circulation [6]. EPA fatty acids are precursors of prostaglandin-3 (which inhibit platelet aggregation), thromboxane-3, and leukotin-5 groups (all eicosanoids). EPA fatty acids also serve as builders of most parts of the cerebral cortex of the brain and the normal growth of other organs [7]. In addition, according to [8], an important role of omega-3 fatty acids is that they can help improve memory for people with Alzheimer's.

The use of biofloc systems is considered effective and able to increase productivity. Because of a narrow biofloc pond, it can produce more fish. Biofloc technology is carried out by adding organic carbohydrates to the maintenance medium to increase the C/N ratio and stimulate the growth of heterotrophic bacteria that can assimilate inorganic nitrogen into bacterial biomass. Heterotrophic bacteria are believed to be able to convert ammonia - nitrogen discharge from a cultivation system into bacterial biomass that has the potential to be used as a source of feed for fish [9]. Thus, production costs can be reduced, especially in terms of feed. In addition, the time required is relatively shorter than using conventional cultivation systems [10].

The purpose of this study is to determine the effect of the addition of soybean oil in feed on the *eicosapentaenoic acid*/EPA fatty acid content and catfish growth (*Clarias sp*). the results of the study are expected to be able to provide information for farmers and the public about the important role of fat in feed to improve catfish quality by using biofloc systems.

2. METHODS

This research was conducted in August – September 2021 at the Inland Fisheries Area Laboratory, Ciparanje, Padjadjaran University. The test fish used was an enlarged catfish obtained from resident catfish farmers in the village of Cileunyi Kulon, Cileunyi District, Bandung Regency, West Java Province. The feed used is pellet-shaped artificial feed. The test feed contains different fats in each treatment. Feeding is carried out 2 times a day during the maintenance period of 30 days in a fiber tub pond with a size of 68x68 cm. Other adjuvants include probiotics, molasses, and adhesives.

The stage of making feed is to mix feed and soybean oil with the addition of progol as an adhesive, then stir until homogeneous and the feed is dried by aerating.

The study was conducted experimentally with a Complete Randomized Design (CRD) consisting of four treatments and four replays. The treatment used is the addition of soybean oil to the feed with the following concentrations.

- A: Control (without the addition of soybean oil)
- B: Feed with the addition of soybean oil 2%
- C: Feed with the addition of soybean oil 4%
- D: Feed with the addition of soybean oil 6%

Acclimatization in this study consists of acclimatization of cultivation media and test feed given during the maintenance period to assist in good growth and be able to survive during the maintenance process. The water medium is carried out to precipitate with the addition of aeration to remove particles or substances that can inhibit the sustainability of fish. During the acclimatization process, the test fish were given artificial feed mixed with soybean oil little by little with the aim that the fish would be able to receive the feed given during the study.

The data observed in this study are the content of omega-3 fatty acids EPA, Hepatosomatic Index (HSI), Specific Growth Rates (SGR), and Average Daily Growth (ADG).

2.1 Omega-3 EPA

Comment [s5]: Add references to research results on biofloc systems, increasing Growth, EPA, and HSI in Biofloc

Comment [s6]: The artificial feed has not been written clearly, what artificial feed consists of ingredients

EPA's omega-3 fatty acids were tested using the gas chromatography method at the Saraswanti Indo Genetech Laboratory and analyzed descriptively.

Comment [s7]: Reference not written yet

2.2 Hepatosomatic Index (HSI)

According to [11], the value of the *hepatosomatic index* (HSI) can be calculated using the following formula.

$$HSI = \frac{W_h}{W_t - W_h} \times 100$$

Information:

HSI = Hepato Somatic Index
W_h = Heavy heart (g)
W_t = Weight (g)

2.3 Specific Growth Rates (SGR)

To see the growth of fish specifically, the formula used to calculate *specific growth rates* (SGR) according to [12] is.

$$SGR = \frac{\ln W_t - \ln W_o}{t} \times 100\%$$

Information:

W_t = Average weight of fish at the end of maintenance (g)
W_o = Average weight of fish at the beginning of maintenance (g)
t = Length of maintenance time

2.4 Average Daily Growth (ADG)

The daily growth rate using the formula according to [13] is.

$$ADG = \frac{ABW_2 - ABW_1}{T}$$

Information:

ABW₂ = Average weight of the second or subsequent sampling (g/fish)
ABW₁ = Average weight of the first or previous sampling (g/fish)
T = Sampling time distance (days)

3. RESULTS AND DISCUSSION

3.1 Omega-3 EPA

Result of omega-3 eicosapentaenoic acid analysis is shown in table below.

Table 1. EPA fatty acid in catfish

Treatment	EPA fatty acid (mg/100g)
Without adding soybean oil	16.4
With added soybean oil 4%	25

Note: The result of chromatography gas test by Saraswanti Indo Genetech

Comment [s8]: Why is the treatment added 4% soybean oil? in the method using 2% soybean oil [B], 4% soybean oil [C], and 6% soybean oil [D], all treatments were 4

Table 1 above shows that feed with the addition of 4% soybean oil and without addition produces different omega-3 levels due to the addition of different soybean oil to the feed. The treatment without the addition of soybean oil only obtained omega-3 fatty acid S EPA 16.4 mg / 100g, while the addition of soybean oil as much as 4% was able to produce a higher value of omega-3 fatty acids EPA of 25 mg / 100g. The yield of EPA fatty acid content in the addition of soybean oil is 4% increased when compared to without the addition of

soybean oil. This is in line with the research of [14] showing that the application of soybean oil to feed produces a relatively high EPA of 5.05% increased from a concentration of 5% to 10%. Meanwhile, the research [15] on the addition of feed with 9% squid oil and 3% soybean oil was able to produce a proportion of EPA fatty acids of 5.10%.

The difference in the EPA fatty acid content of catfish is thought to be due to the composition of fats and fatty acids in fish depending on the type of food consumed by fish [16]. The nutritional balance of catfish feed is very important in supporting the growth and quality of catfish. Fatty acid nutrients from soybean oil are used as additives that play a role in growth. Lack or deficiency of fatty acids in fish can lead to disruption of growth function, feed efficiency till death [17] .

Increased levels of omega-3 fatty acids EPA catfish can provide benefits for those who consume catfish. These omega-3s play a role in vision function, connecting most of the cerebral cortex of the brain and other normal growth [18] as well as helping in the formation of blood cells and the heart, nourishing the blood, and also helping with good blood circulation [6]. Lack of omega-3 fatty acids can inhibit the process of growth, brain development, nerve cell problems, and decrease vision function [19].

3.2 Hepatosomatic Index (HSI)

The results in table 2 showed that the treatment given to soybean oil in catfish feed showed a noticeable difference ($P < 0.05$) with the highest HSI value of 1.52 ± 0.11 in the C treatment (4%). This is following [20] on the addition of soybean substitutions, which influences HSI values ranging from 1.06 ± 0.01 to 1.44 ± 0.01 .

Table 2. Hepatosomatic Index (%)

Component	HSI			
	A (0%)	B (2%)	C (4%)	D (6%)
HSI	1.15 ± 0.15^a	$1.42 \pm 0.21^{\text{from}}$	1.52 ± 0.11^b	$1.41 \pm 0.20^{\text{from}}$

Note: a.b = similar letter notation means that there is no noticeable difference in the level of the test Duncan has a value of 5%.

An increase in hepatosomatic values shows that there is glycogen and fat storage in the liver and muscles as a source of energy. High hepatosomatic values indicate the amount of fat content and fatty acids that can be used as a source of energy to support their growth. Low hepatosomatic values indicate the ability of fish to utilize poor feed, and low nutrient balance so that protein is used as an energy source and causes the growth rate to be slow [21].

According to [22], an increase in HSI values indicates an increase in the amount of nutrients absorbed and causes the amount of accumulated nutrients in the liver to increase as well. According to [23], the liver is one of the places where fat is stored. The liver also has a big role in the occurrence of metabolism and fat synthesis. Fat stored in the liver will cause an increase in size in the liver. The amount of fat content in the fish liver is thought to be directly proportional to the amount of fatty acids contained in the fish.

Comment [s9]: Too little discussion, add more recent references for discussion.

3.3 Specific Growth Rates (SGR)

Growth is defined as a change in fish in units of weight, size, or volume over time. The specific growth rate is calculated to find out the weekly growth of fish expressed in percentages (%).




Fig 1. Specific Growth Rates (SGR) in Catfish

In figure 1 above, it shows that the application of 4% soybean oil to catfish feed shows the highest growth rate value of 1.1%, and the application of 2% soybean oil shows a growth rate value of 1.05%, without giving soybean oil or 0% shows a growth rate value of 1.04% and giving soybean oil 6% shows a growth rate value of 0.9%. The decrease in the growth rate value in the D treatment (6%) is suspected to be because the added soybean oil causes a decrease in the quality value of the feed and the nutritional content becomes unbalanced for catfish.

In contrast to [24] using 4% fat supplementation consisting of a mixture of 2% coconut oil and 2% pecan oil in tilapia fry, produced a higher growth rate value of 2.09%, as well as [20] study on the addition of soybean substitution in barley fish feed, showed the highest growth rate value of 2.60%.

According to [25], The use of oil as a source of fat in fish feed is very important to support fish growth. The use of oil in feed can act as a substitute for protein as an energy source so that energy use can be saved and protein can be absorbed to be optimally utilized in growth. The high-fat content due to the addition of soybean oil causes the activity of lipogenic enzymes to decrease, thus inhibiting the synthesis of fatty acids [26] and can increase the chances of fat peroxidase and affecting the artibut sensors in muscles [27-28] .

Fat peroxidation is thought to reduce the efficiency value of feed and HSI (treatment D). This proves that catfish are able to take advantage of the addition of fat from soybean oil up to the limit of 4%. Existence of this decrease is suspected that the feed given has an unbalanced nutritional value. According to [29], the composition of feed with high-fat content will affect the activity of the enzymes amylase, lipase, and protease which can cause an increase in the occurrence of fat peroxidase resulting in a low growth rate.

3.3 Average Daily Growth (ADG)

Based on the ADG chart (Figure 2) catfish growth during maintenance fluctuates at each treatment. This is thought to be due to competition for feed which causes fish growth to increase and decrease significantly. The average daily growth ranges from 0.21 to 0.27%. The highest value was found in the C (4%) treatment of 0.27% and the smallest value was found in the D (6%) treatment of 0.21%. This is different from the research of [30] shows an ADG value of 0.9% for catfish in biofloc ponds.

There is a decrease in the average daily growth due to the use of protein feed in catfish magnification size allocated for metabolic processes, breeding, energy sources, growth, reproduction, and activity [32].




Fig 2. Average Daily Growth in Catfish

The use of feed with the addition of soybean oil can increase growth at a dosage of up to 4%. Catfish need fat with a content ranging from 4-5% of the weight of the feed. The use of excess fat can reduce daily feed consumption and reduce fish growth [32]. The maximum use of fat in the feed can encourage fish to grow faster and produce a *protein-sparing effect*. The *protein sparing effect* can balance the use of fats and carbohydrates for metabolic activities so that the protein contained in fish can be used for growth [33].

Comment [s10]: Too little discussion, add more recent references for discussion.

4. CONCLUSION

Based on this study, it can be concluded that the addition of 4% soybean oil to the optimum feed in increasing the EPA omega-3 value by 25 mg/100g, the highest HSI value of 1.52 ± 0.11 , the best SGR value of 1.1% and the ADG value of 0.27%.

REFERENCES

Comment [s11]: Check writing references

1. Isa I. Determination of Linoleic and Linolenic Fatty Acids in Soybean Oil by Gas Chromatography. Saintek, 2011:1–6.
2. Puspita YES. The Process of Taking Soybean Oil (Glycine Max) Using A Hydraulic Press With Variable Preheating Temperature And Pressing Pressure.;2016.
3. DPUB. Biofloc System Catfish Cultivation. Directorate of Production and Cultivation Business; 2017.
4. Suryaningrum. Optimizing the Utilization of Dumbo Catfish (*Clarias gariepinus*) to Support Food Security and Aquaculture; 2015.
5. Greatalya LAD. Omega 3 in Sardines as a Memory Enhancer. General Medicine Study Program, Faculty of Medicine, Sebelas Maret University, 2019:5.
6. Georgieff M. Nutrition and the developing brain: nutrient priorities and measurement. American Society for Nutrition; 2007.
7. Sumisih. Extraction of Omega-3 Fatty Acids from Fish Oil by-product of Canning Lemuru with Supercritical CO₂ Fluid Technology. Bogor Agricultural Institute; 2012.
8. Candidate F, Lim GP, Morihara T, Yang F, Ubeda O, Salem N, et al. Dietary n-3 polyunsaturated fatty acid depletion activates caspases and decreases NMDA receptors in the brain of a transgenic mouse model of Alzheimer's disease. *European Journal of Neuroscience*, 2005;22(3), 617–626. <https://doi.org/10.1111/j.1460-9568.2005.04253.x>
9. Wijaya M, Rostika R, Andriani Y. Effect of Different C/N Ratios on Biofloc Formation and Growth of Dumbo Catfish (*Clarias gariepinus*). *Journal of Marine Fisheries* 2016;8(1), 41–47.
10. Rusherlistyani, Sudaryati D, Heriningsih S. Cultivation of catfish with a biofloc pond system. LPPM UPN VY; 2017.
11. Effendie MI. Yogyakarta Fishery Biology. Nusantara Library Foundation; 1997.
12. Steffens W. Principle of fish nutrition. Ellis Horwood Limited; 1989.
13. Jensen G. Handbook for Common Calculation in Finfish Aquaculture. LSU AgCenter Research & Extension; 1992.
14. Salasah R, Nilawati MJ. Study of increasing omega-3 fatty acids EPA and DHA in catfish oil fed soybean oil. *Journal of Science Partners*, 2016;4, 1–12.
15. Marzuqi M, Rusdi I, Giri NA, Suwiry K. Effect of Proportion of Squid Oil and Soybean Oil as a Source of Fat in Feed on Juvenile Growth of Mangrove Crab (*Scylla Paramamosain*). Effect of Proportion of Squid Oil and Soybean Oil as a Source of Fat in Feed on Juvenile Growth of Mangrove Crab (*Scylla Paramamosain*), 2006;8(1), 101. <https://doi.org/10.22146/jfs.170>
16. Ackman R. Fatty acid composition of fish oil. In the Nutritional Evaluation of Long-Chain Fatty Acid In Fish Oil. Press Ltd; 1982.

17. Sumiyati. Effect of Giving Artemia Bioencapsulated Results With Omega-3 Fatty Acids HUFA and Vitamin C on Post Larvae Survival of Windu Shrimp (*Penaeus monodon* Fab.); 2003.
18. Felix. Current status of lipid nutrition white shrimp, *Litopenaeus vannamei*. Food Chemistry, 2002;96, 36–45.
19. Astawan M. Appropriate Animal Food Processing Technology. Presindo Academy; 1998.
20. Yanto H. Effect of Feeding with Different Levels of Fine Bran and Fermented Yellow Corn on Growth Performance of Jelawat Fish (*Leptobarbus hoevenii* Bleeker). Ruaya Journal, 2019;7(1), 8–16.
21. Booth A, Moses D, Allan L. Utilization of Carbohydrate by Yellowtail Kingfish, *Seriola lalandi*. Aquaculture, 2013(376-379):151–161.
22. Yandes Z, Affandi R, Mokoginta I. Effect of cellulose in feed on the biological condition of gouramy (*Ospronomus gourami* Lac) fry; 2003.
23. Deniel C. Les Poissons plats (Téléostéens, Pleuronectiformes) en baie de Douarnenez: reproduction, croissance et migration des Bothidae, Scopthalmidae, Pleuronectidae et Soleidae. Université de Bretagne occidentale-Brest; 1981.
24. Haetami K. Effectiveness of Fat in Formulation on Pellet Quality and Growth of Tilapia. Journal of Community Service, 2018;2(1), 6–11.
25. Masumoto T, Hosokawa H, Shimeno S. Protective Effect Treatment on Endothelial Function of Apolipoprotein E Deficient Mouse Carotid Artery. Nutrient Research, 1991;3(1), 103–108.
26. Wang JT, Liu YJ, Tian LX, Mai K Sen, Du ZY, Wang Y, et al. Effect of dietary lipid level on growth performance, lipid deposition, hepatic lipogenesis in juvenile cobia (*Rachycentron canadum*). Aquaculture, 2005;249(1–4), 439–447. <https://doi.org/10.1016/j.aquaculture.2005.04.038>
27. Bureau DP, Hua K, Harris AM. (2008). The effect of dietary lipid and long-chain n-3 PUFA levels on growth, energy utilization, carcass quality, and immune function of rainbow trout, *Oncorhynchus mykiss*. Journal of the World Aquaculture Society, 2008;39(1), 1–21. <https://doi.org/10.1111/j.1749-7345.2007.00146.x>
28. Jobling M. National Research Council (NRC): Nutrient requirements of fish and shrimp. Aquaculture International, 2012;20(3), 601–602. <https://doi.org/10.1007/s10499-011-9480-6>
29. Mudjiman A. Nutrient Research. Self-Help Disseminator; 2004.
30. Tasyah NN, Mulyono M, Farchan M, Thaib ASPEA. Land Optimization in Sweet Corn Intercropping System. Agroqua Journal, 2020;18(2), 168–174. <https://doi.org/10.32663/ja.v>
31. Niode AR, Nasriani N, Irdja AM. Growth and Survival of Tilapia (*Oreochromis Niloticus*) Seeds on Different Artificial Feeds. Academics: Scientific Journal of Media Publication of Science and Technology, 2017;6(2), 99–112. <https://doi.org/10.31314/akademika.v6i2.51>

32. Williams IH, Williams KC, Smith DM, Jones M. How do polka dot grouper *Cromileptes altivelis*, use dietary fat? Abstract Book (10th International Symposium on Nutrition and Feeding in Fish (ed.)); 2002.
33. Sanjayasari D, Kasprijo. Estimated Protein-Energy Ratio of Fish Feed Senggaringan (*Mystus nigriceps*) Nutritional Basis for Successful Domestication. Journal of Fisheries and Marine Affairs, 2010:2, 89–97.

UNDER PEER REVIEW