

Institutional Analysis of **add the** Bacteriological Profile and Antibiotic Susceptibility of **add the** Neonatal Sepsis in Eastern Nepal

Abstract

Introduction: In **add the** developing nations, neonatal sepsis account for the larger portion of **add the** neonatal morbidity and mortality. Blood culture is considered as the gold standard for the diagnosis. Both **add comma** the conformation and management is difficult for clinicians in **add the** resource limited nations like Nepal.

Method: It is a prospective observational study conducted at one of the tertiary care hospital in eastern Nepal over a year.

Result: Incidence of neonatal sepsis was 12.9%. The incidence of blood culture **positive(correct as positivity in the)** neonatal sepsis was 15%. Coagulase **Negative(small letter)** *Staphylococcus aureus*(**aureus is coagulase positive and write it as Staphylococcus species**) and *Escherichia coli* were two most predominant organism isolated. All of the isolated *Klebsiella* **add spp** and *Pseudomonas* **add spp** showed resistance to **Ampicillin(small letter)**(**ampicillin should not be tested against Pseudomonas and klebsiella as per CLSI guidelines as intrinsically resistant**) whereas **(???majority of)**Coagulase Negative *Staphylococcus aureus* **add spp** showed sensitivity to **Vancomycin.(small letter)**

Conclusion: **add isolated** Bacteria **are showing**(showed) resistance to multiple antibiotics. This is an alarming moment for the pediatrician. Antibiotics should be used judiciously and continuous

surveillance should be done to monitor the changing epidemiology of organisms and antibiotic sensitivity as the emergence of resistance to commonly used antibiotics is high.

Keywords

Neonatal sepsis, antibiotics, blood culture, resistance

Introduction

Septicemia in neonates refers to generalized bacterial infection documented by positive blood culture. It is a clinical syndrome characterized by the systemic sign of infections accomplice by bacteremia[1]. Septicemia in neonates can lead to sepsis. (rewrite the paragraph, avoid repetitions of words and mention the clinical definition as per standard guidelines) Delay in diagnosis and commencement of appropriate treatment may result in high morbidity and mortality rates. Changing bacterial flora and emergence of resistant strains leads to the problem thus neonatal septicemia requires accurate and timely clinical and laboratory diagnosis and proper management for a better outcome[2].

Some commonly found risk factors (mention as the most common risk factors) leading to neonatal sepsis are low birth weight, preterm delivery, maternal infection within two weeks of delivery, meconium stained liquor and premature rupture of membrane for more than twenty-four hours.

The presence of foul-smelling liquor or three of the abovementioned (space bet words) risk factors warrants initiation of antibiotic treatment. Infants with two risk factors should be investigated and then treated accordingly[3]. The incidence of sepsis among preterm neonates is six times greater than of full-term infants which can be attributed to the more immature immune system of preterm infants and their prolonged periods of hospitalization which increase the risk of nosocomial contracted infection[4].

According to Nepal Demographic and Health Survey 2016, the national neonatal mortality rate was 21/1000 live births. Infections including sepsis contributed to 16% of neonatal mortality[1]. Currently, the emergence of multidrug-resistant bacteria imposes challenges in the treatment of

neonatal sepsis[5]. The incidence is higher in late preterm than in term infants. The reported incidence of early and late-onset sepsis in late preterm neonates were 4.4 and 6.3 per 1000 respectively[6]. Though the exact incidence of early-onset neonatal sepsis in Nepal is not available, studies have shown that neonatal sepsis accounts for 17% of total neonatal admission and 5% of death in neonates. Neonates are immunocompromised and defend weakly against bacterial infections. Despite the **advance (add s)** **add the** in health care, neonatal sepsis is **add a** burden for **add the** developing countries. The bacterial agents implicated in early-onset sepsis include *Group B Streptococcus* (GBS), *Escherichia coli*, *coagulase-negative Staphylococcus* **add species(CONS)** , *Haemophilus influenza* (**add influenzae**), and *Listeria monocytogenes*. The organisms commonly associated with late-onset sepsis include *coagulase-negative staphylococci* (CONS), *Staphylococcus aureus*, *Klebsiella pneumonia*, *Escherichia coli*, *Enterobacter spp.*, *Pseudomonas aeruginosa*, and *Acinetobacter* species[7]. Clinical presentation of neonatal sepsis varies and there are no pathognomonic features[8]. Most observed clinical signs and symptoms are decreased feeding, lethargy, inactivity, fast breathing, hyperthermia or hypothermia, etc. Surveillance is needed to identify the common pathogens of the disease as well as the antibiotic susceptibility profile of the pathogens in a particular area. The objective of this study is to find the common bacteriological profile and antibiotic resistance for **add the** neonatal sepsis in eastern Nepal.

Methods

This study was done to determine the prevalence of neonatal septicemia, identify the bacterial isolates and study their antimicrobial susceptibility pattern in neonates admitted to the neonatal intensive care unit of Nobel Medical College and Hospital, Biratnagar, Nepal. This descriptive study was done over a one-year duration from (Jan2018 -Jan2019). Blood culture of all **add the**

neonates who were suspected of neonatal sepsis was performed. (were commercial blood culture bottles used and did u monitor by automated blood culture system) Bacterial isolation, identification, and antimicrobial susceptibility testing were done by the standard microbiological method.

All clinically suspected cases of neonatal sepsis admitted in NICU of Nobel medical college were enrolled in the study. Cases with major congenital malformation, birth weight <1Kg were excluded from the study. Ethical approval was taken from the Institutional Ethical Review Board of Nobel Medical College and Hospital, Biratnagar, Nepal.

Results

Total 1080 neonates admitted in NICU of Nobel medical college over one year of duration (January 2018 to January 2019) out of which 140 cases were neonatal sepsis were included in the study meeting the inclusion criteria. Out of 140 cases, 28(20%) cases were born outside the hospital and 112(80%) cases were born in the hospital maternity ward. Out of total cases, 35 (25%) were preterm and 105(75%) cases were term neonates. The most common clinical presentations in this study were tachypnea, lethargy, refusal to suck, and (???body /rectal)temperature instability.

In this study, out of 140 neonates, blood culture was positive in 21(15%) neonates and 119(85%) were culture negative.

Table 1: Frequency of growth in blood culture

Bacteria	Frequency	Percentage
Coagulase-negative <i>staphylococcus aureus</i> (s capital and not aureus .correct it as spp)	6	28

<i>Pseudomonas aerogenosa</i> spelling	3	14
<i>Klebsiella pneumoniae</i> spelling	3	14
<i>Escherichia coli</i>	4	19
<i>Staphylococcus aureus</i>	2	9
<i>Citrobacter freundii</i> space bet word	1	5
<i>Acinetobacter anitratus</i> spelling	1	5
<i>Enterococcus faecalis</i>	1	5
TOTAL	21	100%

Coagulase-negative *Staphylococcus aureus* add spp or mention as coagulase negative staphylococci (CONS) was found in 6(28%) as shown in table 1.

Above sentence Mention as CONS ACCOUNTED FOR majority of the infection followed by optional name of the other organisms with the % as shown in table no 1.....

Table 2: Antibiotic sensitivity pattern for CONS

Antibiotic	Resistance Frequency	Sensitive Frequency
Cloxacillin	4(67%)	2(33%)
Vancomycin	0	6(100%)
Amikacin	0	6(100%)
Linezolid	0	6(100%)
Teicoplanin	0	6(100%)
Cotrimoxazole	3(50%)	3(50%)
Clindamycin	4(67%)	2(33%)
Penicillin	6(100%)	0
Chloramphenicol	1(17%)	5(83%)

Among CONS, 67% were resistant to Cloxacillin, 100% were resistant to Penicillin, 50% were resistant to Cotrimoxazole, 67% were resistant to Clindamycin and 17% were resistant to Chloramphenicol. All were sensitive to Vancomycin, Amikacin, Linezolid, and Teicoplanin.

Was cefoxitin used to detect MRSA or MRSE or other spp? if detected, what was %?

Cloxacillin also interpret about methicillin resistance in Staphylococci from my point of view, in your study, 4 strains of CONS showed resistance to cloxacillin might be methicillin resistant staphylococci. which guideline was followed for AST (CLSI, BSAC OR EUCAST)

How sensitivity was done by MIC OR KB disc diffusion method?

Table 3: Antibiotic sensitivity pattern for *Pseudomonas*

Antibiotic	Resistance Frequency	Sensitive Frequency
Ampicillin	3(100%)	0
Cefotaxime	2(67%)	1(33%)
Ceftazidime	1(33%)	2(67%)
Ciprofloxacin	0	3(100%)
Gentamicin	1(33%)	2(67%)
Amikacin	0	3(100%)
Tazobactam-piperacillin spelling	0	3(100%)
Meropenem	1(33%)	2(67%)

Pseudomonas aeruginosa counted for 14% of all organisms detected of which 100% were resistant to Ampicillin, 67% were resistant to Cefotaxime, 67% were sensitive to Ceftazidime, 100% sensitive to Amikacin and Tazobactam-piperacillin, 67% sensitive to Meropenem

Table 4: Antibiotics sensitivity pattern of *Escherichia coli*

Antibiotic	Resistance Frequency	Sensitive Frequency
Ampicillin	3(75%)	1(25%)
Cefotaxime	2(50%)	2(50%)
Ceftazidime	1(25%)	3(75%)
Ciprofloxacin	3(75%)	1(25%)
Cefuroxime	3(75%)	1(25%)
Gentamicin	0	4(100%)
Amikacin	0	4(100%)
Tazobactam-pipperacillin	1(25%)	3(75%)

19% of the detected organism in blood culture was *E. Coli* of which 75% resistant to Ampicillin, 50% resistant to Cefotaxime, 75% resistant to Ciprofloxacin, and Cefuroxime, 75% were found sensitive to Ceftazidime and Tazobactum-piperacillin, 100% sensitive to Gentamicin and Amikacin.

The outcome in neonatal sepsis:

At the time of admission, a sepsis screen was done for all the neonates enrolled and empirical antibiotic therapy with Ampicillin and Amikacin were administered intravenously (as per hospital antibiotic policy). The neonates were periodically reviewed clinically and with laboratory results. Based on the organism grown in culture (add comma) the antibiotic regimen was changed according to the sensitivity pattern. In this study, 28(20%) neonates expired and 112(80%) neonates recovered.

Discussion

Follow one tense to discuss and compare (past tense better)

Sepsis remains one of the most important causes (correct it as cause) hospital admission for neonates with bad outcomes despite considerable progress in infection control, the introduction of

new antimicrobial agents, and advanced measures for early diagnosis and treatment[1]. Different hospitals have different bacteriological profile for neonatal sepsis. Similarly, the antibiotic resistance pattern also varies from institute to institute. Our institutional incidence of neonatal sepsis is 12.9% as per this study which is largely lower than the other tertiary care hospital (32%)[9].(in your city or which area)

The commonest clinical presentation in this study is tachypnea whereas another study found that the most common presenting clinical feature was respiratory distress followed by fever and feeding problems[10]. As compared to 15% of blood culture positive rate in this study,10.8% were noted in another study from Chitwan Medical College[11]. Also, in a study conducted in Nepal Medical College Teaching Hospital, Kathmandu, the positivity of the blood culture were 30.8%[12]. Coagulase Negative Staphylococcus aureus add spp (CONS) was seen as the most common (28%) growth after 72 hours where as 46.6% growth were seen in a study conducted in Chitwan Medical college in 2013 [1]. Coagulase Negative Staphylococci(staphylococci) were 17.4% where as Klebsiella(italics and add spp) were 16.11% in a tertiary care hospital in North India [4]. Regarding the antibiotics sensitivity pattern of CONS, 33% of isolates were found to be sensitive to Cloxacillin whereas,(remove comma) in a study conducted in Dhulikhel[12], 57.1% were found to be sensitive to Cloxacillin. In our study (add comma)all of the isolated CONS were found to be sensitive to Vancomycin which is similar to the result of a study conducted in South India[1]. Sensitivity to Amikacin was 100% (??? For all organisms)in this study and 88.9% in a study at Dhulikhel, Nepal[12]. All isolates of CONS were found to be 100% sensitive to Ampicillin, where as 40% were resistant to Ampicillin in a study at Dhulikhel [13]. Around 19% of *Escherichia coli* were isolated after 72 hours of growth. Bergin SP et al. showed 19.7% of blood culture-proven sepsis were *Escherichia coli*[14]. Another study conducted showed

Escherichia coli accounted for 15%[14]. Regarding sensitivity pattern, 14% of isolates were found sensitive to Ampicillin similar (16.6%) to other study [15] A study conducted in Paropakar Maternity and Women Hospital showed 42.85% were sensitive to Ampicillin [16].

In current study, 25% of isolates were found to be sensitive to Amikacin and 100% were found to be sensitive to Gentamicin whereas in a study conducted in Propakar Maternity and Women Hospital[16] 42.85% of isolates were found sensitive to both Amikacin and Gentamycin.(gentamicin) In another study in India, 33.3% and 16.6% of isolates were found sensitive to Amikacin and Gentamycin [15].(gentamicin)

After 72 hours of blood culture, *Klebsiella pneumonia* was isolated in 14%.

In a similar study, *Klebsiella* species accounted for 25% of isolates[17]. A study conducted at a tertiary hospital in Kathmandu, Nepal, *Klebsiella pneumonia*(italics) accounted for 20% of isolates[9]

Pseudomonas species(italics) accounted for 14% of total blood culture in this present study which is almost equal (13.4%) to a another study docnduted (spelling) at a tertiary hospital in Nepal[9].

In a study conducted in Peshawar, Pakistan(erase it) *Pseudomonas* (italics) was isolated in 13% of blood culture-proven sepsis[18][19]. In this study add comma all the cases were found to be resistant to amikacin, and all cases were found to be sensitive to Tazobactam-piperacillin and Ciprofloxacin and 37% were found to be sensitive to Cefotaxime, and 67% were sensitive to meropenem. In a similar study conducted in Rama Medical College, 100% sensitivity was seen with Meropenem and Amikacin, and 66.6%were found to be sensitive to Cefotaxime and tazobactam-piperacillin[20].

Isolated *Staphylococcus aureus* were 9% in the present study which is similar to a study where it was 18%.[21] About half (50%) of cases were sensitive to Cloxacillin which is lower than studies conducted at Dhulikhel, Nepal which showed a sensitivity of 91.7%[13]. Whereas similar rate (40%) of growth was seen in a study conducted in Bayelsa state, Nigeria showed [22]. *Enterococcus*, *Acinetobacter anitratus*(spelling), and *Citrobacter rfruenti* (spelling) have been isolated in 5% of cases of blood culture-proven sepsis. In a similar study conducted in Nigeria, 8% of isolates were *Enterococcus* species(italics) among blood culture-proven sepsis. In a study conducted in Chitwan Medical College, Bharatpur[1], *Acinetobacter* species (spp) accounted for 9.5% of blood culture positive.

Improper antenatal care and nutritional deficiencies (add comma) prematurity and low birth weight are (add some of the) common add contributing factors add of neonatal sepsis. So measures to improve maternal nutrition, health education,(remove comma) and prevention of prematurity would bring down the mortality due to neonatal septicemia. Knowledge of the most commonly isolated bacteria in neonatal intensive care units or nurseries, along with the antimicrobial susceptibilities of these organisms are valuable in treating suspected cases of neonatal septicemia.

Conclusion

The incidence of neonatal sepsis was 12.9% in this study which is similar to most of the studies conducted in other institutions of Nepal and southeast Countries. Blood culture proven sepsis were 15% which is less than most of the studies from different institutions of Nepal. Coagulase Negative *Staphylococcus aureus*(add spp) were the most common organism add s isolated

followed by *Escherichia coli*. Gram-negative growth rate was higher than gram-positive growth. All of isolated *Klebsiella* and *Pseudomonas* were found to be resistant to ampicillin(**intrinsically resistant, should not be tested**). Bacteria **are showing**(**showed**) resistance to multiple antibiotics. This is an alarming moment for the pediatrician.

References: please rewrite all the references page no, year of publication, volume, issue no missing

- [1] Ansari S, Nepal H, Gautam R, ... SS-I journal of, 2015 undefined. Neonatal septicemia in Nepal: early-onset versus late-onset. hindawi.com [Internet]. [cited 2022 Mar 7]; Available from: <https://www.hindawi.com/journals/ijpedi/2015/379806/>
- [2] Reddy KV, Sailaja K AA. Chapagain. Neonatal septicemia in Nepal: early-onset versus late-onset. .International Journal Pediatr. 2015;
- [3] Singh M, Narang A, Pediatrics OB-J of T, 1994 undefined. Predictive perinatal score in the diagnosis of neonatal sepsis. academic.oup.com [Internet]. [cited 2022 Mar 7]; Available from: <https://academic.oup.com/tropej/article-abstract/40/6/365/1641978>
- [4] Lamba M, Sharma R, Sharma D, Choudhary M, Maheshwari RK. Bacteriological spectrum and antimicrobial susceptibility pattern of neonatal septicaemia in a tertiary care hospital of North India. J Matern Neonatal Med. 2016 Dec 16;29(24):3993–8.
- [5] Yadav N, ... SS-B, 2018 undefined. Bacteriological profile of neonatal sepsis and antibiotic susceptibility pattern of isolates admitted at Kanti Children's Hospital, Kathmandu, Nepal. bmcresnotes.biomedcentral.com [Internet]. [cited 2022 Mar 7]; Available from: <https://bmcresnotes.biomedcentral.com/articles/10.1186/s13104-018-3394-6>
- [6] Cohen-Wolkowicz M, ... CM-TP, 2009 undefined. Early and late onset sepsis in late preterm infants. ncbi.nlm.nih.gov [Internet]. [cited 2022 Mar 7]; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2798577/>
- [7] Zakariya BP, Bhat V, Harish BN, Arun Babu T, Joseph NM. Neonatal sepsis in a tertiary care hospital in South India: Bacteriological profile and antibiotic sensitivity pattern. Indian J Pediatr. 2011 Apr;78(4):413–7.
- [8] Mhada T V., Fredrick F, Matee MI, Massawe A. Neonatal sepsis at Muhimbili National Hospital, Dar es Salaam, Tanzania; Aetiology, antimicrobial sensitivity pattern and clinical outcome. BMC Public Health. 2012;12(1).
- [9] Chapagain R, Acharya R, Council NS-... R, 2015 undefined. Bacteriological Profile of Neonatal Sepsis in Neonatal Intermediate Care Unit of Central Paediatric Referral Hospital

- in Nepal. europepmc.org [Internet]. [cited 2022 Mar 7]; Available from: <https://europepmc.org/article/med/27005713>
- [10] Lakhey A, Nepal HS-J of P of, 2017 undefined. Role of sepsis screening in early diagnosis of neonatal sepsis. nepjol.info [Internet]. 2017 [cited 2022 Mar 7];7:1103. Available from: <https://www.nepjol.info/index.php/JPN/article/view/16944>
 - [11] Thapa S, pediatrics LS-I journal of, 2019 undefined. Changing trend of neonatal septicemia and antibiotic susceptibility pattern of isolates in Nepal. hindawi.com [Internet]. [cited 2022 Mar 7]; Available from: <https://www.hindawi.com/journals/ijpedi/2019/3784529/>
 - [12] Raghubanshi BR, Sagili KD, Han WW, Shakya H, Shrestha P, Satyanarayana S, et al. Antimicrobial Resistance among Neonates with Bacterial Sepsis and Their Clinical Outcomes in a Tertiary Hospital in Kathmandu Valley, Nepal. Trop Med Infect Dis 2021, Vol 6, Page 56 [Internet]. 2021 Apr 20 [cited 2022 Mar 7];6(2):56. Available from: <https://www.mdpi.com/2414-6366/6/2/56/htm>
 - [13] Shrestha S, Shrestha N, ... SS-K university, 2013 undefined. Bacterial isolates and its antibiotic susceptibility pattern in NICU. nepjol.info [Internet]. [cited 2022 Mar 7]; Available from: <https://www.nepjol.info/index.php/kumj/article/view/11030>
 - [14] Bergin S, Thaden J, Ericson J, ... HC-TP, 2015 undefined. Neonatal Escherichia coli bloodstream infections: clinical outcomes and impact of initial antibiotic therapy. ncbi.nlm.nih.gov [Internet]. [cited 2022 Mar 7]; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4581845/>
 - [15] Vaniya H, Patel N, ... JA-IJ of, 2016 undefined. Antimicrobial culture sensitivity pattern in neonatal sepsis in a tertiary-care hospital. cabdirect.org [Internet]. [cited 2022 Mar 7]; Available from: <https://www.cabdirect.org/globalhealth/abstract/20163162887>
 - [16] Khanal R, Manandhar S, Acharya GP. Bacteriological Profile of Neonatal Sepsis in a Tertiary Level Hospital of Nepal. J Nepal Paediatr Soc [Internet]. 2014 [cited 2022 Mar 7];34(3):175–80. Available from: <https://www.nepjol.info/index.php/JNPS/article/view/9183>
 - [17] Zaidi A, Thaver D, ... SA-TP infectious, 2009 undefined. Pathogens associated with sepsis in newborns and young infants in developing countries. journals.lww.com [Internet]. [cited 2022 Mar 7]; Available from: https://journals.lww.com/pidj/Fulltext/2009/01001/A_prospective_study_of_neonatal_sepsis_and.3.aspx
 - [18] Rijal KR, Adhikari B, Ghimire B, Dhungel B, Pyakurel UR, Shah P, et al. Epidemiology of dengue virus infections in Nepal, 2006–2019. Infect Dis Poverty. 2021 Dec 1;10(1).
 - [19] Ullah O, Khan A, Ambreen A, ... IA-A of I, 2016 undefined. Antibiotic sensitivity pattern of bacterial isolates of neonatal septicemia in Peshawar, Pakistan. aimjournal.ir [Internet]. [cited 2022 Mar 7]; Available from: <http://aimjournal.ir/Article/1104>

- [20] “Neonatal sepsis”: bacteria & their susceptibility pattern towards antibiotics in neonatal intensive care unit. ncbi.nlm.nih.gov [Internet]. [cited 2022 Mar 7]; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3879858/>
- [21] Kalathia M, Shingala P, Neonatology PP-... C, 2013 undefined. Study of umbilical cord blood culture in diagnosis of early-onset sepsis among newborns with high-risk factors. ncbi.nlm.nih.gov [Internet]. [cited 2022 Mar 7]; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3883211/>
- [22] Peterside O, Pondei K, health FA-T medicine and, 2015 undefined. Bacteriological profile and antibiotic susceptibility pattern of neonatal sepsis at a teaching hospital in Bayelsa State, Nigeria. jstage.jst.go.jp [Internet]. [cited 2022 Mar 7]; Available from: https://www.jstage.jst.go.jp/article/tmh/advpub/0/advpub_2015-03/_article/-char/ja/