

The socio-economic importance and sustainability of Non-Timber Forest Products collection in the South West and Littoral Regions of Cameroon

Abstract

Interest in non-timber forest products (NTFPs) has increased due to their role in poverty reduction, conservation, and food security. However, data on the current environmental and socio-economic aspects of NTFPs harvests are sparse. Using participatory rural appraisal tools, information was gathered on NTFPs collectors, and on the relative socio-economic importance of NTFPs collected, collectors' access to NTFPs and sustainability of collection in major collection sites in the South West and Littoral Regions of Cameroon. An estimated 5500 collectors of NTFPs operate in the South West and Littoral Regions of Cameroon. The study revealed that NTFPs are important in the livelihoods of the village communities in the study area constituting 33% to their household income after agriculture (50.6%). The most important NTFPs collected in the study area were *Gnetum spp.*, *Irvingia sp.*, *Ricinodendron heudelotii* and *Piper guineensis*. Increasing harvests, combined with insufficient regulatory and customary control have led to a situation of long-term unsustainable collection. While NTFP collection is essential in providing income to collectors, their exploitation is failing to contribute in meeting environmental sustainability goals. This study concludes that domestication and awareness raising programs could lessen the pressure on the forest resource base and effective regulatory and customary control measures, if implemented and enforced, could limit over-exploitation and enhance sustainable collection and trade in NTFPs.

Key words: *Non-timber forest products, collectors, socio-economic importance, access, sustainability.*

Introduction

According to the World Bank (2006), around 60 million indigenous people are almost entirely dependent on forests. While, over 350 million people living in or near the world's tropical forests depend largely on this ecosystem. Forest resources are complementary to food production households; provide essential nutritional food and products for medicinal purposes. The forests of Central Africa and Cameroon are rich in non-timber forest products (NTFPs), which have long been an important component in the livelihood strategies of forest-dwelling people, providing subsistence needs, employment and cash income (Arnold and Ruiz-Pérez 2001). Most Cameroonians, particularly the rural inhabitants, depend on NTFPs for subsistence and cash income (Ingram, 2014). As a result, there has been increased interest in the collection and marketing of NTFPs as an instrument for sustainable rural development (Tieguhong and Ndoye, 2006). The importance of NTFPs from outside forests is attracting increasing attention, to help meet growing demands and reduce pressure on natural forests and plantations (Holding *et al.*, 2001). Indigenous people have developed their locality specific knowledge on NTFPs use, management, and conservation for the past centuries (Duguma & Mesele, 2019; Teklehaymanot, 2009; Yigezu *et al.*, 2014). The

subistence production of the rural population comprises fishery, agriculture, livestock husbandry, and the collection of forest resources.

Farmers' livelihoods and economic development is hampered by a low level of education, limited income alternatives and poor infrastructure, and the productivity of the cropland is limited by highly unpredictable rainfall and soil fertility constraints very similar to those encountered in the West African Sahel (Batiolo *et al.*, 1998, Buerkert *et al.*, 1998). Therefore, collection of NTFPs provides an important supplementary source of income (Sula, 2011) and an overuse of such resources threatens people's livelihood.

In the last 15 years, a large number of studies have sought to understand how social, economic, cultural, environmental, and geographical factors influence the traditional knowledge about plants at small scales. Factors such as gender, age, ethnicity, birthplace, and level of education have been identified as important on an individual level (Luoga *et al.*, 2000, Byg, 2004, Byg and Balslev 2006, Paniagua Zambrana *et al.*, 2007). Family size, integration into the market economy (e.g., sale of animals and agricultural products), or amount of material goods at the family level (e.g., possessions of farm animals, tools, and transport) have been linked to the household level (Byg and Balslev 2001, 2004, Reyes-García *et al.*, 2007). Access to commercial centers and to health, education, electricity, or water, as well as land tenure systems and settlement history, have shown a greater relevance at the community level (Takasaki *et al.*, 2001 and Vandebroek, 2010).

A few studies on the socio-economic characteristics of NTFPs collectors and their access to forest resources have been carried out in Cameroon covering many parts of the country but left out certain regions despite their richness in plant diversity (Adjanohoum *et al.*, 1996; Mbolo *et al.*, 2002). An example of such a region are the forested areas of Manyu, Ndian and Kupe-Muanenguba divisions of the South West region and the Mungo division in the Littoral region containing the most important protected areas and technical operation units in the area which are rich in plants and animal species. The consumption and sale of NTFPs can be important particularly for women, whose limited access to land, credit and other assets hinder their ability to pursue alternate livelihood opportunities (Hasalkar and Jadhav, 2004). Research has highlighted the role of gender in shaping access, management and use of forest resources and their associated benefits (Mai *et al.*, 2011, Ndumbe, 2013). Due to the high use of forest resources by the locals as food and income generation, the increasing anthropogenic activities which destroys the natural habitat of these plants calls for the urgent need to conduct a socio-economic survey in order to document the different NTFP species collected for food and income generation, determine their socio-economic importance to collectors, as well as access and sustainability of harvests in the South West and Littoral Regions of Cameroon.

Materials and Methods

Location of Study area

The study area as shown in Figure 1 covered four divisions in Cameroon: Manyu, Kupe-Muanenguba, Meme and Ndian, which were purposively sampled as important NTFP collection zones in the Southwest and one division, Mungo, in Littoral region was

purposively sampled. These were selected based on a situational analysis and rapid assessment that was carried in these regions prior to field work. In Manyu division all four sub-divisions were judged to be important in terms of NTFP collection. In Kupe Muanenguba, Nguti sub-division and in Ndian, the Bamusso sub-division were selected. Two villages were then selected in each sub-division based on their access to markets (easy and difficult).




Figure1: Map of study area

In the Mungo division, Bonalea, Dibombari and Mbanga sub-divisions were selected as productively important. As shown in Table 1, two villages were selected in each sub-division according to their accessibility to markets (ease of access - determined by distance, state of the roads and availability of transportation) with 50% sampled with 'easy' and 50% 'difficult' access. In each village, 25% of estimated NTFP collector population present (after a rapid survey of those present) was interviewed using a questionnaire.

Data collection and analysis

Data was collected on the socio-economic characteristics, NTFPs contribution to collector's household income, tenure, access and sustainability of harvest, using open-ended conversations and semi-structured questionnaires from January to April 2017. Respondents were selected randomly in the villages following age groups. The collected data was analyzed using Statistical Package for Social Sciences software version 20 (SPSS software V.20) using descriptive statistics.

UNDER PEER REVIEW

112 **Table 1: Sampled villages in Study Area**

Region	Division	Sub-division	Village	Access D- Difficult E= easy	Estimated collectors population	Estimated number present	Percentage interviewed of those present	Proportion population Interviewed
South West	Manyu	Akwaya	Bache	E	30	16	25	13
			Tapkwe	D	20	20	25	25
		Mamfe Central	Nchang	D	42	24	25	14
			Okoyong Native	E	25	12	25	12
		Eyumojock	Kembong	E	70	32	25	11
			Eyumojock	D	40	20	25	12
		Upper Bayang	Bachuo-akagbe	E	45	16	25	9
			Etoko	D	40	16	25	10
	Kupe- Muanenguba	Nguti	Ekenge	E	25	12	25	12
			Moungo-Ndor	D	14	8	25	14
	Ndian	Bamusso	Ekombe Liongo	E	20	12	25	15
			Mofako	D	15	8	25	13
	3	6	12	D=50% E=50%	386	196	25	13
Littoral	Mungo	Dibombari	Nkapa camp	E	30	16	25	13
			Bonamateke	D	18	16	25	22
		Bonalea	Souza	E	200	28	25	4
			Mbonjo II	D	30	20	25	17
		Mbanga	Mbanga	E	50	20	25	10
			Mojuka	D	45	16	25	9
	1	3	6	D=50% E=50%	373	116	25	8
Total	2	9	18	D=50% E=50%	759	312	25	10

113
114
115
116
117
118

Table 2: Population statistics of study area

Country	Region/State Division	Surface Area km ²	Population	Density per km ²	Capital	Ethnic groups
Cameroon	<i>Southwest</i>	24,571	838,042 ¹	34	Buea	Bakweri, Anyang,
	Ndian	6,626	129,659 ²	20	Mundemba	Ejaham, Balong,
	Kupe Manengouba	3,404	123,011	36	Bangem	Bassossi, Upper
	Manyu	9,565	177,389	19	Mamfe	Banyang, MboKorup, Isangele, Oroko
	<i>Littoral</i>	20,239	2,202,340	109	Douala	Bassa, Duala (Douala)
	Mungo	3,723	452,722 ²	122	Nkongsamba	Abo, Bankon

¹ 1987 Census, ² 2001 estimate: n "Departments of Cameroon". Statistics from Institut national de la statistique (Cameroun) - Annuaire statistique du Cameroun 2004. <http://www.statoids.com/ycm.html> (old data/2005). Retrieved April 6, 2009., ³ 2005 estimate from 1991 census

3. Results and Discussion

3.1. Socio-economic Characteristics of Collectors

An estimated 5500 collectors of NTFPs operate in the South West and Littoral Regions of Cameroon. The ages of 94.5% collectors were recorded with more than half of them aged above 41.5 years. However, the youngest NTFP collector sampled was 22 years old while the eldest was 70 years old. The average age for NTFP collectors was 43.66 years. Weighing by age group, collectors aged between 31 and 40 had the highest weight 30.8%, followed by collectors between the ages of 41 and 50 years with 25.0%. The age group with the least number of collectors was “61 years and above” making 9.6% of the sample. The majority of collectors in the South West are married. Women constitute the majority of NTFP collectors in the study area (79%) and the majority of them are married (65%). Manyu division had the highest average number of persons 6.47 ± 0.3 living in a NTFP collector’s household. Ndian division had the lowest figure 2 in terms of household size. A majority of collectors 53.6% had attained just primary education, while just 2.7% had attained tertiary education.

3.2. Collectors sources of household Income

The various sources of collectors’ household income are indicated on figure 2. The majority of collectors (50.6%) main source of household income is Agriculture (including market-gardening) followed by the harvesting and sale of NTFPs (33%).




Figure 2: Collectors sources of household income

3.3. Forest products collected and their contribution to collectors' household income

In the South West region thirteen NTFPs were named by collectors as being important in providing income and/or food in addition to other forest products. In Littoral, nine products were named. The most important NTFPs collected are *Gnetum spp.* (Eru), *Irvingia sp.* (bush mango), *Ricinodendron heudelotii* (njansang) and *Piper guineensis* (bush pepper) as indicated on Table 3.

Table 3: The types of forest products collected and their contribution to collectors' household income

NTFPs* % contribution to household income	Divisions and Sub-divisions											Total
	Manyu				Total Manyu	Kupe M	Ndian	Mungo			Total Mungo	Average
	Eyumoc k	Akway a	Upper Bayan g	Mamfe Central		Nguti	Bamuss o	Bonalea	Dibomba ri	Mbang a		
Gnetum (Eru)	33%	26%	19%	24%	26%	20%	22%	92%	57%	47%	63%	33%
Bush Mango	28%	27%	19%	21%	24%	20%	13%	8%	21%	18%	16%	18%
Njangsang	18%	6%	16%	24%	16%	12%	22%	0%	7%	12%	7%	14%
Bush Pepper	3%	12%	11%	8%	8%	20%	22%	0%	0%	6%	2%	13%
Bush Onion	-	9%	11%	8%	7%	12%					-	5%
Native Cola	3%	3%	0%	0%	1%	8%				6%	2%	3%
Hot leaf	8%	12%	19%	16%	14%					-	-	3%
Snails							13%			-	-	3%
Bitter Cola	3%	3%			1%		4%			6%	2%	2%
Monkey Cola			5%		1%	8%	-			-	-	2%
Mushrooms							4%			-	-	1%
Bush meat	3%				1%					6%	2%	1%
Cassa mango									7%	0%	2%	1%
Pepper									7%	0%	2%	1%
Eboya	3%	3%			1%			-	-	-	-	0%
TOTAL	33%	26%	19%	24%	26%	20%	22%	92%	57%	47%	63%	33%

*See Table for English, local and scientific names of the NTFPs

148

149

150

UNDER PEER REVIEW

151 **Table 4: Local and Scientific names of some identified NTFPs**

English name	Local name	Scientific name
Mushroom	Essok	Several species
Bitter Cola	<i>Bassa</i> : wè ; <i>Boulou</i> : onié ; <i>Douala</i> : ebongagnagne ; <i>Ejagham</i> : ejare, nya ; <i>Ewondo</i> : onié ; <i>Ibo</i> : adi ; <i>Pygmée Baka</i> : ngbwel.	<i>Garcinia kola</i> Heckel
Cola	Cola, Cola nut	<i>Cola nitida</i>
Eru	Eru (Efik); eru (Ibibio); ukasi (Igbo); ikokoh, (Ovande); gelu (Anyang); ecole (Boki)	<i>Gnetum africanum</i> and <i>Gnetum buchholzianum</i>
Bush mango	Bush mango (vern.); ogbono (Igbo); bojep (Boki); eloweh (Ovande); kelua (Basho); gluea (Anyang)	<i>Irvingia gabonensis</i> and <i>I. wombolu</i>
Njansang	Njansang (vern.); ngoku (Basho); itche (Becheve); ngoge (Boki); ngongeh (Anyang)	<i>Ricinodendron heudelottii</i>
Bush Onion	Felou (Basho); elongé (Becheve); eloweh (Ovande); elu (Anyang)	<i>Afrotyrax kamerunensis</i>
Bush Pepper	Kakwale (Ovande); iyeyeh (Becheve); ashoesie (Boki); taquale (Basho); acachat (Anyang)	<i>Piper guineensis</i>
Bush meat		Many species of duikers, antelopes, monkeys, wild pigs, rats, snakes, porcupines, cane rats etc.
Faux muscadier	<i>Douala</i> : pebé ; <i>Ewondo</i> : ding ; <i>Pygmée Baka</i> : dengo. <i>Bakoko</i> : gangat ; <i>Bassa</i> : ikoma; <i>Baya</i> : biko ; <i>Boulou</i> : ozek	<i>Monodora myristica</i> (Graertm.) Dunal
Monkey Cola	Monkey Cola	<i>Cola pachycarpa</i> K. Schum.
Snails		Several species

152 Sources: (Eyog Matig *et al.*, 2006; Sunderland *et al.*, 2003; Sunderland *et al.*, 1999)

154 3.4. Tenure and access

155 The situation of access in different forest categories in the South West and Littoral regions is
 156 indicated on table 5. The majority of collectors (75%) have free access to collection sites
 157 while 25% indicated that they do not have free access to community forests, private forests,
 158 national parks and private plantations for different reasons as indicated on Table 5. Under the
 159 1994 Forestry Law (94/01), all forested resources in Cameroon belongs to the state, with
 160 adjacent communities granted forest user rights for normal (not commercial) use of forest
 161 resources.

162 **Table 5: Tenure and access situation per region**

Tenure & Access	Response	South-West %	Littoral %	Total %
Are there areas where you are <u>not</u> allowed to collect eru?	No	75	100	88
	Yes	25		13
If yes, what type of areas?	Community forests	46	50	48
	Private forests	20		10
	National park	33		17
	Private plantations		50	25
Why are you not allowed to collect from these areas?	It is a protected area.			0
	Forests belonging to other village communities who do not allow non-indigenes to exploit.	56	30	43

	Forests/Plantations belonging to private individuals/companies who prohibit harvesters as some also steal crops.	44	69	57
--	--	----	----	----

3.5. Environmental sustainability

The various sites where collectors harvest NTFPs are indicated on Table 6. The majority of harvest is from non-protected areas while 2.9% of respondents collect from national park (protected area). The vast majority, 97% of the total collector's population sampled, responded that the distance travelled to collect NTFPs have increased in the past decade and that distances currently travelled are further than in the past. A large majority (97%) of respondents observed that the forest area around their village area has diminished, and only 2.7% had not yet observed any changes. 68% of respondents attributed this reduction to forest clearance for farmlands and 25% attributed it to the creation of palm plantations. In the Mungo division 7.4% attributed it to logging.

Table 6: Collection sites

Area of harvest		Percent
Protected Area	National Park	2.9
	<i>Sub-Total</i>	2.9
Non-protected Area	Open access forest (Primary and Secondary)	41.9
	Village forest (Primary and Secondary)	20.9
	Farmlands(Group and family)	5.7
	Private forest	17.3
	Palm plantations	11.3
	<i>Sub-Total</i>	97.1
TOTAL		100

The results suggest that the majority of NTFP collectors are youths being in their economically active stage that could drive productivity if supported within an enabling environment. Given that majority of collectors are married, these communities therefore can be more stable and suitable for a consolidated family unit. This offers an opportunity for stakeholders to easily integrate them into management programmes. The collection of NTFPs in the South West and Littoral Regions is an activity that involves adult individuals of which many have not received formal education. This may have a strong impact on introduction of innovations in sustainable harvesting techniques. Lack of education also suggest lack of ability to organise themselves into groups, cooperatives or organisations. Emphasis in this direction will improve the ability of the inhabitants to organise the marketing process of NTFPs to their advantage. The level of education sheds light on the ability to read and write, and how formalised trading is, whether written contracts are required or used for transactions between harvesters and traders (Ndoye 1995).

Collectors in the study area have varying strategies in generating income. Agriculture is their main livelihood activity contributing to 50.6% of household income followed very closely by NTFPs collection which contributes up to 33% of a collector's household income. This

finding matches studies of NTFPs incomes in Takamanda National Park (Sunderland *et al.* 2003; Tajoacha, 2008), Mone Forest Reserve (Mdaihli *et al.*, 2002; Tajoacha, 2008), Korup national park (Fuashi, 1997; Lingondo *et al.*, 2006; Malleson, 2001), Banyang Mbo Forest Reserves (Nkembi *et al.*, 2001; Nkembi, 2003) and Ejaham (Nkwatoh, 2000), and reinforces the importance of NTFPs in the livelihoods of the village communities in the study area.

In the South West region thirteen NTFPs were named by collectors as being important in providing income and/or food in addition to other forest products. In Littoral, nine products were named. The most important NTFPs collected are *Gnetum spp.* (Eru), *Irvingia sp.* (bush mango), *Ricinodendron heudelotii* (njansang) and *Piper guineensis* (bush pepper). This corroborates with the findings of Ingram *et al.*, 2012, Ingram *et. al.* 2016, and Ndumbe *et. al.*, 2018. For example, Ingram *et al.*, (2012) found that *Gnetum* contributes on average to 62% of a harvester's annual income (562, 500FCFA) in the South West and Littoral regions and that 2,324 tons of *Gnetum* was harvested in 18 villages in the South West Region from 2007-2009. The results also corroborate the findings of Ingram *et. al.* 2016 in which bush mango incomes contribute on average to 31% of harvester's annual income. The study revealed that an average annual quantity of bush mango harvested in the South West region was 113 tons while an estimated 4109 tons of bush mango was harvested annually in the period 2007 to 2010 in Southern Cameroon with estimated an estimated value of 1,175,121,208 FCFA while the market value was estimated at 4,801,062,134 FCFA based on average market prices. Similarly, Ndumbe *et al.*, 2018 found out that income from the sale of NTFPs contributes on average 19.8% for those collectors whose main activity is njansang collection, of which the average contribution of njansang was 10%. The quantity of njansang collected by collectors in the South West from 2013 to 2015 was 65.8 tons.

Under the 1994 Forestry Law (94/01), all forested resources in Cameroon belongs to the state, with adjacent communities granted forest user rights for normal (not commercial) use of forest resources. Many respondents in the study area however believe that the forests belong to them by right of inheritance. This is a common misconception in Cameroon (Oyono *et al.*, 2009). Customary rules of land tenure in forested village communities in Cameroon vary from one community to another according to the customs of the people regarding inheritance and who own land. Traditional authorities (village chiefs and councils) in many areas, such as Takamanda, are generally the custodians of forests (i.e lands that are forested and neither family forest or farms) and responsible for forest management in villages. In the communities in the study area, only men have the right to inherit land property as it is held that women are given land when they marry. Women however in most communities interviewed have free access to village and family forest and family farmlands. Women constitute the majority of NTFP collectors in the study area (79%) and the majority of them are married (65%). They are also free to harvest from and cultivate NTFPs on their husband's lands. But if the men (husbands of those married and in-laws of those who are widows) want to create cocoa farms and plantations on the land, women do not have any say.

The majority of respondents (89% in the Southwest and 96% in Littoral) indicated that they do not have to pay for entering the forest or harvest. If payment was required, this was to either the village traditional council in the southwest), or the forest owner (in Littoral). All respondents in Littoral indicated that there had been changes in forest access. One third indicated that local communities had not previously been aware of the economic importance

of some NTFPs and their forests, but now that they are aware, those who are not part of the community no longer have free access and payments were therefore demanded from the 'Forest owner'. The majority (70%) indicated however that forested land that is now being bought and owned by individuals, who prohibit harvester's free access. In the South West, 18% of all respondents indicated that there had been access changes. One third of these changes related to the establishment of protected areas and the restriction of rights in not being able to harvest freely from the National park. Two thirds of respondents who noted access changes indicated that with the increase in the knowledge of the value of some NTFPs, communities now restricted harvest from their own forests. This shows that if land tenure is defined and collectors 'own' or manage land, collection will be controlled.

NTFPs are collected most often from primary or secondary forest or in forest seen as belonging to village (although the majority of this is not legally classed as community forests) where access is free for everyone in the community. The second major source of NTFPs is from private forests and plantations. A small proportion is reported to be collected in protected areas and from farmlands. One third of the therefore harvest originates from privately held lands (farm, private forest or plantations). This differs slightly from CIFOR's 1997/1998 surveys when in the production zones of Lékié and zone Bassa, Mbanga (Centre), Souza (Littoral), Kumba and Mamfe (Southwest) respondents indicated the majority of NTFPs were harvested from long fallow areas, secondary forest, short fallows and primary forest (in that order). However, the focus of this survey was particularly the main collection areas in the Centre province. There are no tenure arrangements specific to NTFPs. Communities legal user rights to use NTFPs in their area for own consumption are however misappropriated with many individuals harvesting for commercial reasons. Collectors tend to act individually and independently, and rationally consulting their own self-interests. An indicator of unsustainable harvest of NTFPs is indicated increased distance to harvest. The vast majority, 97% of the total collector's population sampled, responded that the distance travelled to collect NTFPs have increased in the past decade and that distances currently travelled are further than in the past. This indicates that NTFPs are becoming scarcer and that the rates of harvesting are above the natural regeneration rates for many NTFPs. This matches the experiences in the Centre, East and Littoral regions (Awono *et al.*, 2002; Blackmore and Nkefor, 1998; Fondoun and Tiki-Manga, 2000), where increased NTFPs like *Gnetum spp.* has been found in secondary forest where it thrives after primary forest is disturbed, but is also highly prone to over harvesting with most of the methods used in collecting NTFPs observed to be unsustainable.

Conclusion

An estimated 5500 collectors of NTFPs operate in the South West and Littoral Regions of Cameroon. NTFPs are important in the livelihood's strategies of the rural communities in the South West and Littoral regions of Cameroon constituting 33% to their household income after agriculture (50.6%). The most important NTFPs collected in the study area were *Gnetum spp.*, *Irvingia sp.*, *Ricinodendron heudelotii* and *Piper guineensis*. Increasing harvest, combined with insufficient regulatory and customary control have led to a situation of long-term unsustainable collection. While NTFP collection is essential in providing income to

collectors, their exploitation is failing to contribute in meeting environmental sustainability goals. Findings suggests that distances travelled to collect NTFPs in the forest have increased. Recognising the socio-economic importance of NTFPs and insufficient controls to manage forest resources, domestication of economically important NTFPs on farms is a solution to not only increase production, but also reduces the pressure exerted on the forest resource base. Clarification of land tenure arrangements and the overlaps between unenforced and largely unknown official land tenure rules and customary rules could enhance the management and sustainability of NTFPs.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

References

References must be listed at the end of the manuscript and numbered in the order that they appear in the text. Every reference referred in the text must also present in the reference list and vice versa. In the text, citations should be indicated by the reference number in brackets [3].

World Bank (2006). Forests sourcebook : practical guidance for sustaining forests in development cooperation / World Bank. p. cm. – (Agriculture and rural development). ISBN 978-0-8213-7163-3 – ISBN 978-0-8213-7164-0 (electronic)

Arnold, M. and Ruiz-Pérez, M.(2001): “Can non-timber forest products match tropical forest conservation and development objectives?” *Ecological Economics* 39: 437 – 447.

Ingram, V. (2014). Win-wins in forest product value chains? How governance impacts the sustainability of livelihoods based on non-timber forest products from Cameroon. Dissertation, Faculty of Social and Behavioural Sciences, University of Amsterdam, Amsterdam.

Tieguhong, J.C. and Ndoeye, O. (2006). Africa-Escaping the Primary Commodities Dilemma. *African Development Perspective Yearbook* Vol. 11., chapter Transforming subsistence products to propellers of sustainable rural development: Non-timber forest products (NTFPs) production and trade in Cameroon, pages 107-137. VERLAG, Berlin, Germany.

Holding CP, Njuguna , Gatundu C (2001). Farm sourced timber: The restructuring of the timber industry in Kenya-opportunities and challenges. Forest Extension, IUFRO, Vienna, Austria.

Duguma, I. O., & Mesele, M. A. (2019). Use and management of medicinal plants by indigenous people in Boji Birmeji district, Western Ethiopia. *Ghana Journal of Science*, 60(1), 37–49. <https://doi.org/https://dx.doi.org/10.4314/gjs.v60i1.4>

Teklehaymanot, T. (2009). Ethnobotanical study of knowledge and medicinal plants use by the people in Dek Island in Ethiopia. *Journal of Ethnopharmacology*, 124(1), 69–78. <https://doi.org/10.1016/j.jep.2009.04.005>

- Yigezu, Y., Haile, D. B., & Ayen, W. Y. (2014). Ethnoveterinary medicines in four districts of Jimma zone, Ethiopia: Cross sectional survey for plant species and mode of use. *BMC Veterinary Research*, 10 (1), 76. <https://doi.org/10.1186/1746-6148-10-76>
- Bationo A, Lompo F, Koala S (1998). Research on nutrient flows and balances in West Africa: State-of-the-art. In Nutrient balances as indicators of production and sustainability in sub-Saharan African agriculture, Agriculture, Ecosystems and Environment, Volume 71. Edited by Smaling EMA; 1998:19–36.
- Buerkert A, Hiernaux P (1998). Nutrients in the West African Sudano-Sahelian zone: losses, transfers and role of external inputs. *J Plant Nutr Soil Sci* 1998, 161:365–383.
- SuLa Ma (2011). Sustainable Landmanagement in south-western Madagascar: Recherche participative pour appuyer la gestion durable des terres du Plateau Mahafaly dans le sud-ouest de Madagascar. In Diagnostic participatif de la gestion des ressources naturelles sur le plateau Mahafaly Commune Rurale de Beheloka, Toliara. Madagascar: Rapport Final. Project
- Luoga, E. J., E. T. F. Witkowski, and K. Balkwill. (2000). Differential utilization and ethnobotany of trees in Kitulanghalo Forest Reserve and surrounding communal lands, eastern Tanzania. *Economic Botany* 54(3):328–343. <http://dx.doi.org/10.1007/BF02864785>.
- Byg, A. (2004). Humans and plants of the rain forest: factors affecting local knowledge and use of plants. Dissertation. Department of Systematic, University of Aarhus, Denmark.
- Byg, A., and H. Balslev. (2006). Palms in indigenous and settler communities in southeastern Ecuador: farmers' perceptions and cultivation practices. *Agroforestry Systems* 67:147–158. <http://dx.doi.org/10.1007/s10457-005-1704-1>
- Paniagua Zambrana, N. Y., A. Byg, J.-C. Svenning, M. Moraes, C. Grandez, and H. Balslev. (2007). Diversity of palm uses in the western Amazon. *Biodiversity and Conservation* 16:2771–2787. <http://dx.doi.org/10.1007/s10531-007-9218-y>
- Byg, A., and H. Balslev. (2001). Diversity and use of palms in Zahamena, eastern Madagascar. *Biodiversity & Conservation* 10:951–970. <http://dx.doi.org/10.1023/A:1016640713643>
- Byg, A., and H. Balslev. (2004). Factors affecting local knowledge of palms in Nangaritza Valley in South-Eastern Ecuador. *Journal of Ethnobiology* 24(2):255–278.
- Reyes-García, V., N. Marti, T. McDade, S. Tanner, and V. Vadez. (2007). Concepts and methods in studies measuring individual ethnobotanical knowledge. *Journal of Ethnobiology* 27(2):182–203. [http://dx.doi.org/10.2993/02780771\(2007\)27\[182:CAMISM\] 2.0.CO;2](http://dx.doi.org/10.2993/02780771(2007)27[182:CAMISM] 2.0.CO;2)
- Takasaki, Y., B. L. Barham, and O. T. Coomes. (2001). Amazonian peasants, rain forest use and income generation: the role of wealth and geographical factors. *Society and Natural Resources* 14:291–308. <http://dx.doi.org/10.1080/08941920151080237>.
- Vandebroek, I. (2010). The dual intracultural and intercultural relationship between medicinal plant knowledge and consensus. *Economic Botany* 64(4):303–317. <http://dx.doi.org/10.1007/s12231-010-9135-y>
- Adjanohoun, J.E., Aboubaka, N., Dramane, K., Ebot, N.E., Ekpere, J.A. and Enow-Orock, E.G. (1996). Traditional medicine and pharmacopoeia. Contribution to ethnobotanical and floristic studies in Cameroon.

- Mbolo, M., Walter, S., Lejeune, J. (2002). La collecte et l'analyse des données statistiques sur les produits forestiers non ligneux. Une étude pilote au Cameroun. FAO, Rome (Italy). Dept. des Forêts.
- Hasalkar, S. and Jadhav, V. (2004). Role of women in the use of non-timber forest produce: A review. *Journal of Social Science* 8(3):203-206.
- Mai, Y.H., Mwangi, E., and Wan, M. (2011). 'Gender analysis in forestry: looking back and thinking ahead.' *International Forestry Review* 13(2): 1465-5489.
- Ndumbe, L.N. (2013). *Unshackling women traders: Cross-border trade of Eru from Cameroon to Nigeria*. Africa Trade Policy Notes.
- Eyog Matig, O., O. Ndoye, J. Kengue and A. Awono, Eds. (2006). Les fruitiers forestiers comestibles du Cameroun, International Plant Genetic Resources Institute.
- Sunderland, T. C. H., S. Besong and J. S. O. Ayeni (2003). Distribution, Utilization and Sustainability of Non-timber Forest Products from Takamanda Forest Reserve, Cameroon. SI/MAB Series. 8: Chapter 11: 155-172.
- Sunderland, T. C. H. and P. Tchouto (1999). A Participatory Survey and Inventory of Timber and Non-Timber Forest Products of the Mokoko River Forest Reserve, SW Province, Cameroon. A report for IR1/CARPE, African Rattan Research Programme and Mount Cameroon Project: 45
- Ndoye, O. (1995). The markets for non-timber forest products in the Humid Forest zone of Cameroon and its borders structure, conduct, performance and policy implications. Unpublished Report.
- Tajoacha, A. (2008). Market chain analysis of the main NTFPs in the Takamanda/Mone forest reserves, South West of Cameroon and the Cross River State of Nigeria. Dschang, Cameroon, University of Dschang. **DEA: 96.**
- Mdaihli, M., K. Schmidt-Soltan and J. S. O. Ayeni (2002). Socioeconomic Baseline Survey of the Villages in and around the Takamanda Forest Reserve. Profor, PROFOR: 54.
- Fuashi, N. A. (1997). Production and Marketing of NTFP's in the Korup Project Area Cameroon, and Cross Border Trade with Nigeria. International Workshop on the Domestic Market Potential of Non Tree Timber Products (NTTPs). Eyumojock, Cameroon, Limbe Botanical Garden, Cameroon.
- Ligondo, P.E. ; Atanga, W. ; Tsianhang, D. ; Fru, M. ; Choh, L. ; Meliko, O.M. (2006). A Rapid Market Appraisal on Eru in the Support Zone of the Korup National Park. CENDEP Report.
- Malleson, R. (2001). "Opportunities and Constraints for 'Community-based' forest management: findings from the Korup Forest, Southwest Province, Cameroon." Rural Development Forestry Network.
- Nkembi, L. and D. Hoyle (2001). The Non-Timber Forest Products Status *In* The Banyang-Mbo Wildlife Sanctuary: A Survey Of Household Use, Options For Adding Value And Economic Viability. For the WCS Banyang-Mbo Wildlife Sanctuary Project; The Ministry of Environment and Forestry (MINEF) Cameroon, and the Ministry of

Scientific Research (MINREST), Cameroon. N. T. W. C. Society. Cameroon, NYZS / THE WILDLIFE CONSERVATION SOCIETY: 89.

Nkembi, L. N. (2003). Participatory Forest Conservation and Sustainable Livelihoods: Banyang-Mbo Wildlife Sanctuary. WFC XII, Quebec, Canada, FAO.

Nkwatoh, A. F. (2000). Evaluation of Trade in Non-Timber Forest Products in the Ejagham Forest Reserve of Southwest Cameroon. Unpublished PhD Thesis.

Ingram, V., M.E. Ewane, L.N. Ndumbe and A. Awono (2016).. Challenges to governing sustainable forest food and landscapes: *Irvingia* spp. from southern Cameroon. *Forest Policy and Economics* 84 29–37.

Ingram, V., L.N. Ndumbe and M. E. Ewane (2012). Small Scale, High Value: The *Gnetum* spp. Value chains from Cameroon. *Small-scale Forestry*. DOI 10.1007/s11842-012-9200-8.

Ndumbe, L.N., V. Ingram, M. Tchamba and S. Nya (2018). From trees to money: the contribution of njansang (*Ricinodendron heudelotii*) products to value chain stakeholders' financial assets in the South West Region of Cameroon. *Forest, Trees and Livelihood Journal*. DOI:10.1080/14728028.2018.1559107.

Oyono, P. R., M. B. Biyong and S. Kombo (2009). Les Nouvelles Niches de Droits Forestiers Communautaires au Cameroun: Effets Cumulatifs sur les Moyens de Subsistance et Les Formes Locales de Vulnérabilité. RRI Project. Cifor. Yaoundé, CIFOR: 101

Awono, A., D. L. Ngono, O. Ndoeye, J. Tieguhong, A. Eyebe and M. T. Mahop (2002). Etude Sur La Commercialisation De Quatre Produits Forestiers Non-Ligneux Dans La Zone Forestiere Du Cameroun : *Gnetum* Spp., *Ricinodendron heudelotii*, *Irvingia Spp.*, *Prunus africana*. Fao. Yaounde, FAO : 96.

Blackmore, P. and J. T. Nkefor (1998). The Transfer of the Eru (*Gnetum africanum*, *G. buchholzianum*). Domestication Model to Village-Based Farmers on and around Mount Cameroon. L. B. Garden. Limbe: 8.

Fondoun, J. M. and T.Tiki-Manga (2000). "Farmers indigenous practices for conserving *Garcinia kola* and *Gnetum africanum* in Southern Cameroon." Agroforestry Systems 48: 289-302.