Original Research Article

Influence of COVID-19 Vaccination Coverage on Case Fatality Risk

Abstract

Background: It is well known that COVID-19 vaccines demonstrated higher efficacy against mortalities than mild acute respiratory syndrome coronavirus 2 (SARS-COV2). The estimation of the proportion of mortalities among these morbidities is a measure of case fatality risk (CFR). This study aims to evaluate change in CFR estimates among different countries after the introduction of COVID-19 vaccines and to shed light on the influence of the attack rate (AR) on CFR after the introduction of these vaccines.

Material and methods:

We collected publically available data concerning all countries/territories that implement COVID-19 vaccination at least for a hundred days ending on 3d of April 2021. They were sixteen in number. CFRs were measured as deaths per 100 COVID-19 confirmed cases; vaccine coverage was defined as the number of doses of vaccine per 100 people in the total population.

Descriptive data analyses were used including mean value, standard deviation, and graphical presentation by using Stem-Leaf charts and bar charts.

Inferential data analyses used included the One-Sample Kolmogorov-Smirnov (K-S) test and general linear model procedure (GLM).

Results: Findings showed that in a highly significant association the mean CFR decreased in countries with > 18 COVID-19 vaccine doses per 100 inhabitants. For the time period of the date of 1st day of vaccination till April 3, 2021, the total mean CFR is decreased with a surprising decrease in proportional total deaths and total cases, this decrement is more among total cases.

Conclusion: CFR monitoring may constitute a parameter for measuring vaccination effectiveness and progress of pandemic.

Comment [U1]: Rev. Note 1: No line

Keywords: COVID-19 disease; Vaccination; Case Fatality Risk

Introduction:

The ongoing global pandemic of coronavirus 2019 (COVID-19) was initially reported from Wuhan, China, in December 2019. After few weeks, it has been involved in several countries and became a significant public health problem ^{1,2,3} The rapid spread of this disease has caused substantial burden on morbidity with a variable case fatality risk (CFR). CFR is an important parameter to understand the epidemiological features of an outbreak or epidemic.^{4,5}

Late in 2020 COVID-19 vaccines become crucial tools in the pandemic response and protect against transmission of the disease, sever disease, and death.⁶ Dozens of countries now have advanced vaccination campaigns as they rush to protect their people and get their economies back up and running.

Measures used to monitor countries' vaccination progress included measuring daily or 7- day average decrease in the number of cases, measuring daily or 7- day average decrease in the number of deaths, and measuring the number of confirmed COVID-19 hospital admissions.⁷

COVID-19 vaccine has a substantial impact in reducing the incidence, hospitalizations, and deaths, especially among vulnerable individuals with comorbidities and risk factors associated with severe COVID-19. 8

Although numerous primary studies conducted before the implementation of COVID-19 mass vaccination programs reported variable (CFR)s of COVID-19 across different countries, information about (CFR)s is scares after the implementation of COVID-19 mass vaccinations.

Several factors were suggested to be associated with temporal and spatial variances in COVID-19 CFR. Among these factors comorbidity risk, demographic, socio-economic, and political variables, the age distribution of the community.⁹

Comment [U2]: Rev. Note 2: CFRs maybe?

Comment [U3]: Rev. Note 3: Please check for spelling errors in the whole text.

We conducted this study to look for the influence of COVID-19 vaccines on CFR in different countries and to shed light on the vaccine influence on disease transmission among different countries.

As a global real world study conducted early at mid of 1st week of April 2021 this study is an important one to evaluate the influence of COVID-19 vaccines at such early time of vaccine adminstration.

Material and methods:

We selected all countries / territories that implement COVID-19 vaccination for at least the last hundred days ending on 3d of April 2021. They were sixteen in number. Publically available data derived include total doses, vaccine doses / 100 people, total deaths, and accumulative COVID-19 cases. supplementary file contains this data.

CFRs were computed as total accumulative deaths divided by accumulative total cases x 100

Statistical Analysis: The statistical data analysis approaches were used with (SPSS) ver. (21).

- 1- Descriptive data analysis which included mean value, standard deviation, and Graphical presentation by using Stem-Leaf charts and bar charts.
- 2- Inferential data analyses: These were used to accept or reject the statistical hypotheses, which included the following:
- a- The One-Sample Kolmogorov-Smirnov (K-S) test.
- b- General linear model procedure (GLM).

Results and Findings:

Table (1): The general characteristics of the sample

Total number of countries	16	%
Total doses	385,795,246	

Comment [U4]: Rev. Note 4: No need to mention them as you don't present them.

Comment [U5]: Rev. Note 5: No need for the word Findings.

Comment [U6]: Rev. Note 6: Table 1 should be referred to in short in the text.

Comment [U7]: Rev. Note 7: It should not be here, but next to every number individually.

Total population	2,207,211,884	
Mean COVID-19 doses /100 inhabitant	17.478	
Total deaths 1 (at 1 st day of initiating	574,826	54.178
vaccination)		
Total deaths 2 (At 3 April 2021 including deaths at	1,060,983	100
1 st day of initiating vaccination)		
Total deaths 3(At 3 April 2021 excluding deaths at	486,157	45.82
1 st day of initiating vaccination)		
Total cases 1 (at 1st day of initiating	24,326,745	52.34
vaccination)		
Total cases 2 (At 3 April 2021 including cases at 1st	46,477,803	100
day of initiating vaccination)		
Total cases 3 (At 3 April 2021 excluding cases at $1^{\rm st}$	22,151,058	47.66
day of initiating vaccination)		
Mean CFR1 (at 1st day of initiating	2.362	-
vaccination)		
Mean CFR 2 (At 3 April 2021) total	2.283	-
Mean CFR3 (At 3 April 2021 excluding data at 1st	2.195	-
day of initiating vaccination)		
Mean change (difference) in magnitude of	-0.167	-
CFR (CFR3-CFR1)		
AR1 (at 1st day of initiating vaccination)	11.021	52.34
AR2 (At 3 April 2021)	21.057	100
AR3 (At 3 April 2021 excluding encountered cases	10.035	47.66
at 1st day of initiating vaccination)		
Change (difference) in magnitude of (AR3-	-0.986	-
AR1)		

Table 3 shows a total a highest initial CFR mean value than other values. The results also show that the lowest AR, the number of cases, and the number of deaths values were at 3 April 2021 excluding encountered cases at 1^{st} day of initiating vaccination.

Comment [U8]: Rev. Note 8: With two decimal points maybe uniformly.

Comment [U9]: Rev. Note 9: With two decimal points uniformly.

Comment [U10]: Rev. Note 10: It should be written after Table 2.

Table (2): Normal distribution function test due to different groups in relation to CFR marker

	One-Sample Kolm	ogorov-Smirnov Test		
Groups	Test Statistic	At 12:37pm CEST, 03/04/2021	At day 1 of starting vaccine	
	No.	9	7	
> 18 Doses /	Kolmogorov-Smirnov Z	0.506	0.921	
100 people	Asymp. Sig. (2-tailed)	0.96	0.364	
-	C.S. (*)	NS	NS	
	Test distribution of da	ta follows Normal Shape		
	No.	9	7	
≤ 18 Doses /	Kolmogorov-Smirnov Z	0.566	0.749	
100 people	Asymp. Sig. (2-tailed)	0.906	0.630	
_	C.S. (*)	NS	NS	
	Test distribution of da	ta follows Normal Shape		

(*) NS: Non Sig. at P>0.05

Comment [U12]: Rev. Note 12: (P≥0.05).

Comment [U11]: Rev. Note 11: C.S. should be explained by a note.

Table (2) shows the normal distribution function (goodness of fit test). It represents a one-sample "Kolmogorov-Smirnov" test procedure comparing the observed cumulative distribution function for studied readings with a specified theoretical distribution, which proposed a normal shape (i.e. bell shape).

The results show that the distribution of studied readings regarding CFR marker distribution function in relation to different locations. Since (P-value) is accounted at (P>0.05), this enabled us for applying the convention statistical methods (the parametrical methods).

Table (3): mean values, and standard deviation for the (CFR) marker, according to the assignable factors

Depender					
Countries according to COVID-19 vaccination doses	date	No.	Mean	Std. Deviation	95% confidence interval
> 18 doses / 100 people	At day 1 of starting vaccine (CFR1)	9	1.875	1.474	074202-3.00574

	At 03/04/2021(accumulative)	9	1.449	0.950	0. 71835-2.17898
	At day 1 of starting	7	3.315	2.796	072865-5.90077
≤ 18 doses / 100 people	vaccine (CFR1)				
= 10 doses / 100 people	At 03/04/2021 (7	7 2 202	2.500	0.7034586286
	accumulative)	,	3.283	2.789	

In table (3) results shows that mean CFR is less in countries with > 18 vaccine doses / 100 people compared to countries ≤ 18 vaccine doses / 100 people.

We found that countries and territories that have a level of coverage of > 18 doses/ 100 person showed decreased mean CFR compared to the countries' corresponding CFRs at the time of initiating the vaccine. The mean CFR was also decreased from 1.875 to 1.449. On the other hand, CFR for countries with a coverage rate of ≤ 18 doses per 100 inhabitants showed a lesser extent of decrease in mean CFR from 3.315 to 3.283.

Table (4): General linear model of fixed effects model with interaction for testing Marginal mean values for different Source of Variation in a compact form

Dependent variable CFR											
Source of Variation (SOV)	Type III Sum of Squares	d.f.	Mean Square	F	Sig.	C.S. (*)					
Intercept	193.788	1	193.788	45.91	0.000	HS					
Vaccine dose category/100 people	21.109	1	21.109	5.001	0.033	S					
Time starting the vaccine	0.412	1	0.412	0.098	0.757	NS					
Interaction	0.306	1	0.306	0.073	0.790	NS					
Error	53.71	28	4.221								
Total	2246.3	32									
	R - Squared = 0.1	R - Squared = 0.157									

^(*) HS: Highly Sig. at P<0.01; S: Sig. at P<0.05; NS: Non Sig. at P>0.05

Table (4) shows testing and analyzing the studied marker CFR with different sources of variation (SOV), such as the two different dose categories, countries starting vaccine time, interaction factor represented by applying the GLM of fixed effects model, and testing

effectiveness of the other source of variations which were not included in the studied model (i.e. the intercept). The R – Squared value was 0.157 which determines the proportion of variance in the dependent variable that can be explained by the independent variable .

Results show significant differences accounted at P<0.05 related to studied vaccine dose categories /100 people, while no significant differences at P>0.05 were accounted for both the time that countries starting the vaccine, and the interaction factor. In addition to that, the intercept (the other sources of variations not included in the studied model) recorded highly significant effectiveness at P<0.01.

Discussion:

Since CFR was significantly decreased within countries (as a function of number of COVID-19 doses per 100 population inhabitant) and decreased mean CFR, it is clear that that the decrease in deaths is proportionally more than the decrease in number of cases which is evident as a decrease in AR (tables 1,3, and 4). Our results show that 18 doses of COVID-19 vaccine/ 100 population inhabitant is the cut point for turning mean CFR value down. Usually CFR estimation errors or variances were largely related to testing coverage and detection of cases. In this study a decrease in CFR cannot explained by increase in denominator (cases) alone since increase in cases did not lead to proportional increase in nominator (deaths). Total deaths during 100-116 days since starting vaccination constitutes 45.8% of total deaths (since number of total deaths was estimated to be 486,157 during this period while total deaths were 1,060,983 for a period of more than one year). In contrast total cases during 100-116 days since starting vaccination constitutes 47.7%. Relatively higher total cases than total deaths 47.7 vs 45.8 shifts CFR value lower down.

We suggest her that the effect of COVID-19 vaccine on deaths outweighs its effect on cases. this leads to decrease in CFR. Vaccines provide at least some protection from infection and transmission, but not as much as the protection they provide against serious illness and death. This study gives evidence how vaccination limit infection and transmission on one hand and deaths on other hand.

Vaccination of certain share of population is essential for the reduction of epidemic transmission in a society and protection the unvaccinated individuals. ^{11,12}

Our findings support the findings of a positive association between the COVID-19 AR and CFR raised in recent literature. An increase in attack rate (AR) was suggested by these literature to be correlated to disease severity. The suggested hypothesis is that clustering of cases and viral overload lead to increased mortality rate and CFR. We think that vaccinations can inverse this phenomenon. It is clear that the relative reduction in mortality overcomes the relative reduction in morbidity. This might indicate that the AR has a role in mortality per se as stated in these literature.

In one study COVID-19 vaccination reduced the overall AR from 9.0% to 4.6% over 300 days, which constitute about a 50% reduction. Vaccination markedly reduced adverse outcomes, through decreasing non-intensive care unit (ICU) hospitalizations, ICU hospitalizations, and deaths. ⁸

It was suggested that an increase in fatality rate as the number of infected people increases is related to the overwhelming of the healthcare system. ^{9,16} This should be tested deeply as far as clusters of COVID-19 infections are associated with an increase in fatalities. ^{17,18}

Furthermore, although the number of hospital beds per 1000 people had a negative association with COVID mortality in certain countries including European countries, North America, Mexico, Brazil, Bolivia and USA, these findings were not global. The number of hospital beds per 1000 people did not have such a negative association in many Asian countries (excluding Japan) and in African countries. ¹⁹ They displayed comparatively low mortality regardless of their limited bed capacity. The controversy in these findings might be biased by a high AR in some countries which makes these beds insufficient. On the other hand low attack in other countries probably led to low CFRs regardless of the bed capacity.

Compatible with this study a 10% increase in vaccine coverage was observed with a 7.6% reduction in the CFR (95% confidence interval (CI = -12.6 to -2.7%, P = 0.002) according to a study which evaluated effectiveness of COVID -19 vaccine at third week of April 2021 (rather than mid of 1st week as in this study). ²⁰ Another compatible study which evaluated effectiveness of COVID-19 vaccine on AR as it was in 2 May 2021 showed that when the

accumulated vaccination rate reaches 1.46–50.91 doses per 100 people the infection of disease is reduced. .²¹ Adopted local strict measures of nonpharmaceutical interventions (NPIs) greatly affect AR in addition to vaccination coverage .²¹

This study shed a light on the importance of COVID-19 vaccination coverage in decreasing CFR, a missed parameter before in evaluation of the pandemic and effectiveness of COVID-19 vaccines. Although vaccines protect from severe disease²² which can decrease the CFR among vaccinated people, the finding of low CFR in relatively low vaccination coverage countries might give clue that CFR can be decrease by other mechanism i,e through lowering AR. Anyhow the low R – Squared value and the presence of a highly significant intercept calls for further studies to study the effect of other possible responsible factors in decreasing CFR.

The possible limitations in this study include: (1) The COVID-19 vaccine doses administered per 100 people may not equal the number of people that are vaccinated if the vaccine requires two doses, (2) Change in testing coverage within a country or across countries, (3) difficulty in estimating asymptomatic cases, (4) difficulty in estimation of actual COVID-19 deaths for a variety of reasons, (5) differed COVID-19 preventive approaches across countries and within the same country from time to time, (6) COVID-19 pandemic stage difference across countries, and (7) the contact-reducing interventions in place.

Conclusion:

It was concluded that countries with a higher dose of COVID-19 vaccine indexed as >18 doses /100 people reported the significantly associated lower (CFR)s on April 3,2021 than (CFR)s on day 1 of starting vaccine. Furthermore, data suggested that CFRs reduction is associated with concomitant reduction in (AR)s.

CFR estimate is a parameter for measuring vaccination effectiveness and progress of pandemic.

Comment [U13]: Rev. Note 13: seems to be

Data resources: We used publically available data. Patients were not involved.

Availability of data and materials: attached as supplementary file.

Comment [U14]: Rev. Note 14: No need to mention that in the final text.

References

https://doi.org/10.30491/JMM.22.1.9610.30491/JMM.22.1.96

¹ Nikpouraghdam M, Farahani AJ, Alishiri G, et al. Epidemiological characteristics of coronavirus disease 2019 (COVID-19) patients in IRAN: a single center study. *J Clin Virol* 2020;127:104378. https://doi.org/10.1016/j.jcv.2020.10437810.1016/j.jcv.2020.104378

² Wu JT, Leung K, Bushman M, Kishore N ,et al. Estimating clinical severity of COVID-19 from the transmission dynamics in Wuhan, China. *Nat Med* 2020;26:506-510. https://doi.org/10.1038/s41591-020-0822-710.1038/s41591-020-0822-7

³ Sun P, Lu X, Xu C, Sun W, Pan B. Understanding of COVID-19 based on current evidence. *J Med Virol* 2020;92:548-51. https://doi.org/10.1002/jmv.2572210.1002/jmv.25722

⁴ Wallinga J, Teunis P.Different epidemic curves for severe acute respiratory syndrome reveal similar impacts of control measures. Am J Epidemiol. 2004 Sep 15; 160(6):509-16.

⁵ Alimohamadi Y, Sepandi M: Basic reproduction number: an important indicator for the future of the COVID-19 epidemic in Iran. Journal Mil Med 2020;22:96-7.

⁶ Al-Amer, R., Maneze, D., Everett, B., Montayre, J., Villarosa, A. R., Dwekat, E., & Salamonson, Y. (2022). COVID-19 vaccination intention in the first year of the pandemic: A systematic review. Journal of Clinical Nursing, 31, 62–86. https://doi.org/10.1111/jocn.15951

⁷ CDC. Interpretive Summary for April 2, 2021. The Race to Vaccinate. https://www.cdc.gov/coronavirus/2019-ncov/covid-data/covidview/index.html accessed: April 5,2021

⁸ Moghadas SM ,Vilches TN , Zhang K ,etal . The impact of vaccination on COVID-19 outbreaks in the United States. Version 2. medRxiv. Preprint. 2020 Nov 30 [revised 2021 Jan 2]. doi: 10.1101/2020.11.27.20240051

⁹ Sorci G, Faivre B, Morand S. Explaining among-country variation in COVID-19 case fatality rate. *Sci Rep.* 2020;10(1):18909. Published 2020 Nov 3. doi:10.1038/s41598-020-75848-2

¹⁰ WHO.Vaccine efficacy, effectiveness and protection .14 July 2021

https://www.who.int/news-room/feature-stories/detail/vaccine-efficacy-effectiveness-and-protection

Lancet, 2020 Jun 13; 395(10240):1845-1854

SSRN: https://ssrn.com/abstract=3806123 or http://dx.doi.org/10.2139/ssrn.3806123

medRxiv 2020.12.22.20248696; doi: https://doi.org/10.1101/2020.12.22.20248696

¹¹ Fine P, Eames K, Heymann DL."Herd immunity": a rough guide.Clin Infect Dis. 2011 Apr 1; 52(7):911-6.

¹² Zhu FC, Li YH, Guan XH, et al. Safety, tolerability, and immunogenicity of a recombinant adenovirus type-5 vectored COVID-19 vaccine: a dose-escalation, open-label, non-randomised, first-in-human trial.

¹³ Raham TF. Covid-19 High Attack Rate Can Lead to High Case Fatality Rate. American J Epidemiol Public Health. 2021 April 27;5(2): 045-049. doi: 10.37871/ajeph.id49

¹⁴ Raham, TF. Epidemiological Philosophy of Pandemics (February 20, 2021). Available at SSRN: https://ssrn.com/abstract=3789738 or http://dx.doi.org/10.2139/ssrn.3789738

¹⁵ Al-Naqeeb A. A. A., Raham TF, Case Fatality Rate Components Based Scenarios for COVID-19 Lockdown (March 16, 2021). Available at

¹⁶ Ji Y, Ma Z, Peppelenbosch MP, Pan Q. Potential association between COVID-19 mortality and health-care resource availability.Lancet Glob Health. 2020 Apr; 8(4):e480

¹⁷ McMichael T.M,Currie D.W,Clark S, et al.Epidemiology of COVID-19 in a long-term care facility in King County, Washington.*N Engl J Med.* 2020; 382: 2005-2011

¹⁸ Hashan MR, Smoll N, King C, Ockenden-Muldoon H, Walker J, Wattiaux A, Graham J, Booy R, Khandaker G. Epidemiology and clinical features of COVID-19 outbreaks in aged care facilities: A systematic review and meta-analysis. EClinicalMedicine. 2021 Mar;33:100771. doi: 10.1016/j.eclinm.2021.100771. Epub 2021 Mar 1. PMID: 33681730; PMCID: PMC7917447.

¹⁹ Vineet Jain, Nusrat Nabi, Kailash Chandra, etal. A comparative analysis of COVID-19 mortality rate across the globe: An extensive analysis of the associated factors.

²⁰ Liang LL, Kuo HS, Ho HJ, Wu CY. COVID-19 vaccinations are associated with reduced fatality rates: Evidence from cross-county quasi-experiments. *J Glob Health*. 2021;11:05019. Published 2021 Jul 17. doi:10.7189/jogh.11.05019

²¹ Chen YT. The Effect of Vaccination Rates on the Infection of COVID-19 under the Vaccination Rate below the Herd Immunity Threshold. *Int J Environ Res Public Health*. 2021;18(14):7491. Published 2021 Jul 14. doi:10.3390/ijerph18147491

²² Adil Mahmoud Yousif N, Tsoungui Obama HCJ, Ngucho Mbeutchou YJ, Kwamou Ngaha SF, Kayanula L, Kamanga G, et al. (2021) The impact of COVID-19 vaccination campaigns accounting

for antibody-dependent enhancement. PLoS ONE 16(4): e0245417. doi:10.1371/journal.pone.0245417

Appendices

Appendix (1): references for data

- 1-WHO Coronavirus Disease (COVID-19) Dashboard With Vaccination Data | WHO Coronavirus (COVID-19) Dashboard With Vaccination Data
- 2- Coronavirus (COVID-19) Vaccinations Statistics and Research Our World in Data
- **3-**Covid-19 vaccine tracker: View vaccinations by country (cnn.com)
- 4- Information and public services for the Island of Jersey

Coronavirus (COVID-19) (gov.je)

Appendix (2)
Initial data including CFRs At 12:37pm CEST, 3 April 2021 and at At day 1 of starting vaccine

			Days		pm CEST,		At day 1 of		
			since	3 April 2	021		starting vaccine		
		Doses	first	Deaths	cases	CF	Death	Cases	CF
Location/		/ 100	dose			R	s		R
1000	Total	peopl	vaccine						
population	doses	e	: date						
Locations wi	Locations with > 18 Doses / 100 people								
				6,216	833,269		3069	368617	0.83
Israel	10,057,609	116	105			0.74			3
9,216.90	10,037,009	110	: 19/12			6			
al !!			100	23,421	1,011,485		16228	590914	2.74
Chile	10,780,609	56	100			2.31			6
19,116.21			24/12			6			

				,					
Jersey	59,132	59	111:	69	3,228	2.13	32	1637	1.95
108.809	55,152		13/12			8			4
United			446	126,764	4,350,270	2.91	61434	1737694	3.53
Kingdom	36,249,902	56	116:			4			5
67,215.29	, ,		8/12			-			
Guernsey			40=	14	821	1.70	13	291	4.46
-	33,400	50	107:	14	021		13	291	
63.385			17/12			5			7
Bahrain	5 02 5 20	46	107;	527	146,454	0.36	349	89600	0.38
1,701.58	782,530	46	17/12			0			9
United				547,884	30,238,69	1.81	29684	1586067	1.87
States	157,606,46	47	110:	,	2	2	0	5	1
	3	47	14/12			2	U	3	1
329,484.12									
Serbia	2,521,863	37	100:	5,345	605,406	0.88	2833	312253	0.90
6,908.22	2,321,003	37	24/12			3			7
Qatar			101:	298	181,678	0.16	243	142308	0.17
2,881.06	867,209	30	23/12			4			1
Locations ≤ 18	doses / 100 pe	ople							
Switzerland			101:	9,654	600,331	1.60	6723	423731	1.58
8,636.90	1,536,186	18	23/12			8			6
Canada			110:	23,002	987,918	2.32	13413	454851	2.95
38,005.24	5,968,907	17	14/12		,	8			2
Saudi Arabia			107:	6,684	391,325	1.70	6080	360353	1.68
34,813.87	4,722,340	14	17/12		ŕ	8			7
Mainland			17712	4,851	102,838	4.71	4758	95064	5.00
	133,801,00		100.	4,031	102,030		4730	73004	
China		9	109:			7			5
1,410,929.3	0		14/12						
6									
Russia	11,779,295	8	119:	99,633	4,563,056	2.18	42176	2402949	1.75
144,104.08	11,//9,495	0	4/12			3			5
Costa Rica	204 255	o	100:	2,957	216,764	1.36	2037	159893	1.27
5,094.11	384,355	8	23/12			4			4
Mexico	0.644.446	_	101:	203,664	2,244,268	9.07	11859	1325915	8.94
128,932.75	8,644,446	7	23/12			4	8		4
								1	

Total	46,477,80	17.478	1,060,98	46,477,80	2.28	574,82	24,326,74	2.36
2,207,211.884	3	17.470	3	3	3	6	5	2

Appendix 3

Population data

For Jersey the most recently produced estimate was for year-end 2019. Due to the running of the 2021 census an estimate has not been produced for 2020. Up to date population figures will next be published by Statistics Jersey as part of the upcoming census reports in the first quarter of 2022.

https://www.gov.je/Government/JerseyInFigures/Population/pages/population.aspx

So we took year-end 2019* growth rate for last year.

For Guernsey: GOV.GG The official website for the States of Guernsey:

https://www.gov.gg/population

For Other regions / countries : Population, total | Data (worldbank.org)