

# **Effect of Sub-Lethal Concentrations of *Vernonia amygdalina* (Bitter Leaf) on Testes of *Clarias gariepinus* (African Catfish) Juveniles**

## **ABSTRACT**

*Vernonia amygdalina* has recently been linked to insecticidal and pesticidal properties that could replace the harmful agrochemical pesticide usage around the aquatic environment and to aquatic inhibitors such as fish. This study aimed to determine the effect of sub-lethal concentrations of *Vernonia amygdalina* (bitter leaf) on testes of *Clarias gariepinus* (African catfish) juveniles. The fishes were exposed to 0.00, 0.40, 0.80 and 1.60 g/L graded concentrations of aqueous crude leaves extract of *Vernonia amygdalina* for two months. The organ (testis) changes in antioxidant biomarkers' histology, histomorphometry, and somatic indices compared with the control. There was the reduction in SOD, MDA and GSH activity as the concentration of the toxicant increased compared with the control. Mild interstitial oedema, mild tubular germ cell, interstitial cell depletions, severe depletion of seminiferous luminal content and sloughing off of the seminiferous tubular boundary tissue were observed in various concentrations of *Vernonia amygdalina* compared with the control. There was an increase in testis somatic indices as the plant extract concentration increased compared with the control. Bitter leaf extract was shown to have a toxic effect on *Clarias gariepinus* juveniles. As such, the pesticide use of the plants near the aquatic environment should be discouraged.

**Keywords:** Sub-Lethal Concentrations; *Vernonia amygdalina*; *Clarias gariepinus*; Testes; Juveniles.

## **1. INTRODUCTION**

With the world's population expected to reach 8.2 billion people by 2030, and 842 million people estimated to have been undernourished in 2011–13, the food supply will present a growing challenge in the next two decades (Lem, 2014). With increases in income and demographic changes related to family size, population ageing and urbanization, and consumer trends such as concerns for healthy eating and sustainable production, there will be significant shifts in demand and major changes in the composition of demand (Lem, 2014). This scenario will, in turn have an impact on food supply, which will need to increase and become more efficient if it is to grow within the constraints presented by the availability of natural resources and existing technology (FAO, 2014).

Currently, fish consumption by population has been increasing worldwide, mainly due to the availability, access and price of meat consumption, such as beef, pork, and poultry (Lem, 2014).

Consequently, some concerns begin to emerge, primarily regarding harvesting fishes from natural water bodies and fish treatment by using piscicides by aquaculturists which may affect the aquatic environment and quality of fish available in the market. The bioactive ingredient could be present in any product of animal origin, causing economic losses and putting human and animal health at risk. According to Bostock *et al.* (2010), aquaculture contributes nearly half of all food of aquatic origin intended for human consumption as a vital part of the global food industry.

The African catfish (*Clarias gariepinus*) is a species of catfish of the family Clariidae and comprises the most cultivated fishes in Nigeria. It is the most highly demanded freshwater fish globally due to its resistance to stress, ability to tolerate a wide range of environmental conditions and high stocking densities under culture conditions, and relatively fast growth (APHA, 1988). They are found throughout Africa and the Middle East, and live in freshwater lakes, rivers, swamps and human-made habitats, such as oxidation ponds or even urban sewage systems. The African catfish can never mentioned when talking about traditional fish capture. The current aquaculture practices, are similar to tilapia in terms of culture (Ayotunde *et al.*, 2011). The *C. gariepinus* is a native of Algeria, Angola, Benin, Botswana, Burkina Faso and an excellent sentinel to study of plant crude extracts. It feeds on detritus. It inhabits the streams, rivers, lakes, estuaries and brackish lagoons, found all year round, and its feeding habits reflect the local contamination simply because it is a bottom dweller.

Over the last several decades, botanical extracts have shown much interest in aquaculture to control fish parasites, fish fry predators and unwanted fishes from aquaculture ponds to replace chemical pesticides and pesticides. Since extensive and indiscriminate use of these non-biodegradable synthetic chemicals results in a harmful impact on the aquatic environment and presents a high risk to the non-targeted organisms (Kumar *et al.*, 2010; Das, 2013).

Various plants are reputed for their medicinal and antimicrobial values (Nascimento *et al.*, 2000; Adamu *et al.*, 2005), including their pesticidal, acaricidal and trypanocidal properties (Jhingran, 1975; Mgbojikwe and Okoye, 1998; Atawodi, 2005). However, some are known as potent arrow and fish poisons (Geidam *et al.*, 2007; Kamalkishor and Kulkarni, 2009), depending upon the type and the concentrations of their bioactive constituents. This is because plants contain structurally diverse biological substances with varying properties (Istvan, 2000). Plant extracts are considered promising agents because of their eco-friendliness, ease of availability, high efficiency, rapid biodegradability and reduced toxicity to non-targeted animals (Yunis, *et al.*, 2014). To date, a good number of plants have been investigated in different countries to evaluate their pesticidal (Rahuman *et al.*, 2008; Dubey *et al.*, 2010; Miresmailli and Isman, 2014) and piscicidal activities (Kulakkattolickal, 1989; Chiayvareesajja *et al.*, 1997; Neuwinger, 2004; Tiwari and Singh, 2004; Obomanu *et al.*, 2007; Murthy *et al.*, 2010; Ramanujam and Dominic, 2012). However, commercially available plant products are still limited, and hence, efforts should be made to find new sources of botanical pesticides and piscicides for rapidly growing pisciculture (Ramanujam and Ratha, 2008). Plant extracts are referred as piscicides if they exert toxicological effects on fishes and cause death to these aquatic animals (Burkill, 1995). Plant piscicides are obtained from a variety of plants belonging to different families and species that may vary considerably not only for their taxonomic variations but also for the plant parts used (leaves, barks, fruits and seeds), mode of use, mode of extraction and species of target fishes (Van Andel, 2000; Neuwinger, 2004). Botanical materials contain several bioactive compounds

that work either individually or synergistically as piscicides (Obomanu *et al.*, 2007; Ramanujam and Ratha, 2008). The degree of toxicity and piscicidal activity of any plant extract can be assessed by exposing fishes to it and subsequent estimation of the median lethal concentration (LC50) (Ramanujam and Dominic, 2012). One of the common toxicological effects of plant extract is to bring about changes in haematological, biochemical parameters and antioxidants of fishes (Winkaler *et al.*, 2007; Kavitha *et al.*, 2012), affect biological and physiological activities of fish. Adeyemo (2005) observed the haematological and histopathological effects of Cassava mill effluent in *Clarias gariepinus*.

Man's activities have been responsible for introducing contaminants into the environment in the last few decades (Awoyinka *et al.*, 2019). Environmental factors such as pH, turbidity, alkalinity, dissolved oxygen, temperature and conductivity influence the reaction rate of the pollutants entering the water or the lethal effects on the aquatic organisms (Fagbenro, 2002). Nowadays, pollution of natural ecosystems is increasing, so that with increasing human activities is becoming a fundamental problem. Determination of the toxic compounds in aquatic environments and their effects on aquatic organisms are a fundamental issue in ecotoxicology science. Toxic substances present in the environment can be determined by chemical analysis, but their effects on aquatic organisms in aquatic ecosystems cannot be determined by chemical analysis. Hence, to assess the environmental impact of toxic compounds, bioassay experiments are necessary (Bagheri, 2007).

Information regarding the adverse effect of sub-lethal concentrations on the reproductive system of *C. gariepinus* is scanty. Healthy testes of fish are important determinants of its breeding potential, and thus any toxicological factor adversely affecting the histopathology of testes will reduce the gross production of fish.

*Vernonia amygdalina* has been reported to be used as growth enhancer, phyto-additive, antibacterial, milt booster and insecticidal by *C. gariepinus* farmers (Kayode, 2021). However recent studies showed that *V. amygdalina* has negative effect on glucose level of *C. gariepinus* (Alagoa and Osakwe, 2021). Acute toxicity studies showed that the plant is toxic to catfish (Audu *et al.*, 2017). There is scarcity of information on the testis somatic indices, antioxidant biomarkers and histology and histomorphometry of *C. gariepinus* fish exposed to sub-lethal concentrations of *V. amygdalina*. Hence, this study examined the effect of sub-lethal concentrations of *Vernonia amygdalina* (bitter leaf) on testes of *Clarias gariepinus* (African catfish) juveniles.

## 2. MATERIALS AND METHODS

### 2.1 Description of Study Area

*Vernonia amygdalina* was collected from Keffi Local Government Area of Nasarawa State, Nigeria. This study was carried out at the Department of Zoology Laboratory, Faculty of Natural and Applied Sciences, Nasarawa State University, Keffi.

### 2.2 Procurement and Processing of Experimental Plant (*Vernonia amygdalina*)

The leaves were [air-dried](#) at room temperature ( $27 \pm 2^{\circ}\text{C}$ ) in the Laboratory of Zoology Department, Faculty of Natural and Applied Sciences, Nasarawa State University, Keffi. The dried leaves were pounded with laboratory mortar and pestle into powder, sieved with a 30  $\mu\text{m}$  mesh [sieve](#).

## 2.3 Collection, Transportation and Acclimatization of Experimental Animal

*C. gariepinus* was purchased from Rayuwa Fish Farm, Karu Local Government Area, Nasarawa State, Nigeria. Experimental animals were transported in tanks containing water from the pond to the Laboratory Unit of [the](#) Zoology Department, Nasarawa State University, Keffi. The fish was transferred into aquaria containing dechlorinated municipal tap water and [acclimatized](#) to the laboratory conditions for [two](#) weeks. During this period, the fish was fed to satiation at 8:00 a.m. and 6:00 p.m. with commercial fish feed (Vital feed®).

## 2.4 Sub-Lethal Concentrations of Aqueous Crude Leaf Extract of *V. amygdalina* and Test Procedure

Mortality (0, 1, 2, 5, 7 and 10) of *C. gariepinus* juveniles exposed to acute concentrations of 0.0, 0.19, 0.38, 0.75, 1.50 and 3.00 g/L of crude leaf extract of *A. amygdalina* was reported by [Audu et al.](#) (2018). The  $\text{LC}_{50}$  was calculated as 0.82 g/L by using [the](#) Finney calculator. *V. amygdalina* crude leaves extract [administered](#) at sub-lethal concentrations 1/5, 1/10, 1/20 of the  $\text{LC}_{50}$  ([Abubakar et al.](#), 2018) in a static renewable bioassay system.

The weighed samples were macerated in 1 L of distilled water for 24 hours at room temperature ( $27.0^{\circ}\text{C}$ ) to obtain the stock solution. The obtained stock solution was filtered through a funnel choked with non-absorbent cotton [wool](#), and each filtrate (1.6, 0.8 and 0.4 g/L) was transferred into tanks A1, B1, and [C1, respectively](#), with D1 as the control. Each of the filtrates was diluted by adding 9 L of dechlorinated municipal tap water. The same procedure was repeated in replicate tanks A2, B2 and [C2](#), while tanks D1 and D2 which were not inoculated with the test [materials](#), served as the controls. The sub-lethal toxicity test lasted for two months (56 days or [eight](#) weeks). The test fish was fed 3% body weight at 0800 and 1800hrs with photoperiod being natural.

## 2.5 Experimental Design

The experiments consist of four circular rubber tanks and one hundred and twenty *C. gariepinus* juveniles, mean weight  $23.43 \pm 2.43\text{g}$  and mean total length  $12.501 \pm 0.39\text{cm}$  in a non-randomized block design. Each of the eight tanks was filled with 9 L of dechlorinated municipal tap water. Six of the filled tanks were inoculated with various concentrations (powder earlier macerated in 1 L of water) of aqueous crude leaves extract of *V. amygdalina* and *C. gariepinus* fingerlings were introduced into each.

## 2.6 Assay for Biomarkers of Oxidative Stress

### 2.6.1 Preparation of the Tissue Samples for the Study

At the end of the experiment, fishes were carefully netted to minimize stress, weighed and sacrificed. Tissues such as liver, testes and gills were carefully removed and weighed. The tissues were washed in chilled phosphate buffer saline. Sample homogenates were made of individual tissues with chilled phosphate buffer (0.1 M, pH 7.4), centrifuged in a centrifuge at 9,000 rpm for 20 min at 4°C (Charity *et al.*, 2018). The supernatant obtained was used for further analysis. Samples were transported to Biochemistry Units of National Veterinary Research Institute (NVRI), Vom, Plateau State, Nigeria.

### 2.6.2 Determination of Glutathione Peroxidase

Glutathione peroxidase (GPx) activity was assayed as described by Paglia and Valentine (1967) with modifications according to Lawrence and Burke (1978). The reaction mixture contained 50 mM potassium phosphate buffer (pH 8.3), 1 mM EDTA, 1 mM sodium azide, 0.2 mM nicotinamide adenine dinucleotide phosphate (NADPH), and 1 U/mL glutathione reductase. The reaction was initiated with the addition of 1.5 mM cumene hydroperoxide. The enzyme activity was estimated from the rate of oxidation of NADPH. The reagents were mixed, and the absorbance was measured at 340 nm. Enzyme activity was expressed in mmol/minute/milligram protein.

### 2.6.3 Determination of Superoxide Dismutase Activity

Superoxide dismutase (SOD) activity was assayed as described by Misra and Fridovich (1972). The assay was based on the ability of SOD to inhibit the autoxidation of epinephrine at an alkaline pH. 0.5 ml of tissue homogenate was diluted with 0.5 ml of distilled water, to which 0.25 ml of ice-cold ethanol and 0.15 ml of ice-cold chloroform was added. The mixture was properly mixed using a cyclo mixer for 5 minutes and centrifuged at 2500 rpm. To 0.5 ml of supernatant, 1.5 ml of carbonate buffer (0.05 M, pH 10.2) and 0.5 ml of EDTA solution (0.49 M) were added. The reaction was initiated by the addition of 0.4 ml of epinephrine (3 mM), and the change in optical density/minute was measured at 480 nm against the reagent blank. SOD activity was expressed as units/mg protein. Change in optical density per minute at 50% inhibition of epinephrine to adrenochrome transition by the enzyme was taken as the enzyme unit.

### 2.6.4 Determination of Lipid Peroxidation

The malondialdehyde (MDA) content, a measure of lipid peroxidation, was assayed in the form of thiobarbituric acid reactive substances (TBARS) using the method of Buege and Aust (1978). 1.0 mL of the sample was added to 2 mL of (15% (w/v) trichloroacetic acid, 0.375% (w/v) thiobarbituric acid (TBA), 0.25 M HCl). The solution containing plasma and reagent was heated in a boiling water bath for 15 min and then cooled. Precipitates were removed by centrifuging at 1000 rpm for 10 minutes. The supernatant was removed, and the absorbance read at 535 nm against a blank. MDA was calculated using the molar extinction coefficient  $\epsilon$  for MDA  $1.56 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$ . Results were expressed as  $\mu\text{mol}^{-1}$ .

## 2.7 Extraction of Organs of *C. gariepinus* Juveniles Exposed to Sub-Lethal Concentrations of Aqueous Crude Leaves Extract of *Vernonia amygdalina*

Following the exposure of *C. gariepinus* juveniles to sub-lethal concentrations of crude leaf extract of *V. amygdalina*, four fishes from treatment and control tanks were removed, sacrificed and dissected to obtain the testis, gill, liver and kidney. Each removed organ was rinsed with distilled water to wash off traces of blood. Each organ was preserved in 10 ml specimen bottle containing 5 ml of formal saline before transportation to NVRI, Vom, Plateau State, Nigeria, for histopathological analyses of the tissues.

### **2.7.1 Determination of Organ Somatic Indices of *C. gariepinus* Juveniles Exposed to Sub-Lethal Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina***

The calculation of the testes somatic index (TIS) was done by random selection of four individual fishes from each treatment group and total weight of the individuals were noted (Shalaka and Pragna, 2013). The organs of the fishes like testes were removed carefully and weighed in an electronic weight machine, after removing moisture by blotting paper (Pamela *et al.*, 2018).

Organs Somatic Index was calculated using:

$$\text{OSI} = \frac{\text{Weight of organ}}{\text{Weight of fish}} \times 100$$

### **2.8 Procedure for Histological Examination of Organs of *C. gariepinus* Juveniles Exposed to Sub-Lethal Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina***

The routine Paraffin wax method and haematoxylin-eosin staining techniques of tissue processing described by Drury and Wallington (1967) and Avwioro (2011) were adopted to examine the gills, kidneys and liver of *O. niloticus* fingerlings exposed to aqueous crude leaves extract of *B. aegyptiaca*. The harvested gills, kidneys and liver of *C. gariepinus* were fixed in 10% formalin for three days, cut into thin slices of 5 x 2 x 1 mm thick and then processed with the SPIN Tissue Processor (STP) 120 (Thermo Scientific Model). The tissues were buffered in 10% formalin before passing through the following levels of hydrocarbons for two hours each: 70% Alcohol, 80% Alcohol, 90% Alcohol, 95% Alcohol, Absolute Alcohol I, Absolute Alcohol II, Absolute Alcohol III, Xylene I, Xylene II, Paraffin Wax Oven I, Paraffin Wax Oven II.

Tissues were embedded in molten paraffin wax using embedding moulds. The tissues were embedded using cassettes on a tissue Tek Embedding Centre (SLEE MPS/P2) and cooled rapidly on the cooling component. Tissues were sectioned using a rotary microtome (MICROM HM340E Thermo Scientific)) set at four microns, picked on slides and ready for staining.

Haematoxylin and eosin staining technique was used for the staining of the tissues. Tissue sections were dewaxed and hydrated by passing through two changes of xylene and through descending levels of alcohol (100%, 80%, 70%) for three min each and then into the water before staining in Harris' haematoxylin solution for 5 min and washed in running water. They were differentiated in 1% Acid alcohol and then washed thoroughly in water, blued in Scott's tap water substitute for 5 min and rinsed briefly in distilled water. Each tissue was then counterstained in 1% aqueous eosin for 2 min and then washed in water, dehydrated in descending grades of alcohol before clearing in xylene and mounted in Distrene, Plasticizer and

Xylene (DPX). Sections were then placed in slide carriers and placed in a 40°C oven to dry overnight. Each tissue was read microscopically. Photographs of the prepared sections were taken using a mounted photo-microscopic camera.

## 2.9 Histomorphometrics of Organs of *Clarias gariepinus* Juveniles Exposed to Sub-Lethal Concentrations of Aqueous Crude Leaves Extract of *Vernonia amygdalina*

Ten fishes from each experimental group were randomly selected. In addition, hepatocyte **histomorphometric** measurements such as hepatocyte nuclear diameter and hepatocytes surface area were determined by using Motic image plus 2.0 (Motic Asia, Hong Kong) software (Audu *et al.*, 2017).

## 2.10 Statistical Analysis Method

Data collected were subjected to analysis of variance (ANOVA) using SPSS version 16. Significant means were set at  $p < 0.05$  and separated using Duncan new multiple range test. Results were presented as means  $\pm$  standard errors of the mean.

## 3. RESULTS

There is a reduction in SOD activity as the concentration of the toxicant increases. The highest and lowest value of  $19.4462 \pm 0.98461$  and  $18.1615 \pm 1.10000$  were recorded in control (0.00 g/L) and highest concentration (1.60 g/L). There is also a reduction in GSH activity as the concentration of the toxicant increases, the highest and lowest values of  $10.2500 \pm 9.1250$  were recorded in 0.80 g/L and the highest value recorded in 0.40g/L. There is also reduction in GPx activity as the concentration of the toxicant increases, the highest and lowest values of  $2.9746 \pm 1.9046$  were recorded in 0.40g/L and the highest value recorded in 0.80g/L (Table 1).

There is mild interstitial oedema, mild tubular germ cell, interstitial cell depletions, severe depletion of seminiferous luminal content and sloughing off of the seminiferous tubular boundary tissue observed in testes of the exposed fishes as seen in Plate A-D.

Morphometric parameters such as STD with the highest and lowest value of  $133.79 \pm 9.799$  and  $87.35 \pm 6.211$  shows a significant difference compared to the control recorded in 0.40 g/L and 0.80 g/L. SLD with the highest and lowest value of  $100.45 \pm 7.561$  and  $76.70 \pm 9.241$  shows the significant difference when compared to the control recorded in 0.40 g/L and 0.80 g/L, while SED with the highest and lowest value of  $16.72 \pm 0.740$  and  $9.82 \pm 0.563$  shows significant difference when compared to the control recorded in 0.00 g/L and 1.60 g/L (Table 2).

TSI shows no significant difference compared with the control with the highest and lowest value of  $0.7475 \pm 0.14014$  and  $0.4675 \pm 0.15638$  recorded in the highest concentration (1.60g/L) and (0.80 g/L) (Table 3).

**Table 1: Mean Testes Superoxidase Dismutase, Lipid Peroxidation and Glutathione Peroxidase of *Clarias gariepinus* Juveniles Exposed to Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina***

Concentrations (g/L)	SOD	GSH	GPx
0.00	19.4462 ± 0.98461	9.2500 ± 0.50000	2.153 ± 0.34223
0.40	18.2000 ± 0.41538	10.2500 ± 0.50000	1.9046 ± 0.21081
0.80	18.4615 ± 0.75385	9.1250 ± 0.37500	2.9746 ± 0.64136
1.60	18.1615 ± 1.10000	9.5000 ± 0.75000	2.3272 ± 0.54893
<b>P-value (0.05)</b>	<b>0.7050</b>	<b>0.5370</b>	<b>0.4940</b>

**Key:** SOD = Superoxidase Dismutase; GSH = Glutathione Peroxidase; GPx = Lipid Peroxidation.

**Table 2: Sub-Lethal Effects of Bitter Leaves Crude Extract on Testes Morphometry Parameters**


Concentrations (g/L)	STD	SLD	SED
0.00	88.49±3.174 <sup>A</sup>	84.07±4.902	16.72±0.740 <sup>B</sup>
0.40	133.79±9.799 <sup>B</sup>	100.45±7.561	15.05±0.978 <sup>B</sup>
0.08	87.35±6.211 <sup>A</sup>	76.70±9.241	15.19±0.701 <sup>B</sup>
1.60	129.24±13.068 <sup>B</sup>	99.23±11.098	9.82±0.563 <sup>A</sup>

**Key:** Values with different letters in the same row are significantly different (P<0.05) compared with the control.  
 STD = Seminiferous Tubular Diameter; SLD = Seminiferous Luminal Diameter;  
 SED = Seminiferous Epithelial Diameter.

**Table 3: Mean Testes Somatic Index of *Clarias gariepinus* Juveniles Exposed to Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina***


Concentrations	TSI
1.60	$0.7475 \pm 0.14014$
0.80	$0.4675 \pm 0.15638$
0.40	$0.5100 \pm 0.10141$
0.00	$0.5200 \pm 0.12076$
<b>Total</b>	<b><math>0.5613 \pm 0.06515</math></b>

**Key:** TSI = Testes Somatic Index.




**Plate A-D: Photomicrographs of the Testes of *Clarias gariepinus* Exposed to Graded Concentrations of *Vernonia amygdalina*.**


**Key:** **A. Control (0.00 g/L):** The testes present normal numerous tubular structures housing different spermatogenic cell series (black arrow) and distinct interstitium between the tubules (white arrow). **B. 0.4 g/L:** There was no visible lesion except for mild interstitial oedema (white arrow). **C. 0.8 g/L:** Presence of mild tubular germ cell (black arrow) and interstitial cell depletion (white arrow). **D. 1.6 g/L:** Severe depletion of seminiferous luminal content (star) and sloughing off of the seminiferous tubular boundary tissue (black arrow). Magnification: x400; Stain: Haematoxylin-eosin.




**Figure 1: Mean Testes Somatic Index of *C. gariepinus* Juveniles Exposed to Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina*.**



**Figure 2: Mean Testes Somatic Index of *C. gariepinus* Juveniles Exposed to Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina*.**



**Figure 3: Mean Testes Somatic Index of *C. gariepinus* Juveniles Exposed to Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina*.**



**Figure 4: Mean Testes Somatic Index of *C. gariepinus* Juveniles Exposed to Concentrations of Aqueous Crude Leaves Extract of *V. amygdalina*.**

#### 4. DISCUSSION

Variation in testes antioxidant biomarkers (MDA, GSH, SOD) indicated oxidative stress in testes which is commensurate with the finding of Burak *et al.* (2016) in the testes of fish from the Lake Van and Karasu River of Van fish *Alburnus tarichi*, who reported that there was an unchanged MDA, GSH and SOD level. This research finding is also in line with a report from Alaa and Rania (2016), who opined that there was a significant increase ( $p < 0.05$ ) in activities of serum SOD, glutathione s-transferase (GSH), and malondialdehyde (MDA) when adult *C. gariepinus* were exposed to pure 100  $\mu\text{g/L}$  4-nonylphenol and quince (*Cydonia oblonga*) to fishes exposed for 15 days when compared to the control, but in this research, CAT was not recorded.

The severe sloughing off of the seminiferous tubular boundary tissue and severe depletion of seminiferous luminal content reported in this research work is in line with findings from Priya and Balu (2013) who reported in their work that there was severe deterioration of the testis histology by the 30<sup>th</sup> day of exposure, considerable reduction in spermatozoa in the testis when freshwater fish (*Rasbosa dandia*) was exposed to mercury. However, they reported a congestion of blood vessels and proliferation of interstitial tissues and vacant spaces enlarged which was not seen in this research work. The result shows that there is a significant difference in the testis

somatic indices of *C. gariepinus* concerning the graded concentrations of *V. amygdalina* when compared with the control, and this research work is not in line with the finding of Ada *et al.* (2015), who reported that there were no changes in gonad somatic indices in *C. gariepinus* exposed to Atrazine and coconut water after the fishes were exposed for a period of 14 days. This research work is in line with the findings of Claramma and Radhakrishnan (2016), which reported that there were histopathological changes observed as a distortion of seminiferous tubules, disorganization of spermatogonia, spermatocytes and spermatids with cytoplasmic vacuolization and nuclear pycnosis after exposing *Clarias batrachus* to sub-lethal concentrations of chromium for 45 days. Gupta (2002) also reported distortion of the seminiferous tubules in the testis of freshwater fish *Oreochromis mossambicus* in response to plant nutrients which is in line with the findings of this research work. However, he also recorded an increase in vacuolization, condensation of spermatocytes besides inflammation and inter-tubular vacuolation, which was very prominent.

There is an increment in the value of testis somatic index as the concentration of the plant increases which is not in line with the findings of Malik and Naeem (2020) that reported decline in the value of TSI of the fish with increase in the concentrations. Nassr-Allah (2007) also reported that there was a serious reduction in the GSI of blue tilapia juveniles as concentration levels of phenol increases with no significant increase reported for male fish subjected to 20% 96-h LC<sub>50</sub> of phenol, whereas significant reduction was observed in the case of male fish subjected to 40% and 80% 96-h LC<sub>50</sub> of phenol. Findings from this research work are in line with findings from the work of Shalaka and Pragna (2013), who reported that there was an increase in the GSI value when freshwater fish *Oreochromis mossambicus* were increasingly fed with plant nutrient.

## 5. CONCLUSION

Variation in testis antioxidant biomarkers (MDA, GSH, and SOD) indicated oxidative stress in testes. Alteration in testes histological and histomorphometric parameters also indicated that the bitter leaf is toxic to catfish and that the constant and indiscriminate use of bitter leaves as sperm boosters of catfish should be discouraged.

## 6. RECOMMENDATION

Further toxicity research works on the effect of the plants on the reproductive organs of other commercial fish species should be conducted.

## COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly used in our research area and country. There is no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for litigation but the advancement of knowledge. Also, the research was not funded by the producing company rather, it was funded by the personal efforts of the authors.

## REFERENCES

- Ada, F., Ayotunde, E. O. & William K., B. (2015). Gonado-hepatosomatic Indexes of *Clarias gariepinus* Sub-adult Exposed to Atrazine, *Cocosnucifera* Water and *Phyllanthusmuelarianus* Extract. *Journal of Aquaculture Research & Development*. **6**: 11.
- Adama, K. K., Oberafo, A. A. & Dika, S. (2011). Bitterleaf as Local Substitute for Hops in the Nigerian Brewing Industry. *Scholars Research Library*. **3**(4): 388–397.
- Adamu, H. M., Abayeh, O. J., Agho, M. O., Abdullahi, A. L., Uba, A., Dukku, H. U. & Wufem, B. M. (2005). An Ethnobotanical Survey of Bauchi State Herbal Plants and their Antimicrobial Activity. *Journal of Ethnopharmacology*. **99**: 1–4.
- Adedapo, A. A., Aremu, O. J. & Oyagbemi, A. A. (2014). Anti-oxidant, Anti-inflammatory and Antinociceptive Properties of the Acetone Leaf Extract of *Vernonia amygdalina* in Some Laboratory Animals. *Advanced Pharmaceutical Bulletin*. **4**: 591–598.
- Adesanoye, O. A. & Farombi, E. O. (2010). Hepatoprotective Effects of *Vernoniaamygdalina* (Asteraceae) in Rats Treated with Carbon Tetrachloride. *Experimental and Toxicologic Pathology*. **62**(2): 197–206.
- Adetunji, C. O., Olaniyi, O. O. & Ogunkunle, A. T. J. (2013). Bacterial Activity of Crude Extracts of *Vernoniaamygdalina* on Clinical Isolates. *Journal of Microbiology and Antimicrobials*. **5**(6): 60– 64.
- Adeyemo, O. K. (2005). Haematological and Histopathological Effects of Cassava Mill Effluent in *Clariasgariepinus*. *African Journal of Biomedical Research***8**: 179–183.
- Adiukwu, P., Amon, A., Nambatya, G., Adzu, B., Imanirampa, L., Twinomujuni, S. & Katusiime, B. (2012). Acute Toxicity, Antipyretic and Antinociceptive Study of the Crude Saponin from an Edible Vegetable *Vernoniaamygdalina* leaf. *International Journal of Biological and Chemical Sciences*. **6**(3): 1019–1028.
- Agbogidi, O. M. & Akpomorine, M. (2013). Health and Nutritional Benefits of Bitterleaf (*Vernonia amygdalina* Del.). *International. Journal of Applied Pharmaceutical Sciences and Biological Sciences*. **2**(3): 164-170.

- Akah**, P., Njoku, O., Nwanguma, A. & Akunyili, D. (2004). Effects of Aqueous Leaf Extract of *Vernonia amygdalina* on Blood Glucose and Triglyceride Levels of Alloxan-Induced Diabetic Rats (*Rattus rattus*). *Animal Research International*. **1**: 90–94.
- Akinyele**, B. J., Oladejo, B. O., Akinyemi, A. I. & Ezem, L. O. (2014). Comparative Study of the Antibacterial Effect of Mouth Washes and *Vernonia amygdalina* (Del.) on Some Tooth Decay Causing Bacteria. *British Microbiology Research Journal*. **4**(7): 749–758.
- Alaa**, E. H. S., & Rania, F. K. I. (2016). Endocrine Disruption, Oxidative Stress, and Testicular Damage Induced by 4-nonylphenol in *Clarias gariepinus*: the Protective Role of *Cydonia oblonga*. *Fish Physiology and Biochemistry*. **10**:100-117.
- Alabi**, D. A. & Amusa, N. A. (2005). Chemicals and Nutritional Composition of Four Botanicals with Fungitoxic Properties. *World Journal of Agricultural Sciences*. **1**(1): 84–88.
- Alara**, O. R., Abdurahman, N. H. & Olalere, O. A. (2017c). Ethanolic Extraction of Flavonoids, Phenolics and Antioxidants from *Vernonia amygdalina* Leaf using Two-Level Factorial Design. *Journal of King Saud University – Science*.
- Amodu**, A., Itodo, S. E. & Musa, D. E. (2013). Nigerian Foodstuffs with Tumour Chemosuppressive Polyphenols. *International Journal of Pharmaceutical Science Invention*. **2**(1): 12–17.
- Anil**, K. T. & Kanika, P. (2018). Testicular Oxidative Stress and Cellular Deformities in *Clarias gariepinus* (Burchell) from River Yamuna in Delhi Region, India. *Bulletin of Environmental Contamination and Toxicology*. Retrieved from; <https://doi.org>.
- Ankit**, S., Chetan, S. K. R. & Aneja, R. P. (2010). Bitterleaf as Local Substitute for Hops in the Nigerian Brewing Industry. *Pharmacia*. **2**(2): 208–220.
- Asifa**, K. P. & Chitra, K. C. (2016). Induction of Testicular Oxidative Stress by Nonylphenol in Cichlid Fish, *Etroplus maculatus* (Bloch, 1795). *International Journal of Fisheries and Aquatic Studies*, **4**(4): 453-456.
- Atangwho**, I. J., Egbung, G. E., Ahmad, M., Yam, M. F. & Asmawi, M. Z. (2013). Antioxidant Versus Anti-diabetic Properties of Leaves from *Vernonia amygdalina* Del. Growing in Malaysia. *Food Chemistry*. **141**(4): 3428–3434.
- Audu**, B. S., Omirinde, J. O., Gosomji, I. J. & Wazhi, P. E. (2017). Histopathological Changes in the Gills and Liver of *Clarias gariepinus* Exposed to Acute Concentrations of *Vernonia amygdalina*. *Animal Research International*. **14**(1): 2576-2587.
- Awoyinka**, O. A., Omodara, T. R., Oladele, F. C. & Ajayi, D. D., (2019). In-vitro Protein Digestibility of Selected Under-utilized Local Wild Beans and Bio-availability in Rats Model. *Academia journal Biotechnology*. **7**(4): 063-067.
- Ayoola**, G. A., Coker, H., Adesegun, S. A., Adepoju-Bello, A. A., Obaweya, K., Ezennia, E. C. & Atangbayila, T. O. (2008). Phytochemical Screening and Antioxidant Activities of Some Selected Medicinal Plants Used for Malaria Therapy in Southwestern Nigeria. *Tropical Journal of Pharmaceutical Research*. **7**(3): 1019-1024.

- Ayotunde** E. O., Offem, E. O. & Bakeh A. F. (2011). Toxicity of *Carica papaya* Linn: Haematological and Piscicidal Effect on Adult Catfish (*Clarias gariepinus*). *Journal of Fisheries and Aquatic Sciences*.**6**:291–308.
- Bagheri**,F.(2007). Study of Pessticial Residue (Diazinon, Azinphosmethyl) in Rivers of Golestan Province (Gorganround and Gharehsou). *Open Journal of Marine*. **7**(2): 1-125.
- Burak**, K., Ertuğrul, K., Abdulahad, D. & Atilla, D. (2016). Alterations in Histology and Antioxidant Defense System in the Testis of the Lake Van Fish (*Alburnustarichi* Güldenstädt, 1814). *Environment Monitoring Assessment*.**188**:474-547.
- Bostock**, J., Brendan Mc., Randolph, R., Kim J., Trevor, T., Kai L., David, L., Lindsay, R., Neil, H., Iain G. & Richard, C. (2010). Aquaculture: Global Status and Trends. *Philosophical Transactions of the Royal Society Biological Sciences*. DOI: 10.1098/rstb.2010.0170
- Burkill**,H. N. (1995).The Useful Plants of West Africa (tropical) Ed. 2 Vol. I. Families A.D. Royal Botanical Garden, Kew.
- Chiayvareesajja**, S., I., Chiayvareesajja, N. Rittibhonbhun, I. & Wiriyaichitra P. (1997). The toxicity of five native Thai Plants to Aquatic Organisms. *Asian Fisheries Science*. **9**:261–267.
- Claramma**, J. & Radhakrishnan, M. V. (2016). Histopathological Analysis of Chromium Toxicity to Testis of the Catfish *Clarias batrachus* (Linn). *IOSR Journal of Environmental Science, Toxicology and Food Technology*. **10**: 01-04.
- Clement**, E., Erharuyi, O., Vincent, I., Joy, A., Christopher, A., Anthony, A. U., Onyekaba, T., Osakue, J., Iftikhar, A. & Abiodun, F. (2014). Significance of Bitter Leaf (*Vernonia amygdalina*) in Tropical Diseases and Beyond: A Review.
- Dubey**, S. P., Manu, L. &Manu, S. (2013). Green Synthesis and Characterizations of Silver and Gold Nanoparticles using Leaf Extract of *Rosa rugosa*. *International Journal of Colloids and Surfaces A: Physiochemical and Engineering Aspects*. **364**: 34-41.
- Ejike**, C. E. C. C., Awazie, S. O., Nwangozi, P. A. & Godwin, C. D. (2013). Synergistic Postprandial Blood Glucose Modulatory Properties of *Vernonia amygdalina* (Del.), *Gongronem alatifolium* (Benth.) and *Occimum gratissimum* (Linn.) Aqueous Decoctions. *Journal of Ethnopharmacology*. **149**(1): 111–116.
- Erasto**, P., Grierson, D. S. and Afolayan, A. J. (2006). Bioactive Sesquiterpene Lactones from the Leaves of *Vernonia amygdalina*. *Journal of Ethnopharmacology*. **106**: 117–120.
- Ertuğrul K.**, **Abdulahad D.** & **Atilla, D.** (2016). Alterations in Histology and Antioxidant Defense System in the Testis of the Lake Van Fish (*Alburnustarichi* Güldenstädt, 1814). *Environ Monit Assess*. **188**(8):474.
- Ezeadila**, J. O., Nwande, M. O., Ogu, G. I., Aneke, F. A. & Ezeadila, J. O. (2015). Antibacterial Activity of Ethyl Acetate And n-Hexane Extracts of *Vernoniaiaamygdalina* and

*Moringaoleifera* Leaves on *Staphylococcusaureus* and *Escherichiacoli* Isolated from Urine Samples. *Pelagia Research Library*. **6**(11): 23–28.

- Fagbenro**, O. A. (2002). The Dietary Habits of Clarid Catfish *Heterobranchus bidorsalis* (Geoffery st. Hilaria 1809) in Owerri, Southern Nigeria. *Tropical Zoology*. **5**: 11–17.
- Farombi**, E.O. (2003). African Indigenous Plants with Chemotherapeutic Potentials and Biotechnological Approach to the Production of Bioactive Prophylactic Agents. *African Journal Biotechnology*. **2**: 662–671.
- Farombi**, E. O., & Owoeye, O., (2011). Antioxidative and Chemopreventive Properties of *Vernoniaamygdalina* and *Garciniabiflavonoid*. *International Journal of Environmental Research Public Health*. **8**: 2533-2555; DOI 10.3390/ijerph8062533.
- Georgewill**, O.A. & Georgewill, U. O. (2010). Evaluation of the Anti-inflammatory Activity of Extract of *Vernoniaamygdalina*. *Asian Pacific Journal of Tropical Medicine*. **6**:150–151.
- Geidam**, M. A., Dauda, E. & Hamza, H. G. (2007). Effects of Aqueous Stem-bark Extract of *Momordicabalsamina* Linn on some Serum Enzymes in Normal and Ethanol Fed Rats. *Journal of Biological Sciences* **7**: 397–400.
- Gresham**, L. J., Ross, J. & Izevbigie, E. B. (2008). *Vernoniaamygdalina*: Anticancer Activity, Authentication, and Adulteration Detection. *International Journal of Environmental Research and Public Health*. **5**(5): 342–348.
- Hanan**, S. A., Shimaa, A. A., Dina, M. M. A., Alshimaa, A. K., Eman, M. Z. & Doaa, A. El-A. (2020). Optimizing the Growth, Health, Reproductive Performance, and Gonadal Histology of Broodstock Fantail Goldfish (*Carassiusauratus*, L.) by Dietary Cacao Bean Meal. *Animals*. DOI: 10.3390/ani10101808s.
- Igile**, G. O., Oleszek, W., Jurzysta, M. (1995). Vernoniosides D and E, two Novel Saponins from *Vernoniaamygdalina*. *Journal of National Prodruction*. **58**: 1438–1443.
- Igile**, G. O., Wieslaw, O., Jurzysta, M., Stanislaw, B. and Fasanmade, A. (1995). Flavonoids from *Vernoniaamygdalina* and their Antioxidant Activities. *Journal of Agricultural and Food Chemistry*. **42**: 2445–2448.
- Ijeh**, I. I. & Ejike, C. E. C. C (2011). Current Perspectives on the Medicinal Potentials of *Vernoniaamygdalina* Del. *Journal of Medicinal Plants Research*. **5**(7): 1051-1061.
- Imaga**, N. O. A. & Bamigbetan, D. O. (2013). In Vivo Biochemical Assessment of Aqueous Extracts of *Vernoniaamygdalina* (Bitter leaf). *International Journal of Nutrition and Metabolism*. **5**: 22–27.
- Istvan**. U. (2000). Semi-Natural Products and Related Substances as Alleged Botanical Pesticides. *Journal of Pest Management Sciences*. **56**: 703–705.
- Kiplimo**, J. J., Koorbanally, N. A. & Chenia, H. (2011). Triterpenoids from *Vernoniaauriculifera* Hiern Exhibit Antimicrobial Activity. *African Journal of Pharmacy and Pharmacology*. **5**(8): 1150–1156.

- Kumar**, A., Lingadurai, S., Jain A. & Barman, N. R. 2010. *Erythrina variegata* Linn: A Review on Morphology, Phytochemistry, and Pharmacological Aspects. *Pharmacognosy Review*. **4**(8): 147-152.
- Kamalkishor**, H. N. & Kulkarni, HN. (2009). Fish Stupefying Plants used by the Gond Tribal of Mendha Village of Central India. *Indian Journal of Traditional Knowledge*. **8**: 531-534.
- Kavitha**, C., Ramesh, M., Kumaran, S. S., & Lakshmi, S. A., (2012). Toxicity of *Moringa oleifera* Seed Extract on Some Hematological and Biochemical Profiles in a Freshwater Fish, *Cyprinus carpio*. *Exp. Toxicol. Pathol.* **64**(7-8): 681-687.
- Kulakkattolickal**, A. T. (1989). Piscicidal Plants of Nepal: Toxicity to Air Breathing Predatory fish (*Ophicephalus punctatus*, *Clarias barbatrachus* and *Heteropneustes fossilis*) and duration of Risk to Cultivated fish. *Aquaculture*. **78**:285–292.
- Lem**, A., Bjørndal, T. & Lappo, A. (2014). Economic Analysis of Supply and Demand for Food up to 2030 – Special Focus on Fish and Fishery Products. *FAO Fisheries and Aquaculture Circular No. 1089*.
- Luo**, X., Jiang, Y., Fronczek, F. R., Lin, C., Izevbigie, E. B., Lee, S. Lee, K. S. (2017). Isolation and Structure Determination of a Sesquiterpene Lactone (Vernodalinalol) from *Vernonia amygdalina* Extracts. *Pharmaceutical Biology*. **49**(5): 464–470.
- Momoh**, M., Adikwu, M. & Oyi, A. (2010). *Vernonia amygdalina* Extract and CD4+ Cell Counts: An Immune Study. *Global Journal of Biotechnology and Biochemistry*. **5**(2):92-96.
- Murthy**, K. S., Kiran, B. R., Venkateshwarlu, M. (2010). A Review on Toxicity of Pesticides in Fish. *International Journal of Open Science Research*. **1**(1): 15-36.
- Nascimento**, G.G.F., Locatelli, J., Freitas, P. C. & Silva, G. L. (2000). Antibacterial Activity of Plant Extracts and Phytochemicals on Antibiotic Resistant Bacteria. *Brazilian Journal of Microbiology*. **31**: 247–256.
- Neuwinger**, H. D. (2004). Fish Poisoning Plants of Africa. *Botanica Acta*. **107**:263–270.
- Njan**, A.A., Bulus, A., Amos, G.A., Dominic, B., Silvia, & David, R.B. (2008). The Analgesic and Antiplasmodial Activities and Toxicity of *Veronia amygdalina*. *Journal of Medicinal Food*. **11**(3): 574-581.
- Ngatu**, N. R., Okajima, M. K., Yokogawa, M., Hirota, R., Takaishi, M., Eitoku, M. & Suganuma, N. (2012). Anti-Allergic Effects of *Vernonia amygdalina* Leaf Extracts in Hapten-Induced Atopic Dermatitis-Like Disease in Mice. *Allergology International*. **61**(4): 597–607.
- Nwanjo**, H. U. (2005). Efficacy of Aqueous Leaf Extract of *Vernonia amygdalina* on Plasma Lipoprotein and Oxidative Status in Diabetic Rat Models. *Nigerian Journal of Physiological Sciences: Official Publication of the Physiological Society of Nigeria*. **20**(1–2): 39–42.

- Nwaoguikpe**, R. N. (2010). The effect of Extract of Bitter Leaf (*Vernonia amygdalina*) on Blood Glucose Levels of Diabetic Rats. *International Journal of Biological and Chemical Sciences*. **4**(3): 721-729.
- Obomanu**, F. G., Ogbalu, O. K., Gabriel, U. U., Fekarurhobo, S. G. B. & Abadi S. U. (2007). Piscicidal Effects of *Lepidagathisalopecuroides* on Mudskipper, *Periophthalmus papillio* from the Niger Delta. *Research Journal Applied Sciences* **2**(4): 382-387.
- Olowolafe**, T. & Olufayo M. O., (2018). Toxicity of Aqueous Extracts of Bitter leaf (*Vernonia amygdalina*) on Haematological Profile of African Catfish (*Clarias gariepinus*) Juveniles. *International Journal of Fisheries and Aquatic*. **6**(2): 596-600.
- Osinubi**, A. A. A. (2008). Effects of *Vernonia amygdalina* and Chlorpropamide on Blood Sugar. *Medical Journal of Islamic World Academy of Sciences*. **16**(3): 115–119.
- Opata**, M. M. & Izevbigie, E. B. (2006). Aqueous *Vernonia amygdalina* Extracts Alter MCF-7 Cell Membrane Permeability and Efflux. *International Journal of Environmental Research and Public Health*. **3**(2): 174–179.
- Owu**, D. U., Ben, E. E., Antai, A. B., Ekpe, E. A. & Udia, P. M. (2008). Stimulation of Gastric Acid Secretion and Intestinal Motility by *Vernonia amygdalina* Extract. *Fitoterapia*. **79**(2): 97–100.
- Priya**, R., & Balu, T. K. (2013). Mercury Induced Histopathological Changes in the Testis of the Freshwater Fish, (*Rasbora dandia*). *International Journal of Science and Research*. **4**: 2319-7064.
- Rahuman**, A.A, Gopalakrishnan, G, Venkatesan, P. & Geetha K. (2008). Larvicidal Activity of Some Euphorbiaceae Plant Extracts against *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae). *Parasitology Research*. **102**:867-873.
- Ramesh**, H. & Muniswamy, D. (2009). Behavioral Response of the Freshwater Fish, *Cyprinus carpio* (Linnaeus) Following Exposure to Chlorpyrifos. *Turkish Journal of Fisheries and Aquatic Sciences*. **9**: 233-238.
- Samuel**, U., Emmanuel, I. N., Gregory, E. O., Bright, C. I. & Ogonna, C. A. (2021). Evaluation of Lambda-cyhalothrin Oxidative Stress and Gonad Histoarchitecture Toxicity Potency in *Clarias gariepinus*. *Journal of Basic and Applied Zoology*. **82**: 2-11.
- Shalaka**, S. & Pragna, P. (2013). Gonadosomatic and Hepatosomatic Indices of Freshwater Fish *Oreochromis mossambicus* in Response to a Plant Nutrient. *Journal of Zoology*. **8**(1): 110-118.
- Tiwari**, S. & Singh, A. (2004). Control of Common Freshwater Predatory Fish, *Channa punctatus* through *Nerium indicum* leaf Extracts. *Chemosphere*. **53**: 865–875.
- Toyang**, N. J. & Verpoorte R. (2013). A Review on Antimicrobial Potential of Species of the Genus *Veronia*. *A Journal of Medicinal Plant Research*. **9**(31): 838-850.

- Toyang**, N. J. & Verpoorte, R. (2013). A Review of the Medicinal Potentials of Plants of the Genus *Vernonia* (Asteraceae). *Journal of Ethnopharmacology*. **146**(3): 681–723.
- Udochukwu**, U., Omeje, F. I., Uloma, I. S. & Oseiwe, F. D. (2015). Phytochemical Analysis of *Vernoniaamygdalina* and *Ocimumgratissimum* Extracts and their Antibacterial Activity on Some Drug Resistant Bacteria. *American Journal of Research Communication*. **3**(5): 225–235.
- Van**, F., & Andel. T. (2000). The Diverse Use of Fish Poison Plants in Northwest Guyana. *Economic Botany*. **54**:500–512.
- Quasie**, O., Zhang, Y., Zhang, H., Luo, J. & Kong, L. (2016). Four New Steroid Saponins with Highly Oxidized Side Chains from the Leaves of *Vernoniaamygdalina*. *Phytochemistry Letters*. **15**: 16– 20.
- Winkaler**, E. U., Santos, T. R. M., Machado-Neto, J. G. & Martinez, C. B. R. (2013). Acute Lethal and Sublethal Effects of Neem Leaf Extract on the Neotropical Freshwater Fish *Prochilodus lineatus*. *Comp. Biochem. Physiol. Toxicol., Pharmacol.*, **145**(2): 236-244.
- Yeap**, S. K., Wan, Y. H., Boon, K. B., Woon, S. Li., Huynh, K., Abdul, H. N. Y., & Noorjahan B. A. (2010). *Vernoniaamygdalina*, an Ethnoveterinary and Ethnomedical Used Green Vegetable with Multiple Bioactivities. *Journal of Medicinal Plants Research*. **4**(25): 2787-2812.
- Yunis**, J.A., Gustavo, S.C., Paulo, F.M., Cynthia, I., George, G.O., Silas, F.E., Claudine, C.M., Juliet, R.E.M., Flavio, R.M. & Joao, B.K.F. (2014). Acute Toxicity and Determination of the Acute Constituents of Aqueous Extract of *Uncaria tomentosa* Bark in *Hyphessobrycon eques*. *Journal of Toxicology*. **20**(14) 56-87.