

Mini review Article

Virgin Olive Oil: a rich source of functional bioactive compounds in the promotion of human health-An overview

Comment [P1]: Line 12

ABSTRACT

Comment [P2]: Abstract

The prevention of pathologies by diet management is an important public health challenge. The Mediterranean diet (Med diet) is considered one of the healthiest dietary patterns. According to current scientific research, this dietary pattern is linked to a decreased prevalence and incidence of various chronic diseases, including cardiovascular disease, diabetes, cancer, and age-related cognitive decline, as well as a lower overall mortality rate. In the Mediterranean diet, a high consumption of Virgin Olive Oil (VOO), fruit, nuts, vegetables, and grains is linked to a decreased incidence of many diseases and a longer life expectancy. According to numerous studies, VOO is the major food responsible for the health and nutritional benefits of the Mediterranean diet. VOO chemical compounds have been shown *in vitro* and *in vivo* investigations to have potentially favorable health benefits as a result of their biological activities. VOO composition has been linked to a variety of biological activities, including antioxidant, anti-inflammatory, anti-diabetic, anti-cancer, and antiviral characteristics. As a result, increasing attention is being paid to the composition of VOO and the identification of its biologically active components. Because of their antioxidant, anti-inflammatory, and anti-thrombotic properties, phytochemicals, particularly phenolic compounds, have health-promoting effects, especially on cardiovascular and metabolic illnesses. These characteristics are even more important in light of the COVID-19 severe illness's heightened inflammatory and pro-thrombotic consequences. This overview collects and discusses the scattered data available in the literature concerning VOO compounds of biological interest and highlights their possible mechanisms of action and effects on human health.

Keywords: Virgin Olive Oil (VOO), bioactive compounds, biological activities, beneficial health effects

ABBREVIATIONS

Comment [P3]: Small letter

AChe: Acetylcholinesterase, **ACE2:** Angiotensin-Converting Enzyme 2, **BHT:** 2,6-di-tert-butyl-hydroxytoluene, **BuChE:** Butyrylcholinesterase, **CAT:** Catalase, **CNS:** Central nervous system, **COX:** Cyclooxygenase, **DNA:** Deoxyribonucleic acid, **EFSA:** European Food Safety Authority, **GSH:** Glutathione, **HDL:** high density lipoprotein; **HCV:** Hepatitis C virus, **HIV:** human immunodeficiency virus, **4-HNE:** 4- Hydroxynonenal, **HT:** Hydroxytyrosol, **•OH** **sacavenging:** Hydroxyl radical scavenging, **IOC:** International olive council, **5-LOX:** 5-Lipoxygenase, **LDL:** low density lipoprotein, **Med diet:** Mediterranean diet, **OA:** Oleocanthal, **OE:** Oleuropein, **ROS:** Reactive oxygen species, **SOD:** Superoxide dismutase, **TBARS:** Thiobarbituric acid-reacting substances, **T:** Tyrosol, **VOO:** Virgin olive oil.

1. INTRODUCTION

Comment [P4]: Small letter

Worldwide olive oil production was about 3266.5-mile tons in 2019/2020, the majority of them in Mediterranean countries [1]. Spain, Italy, Tunisia, and Greece (followed by Turkey, Morocco, Portugal, and Algeria) are the biggest olive oil-producing countries [1] (Fig. 1).

Comment [P5]: Delete

Comment [P6]: Delete




Fig. 1. Olive oil producing countries [1]

Because of the health benefits of olive oil [2], consumption is not restricted to the Mediterranean basin and is consumed in non-traditional producing countries, such as the United States (402.5-mile tons) [3]. Spain consumes the largest per capita amount of olive oil with over 521.6-mile tons and Italy with around 404.4-mile tons in 2019/2020 [3].

Virgin olive oil is the main fat of the Med diet. Virgin olive oil is oil which is obtained from the fruit of the olive tree (*Olea europaea* L.) solely by mechanical or other physical means under conditions, particularly thermal conditions, that do not lead to alterations in the oil, and which have not undergone any treatment other than washing, decantation, centrifugation and filtration [4].

Several epidemiological studies have shown that Med diet rich in VOO can diminish the incidence of chronic diseases (Fig. 2) [5, 6, 7]. Furthermore, many experiences in human

beings have revealed that the intake of VOO may be protective against these pathologies [8, 9, 10, 11].




Fig. 2. The protective potential of the Mediterranean diet

It is well reported that in living organisms, the balance between reactive oxygen species (ROS) and endogenous antioxidants may be altered in some pathological and physiological conditions, leading to a lower capacity of cellular defense against free radicals and/or increased production of ROS, thus creating cellular oxidative stress [12]. Oxidative stress occurs when cells are unable to eliminate free radicals known as reactive oxygen species (ROS) using the natural antioxidant defense system (Fig. 3). Target molecules for ROS are lipids, DNA, and proteins. Oxidative stress associated with aging appears to be involved in several chronic pathologies like atherosclerosis, Alzheimer's disease, type 2 diabetes mellitus, and cancer [13].




Fig. 3. Oxidative stress

Exploring the chemical composition of our diet, which has enhanced our overall standard of health, is one of the latest themes in food science. Food chemical compounds, in fact, intervene in biological processes to deliver a variety of health benefits. Bioactive chemicals of interest could be examined to see if they have a wide range of biological applications. As a result, there is a massive list of bioactive compounds that have already been published in the literature, and new compounds are being introduced on a regular basis. VOO, on the other hand, is one of the greatest sources of phytochemicals with biological activities known to date, as well as the only food with EFSA-approved health benefits [14]. VOO has been extensively researched for a variety of health advantages over the years. The health-promoting characteristics of VOO have sparked a lot of curiosity. Following research (both *in vivo* and *in vitro*) has shown that VOO bioactive chemical compounds have beneficial effects on physiological parameters, potentially lowering the risk of chronic diseases linked to oxidative stress [15]. As a source of bioactive compounds, it plays an important role as the main food of the Med diet [16, 17], which is associated with a lower incidence of several diseases [18, 19, 20]. With the advancement in scientific research, VOO bioactive compounds were identified, as well as their applications in the biological system [8, 21, 22, 23]. Monounsaturated fatty acids (especially oleic acid), phenolic compounds, and squalene are some of the main active components found in this matrix [24].

Following these trends, this article will focus on the most existing features of virgin olive oil's health benefits, which are mostly due to the presence of phenolic compounds.

2. METHODOLOGY

The current study was based on literature that is already available on various scientific databases and highlighted the health benefits of VOO. PubMed, Medline, PubMed Central, Science Direct, and a few more scientific databases were included in the study. The current work compiles, critically interprets, and presents the data gathered from these various sources.

3. FATTY ACIDS

Comment [P7]: Fatty acids

3.1. FATTY ACIDS PROFILE

Comment [P8]: Fatty acids profile

VOO is primarily composed of mono- and polyunsaturated fatty acids, accounting for more than 98% of the total weight. VOO mainly contains palmitic acid (C16:0), hypogeic acid (C16:1 ω 9), palmitoleic acid (C16:1 ω 7), margaric acid (C17:0), margaroleic acid (C17:1), stearic acid (C18:0), oleic acid (C18:1 ω 9), cis-vaccenic acid (C18:1 ω 7), linoleic acid (C18:2), arachidic acid (C20:0), alpha-linolenic acid (C18:3), eicosenoic acid (C20:1), behenic acid (C22:0), and lignoceric acid (C24:0). Out of the given fatty acids, oleic acid is found in the highest concentration [23]. This fatty acid represents the most abundant fatty acid in VOO. Its concentration ranges from 56% to 84% of total fatty acids [23].

UNDER PEER REVIEW

Table 1. Fatty acid composition of virgin olive oil

Source of Virgin Olive Oil	Localization	GPS coordinate	Fatty acid composition (%)							Reference
			C16:0	C16:1 ω7	C17:0	C18:0	C18:1 ω9	C18:2 ω6	C18:3 ω3	
Tunisian varieties										
Chemlali	Sfax	34°44'28.0N 10°45'35.1"E	15.81 ±0.60	3.61 ±0.15	0.03 ±0.002	1.77 ±0.08	61.92 ±2.7	15.77 ±1.02	0.56 ±0.03	[25]
Chétoui	Oueslatia	35°51'02.7N 9°35'10.4"E	11.53 ±0.80	0.39 ±0.03	n.d	2.99 ±0.12	64.24 ±1.22	19.63 ±0.62	0.68 ±0.15	[26]
Zalmati	Medenine	33°20'46.3N 10°29'24.6"E	17.00 ±1.10	1.93 ±0.38	0.03 ±0.05	2.06 ±0.01	63.43 ±2.31	14.47 ±0.6	0.51 ±0.08	[25]
Italian varieties										
Coratina	Calabria	38°06'37" N 15°39'40" E	12.62 ± 1.32	0.93 ± 0.03	0.02 ± 0.00	1.74 ± 0.24	71.70 ± 4.50	10.42 ± 0.93	0.80 ± 0.02	[27]
Leccino	Calabria	38°06'37" N 15°39'40" E	14.25 ± 1.21	1.31 ± 0.04	0.02 ± 0.00	2.50 ± 0.84	70.52 ± 3.90	10.43 ± 0.62	0.67 ± 0.04	[27]
Ottobratica	Calabria	38°06'37" N 15°39'40" E	15.12 ± 2.22	1.44 ± 0.13	0.02 ± 0.00	2.23 ± 0.40	69.60 ± 6.21	9.92 ± 0.70	0.90 ± 0.02	[27]
Nocellara del Belice	Calabria	38°06'37" N 15°39'40" E	11.14 ± 1.01	0.60 ± 0.01	0.08 ± 0.00	2.24 ± 0.41	76.30 ± 4.52	8.12 ± 0.70	0.58 ± 0.03	[27]
Spanish varieties										
Arbequina	Extremadura	39°31'55.5N 6°23'38.4"W	15.32 ±1.41	1.64 ±0.24	0.11 ±0.01	1.33 ±0.48	67.25 ±4.31	12.66 ±2.60	0.59 ±0.13	[28]
Cornicabra	Extremadura	39°31'55.5N 6°23'38.4"W	12.12 ±2.06	1.00 ±0.32	0.05 ±0.01	2.44 ±1.10	77.36 ±6.07	5.32 ±4.25	0.67 ±0.25	[28]
Picual	Extremadura	39°31'55.5N 6°23'38.4"W	11.62 ±1.18	1.02 ±0.17	0.04 ±0.01	1.96 ±0.88	80.67 ±1.95	3.08 ±0.46	0.65 ±0.12	[28]

Data are presented as the mean of the number ± Standard deviation of different samples analyzed.

Fatty acids: C16:0, palmitic acid (hexadecanoic acid); C16:1ω7, palmitoleic acid (9-hexadecenoic acid); C17:0, margaric acid (heptadecanoic acid); C18:1ω9, oleic acid (9-octadecenoic acid); C18:2ω6, linoleic acid (9,12-octadecadienoic acid); C18:3ω3, linolenic acid (9,12,15-octadecatrienoic acid).

3.2. Biological activities of fatty acids

Many experimental studies on human beings have disclosed that the intake of monounsaturated fatty acids may be protective against chronic diseases. Extensive evidence points to a series of potential health effects of oleic acid in VOO. Indeed, Bermudez et al. [29] demonstrated the relevance of oleic acid in VOO on different nutrition-related issues and discussed the impact of oleic acid in olive oil and its clinical relevance to major risk factors for chronic diseases. In this respect, Soriguer et al. [9] showed that the regular consumption of VOO exerts a healthy effect on several cardiovascular danger factors, particularly in the presence of obesity, a reduced glucose tolerance, or a sedentary way of life. Besides, Kien et al. [30] reported that replacing dietary palmitic acid with oleic acid reduces the blood LDL concentration and whole-body fat oxidation. Montserrat-De La Paz et al. [31] reported that monounsaturated fatty acids present in VOO prevent atherosclerosis by interfering with various inflammatory responses. Lamy et al. [32] demonstrated that oleic acid may be useful in cancer chemoprevention.

4. Phenolic compounds

4.1. Phenolic compounds profile

Secondary metabolites with at least one aromatic ring and one or more hydroxy substituents are known as phenolic compounds [33]. In VOO, more than 36 phenolic compounds have been discovered [34, 35, 36, 37]. The main phenolic compounds identified in VOO are phenolic acids (benzoic acid derivatives, cinnamic acid derivatives, and other phenolic acids and derivatives), phenolic alcohols (tyrosol and hydroxytyrosol), secoiridoids (oleuropein and ligstroside), hydroxy-isocromans (3,4-dihydro-1H-benzo[c]pyran derivatives), flavonoids (flavones and flavanols) [34, 38]. Fig. 4 shows the chemical structures of various phenolic compounds of VOO.




Fig. 4. Chemical structures of phenolic compounds of virgin olive oil

Numerous factors influence the concentration of phenolics in VOO, including agronomic factors such as the area of origin, cultivar, and stage of fruit ripening, as well as agronomic practices and the technological and operational circumstances of the oil extraction process [39, 40, 41]. Table 2 shows the phenolic content of virgin olive oils from Tunisian, Italian, and Spanish varieties.

Table 2. Phenolic content of virgin olive oils from some Tunisian, Italian and Spanish varieties

Source of virgin olive oil	Localization	GPS coordinate	Phenolic content (mg GAE/kg)*	Reference
Tunisian varieties				
Chemlali	Sfax	34°44'28.0"N 10°45'35.1"E	266.00 ± 5.76	[25]
Chétoui	Oueslatia	35°51'02.7"N 9°35'10.4"E	800.00 ± 3.32	[26]
Zalmati	Medenine	33°20'46.3"N 10°29'24.6"E	189.63 ± 4.67	[25]
Italian varieties				
Colozzese	Lecce	40°21'17.32" N, 18°10'20.78" E	251.00 ±12.00	[42]
Barone di Monteprofico	Lecce	40°21'17.32" N, 18°10'20.78" E	202.00 ± 14.00	[42]
Cellina di Nardò	Lecce	40°21'17.32" N, 18°10'20.78" E	253.00 ± 7.00	[42]
Spanish varieties				
Arbequina	Extremadura	39°31'55.5"N 6°23'38.4"W	234.00 ± 2.40	[43]
Manzanilla	Extremadura	39°31'55.5"N 6°23'38.4"W	210.70 ± 13.20	[43]
Picual	Extremadura	39°31'55.5"N 6°23'38.4"W	590.30 ± 2.20	[43]

*Data are presented as the mean of the number of different samples analyzed; *: mg Gallic Acid Equivalent*

4.2 Biological activities of phenolic compounds

VOO phenolics have been shown to have positive effects on specific physiological parameters and decrease the risk of chronic diseases in both *in vitro* and *in vivo* studies [34,

44, 45, 46, 47, 48, 49]. Furthermore, VOO phenolic compounds are highly bioavailable, thus reinforcing their potential health-promoting properties [50, 51, 52].

4.2.1 Antioxidant activity

Interest in phenolic compounds is related primarily to their antioxidant activity, which may contribute to countering many chronic diseases that may be related to oxygen radical formation, oxidative damage, and aging [53]. Phenolic compounds work inside cellular compartments as the first line of defense against free radicals due to their capacity to stabilize free radicals through the formation of intramolecular hydrogen bonds. In this context, many studies disclose the important antioxidant potential of VOO phenolics [42, 54, 55, 56].

The antioxidant properties of phenolic compounds can be achieved in a variety of ways. They can chelate metallic ions in oxidative systems using transition metals, preventing their involvement in Fenton reactions, which can generate significant amounts of hydroxyl radicals [57, 58]. Furthermore, the most essential antioxidant activity is connected to the ability to scavenge free radicals by interrupting the chain of reactions set off by free radicals. Indeed, the potential of phenolics to promote radical stability by creating an intramolecular hydrogen link between the free hydrogens of their hydroxyl groups and the phenoxyl radicals is well recognized [59]. *In vivo* studies confirmed that VOO phenolic compounds appear to regulate the oxidative/antioxidative balance and explained that the mechanisms by which phenolic compounds of VOO can protect lipids and DNA are primarily related to their abilities to counteract metal- and radical-dependent oxidation and to act as chelating agents, thus breaking the chain-like propagation of the lipid peroxides [60, 61]. Moreover, an important claim by the EFSA [14], based on scientific reports, supported the effectiveness of the ingestion of VOO phenolics at a concentration evaluated to be 5 mg/day in protecting against oxidative stress. The most extensively investigated VOO phenolics are hydroxytyrosol (HT) and oleuropein (OE), and their antioxidant potentialities have been extensively reported in the scientific literature. Indeed, many authors have indicated that HT and OE are potent scavengers of free radicals [21]. Also, HT and OE inhibit copper sulfate-induced oxidation of LDL, according to Visioli et al. [63] and Visioli et al. [64]. The protective effects of HT and OE have been demonstrated through the assessment of various markers of LDL oxidation, including the thiobarbituric acid-reacting substances (TBARS), lipid peroxides, and 4-hydroxynonenal (4-HNE) adducts [63, 64].

On the other hand, VOO phenolics enhance the endogen antioxidant system, thereby improving the antioxidant enzymes including catalase, superoxide dismutase (SOD), and glutathione peroxidase (GPx) [65]. According to the previous report by Chashmi et al. [66], these effects are mainly attributed to HT, which increases SOD and glutathione (GSH) activities. In the same context, Pérez-Bonilla et al. [67] found that oleocanthal (OA) could display a higher anti-oxidative effect against the free radical DPPH than the reference 2,6-di-tert-butyl-hydroxytoluene (BHT).

4.2.2 Anti-atherosclerosis activity

Considering the role of oxidative stress in the pathogenesis of atherosclerosis, previous studies evaluated the protective effect of VOO phenolics. Based on various studies performed *in vitro* and *in vivo*, phenolic compounds of VOO play a primordial role in the prevention of atherosclerosis through their inhibition of LDL oxidation [68, 69, 70]. Many studies have shown that VOO phenolic compounds can bind to LDL and thus protect them against oxidation [71]. In this respect, some studies have focused attention on the antioxidant power of VOO phenolics on plasma LDL. Plasma was incubated with various

VOO phenolics to simulate an in vivo situation. LDL was then isolated and submitted to the action of free radicals. The obtained results showed that HT and OE are more effective than tyrosol and ligstroside aglycone [72]. Another important risk factor for the onset of atherosclerosis is a high blood concentration of cholesterol. Benkhalti et al. [73] reported that VOO phenolics can interfere with 3-hydroxy3-methylglutaryl (HMG)-CoA reductase, a key enzyme involved in the synthesis of cholesterol.

4.2.3 Cardioprotective activity

Some clinical trials that have examined the effects of VOO on cardiovascular disease have indicated that the consumption of VOO can protect various aspects of cardiovascular disease [18, 74, 75]. According to Bullota et al. [76], VOO phenolics play a significant role in improving cardiovascular health in adherents to the MED diet. Besides, phenolics have been shown to improve vascular function and reduce inflammation in heart tissues [77, 78, 79].

4.2.4 Anti-cancer activity

Studies have corroborated the protective effect of olive oil against various forms of cancer. VOO consumption and the incidence of many forms of cancer, including colon, breast, and skin cancer, have been inversely correlated [47]. Recent studies attribute this effect to the inhibitory potential of phenolic compounds on tumor growth [80, 81, 82, 83]. These bioactive substances appear to be able to influence gene expression, reduce DNA damage caused by ROS, and alter tumor eicosanoid biosynthesis and cell signaling pathways. Among the phenolic compounds possessing anticancer activity, HT is of special interest. Deiana et al. [84] highlighted that the presence of HT reduced the biochemical effects of peroxynitrites, such as oxidizing methionine and protecting DNA against damage. Della Ragione et al. [85] studied the anticancer effect of HT on the proliferation of human promyelocytic leukemia cells and highlighted its anti-cancer potential. HT delivered orally was shown to decrease breast tumor volume [86]. According to García-Villalba et al. [87], HT and luteolin have been shown to exhibit anti-cancerous effects against breast cancer cell lines. In the same context, some scientific studies have found that lignans inhibit the growth of various types of tumors, including cutaneous, mammary, colonic, and pulmonary tumors [88]. According to Akl et al. [89], OA was found to suppress tumor cell growth in a model of breast cancer. More recently, López de las Hazas et al. [90] reported the potential apoptotic and antiproliferative effects of VOO phenolics.

4.2.5 Anti-inflammatory activity

Natural inhibitors of cyclooxygenase (COX), important enzymes involved in the management of inflammation, can be an effective therapeutic approach. Some studies have found that VOO phenolics have an inhibitory effect on COX [91, 92, 93]. The anti-inflammatory activity of HT is similar to that of aspirin, a drug that is well known for its inhibitory potential on COX activity [91]. As reported by Facchini et al. [94], HT has also been shown to decrease the activity of COX-2. OA was found to have a similar anti-inflammatory role to that of the drug Ibuprofen [95]. Procopio et al. [96] explained that the anti-inflammatory activity of OA could be achieved by inhibiting COX enzymes. Therefore, OA can play an effective role in anti-inflammatory activities. According to the studies of De La Puerta [92] and Barbaro et al. [97], OE has been found to inhibit 5-lipoxygenase (5-LOX) activity, an enzyme with an important role in inflammation.

4.2.6 Antimicrobial activity

VOO is a promising source of bioactives with antimicrobial properties; the majority of them act as antimicrobial agents against a wide range of pathogenic microorganisms (bacteria, fungi, viruses, and protozoa) [98]. Phenolic compounds found in VOO have been shown to possess antimicrobial potential against several strains of bacteria [37]. OE can inhibit the development and production of enterotoxin B by *Staphylococcus aureus*, the development of *Salmonella enteritidis*, and the germination of spores of *Bacillus cereus* [99, 100, 101]. OE and other olive oil phenolic compounds (*p*-hydroxybenzoic, vanillic, and *p*-coumaric acids) inhibit the proliferation of *Klebsiella pneumoniae*, *Escherichia coli*, and *Bacillus cereus* [102]. HT shows a highly antibacterial effect against *Pseudomonas syringae* pv *savastanoi* and *Corynebacterium michiganense* [103]. Hydroxytyrosol and oleuropein have been shown to inhibit the growth of *Staphylococcus aureus* and *Candida albicans* [98]. The dialdehydic form of the decarboxymethyl oleuropein aglycon, OA, HT, and T have been shown to possess potent activity against several strains of bacteria responsible for intestinal and respiratory infections *in vitro* [104]. As reported by Romero et al. [105], Oleocanthal has also been found to inhibit the growth of *Helicobacter pylori*, a bacterium associated with peptic ulcer and gastric cancer development. According to Karaosmanoglu et al. [106], the antimicrobial activities of VOO phenolics were reported against three foodborne pathogenic bacteria: *Escherichia coli* O157: H7, *Listeria monocytogenes*, and *Salmonella enteritidis in vitro*.

4.2.7 Anti-diabetic activity

VOO bioactive compounds have been investigated for their contribution to the prevention and/or treatment of type 2 diabetes mellitus. Rigacci et al. [74] concluded that OA could prevent or retard the development of type 2 diabetes. Loizzo et al. [107] reported that phenol-rich VOO extracts from "Frantoio," "Ortice," and "Ortolana" produced in Campania (Italy) showed antidiabetic activity. HT has been found to prevent high-fat diet adverse effects such as obesity, diabetes, and insulin resistance. It also increases glucose consumption and inhibits adipogenesis [72, 108]. In fact, HT and OE could act as α -glucosidase and α -amylase inhibitors, according to Hadrach et al. [109]. Besides, Santangelo et al. [110] indicated that VOO phenolic compounds induced a direct effect on sugar metabolism and blood glucose levels. OE has been shown to protect β cells from cytotoxicity induced by amylin amyloids that are characteristic of type 2 diabetes [111]. In the same context, Collado-González et al. [24] reported that luteolin is the main phenolic compound that contributes to the antidiabetic activity of VOO.

More recently, Figueiredo-González et al. [112] revealed the antidiabetic potential of phenol-rich extracts from monovarietal VOOs (obtained from 'Cornicabra' and 'Picual' varieties, two of the most representative Spanish cultivars), through inhibition of α -amylase and α -glucosidase, key enzymes involved in the treatment of diabetes mellitus. In this respect, it is interesting to mention that many phenolic compounds have been described as inhibitors of key enzymes in the management of diabetic complications [24, 109, 113, 114, 115].

4.2.8 Neuroprotective activity

VOO phenolics are increasingly being investigated for their potential to manage the devastating central nervous system (CNS) [116]. HT has been shown to protect brain cells from free radical damage that contributes to neurodegenerative processes [117]. HT has also been shown to protect against mitochondrial dysfunction in rat brains [118]. It induces neuroprotection by increasing the activity of detoxifying enzymes, such as the GSH enzyme [119]. OE has demonstrated protective effects on dopaminergic neurons, implying possible roles in preventing or treating Parkinson's disease [97]. OE interacts with some of the biochemical processes behind the formation and deposition of β -amyloid plaques associated with Alzheimer's disease [97]. OA also promotes neuroprotection in cultured neuronal cells,

according to Barbaro et al. [97], by interfering with Amyloid aggregation, diminishing aggregate cytotoxicity, and counteracting related neuroinflammation. Another noteworthy study shows that VOO restores cholinergic deficits by restoring acetylcholine levels by inhibiting the two main forms of cholinesterases (BuChE and AChE) [112].

4.2.9 Protective activity against COVID-19

Comment [P9]: Covid

SARS-CoV-2 (severe acute respiratory syndrome coronavirus-2) is a hazardous infection that is causing global health problems. It causes coronavirus disease 2019 (COVID-19), which has a high rate of morbidity and fatality. The coronavirus disease (Covid-19) is a pathogenic virus caused by SARS-CoV-2 that has spread fast around the world. Antiviral bioactivities of phenolics have been identified against HCV, HIV, and other viruses. Protease inhibitors, such as phenolic compounds, have already been demonstrated to be particularly effective in suppressing virus-induced infections in multiple investigations [120, 121, 122, 123, 124]. There is currently insufficient data from in vivo studies indicating correlations between phenolics and ACE2 expression downregulation. Nonetheless, phenolics have recently been discovered to interact with SARS-CoV-2 viral proteins and their cellular targets [125]. As a result, the potential modification of immune response by these natural bioactive chemicals might be viewed as a positive element of the human body's defence against COVID-19. According to Paraiso et al. [125], phenolic compounds have been shown to have an effect on various stages of the SARS-CoV-2 life cycle. According to El-Missiry et al. [126], phenolics could be used as dietary adjuvants to target COVID-19. The health-promoting benefits of these natural substances on COVID-19 may be attributable to the body's anti-inflammatory and antioxidant defenses against viral infection, according to the researchers. Other approaches, such as targeting virus proteins and/or inhibiting cellular receptors, can still be effective in preventing virus entry and multiplication in host cells [126]. Although robust in vivo evidence for the efficacy of phenolics against the COVID-19 pandemic is still lacking, the preliminary findings can pave the way for more systematic and advanced experimental research into the efficacy of phenolic compounds derived from natural sources in the prevention and/or treatment of COVID-19. COVID-19 patients may benefit from natural bioactive compounds in their dietary control [127].

Since ancient times, virgin olive oil (VOO) has been used for therapeutic purposes as a valuable component of MED human diets. During COVID-19, VOO is regarded as a nutritional therapeutic immuno-enhancer since it can help keep the immune system healthy and minimize inflammation and oxidative stress, which can lead to a number of diseases and conditions, raising the likelihood of major COVID-19 instances [128].

Table 3. Some biological activities and health benefits of virgin olive oil and its phytochemicals

Source	Biological activity	Mechanisms	References
Phenolic extract of VOO	Antioxidant activity	Reduction of reactive oxygen species, lipid peroxidation, and NO [•] overproduction through modulating inducible nitric oxide synthase protein levels after H ₂ O ₂ induced oxidative stress	[129]
OE		Protection of L-O ₂ cells from H ₂ O ₂ -induced oxidative stress by boosting SOD, GPx and CAT expression	[130]
VOO		Improvement of antioxidant status in plasma and cell blood	[65]
VOO	Anti-obesity activity	Dietary 5% VOO increased serum HDL concentrations while decreasing triglyceride levels. Hydroxytyrosol modulated adipocyte gene expression.	[131] [132]
VOO		Increased immune system activity aids in immune resistance.	[133]
Phenolic compounds	Immunological resistance activity	White blood cell proliferation and the production of cytokines, which enhance immune resistance	[134]
VOO		Ethopone-induced nephrotoxicity was reduced by 2mL/Kg/day (150 mg per kg daily).	[135]
VOO		Inhibition of amikacin-induced	[136]

VOO + naringenin		nephrotoxicity. During a 45-day treatment period, the combined use of olive oil (1.25 ml/kg/day virgin olive oil) and naringenin (100 mg/kg/day naringenin) reduced the cyclosporine-induced (25 mg/kg/day cyclosporine) nephrotoxicity by improving renal functionality and lowering serum urea and creatinine concentrations in rats.	[137]
VOO	Anti-cancer activity	Protection against mammary carcinogenesis in both <i>in vivo</i> and <i>in vitro</i> studies.	[138]
VOO + phenolic compounds		VOO and its phenolic components, such as HT, OE, and OA, have been shown to have anti-cancer properties.	[139]
Secoiridoids		Secoiridoids found in olive oil have been demonstrated to have anti-cancer and anti-preventive properties against a variety of human malignancies.	[140]
VOO		VOO helps to prevent cancers of the breast, prostate, and digestive system.	[133]
MUFAs		MUFAs can potentially inhibit tumor growth and proliferation.	[138]

VOO		Olive oil's phenolic components, which scavenge free radicals and other reactive oxygen species, may help to slow or stop cancer progression.	[141]
HT			[142]
Secoiridoids		HT exerted an anti-proliferative impact on human colon cancer growth.	[140]
		In several cancer cell lines, VOO secoiridoids suppressed cell proliferation and triggered apoptosis.	[143]
OE		The mammary tumor MCF-7 cell line was used to demonstrate that OE had powerful anti-breast cancer effects.	[144]
VOO+ diallyl sulfide	Hepatoprotective activity	In mice, VOO combined with diallyl sulfide protects them from CCl ₄ induced acute liver damage.	[145]
VOO		In animal studies and cellular cultures, VOO and its phytochemicals exert a protective effect on the liver.	[146]
HT		HT helps to avoid liver steatosis and the mitochondrial dysfunction that comes with it, both of which are caused by a high-fat diet.	[147]
		Because of its strong antioxidant properties, a combination of VOO and	[148]


VOO + camel milk		camel milk showed hepatoprotective effects in mice against acetaminophen-induced hepatotoxicity at a single dose (500 mg/kg).	
VOO		Deltamethrin-induced hepatotoxicity was minimized when VOO was consumed.	[149] [150]
VOO	Anti-inflammatory activity	VOO (0.3 g/animal/day) showed cell membrane protecting and anti-inflammatory properties.	[151]
HT		In a mouse model of systemic inflammation, HT has anti-inflammatory and anti-oxidant properties.	[152]
OE		By lowering the secretion of COX inflammatory enzymes, OE has the power to break the inflammatory cascade.	[153]
MUFAs	Cardioprotective activity	A higher concentration of MUFAs (such as oleic acid) benefited the male albino rats' cardiovascular systems.	[154]
VOO		Olive oil is linked to a lower risk of cardiovascular disease and stroke. With a daily intake of 20 to 30 g, the greatest effect might be reached.	[155]
HT	Anti-Neurodegenerative activity	In a mouse model of amyloid deposition, HT supplementation improves brain disease and recovers cognitive skills.	[156]

Phenolic compounds		Polyphenols from VOO can affect a different cellular pathway associated with the disease's onset and progression.	[157]
OE		In an <i>in vitro</i> model of Parkinson's disease, oleuropein has been proven to diminish cell damage, apoptosis, and oxidative stress in PC12 cells generated by 6-OHDA.	[158]
OE		On PC12 cells exposed to 150 mM 6-OHDA, OE (20 and 25 g/ml) showed neuroprotective effects.	[159]
VOO	Antidiabetic activity	In type 1 diabetes, VOO has been linked to reduced postprandial glycemia.	[160]
VOO		In patients with impaired fasting glucose, VOO improves post-prandial glycemic and lipid profiles.	[161]
HT	Antimicrobial activity	Inactivation of <i>S. aureus</i> as well as the enterotoxin A	[162]
VOO		<i>In vitro</i> antimicrobial activities of the <i>Listeria monocytogenes</i> DSM 20600, <i>Staphylococcus aureus</i> DSM 20231, <i>Escherichia coli</i> DSM 30083, <i>Salmonella bongori</i> DSM13772, <i>Lactocaseibacillus</i>	[163]

		casei <i>Shirota</i> , <i>Limosilactobacillus reuteri</i> DSM 17938, <i>Lactacaseibacillus casei</i> RI4, and the yeast <i>Candida albicans</i> 3393	
--	--	--	--

VOO: virgin olive oil; OE: oleuropein; MUFAs: Monounsaturated fatty acids; HT: hydroxytyrosol; H₂O₂: Hydrogen peroxide; NO: nitric oxide; L-0₂: cells Human Normal Liver Cell Line; SOD: Superoxide dismutase; GPx: glutathione peroxidase; CAT: catalase; OA: oleocanthal; MCF-7: human breast cancer cell line; COX: cyclooxygenase; PC12 cells: pheochromocytoma cells; 6-OHDA: 6-Hydroxydopamine.

UNDER PEER REVIEW



4.3. Bioavailability

The bioavailability of VOO bioactive compounds refers to the degree to which they are extracted from this food matrix and absorbed by the body [164]. Bioavailability is essential in determining the beneficial health effects. Indeed, achieving an effect in specific tissues or organs depends on the bioavailability of VOO bioactive compounds. The scientific community has looked at the metabolic fate of phenolic substances after consumption. Phenolics are known to have strong metabolism in phases I and II, during which they are hydrolyzed (phase I) and then conjugated (phase II) into their glucuronidated, methylated, and sulfated forms, which are then absorbed [165].

The majority of studies regarding the bioavailability of VOO bioactive compounds have focused on phenolics. Some studies have shown that HT and T are absorbed after ingestion in a dose-dependent manner [62, 63, 166]. Vissers et al. [167] reported that the absorption of administered ligstroside-aglycone, HT, T, and OE aglycone was as high as 55–66% in human subjects. According to Vissers et al. [167], the polarities of the phenolics seemed to play a key role in the absorption of these compounds. In this context, Manach et al. [165] explained that the absorption of T and HT occurs via passive diffusion. However, OE-glycoside may be absorbed via a different mechanism to T and HT. Edgecombe et al. [168] explained that OE-glycoside may diffuse through the lipid bilayer of the epithelial cell membrane and be absorbed via a glucose transporter.

On the other hand, some work has been conducted to evaluate the quantity of VOO phenolics excreted. A smaller quantity of HT and T was recovered in urine (5%–16%) [167]. Casas et al. [18] revealed that approximately 24% of administered T was excreted. Visioli et al. [62] reported that the excretion of administered HT and T was evaluated at 30–60% and 20–22% of the total ingested by human subjects, respectively. These findings demonstrate that humans absorb a significant quantity of the dietary VOO phenolics.

5. Other bioactive compounds and their biological activities

5.1. Tocopherols

Tocopherols are hetero acid compounds with a high molecular weight that have been designated as α -, β -, γ - and δ -tocopherols. 90 % of the tocopherols in VOO are α -tocopherol, which has a concentration of up to 300 mg/kg, while others (β -, γ -, and δ -) are present in small amounts, up to 25 ppm [44, 169]. Tocopherols are particularly important bioactive constituents in VOO mainly due to their antioxidative effects [170]. They can act as antioxidants by two mechanisms. The first is a chain-breaking electron donor mechanism in which tocopherol donates its hydrogen atom to lipid radicals. The second is a chain-breaking acceptor mechanism that includes singlet oxygen scavenging or quenching and inhibits oxidation induced by excited singlet oxygen. High levels of α -tocopherol prevent the oxidation of polyunsaturated fatty acids in plasma lipoproteins, which causes atherosclerosis and modifies numerous pathways implicated in atherogenesis prevention [171]. Furthermore, α -tocopherol has been shown to inhibit prostate cancer cell growth in animal model experiments [172], decrease the progression of Alzheimer's disease [173], and protect against upper respiratory tract infections [174]. Fig. 5 shows the chemical structure of α -tocopherol.




Fig 5. Structure of α -tocopherol

5.2. Carotenoids

Carotenoid pigments are responsible for the color of olive oil. Their content does not usually exceed 10 mg/kg, with lutein being the major carotenoid, followed by β -carotene. They are substances with antioxidant activity [170]. In this context, β -Carotene is known to be a potent singlet oxygen quencher and free radical scavenger.

5.3. Hydrocarbons

Hydrocarbons are always present in VOO. Squalene is the most important hydrocarbon of VOO. Therefore, VOO can be considered as significant dietary source of squalene [170]. It is composed of approximately 0.7% squalene [171]. The biological activities of squalene are well-known. Indeed, the high squalene level of VOO, when compared with other foods, may account for the low incidence of certain tumors in the Mediterranean area [172, 173]. In vitro and in vivo animal model studies have shown inhibition of chemically induced colon and lung cancer formation by squalene [175, 176]. Squalene appears to be critical for reducing free radical oxidative damage. According to Kelly [177] and Huang et al. [178], it seems to function in the skin as a quencher of singlet oxygen, thus inhibiting lipid peroxidation due to exposure to UV light and other factors of oxidative stress. The chemical structure of squalene is provided in Fig. 6.




Fig .6. Structure of squalene

5.4. Phytosterols

VOO has between 1000 and 2300 ppm of sterols, with β -sitosterol being the most abundant [169]. According to several epidemiological investigations [179, 180], phytosterols have been shown to have immunological and anticancer properties in animal models, both in vitro and in vivo. Beta-sitosterol substantially suppressed the tumor growth of a human colon cancer cell line [181]. Epidemiological investigations linking dietary phytosterol intake to a lower risk of a variety of diseases corroborated these findings. In the same vein, multiple epidemiological studies suggest that having them in one's diet is linked to a lower risk of common malignancies like colon, breast, and prostate cancer [170].

Furthermore, phytosterols act as cholesterol-lowering agents by reducing cholesterol absorption [170]. Consumption of phytosterol may also increase the activity of antioxidant enzymes, reducing oxidative stress [180]. Fig. 7 shows the chemical structure of β -sitosterol.




Fig. 7. Structure of β -sitosterol

5.5. Triterpenic Acids

Oleanolic acid, maslinic acid, ursolic acid, and betulinic acid are triterpenic acids that are present in small amounts. Prades et al. [182] indicated that these acids could act as anti-inflammatory, antioxidant, antihypertensive, antihyperlipidemic, antidiabetic, antiviral, and antitumoral agents. Additionally, some studies have reported their neuroprotective effect. Additionally, it has been described that oleanolic acid protects the brain during hypoxic injury in rats and that ursolic acid protects the brain against ischemic injury in mice via Nrf2 activation [183, 184, 185]. According to Andrikopoulos et al. [186], oleanolic acid was described as a protective agent against LDL oxidation. Oleanolic acid is also involved in atherosclerosis protection, with antihyperlipidemic effects in animal models. Previous studies of isolated oleanolic acid describe its antitumoral activity.

Maslinic acid strongly inhibits *in vitro* LDL oxidation. Maslinic acid has previously been shown to have cardioprotective properties. This chemical may also protect against atherosclerosis by lowering insulin resistance in a mouse model, with potential antioxidant and hypoglycemic effects. Different research has focused on different aspects of the role that maslinic acid appears to play in cancer [187, 188, 189].

6. Critical perspective

The data allows us to show that VOO has a preventive impact on a variety of diseases. An overview of the literature reveals that the vast majority of the stated research works on VOO biological activities were carried out *in vitro* and *in vivo* models. Similarly, while research on VOO phytochemical substances is promising, the majority of their protective properties have only been established *in vitro* and *in vivo* models. However, it is critical that the investigations of this matrix and its phytochemicals continue to follow the proper phases of efficacy and safety testing in more *in vivo* models and eventually clinical trials in order to

validate this knowledge for the development of functional foods and nutraceuticals. Furthermore, more research is required to assess the precise situation. Furthermore, more research is required to determine the precise mechanism of action of these chemicals. According to the experimental investigations described in this study, hydrocarbons and triterpenic acids have not been researched as completely as phenolic chemicals and fatty acids.

7. Conclusion

In terms of culture and health, olive oil consumption is a significant advantage for many countries. Consuming VOO, which is high in phenolic compounds and other physiologically active substances, may help to lower the risk of developing a variety of diseases. The quest for natural, biologically active substances is becoming increasingly popular, and academics are becoming increasingly interested in the subject. Dietary management as a means of preventing disease is a major public health concern around the world. VOO has been proven to be a significant source of bioactive chemicals with a variety of biological effects in both in vitro and in vivo experimental models. The health benefits outlined in this review lead us to believe that virgin olive oil is one of the healthiest foods available.

The potential health benefits gained from VOO as a possible candidate in developing pharmaceutical preparations and nutraceutical or functional foods for numerous pathological conditions are supported by the health benefits outlined in this article. This hypothesis could lead to further research concentrating on more *in vivo* and, eventually, clinical trials. Furthermore, more thorough research on the mechanisms of action and efficacy is required.

REFERENCES

1. IOC, Production data. 2019/2020.
2. Perez-Martinez P, Garcia-Rios A, Delgado-Lista J, Perez-Jimenez F, Lopez-Miranda J. Mediterranean Diet Rich in Olive Oil and Obesity, Metabolic Syndrome and Diabetes Mellitus. *Curr. Pharm. Des.* 2011;17(8):769-777.
3. IOC, Consumption data. 2019/2020.
4. IOC, 2019. Trade standard applying to olive oils and olive pomace oils.
5. Owen RW, Giacosa A, Hule WE, Spiegelhader B, Bartsch H. Phenolic compounds and squalene in olive oils: the concentration and antioxidant potential of total phenols, simple phenols, secoroids, lignans and squalene. *Food Chem. Toxicol.* 2000;38(8):647-659.
6. Luisi MLE, Lucarini L, Biffi B, Rafanelli E, Pietramellara G, Durante M, Vidali S, Provensi G, Madiati S, Gheri CF, Masini E, Ceccherini MT. Effect of Mediterranean diet enriched in high quality extra virgin olive oil on oxidative stress, inflammation and gut microbiota in obese and normal weight adult subjects. *Front. Pharmacol.* 2019;10:1366.
7. Mazzocchi A, Leone L, Agostoni C, Pali-Schöll I. The secrets of the Mediterranean diet. Does [only] olive oil matter? *Nutrients*; 2019;11,2941.

8. Morello JR, Romero MP, Motilva MJ. Effect of the maturation process of the olive fruit on the phenolic fraction of drupes and oils from Arbequina, Farga and Morrut cultivars. *J. Agr. Food Chem.* 2004;52:6002–6009.
9. Soríguer F, Rojo-Martínez G, Goday A, Bosch-Comas A, Bordiù E, Caballero-Díaz F, et al. Olive oil has a beneficial effect on impaired glucose regulation and other cardiometabolic risk factors. Di@bet.es study. *Eur. J. Clin. Nutr.* 2013;67 (9):911–916.
10. Estruch R, Ros E, Salas-Salvadó J, Covas MI, Corella D, Arós F, Gómez-Gracia E, Ruiz-Gutiérrez V, Fiol M, Lapetra J, Lamuela-Raventos RM, Serra-Majem L, Pintó X, Basora J, Muñoz MA, Sorlí JV, Martínez JA, Fitó M, Gea A, Hernán MA, Martínez-González MA. Primary prevention of cardiovascular disease with a Mediterranean diet supplemented with extra-virgin olive oil or nuts. *N. Engl. J. Med.* 2018;378:e34.
11. Olalla J, García de Lomas JM, Chueca N, Pérez-Stachowski X, De Salazar A, Del Arco A, Plaza-Díaz J, De la Torre J, Prada JL, García-Alegria J, Fernández-Sánchez F, García F. Effect of daily consumption of extra virgin olive oil on the lipid profile and microbiota of HIV-infected patients over 50 years of age. *Medicine.* 2019;98:e17528.
12. Uttara B, Singh AV, Zamboni P, Mahajan RT. Oxidative stress and neurodegenerative diseases: a review of upstream and downstream antioxidant therapeutic options. *Curr. Neuropharmacol.* 2009;7(1):65-74.
13. Tan BL, Norhaizan ME, Liew W-P, Rahman HS. Antioxidant and oxidative stress: A mutual interplay in age-related diseases. *Front. Pharmacol.* 2018;9:1162.
14. EFSA. Scientific Opinion on the substantiation of a health claim related to polyphenols in olive and maintenance of normal blood HDL-cholesterol concentrations (ID 1639, further assessment) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. *EFSA J.* 2012;10(8):2848.
15. Jimenez-Lopez C, Carpena M, Lourenço-Lopes C, Gallardo-Gomez M, Lorenzo JM, Barba FJ, Prieto MA, Simal-Gandara J. Bioactive compounds and quality of extra virgin olive oil. *Foods.* 2020;9:1014.
16. Caramia G, Gori A, Valli E, Cerretani L. Virgin olive oil in preventive medicine: From legend to epigenetics. *Eur. J. Lipid Sci. Technol.* 2012;114:375-388.
17. European Commission Regulation EC No. 432/2012 establishing a list of permitted health claims made on foods, other than those referring to the reduction of disease risk and to children's development and health. *Off. J. Eur. Union.* 2012;L136:1–40.
18. Casas R, Sacanella E, Urpí-Sardà M, Chiva-Blanch G, Ros E, Martínez-González MA, Covas MI, Lamuela-Raventos RM, Salas-Salvadó J, Fiol M, Arós F, Estruch R. The effects of the Mediterranean diet on biomarkers of vascular wall inflammation and plaque vulnerability in subjects with high risk for cardiovascular disease. A randomized trial. *PLoS One.* 2014;9(6):e100084.
19. Hervet-Hernández D, García OP, Rosado JL, Goñi I. The contribution of fruits and vegetables to dietary intake of polyphenols and antioxidant capacity in a Mexican rural diet: Importance of fruit and vegetable variety. *Food Res. Int.* 2011;44:1182-1189.

20. Mayneris-Perxachs J, Sala-Vila A, Chisaguano M, Castellote AI, Estruch R, Covas MI, Fitó M, Salas-Salvadó J, Martínez-González MA, Lamuela Raventós R, Ros E, López-Sabater MC. Effects of 1-year intervention with a Mediterranean diet on plasma fatty acid composition and metabolic syndrome in a population at high cardiovascular risk. *PLoS One*. 2014;9:e85202.
21. Sabatini N. Recent Patents in Olive Oil Industry: New Technologies for the Recovery of Phenols Compounds from Olive Oil, Olive Oil Industrial by-Products and Waste Waters. *Recent Pat. food, Nutr. Agric.* 2010;2(2):154-159.
22. Reboredo-Rodríguez P, Figueiredo-González M, González-Barreiro C, Simal-Gándara J, Salvador MD, Cancho-Grande B, Fregapane G. State of-the-art on functional virgin olive oils enriched with bioactive compounds and their properties. *Int. J. Mol. Sci.* 2017;18(3):668-195.
23. Sabatini N, Perri E, Rongai D. Olive Oil Antioxidants and Aging. In *Handbook of Food Bioengineering, Food Quality: Balancing Health and Disease*, Academic Press, 2018;145-157.
24. Collado-González J, Grosso C, Valentão P, Andrade PB, Ferreres F, Durand T, Gil-Izquierdo Á. Inhibition of α -glucosidase and α -amylase by Spanish extra virgin olive oils: The involvement of bioactive compounds other than oleuropein and hydroxytyrosol. *Food Chem.* 2017;235,298–307.
25. Ennouri K, Ben Hlima H, Ben Ayed R, Ben Braïek O, Mazzarello O, Ottaviani E, Mallouli L, Smaoui S. Assessment of Tunisian virgin olive oils via synchronized analysis of sterols, phenolic acids, and fatty acids in combination with multivariate chemometrics. *Eur. Food Res. Technol.* 2019;245(6):1811–1824.
26. Ben Youssef N, Abaza L, Ouni Y, Salma Naeit M, Debbech N, Abdelly C, Zarrouk M. Influence of the Site of Cultivation on Chétoui Olive (*Olea europaea* L.) Oil Quality, *Plant Prod. Sci.* 2002;15(3):228-237.
27. Sicari V, Leporini M, Giuffré AM, Aiello F, Falco T, Pagliuso MT, Ruffolo A, Reitano A, Romeo R, Tundis R, Loizzo MR. Quality parameters, chemical compositions and antioxidant activities of Calabrian (Italy) monovarietal extra virgin olive oils from autochthonous (Ottobratica) and allochthonous (Coratina, Leccino, and Nocellara Del Belice) varieties. *J. Food Meas. Charact.* 2020;15:363-375.
28. Montaña A, Hernández M, Garrido I, Llerena JL, Espinosa F. Fatty Acid and Phenolic Compound Concentrations in Eight Different Monovarietal Virgin Olive Oils from Extremadura and the Relationship with Oxidative Stability. *Int. J. Mol. Sci.* 2016;17(11):1960.
29. Bermudez B, Lopez S, Ortega A, Varela LM, Pacheco YM, Abia R, Muriana FJG. Oleic Acid in Olive Oil: From a Metabolic Framework Toward a Clinical Perspective. *Curr. Pharm. Des.* 2011;17(8):831-843.
30. Kien CL, Bunn JY, Stevens R, Bain J, Ikayeva O, Crain K, Koves TR, Muoio DM. Dietary intake of palmitate and oleate has broad impact on systemic and tissue lipid profiles in humans. *Am. J. Clin. Nutr.* 2014. 99(3),436-45.

31. Montserrat-De La Paza S, Naranjo MC, Lopez S, Abia R, Muriana F-JG, Bermudez B. Olive oil, compared to a saturated dietary fat, has a protective role on atherosclerosis in niacin-treated mice with metabolic syndrome. *J. Funct. Foods.* 2016;26:557-564.
32. Lamy S, Ben Saa A, Zgheib A, Annabi B. Olive oil compounds inhibit the paracrine regulation of TNF- α -induced endothelial cell migration through reduced glioblastoma cell cyclooxygenase-2 expression. *J. Nutr. Biochem.* 2016;27:136-145.
33. Ryan D, Robards K. Phenolic compounds in olives. *Crit. Rev.* 1998;123:31R-44R.
34. Cicerale S, Conlan XA, Sinclair AJ, Keast SJ. Chemistry and health of olive oil phenolics. *Crit. Rev. Food Sci. Nutr.* 2009;49(3):18-36.
35. Young DP, Kong DX, Zhang HY. Pharmacological effects of olive oil phenols. *Food. Chem.* 2007;104:1269-1271.
36. Boskou D. Olive oil—Constituents, Quality, Health Properties and Bioconversions. 1st ed InTech, Croatia. 2012.
37. Cicerale S, Lucas LJ, Keast RSJ. Antimicrobial, antioxidant and anti-inflammatory phenolic activities in extra virgin olive oil. *Curr. Opin. Biotechnol.* 2012;23 (2):129-135.
38. Brenes M, Hidalgo FJ, García A, Rios JJ, García P, Zamora R, Garrido A. Pinoresinol and 1-acetoxypinoresinol, two new phenolic compounds identified in olive oil. *J. Am. Oil Chem. Soc.* 2000;77(7):715-720.
39. Vierhuis E, Servili M, Baldioli M, Schols HA, Voragen AGJ, Montedoro GF. Study of the effect of enzyme treatment during the mechanical extraction of olive oil on phenolic compounds and polysaccharides. *J. Agric. Food Chem.* 2001;49(3):1218-1223.
40. Inglese P, Famiani F, Galvano F, Servili M, Esposito S, Urbani S. Factors Affecting Extra-Virgin Olive Oil Composition. In *Horticultural Reviews*, J. Janick (Ed.). 2010
41. El Riachy M, Priego-Capote F, León L, Luque De Castro MD, Rallo R. Virgin olive oil phenolic profile and variability in progenies from olive crosses. *J. Sci. Food Agric.* 2012;92(12):2524-2533.
42. Negro C, Aprile A, Luvisi A, Nicoli F, Nutricati E, Vergine M, Miceli A, Blando F, Sabella E, De Bellis L. Phenolic Profile and Antioxidant Activity of Italian Monovarietal Extra Virgin Olive Oils. *Antioxidants.* 2019;8(161):1-14.
43. Franco MN, Galeano-Díaz T, Sánchez J, De Miguel C, Martín-Vertedor D. Total Phenolic Compounds and Tocopherols Profiles of Seven Olive Oil Varieties Grown in the South-West of Spain. *J. Oleo Sci.* 2014;63(2): 1-11.
44. Boskou D, Blekas G, Tsimidou M. In: Boskou, D. (Ed.), *Olive Oil Chemistry, Properties, Health Effects*. AOCS Press, Thessaloniki, 2006;41-72.
45. Covas M.-I, Ruiz-Gutierrez V, De La Torre R, Kafatos A, Lamuela-Raventos RM, Osada J, Owen RW, Visioli F. Minor components of olive oil: evidence to date of health. *Nutr. Rev.* 2006;64(10):20-30.

46. Visioli F, Poli A, Galli C. Antioxidant and other biological activities of phenols from olives and olive oil. *Med. Res. Rev.* 2002;22(1):65-75.
47. Parkinson L, Cicerale S. The health benefiting mechanisms of virgin olive oil phenolic compounds. *Molecules.* 2016;21(12):1734.
48. Reboredo-Rodríguez P, Varela-López A, Forbes-Hernández TY, Gasparri M, Afrin S, Cianciosi D, Zhang J, Manna PP, Bompadre S, Quiles JL, Battino M, Giampieri F. Phenolic Compounds Isolated from Olive Oil as Nutraceutical Tools for the Prevention and Management of Cancer and Cardiovascular Diseases. *Int. J. Mol. Sci.* 2018;19(8):2305.
49. Romani A, Ieri F, Urciuoli S, Noce A, Marrone G, Nediani C, Bernini R. Health Effects of Phenolic Compounds Found in Extra-Virgin Olive Oil, By-Products, and Leaf of *Olea europaea* L. *Nutrients.* 2019;11(8):1776.
50. Miro Casas E, Albadalejo MF, Covas Planells MI, Colomer FM, Lamuela Raventos RM, De La Torre Fornell R. Tyrosol bioavailability in humans after ingestion of virgin olive oil. *Clin. Chem.* 2001;47:341–343.
51. Tuck KL, Hayball PJ. Major phenolic compounds in olive oil: metabolism and health effects. *J. Nutr. Biochem.* 2002;13:636–644.
52. De la Torre R. Bioavailability of olive oil phenolic compounds in humans. *Inflammopharmacology.* 2008;16(5):245-247.
53. Notarnicola M, Pisanti S, Tutino V, Bocale D, Rotelli MT, Gentile A, Memeo V, Bifulco M, Perri E, Caruso MG. Effects of olive oil polyphenols on fatty acid synthase gene expression and activity in human colorectal cancer cells. *Genes Nutr.* 2010;6:63-69.
54. Ricciutelli M, Marconi S, Boarelli MC, Caprioli G, Sagratini G, Ballini R, Fiorini D. Olive oil polyphenols: A quantitative method by high-performance liquid-chromatography-diode-array detection for their determination and the assessment of the related health claim. *J. Chromatogr. A.* 2017;20(148):53-63.
55. Gavahian M, Khaneghah AM, Lorenzo JM, Munekata PE, Garcia-Mantrana I, Collado M, Meléndez-Martínez AJ, Barba FJ. Health benefits of olive oil and its components: Impacts on gut microbiota antioxidant activities, and prevention of non communicable diseases. *Trends Food Sci. Technol.* 2019;88:220-227.
56. Mosca A, Crudele A, Smeriglio A, Braghini MR, Panera N, Comparcola D, Alterio A, Sartorelli MR, Tozzi G, Raponi M, Trombetta D, Alisi A. Antioxidant activity of Hydroxytyrosol and Vitamin E reduces systemic inflammation in children with paediatric NAFLD. *Dig. Liver Dis.* 2020;53(9):1154-1158.
57. Halliwell B, Gutteridge JM. Role of free radicals and catalytic metal ions in human disease: an overview. *Methods Enzymol.* 2010;186:1-85.
58. Halliwell B, Murcia MA, Chirico S, Aruoma OI. Free radicals and antioxidants in food and in vivo: what they do and how they work. *Crit. Rev. Food Sci. Nutr.* 1995;35(1-2):7-20.
59. Rice-Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic. Biol. Med.* 1996;20(7):933-956.

60. Carluccio MA, Siculella L, Ancora MA, Massaro M, Scoditti E, Storelli C, Visioli F, Distante A, De Caterina R. Olive oil and red wine antioxidant polyphenols inhibit endothelial activation: antiatherogenic properties of Mediterranean diet phytochemicals. *Arterioscler. Thromb. Vasc. Biol.* 2003; 23(4):622-629.
61. Weinbrenner T, Fitó M, De la Torre R, Saez GT, Rijken P, Tormos C, Coolen S, Albaladejo MF, Abanades S, Schroder H, Marrugat J, Covas M-I. Olive oils high in phenolic compounds modulate oxidative/antioxidative status in men. *J. Nutr.* 2004;134(9):2314-2321.
62. Visioli F, Galli C, Bornet F, Mattei A, Patelli R, Galli G, Caruso D. Olive oil phenolics are dose-dependently absorbed in humans. *FEBS Lett.* 2000;468:159-160.
63. Visioli F, Galli C, Plasmati E, Viappiani S, Hernandez A, Colombo C, Sala A. Olive phenol hydroxytyrosol prevents passive smoking-induced oxidative stress. *Circulation.* 2000;102(18):2169-2171.
65. Oliveras-López M-J, Bernáa G, Jurado-Ruiza E, López-García de la Serrana H, Martín F. Consumption of extra-virgin olive oil rich in phenolic compounds has beneficial antioxidant effects in healthy human adults. *J. Func. Foods.* 2014;10:475-484.
66. Chashmi NA, Emadi S, Khastar H. Protective effects of hydroxytyrosol on gentamicin induced nephrotoxicity in mice. *Biochem. Biophys. Res. Commun.* 2016;482(4):1427-1429.
67. Pérez-Bonilla M, Salido S, Van Beek TA, Altarejos J. Radical scavenging compounds from olive tree (*Olea europaea* L.) wood. *J. Agric. Food Chem.* 2014;62:144-151.
68. Marrugat J, Covas MI, Fitó M, Schroder H, Miro-Casas, Gimeno E, Lopez-Sabater MC, De la Torre R, Farre M. Effects of differing phenolic content in dietary olive oils on lipids and LDL oxidation — a randomized controlled trial. *Eur. J. Nutr.* 2004;43(3):140-147.
69. De la Torre-Carbot K, Chavez-Servin JL, Jauregui O, Castellote AI, Lamuela-Raventos RM, Fitó M, Covas MI, Munoz-Aguayo D, Lopez-Sabater MC. Presence of virgin olive oil phenolic metabolites in human low density lipoprotein fraction: determination by high-performance liquid chromatography–electrospray ionization tandem mass spectrometry. *Anal. Chim. Acta.* 2007;583(2):402-410.
70. De la Torre-Carbot K, Chavez-Servin JL, Jauregui O, Castellote AI, Lamuela-Raventos RM, Nurmi T, Poulsen HE, Gaddi AV, Kaikkonen J, Zunft HF, Kieseewetter H, Fitó M, Covas M-I, López-Sabater M-C. Elevated circulating LDL phenol levels in men who consumed virgin rather than refined olive oil are associated with less oxidation of plasma LDL. *J. Nutr.* 2010;140(3):501-508.
71. González-Santiago M, Fonollá J, Lopez-Huertas E. Human absorption of a supplement containing purified hydroxytyrosol, a natural antioxidant from olive oil, and evidence for its transient association with low-density lipoproteins. *Pharmacol. Res.* 2010;61(4):364-370.
72. Cao K, Xu J, Zou X, Li Y, Chen C, Zheng A, Li H, Li H, Szeto IM-Y, Shi Y, Long J, Liu J, Feng Z. Hydroxytyrosol prevents diet-induced metabolic syndrome and attenuates mitochondrial abnormalities in obese mice. *Free Radic. Biol. Med.* 2014;67:396-407.
73. Benkhalti F, Prost J, Paz E, Perez-Jimenez F, El modafar C, El Boustani E. Effects of feeding virgin olive oil or their polyphenols on lipid of rat liver. *Nutr. Res.* 2002;22(9):1067-1075.

74. Rigacci S, Stefani M. Nutraceutical properties of olive oil polyphenols. An itinerary from cultured cells through animal models to humans. *Int. J. Mol. Sci.* 2016;17(6):843.
75. Mehmood A, Usman M, Patil P, Zhao L, Wang C. A review on management of cardiovascular diseases by olive polyphenols. *Food Sci. Nutr.* 2020;8:4639-4655.
76. Bulotta S, Celano M, Lepore SM, Montalcini T, Pujia A, Russo D. Beneficial effects of the olive oil phenolic components oleuropein and hydroxytyrosol: focus on protection against cardiovascular and metabolic diseases. *J. Transl. Med.* 2014;12:219-241.
77. Poudyal H, Campbell F, Brown L. Olive Leaf Extract Attenuates Cardiac, Hepatic, and Metabolic Changes in High Carbohydrate-, High Fat-Fed Rats. *J. Nutr.* 2010;140(5):946-953.
78. Nocella C, Cammisotto V, Fianchini L, D'Amico A, Novo M, Castellani V, Stefanini L, Violi F, Carnevale R. Extra Virgin Olive Oil and Cardiovascular Diseases: Benefits for Human Health Endocrine, Metabolic & Immune Disorders. *Curr. Drug Targets.* 2018;18(1): 4-13.
79. Pintó X, Fanlo-Maresma M, Corbella E, Corbella X, Mitjavila MT, Moreno JJ, Casas R, Estruch R, Corella D, Bulló M, Ruiz-Canela M, Castañer O, Martínez JA, Ros E. A Mediterranean diet rich in extra-virgin olive oil is associated with a reduced prevalence of nonalcoholic fatty liver disease in older individuals at high cardiovascular risk. *J. Nutr.* 2019;149(1):1920-1929.
80. Budiyo A, Ahmed NU, Wu A, Bito T, Nikaido O, Osawa T, Ueda M, Ichihashi M. Protective effect of topically applied olive oil against photocarcinogenesis following UVB exposure of mice. *Carcinogenesis.* 2000;21(11):2085-90.
81. Thuy NT, He P, Takeuchi H. Comparative effect of dietary olive, safflower, and linseed oils on spontaneous liver tumorigenesis in C3H/He mice, *J. Nutr. Sci. Vitaminol.* 2001;47: 363-366.
82. Fabiani R, De Bartolomeo A, Rosignoli P, Servili M, Selvaggini R, Montedoro GF, Di Saverio C, Morrozi G. Virgin olive oil phenols inhibit proliferation of human promyelocytic leukemia cells (HL60) by inducing apoptosis and differentiation, *J. Nutr.* 2006;136(3):614-619.
83. Calahorra J, Martínez-Lara E, Granadino-Roldán JM, Martí JM, Cañuelo A, Blanco S, Oliver FJ, Siles E. Crosstalk between hydroxytyrosol, a major olive oil phenol, and HIF-1 in MCF-7 breast cancer cells. *Sci. Rep.* 2020;10:6361.
84. Deiana M, Aruoma OI, De Lourdes P Bianchi M, Spencer JPE, Kaur H, Halliwell B, Aeschbach R, Banni S, Assunta Dessi M, Corongiu FP. Inhibition of peroxynitrite dependent dna base modification and tyrosine nitration by the extra virgin olive oil-derived antioxidant hydroxytyrosol. *Free Radic. Biol. Med.* 1999;26(5/6):762-769.
85. Della Ragione F, Cucciolla V, Borriello A, Della Pietra V, Pontoni G, Racioppi L, Manna C, Galletti P, Zappia V. Hydroxytyrosol, a Natural Molecule Occurring in Olive Oil, Induces Cytochrome c-Dependent Apoptosis. *Biochem. Biophys. Res. Commun.* 2000;278:733-739.

86. Granados-Principal S, Quiles JL, Ramirez-Tortosa CL, Sanchez-Rovira P, Ramirez-Tortosa MC. Hydroxytyrosol: from laboratory investigations to future clinical trials. *Nutr. Rev.* 2006;68(4):191-206.
87. García-Villalba R, Carrasco-Pancorbo A, Oliveras-Ferraro C, Menendez JA, Segura-Carretero A, Fernandez-Gutierrez A. Uptake and metabolism of olive oil polyphenols in human breast cancer cells using nano-liquid chromatography coupled to electrospray ionization-time of flight-mass spectrometry. *J. Chromatogr.* 2012;B898:69-77.
88. López-Biedma A, Sánchez-Quesada C, Delgado-Rodríguez M, Gaforio JJ. The biological activities of natural lignans from olives and virgin olive oils: A review. *J. Funct. Foods.* 2016;26:36-47.
89. AKL MR, Ayoub NM, Mohyeldin MM, Busnena AA, Foudah AL, Liu Y-Y, El Sayed KA. Olive phenolics as c-met inhibitors: (-)-oleocanthal attenuates cell proliferation, invasiveness, and tumor growth in breast cancer models. *PLoS ONE* 2014;9(5): e97622.
90. López de las Hazas M-C, Piñol C, Macià A, Motilva M-J. Hydroxytyrosol and the colonic metabolites derived from virgin olive oil intake induce cell cycle arrest and apoptosis in colon cancer cells. *J. Agric. Food. Chem.* 2017;65(31):6467-6476.
91. Petroni A, Blasevich M, Salami M, Papini N, Montedorol GF, Galli C. Inhibition of platelet aggregation and eicosanoid production by phenolic components of olive oil. *Thromb. Res.* 1995;78(2):151-160.
92. De La Puerta R, Ruiz Gutierrez V, Hoult JR. Inhibition of leukocyte 5-lipoxygenase by phenolics from virgin olive oil. *Biochem. Pharmacol.* 1999;57(4):445-449.
93. Serrelli G, Deiana M. Extra virgin olive oil polyphenols: Modulation of cellular pathways related to oxidant species and inflammation in aging. *Cells.* 2020;9:478.
94. Facchini A, Cetrullo S, D'Adamo S, Guidotti S, Minguzzi M, Facchini A, Borzi RM, Flamigni F. Hydroxytyrosol prevents increase of osteoarthritis markers in human chondrocytes treated with hydrogen peroxide or growth-related oncogene α . *PLoS One.* 2016;11(10):e019724.
95. Beauchamp GK, Keast RSJ, Morel D, Lin J, Pika J, Han Q, Lee C-H, Smith AB, Breslin PAS. Phytochemistry: ibuprofen-like activity in extra-virgin olive oil. *Nat.* 2005;437(7055):45-46.
96. Procopio A, Alcaro S., Nardi, M., Oliverio, M., Ortuso, F., Sacchetta, P., et al. (2009). Synthesis, biological evaluation, and molecular modeling of oleuropein and its semisynthetic derivatives as cyclooxygenase inhibitors. *J. Agric. Food Chem.* 57(23), 11161–11167.
97. Barbaro B, Toietta G, Maggio R, Arciello M, Tarrocchi M, Galli A, Balsano C. Effects of the olive-derived polyphenol oleuropein on human health. *Int. J. Mol. Sci.* 2014;15(10): 18508-18524.
98. Bisignano G, Tomaino A, Lo Cascio R, Crisafi G, Uccella N, Saija A. On the *in-vitro* antimicrobial activity of oleuropein and hydroxytyrosol. *J. Pharm. Pharmacol.* 1999;51:971–974.

99. Fleming HP, Walter WM, Etchells JL. Antimicrobial Properties of Oleuropein and Products of Its Hydrolysis from Green Olives. *J. Appl. Microbiol.* 1973;26(5):777-782.
100. Walter WMJR, Fleming HP, Etchells JL. Preparation of Antimicrobial Compounds by Hydrolysis of Oleuropein from Green Olives. *J. Appl. Microbiol.* 1973;26(5):773-776.
101. Tassou CC, Nychas GJ, Board RG. Effect of phenolic compounds and oleuropein on the germination of *Bacillus cereus* T spores. *Biotechnol. Appl. Biochem.* 1991;13(2):231-237.
102. Aziz NH, Frag SE, Mousa LA, Abo-Zaid MA. Comparative antibacterial and antifungal effects of some phenolic compounds. *Microbios.* 1998;93:43-54.
103. Capasso R, Evidente A, Schivo L, Orru G, Marcialis MA, Cristinzio G. Antibacterial polyphenols from olive oil mill waste waters. *J. Appl. Bacteriol.* 1995;79(4):393-398.
104. Medina E, De Castro A, Romero C, Brenes M. Comparison of the concentrations of phenolic compounds in olive oils and other plant oils: correlation with antimicrobial activity. *J. Agric. Food. Chem.* 2006;54(14):4954-4961.
105. Romero C, Medina E, Vargas J, Brenes M, De Castro A. *In vitro* activity of olive oil polyphenols against *Helicobacter pylori*. *J. Agric. Food Chem.* 2007;55(3):680-686.
106. Karaosmanoglu H, Soyer F, Ozen B, Tokatli F. Antimicrobial and antioxidant activities of Turkish extra virgin olive oils. *J. Agric. Food Chem.* 2010;58(14):8238-8245.
107. Loizzo MR, Di Lecce G, Boselli E, Menichini F, Frega NG. Inhibitory activity of phenolic compounds from extra virgin olive oils on the enzymes involved in diabetes, obesity and hypertension. *J. Food Biochem.* 2011;35 (2):381-399.
108. Drira R, Sakamoto K. Modulation of adipogenesis, lipolysis and glucose consumption in 3T3-L1 adipocytes and C2C12 myotubes by hydroxytyrosol acetate: A comparative study. *Biochem. Biophys. Res. Commun.* 2013;440(4):576-581.
109. Hadrich F, Bouallagui Z, Junkyu H, Isoda H, Sayadi S. The α -glucosidase and α -amylase enzyme inhibitory of hydroxytyrosol and oleuropein. *J. Oleo Sci.* 2015;64(8):835-843.
110. Santangelo C, Filesi C, Varí R, Scazzocchio B, Filardi T, Fogliano V, D'Archivio M, Giovannini C, Lenzi A, Morano S, Masella R. Consumption of extra-virgin olive oil rich in phenolic compounds improves metabolic control in patients with type 2 diabetes mellitus: a possible involvement of reduced levels of circulating visfatin. *J. Endocrinol. Invest.* 2016;39(11):1295-1301.
111. Wu L, Velander P, Liu D, Xu B. Olive component oleuropein promotes β -cell insulin secretion and protects β -cells from amylin amyloid-induced cytotoxicity. *Biochemistry.* 2017;56(38):5035-5039.
112. Figueiredo-González M, Reboredo-Rodríguez P, González-Barreiro C, Carrasco-Pancorbo A, Simal-Gándara J, Cancho-Grande B. Nutraceutical Potential of Phenolics from 'Brava' and 'Mansa' Extra-Virgin Olive Oils on the Inhibition of Enzymes Associated to Neurodegenerative Disorders in Comparison with Those of 'Picual' and 'Cornicabra'. *Molecules.* 2018;23(4):722.

113. Proença C, Freitas M, Ribeiro D, Oliveira EFT, Sousa JLC, Tomé SM, Ramos MJ, Silva AMS, Fernandes PA, Fernandes E. α -Glucosidase inhibition by flavonoids: An *in vitro* and silico structure activity relationship study. *J. Enzyme Inhib. Med. Chem.* 2017;32(1):1216–1228.
114. Zhang Y, Li X, Sun WL, Xing Y, Xiu ZI, Zhuang CI, Dong YS. Dietary flavonoids and acarbose synergistically inhibit α -glucosidase and lower postprandial blood glucose. *J. Agr. Food Chem.* 2017; 65(38):8319-8330.
115. Figueiredo-González M, Reboredo-Rodríguez P, González-Barreiro C, Carrasco Pancorbo A, Cancho-Grande B, Simal-Gándara J. The involvement of phenolic-rich extracts from Galician autochthonous extra-virgin olive oils against the α -glucosidase and α -amylase inhibition. *Food Res. Int.* 2019;116:447-454.
116. Salis C, Papageorgiou L, Papakonstantinou E, Hagidimitriou M, Vlachakis D. Olive Oil Polyphenols in Neurodegenerative Pathologies. *Adv. Exp. Med. Biol.* 2020;1195:77-91.
117. González-Correa JA, Navas MD, Lopez-Villodres JA, Trujillo M, Espartero JL, De La Cruz JP. Neuroprotective effect of hydroxytyrosol and hydroxytyrosol acetate in rat brain slices subjected to hypoxia-reoxygenation. *Neurosci. Lett.* 2008;446(2-3):143-146.
118. Soni M, Prakash C, Sehwal S, Kumar V. Protective effect of hydroxytyrosol in arsenic-induced mitochondrial dysfunction in rat brain. *J. Biochem. Mol. Toxicol.* 2017;31(7): e21906.
119. Cabrerizo S, De La Cruz JP, López-Villodres JA, Muñoz-Marína J, Guerrero A, Reyes JJ, Labajos MT, González-Correa JA. Role of the inhibition of oxidative stress and inflammatory mediators in the neuroprotective effects of hydroxytyrosol in rat brain slices subjected to hypoxia reoxygenation. *J. Nutr. Biochem.* 2013;24(12):2152-2157.
120. Liu W, Zhou P, Chen K, Ye Z, Liu F, Li X, He N, Wu Z, Zhang Q, Gong X, Tang Q, Du X, Ying Y, Xu X, Zhang Y, Liu J, Li Y, Shen N, Couban RJ, Ibrahim QI, Guyatt G, Zhai S. Efficacy and safety of antiviral treatment for COVID-19 from evidence in studies of SARS-CoV-2 and other acute viral infections: a systematic review and meta-analysis. *CMAJ.* 2020;192(27):E734-E744.
121. Liu A-L, Liu B, Qin H-L, Lee SM, Wang Y-T, Du G-H. Anti-influenza virus activities of flavonoids from the medicinal plant *Elsholtzia rugulosa*. *Planta Med.* 2008;74(8):847-851.
122. Mohammadi N, Shaghghi N. Inhibitory Effect of Eight Secondary Metabolites from Conventional Medicinal Plants on COVID_19 Virus Protease by Molecular Docking Analysis. *Bio. Med. Chem.* 2020;1:1-11.
123. Shaghghi N. Molecular Docking Study of Novel COVID-19 Protease with Low Risk Terpenoid Compounds of Plants. *Bio. Med. Chem.* 2020;1:1-9.
124. Mukherjee PK, Efferth T, Das B, Kar A, Ghosh S, Singha S, Debnath P, Sharma N, Bhardwaj PK, Haldar PK. Role of medicinal plants in inhibiting SARS-CoV-2 and in the management of post-COVID-19 complications. *Phytomedicine.* 2022;98:153930.
125. Paraiso IL, Revel JS, Stevens JF. Potential use of polyphenols in the battle against COVID-19. *Curr. Opin. Food Sci.* 2020;32:149-155.

126. EL-Missiry M, Fekri A, Kesar LA, Othman AI. Polyphenols are potential nutritional adjuvants for targeting COVID -19. *Phytother. Res.* 2020;35(6):2879-2889.
127. Ruhidas S, Gayen R, Gopal SK. Nutritional Management of Covid-19 Disease. *J. Sci. Technol.* 2021;6(4):84-92.
128. Almoselhy RIM. Extra Virgin Olive Oil as Nutritional Therapeutic Immuno-Enhancer. *Int. Journal of Fam. Studies, Food Sci. and Nutr. Health* 2021;4(2):26- 45.
129. Bartolomei M, Bollati C, Bellumori M, Cecchi L, Cruz-Chamorro I, Santos-Sánchez G, Ranaldi G, Ferruzza S, Sambuy Y, Arnoldi A, Mulinacci N, Lammi C. Extra Virgin Olive Oil Phenolic Extract on Human Hepatic HepG2 and Intestinal Caco-2 Cells: Assessment of the Antioxidant Activity and Intestinal Trans-Epithelial Transport. *Antioxidants* 2021;10:118.
130. Shi C, Chen X, Liu Z, Meng R, Zhao X, Liu Z, Gyu Na. Oleuropein Protects L-02 Cells against H₂O₂-Induced Oxidative Stress by Increasing SOD1, GPx1 and CAT Expression. *Biomed. Pharmacother.* 2017;85:740–748.
131. Scoditti E, Massaro M, Garbarino S, Toraldo DM. Role of Diet in Chronic Obstructive Pulmonary Disease Prevention and Treatment. *Nutrients.* 2019;11(6):1357.
132. Scoditti E, Carpi S, Massaro M, Pellegrino M, Polini B, Carluccio MA, Wabitsch M, Verri T, Nieri P, De Caterina R.. Hydroxytyrosol Modulates Adipocyte Gene and miRNA Expression under Inflammatory Condition. *Nutrients.* 2019;11 (10):2493.
133. Psaltopoulou T, Kostis RI, Haidopoulos D, Dimopoulos M, Panagiotakos DB. Olive Oil Intake Is Inversely Related to Cancer Prevalence: a Systematic Review and a Meta-Analysis of 13800 Patients and 23340 Controls in 19 Observational Studies. *Lipids Health Dis;* 201;10:127.
134. Ding S, Jiang H, Fang J. Regulation of Immune Function by Polyphenols. *J. Immunol. Res.* 2018.;2018:1264074.
135. Mokhtari T, Hussein Osman H.-E, El-Meghawry El-Kenawy A, Dashti N. Ameliorative Effect of virgin Olive Oil against Nephrotoxicity Following Sub-chronic Administration of Ethephon in Male Rats. *J. Traditional Complement. Med.* 2020;10:487-495.
136. Abdelgayoum AA-G, Abdelrahman AA-H, Ibrahim AG, Ibraheem MA. The ameliorative effects of virgin olive oil and olive leaf extract on amikacin-induced nephrotoxicity in the rat. *Toxicol. Rep.* 2015;30(2):1327-1333.
137. Elshama SS, Osman H-EH, El-Kenawy E-MA. Renoprotective Effects of Naringenin and Olive Oil against Cyclosporine- Induced Nephrotoxicity in Rats. *IJT.* 2016;10(5):27-37.
138. Moral R, Escrich E. Influence of Olive Oil and Its Components on Breast Cancer: Molecular Mechanisms, *Molecules.* 2022;27(2):477.
139. Sain A, Sahu S, Naskar D. Potential of olive oil and its phenolic compounds as therapeutic intervention against colorectal cancer: A comprehensive review. *Br. J. Nutr.* 2021;2:1-17.

140. Emma MR, Augello G, Di Stefano V, Azzolina A, Giannitrapani L, Montalto G, Cervello M, Cusimano A. Potential Uses of Olive Oil Secoiridoids for the Prevention and Treatment of Cancer: A Narrative Review of Preclinical Studies. *Int. J. Mol. Sci.* 2021;22(3):1234.
141. Wahle KW, Brown I, Rotondo D, Heys SD. Plant Phenolics in the Prevention and Treatment of Cancer. *Adv. Exp. Med. Biol.* 2010;698:36-51.
142. Terzuoli E, Giachetti A, Ziche M, Donnini S. Hydroxytyrosol, a Product from Olive Oil, Reduces colon Cancer Growth by Enhancing Epidermal Growth Factor Receptor Degradation. *Mol. Nutr. Food Res.* 2016;60 (3):519-529.
143. Hassan ZK, Elamin MH, Omer SA, Daghestani MH, Al-Olayan ES, Elobeid MA, Virk P. Oleuropein Induces Apoptosis via the P53 Pathway in Breast Cancer Cells. *Asian Pac. J. Cancer Prev.* 2014;14(11):6739-6742.
144. Sepporta MV, Fuccelli R, Rosignoli P, Ricci G, Servili M, Morozzi G, Fabiani R. Oleuropein Inhibits Tumour Growth and Metastases Dissemination in Ovariectomised Nude Mice with MCF-7 Human Breast Tumour Xenografts. *J. Funct. Foods.* 2014;8:269-273.
145. Habibi E, Baâti T, Njim L, M'Rabet Y, Hosni K. Antioxidant and protective effects of extra virgin olive oil incorporated with diallyl sulfide against CCl₄-induced acute liver injury in mice. *Food Sci. Nutr.* 2021;9:6818-6830.
146. Han CY, Koo JH, Kim SH, Gardenghi S, Rivella S, Strnad P, Kim SG. Hepcidin Inhibits Smad3 Phosphorylation in Hepatic Stellate Cells by Impeding Ferroportin-Mediated Regulation of Akt. *Nat. Commun.* 2016;7 (1):1-14.
147. Pirozzi C, Lama A, Simeoli R, Paciello O, Pagano TB, Mollica MP, Di Guida F, Russo R, Magliocca S, Berni Canani R, Mattace Raso G, Calignano A, Meli R. Hydroxytyrosol Prevents Metabolic Impairment Reducing Hepatic Inflammation and Restoring Duodenal Integrity in a Rat Model of NAFLD. *J. Nutr. Biochem.* 2016;30:108–115.
148. Ortiz M, Soto-Alarcón SA, Orellana P, Espinosa A, Campos C, López-Arana S, Rincón MA, Illesca P, Valenzuela R, Videla LA. Suppression of High-Fat Diet-Induced Obesity-Associated Liver Mitochondrial Dysfunction by Docosahexaenoic Acid and Hydroxytyrosol Co-administration. *Dig. Liver Dis.* 2020;52 (8):895–904.
149. Ibrahim M, Wani F, Rahiman S. Hepatoprotective Effect of Olive Oil and Camel Milk on Acetaminophen-Induced Liver Toxicity in Mice. *Int. J. Med. Sci. Public Health.* 2017;6(1): 186-194.
150. Khalatbary AR, Ghabaee DNZ, Ahmadvand H, Amiri FT, Lehi ST. Deltamethrin-induced Hepatotoxicity and virgin Olive Oil Consumption: an Experimental Study. *Iran J. Med. Sci.* 2017;42 (6):586-592.
151. Molnar R, Szabo L, Tomesz A, Deutsch A, Darago R, Ghodrattollah N, Varjas T, Nemeth B, Budan F, Kiss I. In Vivo effects of Olive Oil and Trans-fatty Acids on miR-134, miR-132, miR-124-1, miR-9-3 and mTORC1 Gene Expression in a DMBA-Treated Mouse Model. *Plos one.* 2021;16 (2):e0246022.
153. Zarghi A, Arfaei S. Selective COX-2 Inhibitors: A Review of Their Structure-Activity Relationships. *Iran J. Pharm. Res.* 2011;10 (4):655–683.

152. Fuccelli R, Fabiani R, Rosignoli P. Hydroxytyrosol exerts anti-inflammatory and anti-oxidant activities in a mouse model of systemic inflammation. *Molecules*. 2018;23:3212.
154. Farahat AA, Sawiress FAS, Aghwider AA. Effect of virgin Olive Oil Supplementation on Some Hematologic and Thyroid Hormones, Levels in Rats. *J. Vet. Med. Res.* 2019;26(1):41–47.
155. Donat-Vargas C, Sandoval-Insausti H, Peñalvo JL, Iribas MCM, Amiano P, Bes-Rastrollo M, Molina-Montes E, Moreno-Franco B, Agudo A, Mayo CL, Laclaustra M, Arrillaga CDLF, Lopez MDC, Sánchez M-J, Martínez-Gonzalez MA, Pilar G-C. Olive oil consumption is associated with a lower risk of cardiovascular disease and stroke. *Clin. Nutr.* 2021;41(1): 122-130.
156. Nardiello P, Pantano D, Lapucci A, Stefani M, Casamenti F. Diet supplementation with hydroxytyrosol ameliorates brain pathology and restores cognitive functions in a mouse model of amyloid- β deposition. *J. Alzheimer's Dis.* 2018;63:1161–1172.
157. Sarrafchi A, Bahmani M, Shirzad H, Rafieian-Kopaei M. Oxidative Stress and Parkinson's Disease: New Hopes in Treatment with Herbal Antioxidants. *Curr. Pharm. Des.* 2016;22:238-246.
158. Achour I, Arel-Dubeau AM, Renaud J, Legrand M, Attard E, Germain M, Martinoli MG. Oleuropein Prevents Neuronal Death, Mitigates Mitochondrial Superoxide Production and Modulates Autophagy in a Dopaminergic Cellular Model. *Int. J. Mol. Sci.* 2016;17:1293.
159. Pasban-Aliabadi H, Esmaeili-Mahani S, Sheibani V, Abbasnejad M, Mehdizadeh A, Yaghoobi MM. Inhibition of 6-Hydroxydopamine-Induced PC12 Cell Apoptosis by Olive (*Olea Europaea* L.) Leaf Extract Is Performed by its Main Component Oleuropein. *Rejuvenation Res.* 2013;16:134-142.
160. Bozzetto L, Alderisio A, Clemente G, Giorgini M. Gastrointestinal effects of extra-virgin olive oil associated with lower postprandial glycemia in type 1 diabetes. *Clin. Nutr.* 2019;38: 2645–2651.
161. Carnevale R, Loffredo L, Del Ben M, Angelico F, Nocella C, Petruccioli A, Bartimoccia S, Monticolo R, Cava E, Violi F. Extra virgin olive oil improves post-prandial glycemic and lipid profile in patients with impaired fasting glucose. *Clin. Nutr.* 2017;36:782-787.
162. Friedman M, Rasooly R, Do PM, Henika PR. The Olive Compound 4-Hydroxytyrosol Inactivates *Staphylococcus aureus* Bacteria and Staphylococcal Enterotoxin A (SEA). *J. Food Sci.* 2011;76(8):M558-M563.
163. Fancello F, Multineddu C, Santona M, Molinu MG, Zara G, Dettori S, Deiana P, Zara S. Antimicrobial activities of virgin olive oils *in vitro* and on lettuce from pathogen-inoculated commercial quick salad bags. *Food Control.* 2022;133(Part B):108657.
165. Manach C, Scalbert A, Morand C, Rémésy C, Jiménez L. Polyphenols: food sources and bioavailability. *Am. J. Clin. Nutr.* 2004;79(5):727 – 747.
166. Caruso D, Visioli F, Patelli R, Galli C, Galli G. Urinary excretion of olive oil phenols and their metabolites in humans. *Metabolism.* 2001;50(12):1426–1428.

167. Vissers MN, Zock PL, Katan MB. Bioavailability and antioxidant effects of olive oil phenols in humans: a review. *Eur. J. Clin. Nutr.* 2004; 58:955–965.
168. Edgecombe SC, Stretch GL, Hayball PJ. Oleuropein, an Antioxidant Polyphenol from Olive Oil, Is Poorly Absorbed from Isolated Perfused Rat Intestine. *J. Nutr.* 2000;130(12): 2996–3002.
169. Ghanbari R, Anwar F, Alkharfy KM, Gilani A-H, Saari N. Valuable nutrients and functional bioactives in different parts of olive (*Olea europaea* L.)-A review. *Int. J. Mol. Sci.* 2012;13(3):3291-3340.
170. García-González DL, Aparicio-Ruiz R, Aparicio R. Virgin olive oil - Chemical implications on quality and health. *Eur. J. Lipid. Sci. Technol.* 2008;110(7):602-607.
171. Lou-Bonafonte JM, Arnal C, Navarro MA, Osada J. Efficacy of bioactive compounds from extra virgin olive oil to modulate atherosclerosis development. *Mol. Nutr. Food Res.* 2012;56:1043-1057.
164. Martini FH. *Fundamentals of Anatomy and Physiology*, 7 ed.; Pearson Education Inc.: San Francisco, CA, USA, 2006.
171. Traber MG, Packer L. Vitamin E: beyond antioxidant function. *Am. J. Clin. Nutr.* 1995;62(6 Suppl):1501S-1509S.
172. Morley S, Thakur V, Danielpour D, Parker R, Arai H, Atkinson J, Barnholtz-Sloan J, Klein E, Manor D. Tocopherol Transfer Protein Sensitizes Prostate Cancer Cells to Vitamin E. *Lipids.* 2010;85(46):35578-35589.
172. Newmark HL. Squalene, olive oil, and cancer risk. *Ann. N. Y. Acad. Sci.* 1999;889:193-203.
173. Sano M, Ernesto C, Thomas RG, Klauber MR, Schafer K, Grundman M, Woodbury P, Growdon J, Cotman CW, Pfeiffer E, Schneider LS, Thal LJ. A controlled trial of selegiline, alpha-tocopherol, or both as treatment for Alzheimer's disease. The Alzheimer's Disease Cooperative Study. *N. Engl. J. Med.* 1997;336(17):1216-1222.
174. Meydani SN, Lewis ED, Wu D. Perspective: Should Vitamin E Recommendations for Older Adults Be Increased? *Adv. Nutr.* 2018;9(5):533-543.
175. Murakoshi M, Nishino H, Satomi Y, Takayasu J, Hasegawa T, Tokuda H, Iwashima A, Okuzumi J, Okabe H, Kitano H. Potent preventive action of alpha-carotene against carcinogenesis: spontaneous liver carcinogenesis and promoting stage of lung and skin carcinogenesis in mice are suppressed more effectively by alpha-carotene than by beta-carotene. *Cancer Res.* 1992;52(23):6583-6587.
176. Newmark HL. Is oleic acid or squalene the important preventive agent?. *Am. J. Clin. Nutr.* 2000;72(2):502.
177. Kelly GS. Squalene and its potential clinical uses. *Altern. Med. Rev.* 1999;4(1):29-36.

178. Huang Z-R, Lin Y-K, Fang J-Y. Biological and pharmacological activities of squalene and related compounds: potential uses in cosmetic dermatology. *Molecules*. 2009;14(1), 540-554.
179. Jones PJH, Abumweis SS. Phytosterols as functional food ingredients: Linkages to cardiovascular disease and cancer. *Curr. Opin. Clin. Nutr.* 2009;12(2):147-151.
180. Woyengo TA, Ramprasath VR, Jones PJH. Anticancer effects of phytosterols. *Eur. J. Clin. Nutr.* 2009;63:813-820.
181. Awad AB, Fink CS. Phytosterols as Anticancer Dietary Components: Evidence and Mechanism of Action. *J. Nutr.* 2000;130(9): 2127-2130.
182. Prades J, Vögler O, Alemany R, Gomez-Florit M, Funari SS, Ruiz-Gutiérrez V, Barceló F. Plant pentacyclic triterpenic acids as modulators of lipid membrane physical properties. *Biochim. Biophys. Acta*. 2011;1808:752-760.
183. Qian Y, Guan T, Tang X, Huang L, Huang M, Li Y, Sun H. Maslinic acid, a natural triterpenoid compound from *Olea europaea*, protects cortical neurons against oxygen-glucose deprivation-induced injury. *Eur. J. Pharmacol.* 2011;670:148-153.
184. Qian Y, Guan T, Tang X, Huang L, Huang M, Li Y, Sun H, Yu R, Zhang F. Astrocytic glutamate transporter-dependent neuroprotection against glutamate toxicity: An *in vitro* study of maslinic acid. *Eur. J. Pharmacol.* 2011;651:59-65.
185. Caltana L, Rutolo D, Nieto ML, Brusco A. Further evidence for the neuroprotective role of oleanolic acid in a model of focal brain hypoxia in rats. *Neurochem. Int.* 2014;79:79-87.
186. Andrikopoulos NK, Kaliora AC, Assimopoulou AN, Papageorgiou VP. Inhibitory activity of minor polyphenolic and nonpolyphenolic constituents of olive oil against *in vitro* low-density lipoprotein oxidation. *J. Med. Food* 2002;5(1): 1-7.
187. Shaik AH, Shaik SR, Daddam JR, Ali D, SalimManoharadas A, Arafah MD, Kodidhela LD. Maslinic acid and gallic acid protective efficacy on lipids, lipoproteins and lipid metabolizing enzymes against isoproterenol administered cardiotoxicity: An *in vivo* and *in silico* molecular docking evidences. *J. King Saud Univ. Sci.* 2021;33(1):101230.
188. Lozano-Mena G, Sánchez-González M, Juan ME, Planas JM. Maslinic Acid, a Natural Phytoalexin-Type Triterpene from Olives — A Promising Nutraceutical?. *Molecules*. 2014;19:11538-11559.
189. Shaik AH, Rasool SN, Kareem MA, Krushna GS, Akhtar PM, Devi KL. Maslinic Acid Protects Against Isoproterenol-Induced Cardiotoxicity in Albino Wistar Rats. *J. Med. Food*. 2012;15(8):741-746.