

## Original Research Article

### Seasonal incidence of fall armyworm, *Spodoptera frugiperda* (J. E. Smith) on sorghum

#### Abstract:

The study on seasonal incidence of fall armyworm was carried out during *kharif*, 2019 at Agricultural college farm, Bapatla.- The oviposition of fall armyworm was observed from 34<sup>th</sup> SMW and reached its peak during 40<sup>th</sup> SMW (0.03 egg masses plant<sup>-1</sup>). The larval population of fall armyworm was commenced during 35<sup>th</sup> SMW and increased gradually to a peak of 1.67 larvae plant<sup>-1</sup> during 41<sup>st</sup> SMW. The maximum plant infestation (60.00%) and leaf damage severity rating (3.13) of fall armyworm was observed during 41<sup>st</sup> SMW. The peak activity of natural enemies was observed during 41<sup>st</sup> SMW which coincides with the larval population of fall armyworm. The correlation analysis indicated that eggmasses of fall armyworm was positively correlated with minimum temperature ( $r=0.668$ ) and wind speed ( $r=0.529$ ), while, the larval population showed significant positive correlation with maximum temperature ( $r=0.029$ ). The plant infestation caused by fall armyworm had shown significant negative correlation with maximum temperature ( $r=-0.633$ ) ( $r=0.678$ ) and evening relative humidity ( $r=0.664$ ) whereas, the leaf damage severity rating exhibited significant positive correlation with evening relative humidity ( $r=0.691$ ). The multiple regression analysis revealed that the influence of weather parameters on the incidence of fall armyworm and natural enemies in sorghum ecosystem was more than 65% and 40%, respectively.

**Key words:** Fall armyworm, seasonal incidence, correlation, weather parameters, sorghum.

#### 1. INTRODUCTION

Sorghum (*Sorghum bicolor* (L.) Moench) is an important staple food for more than 500 million people after wheat, rice, maize and barley in the world whereas in India, it occupied third place after rice and wheat in terms of production and utilization (Varaprasad and Staggenborg, 2011). It occupied an area of 4.96 m ha with an annual production of 4.80 MT in India during 2018-19 (FAO, 2018-19). The productivity of sorghum in India (967 kg ha<sup>-1</sup>) was low compared to the world productivity (1408 kg ha<sup>-1</sup>) due to abiotic and biotic stress. Among the biotic stresses, the insect pests and diseases are causing significant yield loss (Reddy and Patil, 2015).

**Comment [A1]:** Title is written using Arial, 18

Based on the Author Guidelines, the body of manuscript should follow the structure of Abstract, Introduction, Methodology, Results and Discussion, Conclusion, Acknowledgements, Competing Interests, Authors' Contributions, Consent (where applicable), Ethical approval (where applicable), and References plus figures and/or tables.)

**Comment [A2]:** Almost 90% similarity with <https://www.entomoljournal.com/archives/7?year=2020&vol=8&issue=3&ArticleId=6722>

Please re-paraphrase the title using another sentences or terminology

**Comment [A3]:** ABSTRACT (ARIAL, BOLD, 11 FONT, LEFT ALIGNED, CAPS)

**Comment [A4]:** Define the abbreviation clearly for the first use

**Comment [A5]:** Which value of r describes the sentence in front of it?

**Comment [A6]:** Based on the Author Guidelines, the abstract should be concise and informative. It should briefly describe the purpose of the work, techniques and methods used, major result and conclusions. Different sub-sections, as given below, should be used:

Abstract:

- Aims
- Study design
- Place and Duration of Study
- Methodology
- Result
- Conclusion

Generally non-standard abbreviations should not be used, if necessary they should be clearly defined in the abstract, at first use.

The abstract of this manuscript have not been compiled using sub-sections according to the author guidelines.

There is an abbreviation used in the abstract and not clearly define yet

**Comment [A7]:** The conclusion not stated yet

**Comment [A8]:** Arial, inclined, 10 font, justified, italic

**Comment [A9]:** Sub-section title should be written using Arial, Bold, 11 font, left aligned, caps

It harbours nearly 150 species of insects in different agroecosystems of India (Sharma, 1993). Shoot fly, stem borer and sorghum midge has attained the major pest status in India and accounts to the yield loss of nearly 32 per cent in India (Nwanze *et al.*, 1995). The recently introduced pest, fall armyworm [*Spodoptera frugiperda* (J. E. Smith)] has become a great threat to cereal production in the world (Day *et al.*, 2017) and causing 16 per cent yield loss in sorghum (Abrahmas *et al.*, 2017). The pest status was not static on sorghum even though it is one of the preferred host for fall armyworm and it has been largely influenced by weather parameters like temperature, relative humidity and wind.

In view of the existing situation and importance of sorghum in India, there is a need for the development of economically viable and environmentally safe approach for successful management of fall armyworm. For this, the knowledge on population dynamics fall armyworm in sorghum is required but, the availability of literature on seasonal incidence of fall armyworm in India is scanty. Hence, the present investigation was carried out to find out the influence of weather parameters on incidence of fall armyworm and its natural enemies in sorghum.

## 2. MATERIALS AND METHODS

The field experiment on seasonal incidence of fall armyworm in sorghum was carried out during *kharif*, 2019 at Agricultural college farm, Bapatla. The sorghum variety (CSH-16) was sown during second week of August in an area of 400 m<sup>2</sup> with a spacing of 45× 15 cm. The crop was raised as per the agronomic recommendations of ANGRAU and the plot was kept under unprotected conditions throughout the crop growth period.

The observations on incidence of fall armyworm (No. of egg masses and larvae per plant), pest damage (Plant infestation and leaf damage severity) and population of natural enemies (Coccinellids, spiders and predatory bugs) were recorded on 30 randomly selected plants at weekly interval from 10 DAS to crop maturity.

The leaf damage severity was recorded by using the rating scale of Wiseman *et al.* (1966). 0- no damage, 1- small amount of pinhole-type injury, 2- several pinholes, 3-small amount of shot hole type injury with 1 or 2 lesions, 4- several shot hole type injuries and few lesions, 5- several lesions, 6-several lesions, shot hole injury and portions eaten away, 7-several lesions and portions eaten away and areas drying, 8- several portions of the whorl eaten away

**Comment [A10]:** According to Author Guidelines the manuscript's body using Arial, normal, 10 font, justified

**Comment [A11]:** Sub-section title should be written using Arial, Bold, 11 font, left aligned, caps

**Comment [A12]:** Define the abbreviation clearly for the first time use

and areas drying, 9- the whorl completely eaten away and more areas drying or plant dead was used for rating the fall armyworm damage.

The data on various weather parameters (Maximum temperature, minimum temperature, morning relative humidity, evening relative humidity, rainfall, rainy days and wind speed) during the entire crop growth period was collected from meteorological observatory, Agricultural College Farm, Bapatla. In order to determine the relationship between the weather parameters and incidence of fall armyworm and natural enemies, the mean population of pest and natural enemies were subjected to simple correlation analysis with weather parameters and multiple linear regression equations for pest and natural enemies population with weather parameters were formulated by using SPSS 16.0 software.

### 3. RESULTS AND DISCUSSION

#### 3.1 Seasonal incidence of fall armyworm on sorghum:

The observation on seasonal incidence of fall armyworm and their natural enemies on sorghum were recorded from 34<sup>th</sup> SMW (Standard Meteorological Week) to 49<sup>th</sup> SMW (Table 1). The oviposition of fall armyworm was observed from 34<sup>th</sup> SMW (0.03 egg masses plant<sup>-1</sup>) thereafter, it was increased gradually and attained the peak during 40<sup>th</sup> SMW (0.30 egg masses plant<sup>-1</sup>). The results were in accordance with the reports of Shylesha and Sravika (2018) who reported that the maximum oviposition of FAW was noticed during 39<sup>th</sup> and 40<sup>th</sup> SMW in maize. The moths of FAW preferred to oviposit on 54 to 64 days old sorghum plants compared to 22 to 42 days old plants (Pitre *et al.*, 1983).

The larval population of FAW (0.37 larvae plant<sup>-1</sup>) was noticed initially during 35<sup>th</sup> SMW then population increased gradually and attained the peak population of 1.67 larvae plant<sup>-1</sup> during 41<sup>st</sup> SMW (59 DAS). The findings were in conformation with Paul and Deole (2020) who reported that the maximum larval population of FAW was observed during 39<sup>th</sup> and 40<sup>th</sup> SMW in maize. Kumar *et al.* (2020) observed the peak incidence of FAW during the first fortnight of November in maize.

The infestation of FAW was observed from 35<sup>th</sup> SMW (16.67%) to 49<sup>th</sup> SMW (43.33%). The per cent infestation was more than 50 per cent from 40<sup>th</sup> SWM to 47<sup>th</sup> SWM however, the peak infestation of 60.00 per cent was observed during 41<sup>st</sup> SWM (59 DAS). The findings were in line with the results of Wyckhuys and Neil (2006) who reported that the peak infestation of

**Comment [A13]:** Body of manuscript should be written using Arial, normal, 10 font, justified

**Comment [A14]:** Sub-section title should be written using Arial, Bold, 11 font, left aligned, caps

**Comment [A15]: 2.1 Subheading Subheading (ARIAL, BOLD, 11 FONT, LEFT ALIGNED) - second level heading.**

**2.1.1 Sub-subheading (ARIAL, BOLD, 10 FONT, LEFT ALIGNED, underlined)- third level heading.**

**Comment [A16]:** Describe the abbreviate clearly for the first use

fall armyworm was noticed during the whorl stage of maize and the infestation was decreased gradually during post whorl stage of the crop. The maximum infestation of FAW on plants (53.89%) was observed at 66 days after seedling emergence in maize (Pinango *et al.*, 2001).

The leaf damage severity rating was low during the early crop growth period. Later, the damage of FAW on leaves increased with the increase in larval population. However, the maximum leaf damage severity rating of 3.13 was observed during 41<sup>st</sup> SMW (59 DAS) which was coincided with the peak population of larvae. Similarly, Chamberlin and All (1991) found that the leaf injury rating of fall armyworm on sorghum was low upto 43 days after planting later, it was increased at 51 and 58 days after planting. According to Macharia *et al.* (2019), the severity rating of fall armyworm in maize was very low up to 30 days after emergence (DAE) later, it was increased at 44 DAE and 58 DAE.

### 3.2 Seasonal incidence of natural enemies on sorghum

The coccinellids viz., *Cheilomenes sexmaculata* (Fab.), *Coccinella transversalis*(Fab.), *Brumoides suturalis* (Fab.) and *Cycloneda sanguinea* (L.) were observed from 35<sup>th</sup> SMW to 48<sup>th</sup> SMW. The peak population was noticed during 41<sup>st</sup> SMW (0.43 coccinellids plant<sup>-1</sup>). The results were in agreement with the findings of Ankita *et al.* (2020) who reported that the activity of coccinellids was observed during the first week of August however, the maximum activity of coccinellids was noticed during the fourth week of September in maize.

The population of spiders was low during the initial crop growth period *i.e.* up to 45 DAS (39<sup>th</sup> SMW). Later, the population was raised and attained the peak population of 0.90 spiders plant<sup>-1</sup> during 41<sup>st</sup> SMW. The results were in coherence with the findings of Sidar *et al.* (2017) who recorded the incidence of spiders on maize from 32<sup>nd</sup> SMW to 44<sup>th</sup> SMW but, the peak population was noticed during 39<sup>th</sup> SMW.

The population of predatory bugs such as *Rhynocoris fuscipes* (Fab.), *Orius* sp. and *Eocanthecona furcellata* (Wolff) was observed from 35<sup>th</sup> SMW to 47<sup>th</sup> SMW. The maximum population of 0.26 predatory bugs plant<sup>-1</sup> was recorded during 41<sup>st</sup> SMW. The present observations were in line with the findings of Shylesha and Sravika (2018) who reported that the peak population of predatory bugs was observed from 38<sup>th</sup> SMW to 40<sup>th</sup> SMW in maize. The

**Comment [A17]:** Body of manuscript should be written using Arial, normal, 10 font, justified

**Comment [A18]:** 2.1 Subheading Subheading (ARIAL, BOLD, 11 FONT, LEFT ALIGNED) - second level heading.

**2.1.1 Sub-subheading** (ARIAL, BOLD, 10 FONT, LEFT ALIGNED, underlined)- third level heading.

incidence of *E. furcellata* in maize was noticed during the 36<sup>th</sup> SMW and the peak population was observed during the 40<sup>th</sup> SMW (Keerthi *et al.*, 2020).

### 3.3 Correlation between weather parameters and incidence of fall armyworm and natural enemies in sorghum

The correlation studies worked out between the weather parameters and population of FAW (Egg masses, larval population, per cent plant infestation, leaf damage severity rating) and their natural enemies revealed that the oviposition of FAW had significant positive correlation with the minimum temperature ( $r = 0.668$ ) and wind speed ( $r = 0.529$ ). However, the population showed non-significant positive correlation with other abiotic factors. This observation was in agreement with findings of Kundra *et al.* (2020) who reported that the oviposition of shoot fly had significant and positive correlation with the evening RH and rainfall in little millet (Table 2).

The larval population showed significant positive correlation with maximum temperature ( $r = 0.029$ ) and non-significant positive correlation with minimum temperature ( $r = 0.383$ ), morning RH ( $r = 0.222$ ), evening RH ( $r = 0.274$ ), and wind speed ( $r = 0.362$ ) however, it showed non significant negative correlation with precipitation ( $r = -0.192$ ) and no. of rainy days ( $r = -0.200$ ). The results were in agreement with the findings of Paul and Deole (2020) who found the existence of non significant positive correlation between the larval population of FAW in maize and the weather parameters like minimum temperature and morning RH. According to Barrios *et al.* (2019), the larval population of *S. frugiperda* was positively correlated with the relative humidity in maize ecosystem.

The per cent plant infestation of fall armyworm showed significant positive correlation with morning RH ( $r = 0.678$ ) and evening RH ( $r = 0.664$ ) however, it had negative correlation with the maximum temperature ( $r = -0.633$ ). The results were in partial coherence with Pazhanisamy *et al.* (2019) who observed the significant positive correlation between leaflet damage of *S. litura* in groundnut and relative humidity and non significant positive correlation with the minimum temperature. However, the leaflet damage exhibited negative correlation with maximum temperature, wind speed and rainfall. The leaf damage severity of fall armyworm showed significant and positive correlation with evening RH ( $r = 0.691$ ) and negative correlation with the maximum temperature ( $r = -0.237$ ). The results were partially in coherence with the

**Comment [A19]:** Body of manuscript should be written using Arial, normal, 10 font, justified

**Comment [A20]:** 2.1 Subheading Subheading (ARIAL, BOLD, 11 FONT, LEFT ALIGNED) - second level heading.

**2.1.1 Sub-subheading** (ARIAL, BOLD, 10 FONT, LEFT ALIGNED, underlined)- third level heading.

findings of Dar *et al.* (2019) who observed the significant positive correlation between the pest damage index caused by *Tetranychus turkestani* in mulberry and weather parameters like maximum temperature, minimum temperature and relative humidity.

The coccinellid population showed significant positive correlation with no. of rainy days ( $r = 0.022$ ) and negative correlation with evening RH ( $r = -0.008$ ). The population of spiders showed significant positive correlation with no. of rainy days ( $r = 0.021$ ) and maximum temperature ( $r = 0.033$ ). Whereas, the predatory bugs showed significant positive correlation with the maximum temperature ( $r = 0.033$ ). The present findings were in close conformity with the findings of Ahirwar *et al.* (2015) who stated that the non significant positive correlation associated between the population of natural enemies (Coccinellids, spiders and predatory bugs) in soybean and abiotic factors like maximum temperature and minimum temperature.

### 3.4 Multiple linear regression analysis

The contribution of weather variables *viz.*, maximum temperature, minimum temperature, morning relative humidity, evening relative humidity, rainfall, rainy days and wind speed on the incidence of fall armyworm egg masses, larvae, coccinellids, spiders and predatory bugs on sorghum was 75.77, 65.91, 41.86, 43.64 and 44.52 per cent, respectively ( Table 3). Similarly, Kumar *et al.* (2020) reported that the abiotic factors were responsible for the 76%, 74%, 64% and 71% variation in the incidence of larval population of fall armyworm in the maize fields of Perambalur, Veppanthattai, Alathur and Veppur blocks of Tamil Nadu, respectively. The overall impact of abiotic factors on the incidence of natural enemies like spiders, coccinellids and chrysopids in potato was 26.60, 35.90 and 39.90 per cent, respectively (Natikar *et al.*, 2018). The influence of weather parameters on the incidence of predatory bugs *viz.*, *E. fuscicornis* and *R. fuscipes* on pigeonpea was 81.90 and 69.60 per cent, respectively (Chakravarty *et al.*, 2017).

## 4. CONCLUSION

## ACKNOWLEDGEMENTS

## COMPETING INTERESTS

**Comment [A21]:** Body of manuscript should be written using Arial, normal, 10 font, justified

**Comment [A22]:** 2.1 Subheading Subheading (ARIAL, BOLD, 11 FONT, LEFT ALIGNED) - second level heading.

**2.1.1 Sub-subheading** (ARIAL, BOLD, 10 FONT, LEFT ALIGNED, underlined)- third level heading.

**Comment [A23]:** Body of manuscript should be written using Arial, normal, 10 font, justified

## AUTHORS' CONTRIBUTIONS

## CONSENT (WHERE EVER APPLICABLE)

## ETHICAL APPROVAL (WHERE EVER APPLICABLE)

## References

- Abrahams P, Beale T, Cock M, Corniani N, Day R, Godwin J, Murphy S, Richards, G, Vos J. Fall Armyworm status. impacts and control options in Africa: Preliminary evidence note. International maize and wheat improvement center (CIMMYT), Mexico.2017; **4**.
- Ahirwar R, Devi P, Gupta R. Influence of ambient weather on the incidence of major insect pests and their bioagents of soybean crop (*Glycine max* L. Merrill). International Journal of Plant Protection. 2015; 8 (2): 234- 240.
- Ankita, Sharma PK, Sharma PC. Fall army worm *Spodoptera frugiperda* (J E Smith) on maize in Himachal Pradesh. Indian Journal of Entomology. 2020; 82: 1-4.
- Barrios CIJ, Quijano, BE, Andrade MB. Population of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) cause significant damage to genetically modified corn crops. Revista Facultad Nacional de Agronomia. 2019; 72 (3): 8953-8962.
- Chakravarty S, Agnihotri M, Jagdish J. Seasonal abundance of predatory bugs, *Eocanthecona furcellata* (Wolff.) and *Rhynocoris fuscipes* (F.) and its olfactory responses towards plant and pest mediated semiochemical cues in pigeonpea ecosystem. Legume Research: An International Journal. 2017; 40 (2): 351-357.
- Chamberlin JR, All JN. Grain sorghum response to fall armyworm and corn earworm infestation. Journal of Economic Entomology. 1991;84 (2): 619-624.
- Dar MY, Singh A, Ramegowda GK, Rao RJ. Influence of weather factors on incidence and severity of *Tetranychus turkestani* (Acari: Tetranychidae) on mulberry. Journal of Applied Sciences. 2019; 19: 504-512.
- Day R, Abrahams P, Bateman M, Beale T, Clotley V, Cock M, Yelitza C, Natalia C, Early R, Godwin J, Gomez J, Moreno PG, Murphy ST, Mensah B O, Phiri N, Pratt C, Silvestri, S, Witt A. Fall armyworm: Impacts and implications for Africa. Outlook on Pest Management. 2017; 28 (5): 196-201.
- Food and Agricultural organizations (FAO), 2018-19.Crops. 8 April 2020. <http://www.fao.org/faostat/en/#data/QC>.

**Formatted:** Indent: First line: 0 cm

**Comment [A24]:** Sub-section title should be written using Arial, Bold, 11 font, left aligned, caps. Citation style

- Keerthi MC, Sravika A, Mahesha HS, Gupta A, Bhargavi HA, Ahmed S. Performance of the native predatory bug, *Eocanthecona furcellata* (Wolff) (Hemiptera: Pentatomidae), on the fall armyworm, *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae), and its limitation under field condition. *Egyptian Journal of Biological Pest Control*. 2020; 30: 691-4.
- Kumar VN, Yasodha P, Justin CGL. Seasonal incidence of maize fall armyworm *Spodoptera frugiperda* (J.E. Smith) (Noctuidae; Lepidoptera) in Perambalur district of Tamil Nadu, India. *Journal of Entomology and Zoology Studies*. 2020; 8 (3): 1- 4.
- Kundra KKK, Chakravarty MK, Kumari A. Influence of abiotic factors on the shoot fly (*Atherigona pulla* Wiede) infestation. *International Journal of Current Microbiology and Applied Sciences*. 2020; 9 (6): 4037- 4043.
- Macharia JK, Birech RJ, Njoroge M, Nganga JN. Evaluation of selected organic biopesticides to control fall armyworm (*Spodoptera frugiperda* (J.E.Smith) (Lepidoptera, Noctuidae). *International Organization of Scientific Research Journal of Agriculture and Veterinary Science*. 2019; 12(6):26-29.
- Natarikar PK, Balikai RA, Kambekar DN. Prediction models for insect pests of potato and their natural enemies based on abiotic factors in Karnataka. *Journal of Entomology and Zoology Studies*. 2018; 6 (4): 1151-1155.
- Nwanze KF, Seetharama N, Sharma HC, Stenhouse JW. Biotechnology in pest management: Improving resistance in sorghum to insect pests. *African Crop Science Journal*. 1995; 3: 209-215.
- Paul, N, Deole, S. Seasonal incidence of fall army worm, *Spodoptera frugiperda* (Smith) infesting maize crop at Raipur (Chhattisgarh). *International Journal of Chemical Studies*. 2020; 8 (3): 2644-2646.
- Pazhanisamy M, Sathyaseelan V, Senthilkumar M. Seasonal incidence of leaf eating caterpillar (*Spodoptera litura* Fabricius) in groundnut ecosystem during *rabi* season. *Plant Archives*. 2019; 19 (1): 1159-1163.
- Pinango L, Arnal E, Rodriguez B. Population fluctuation of *Spodoptera frugiperda* (Smith) (Lepidoptera: Noctuidae) in corn cultivation under three tillage systems. *Entomotropica*. 2001; 16 (3): 173-179.
- Pitre HN, Mulrooney JE, Hogg BD. Fall armyworm (Lepidoptera: Noctuidae) oviposition: crop preferences and egg distribution in plants. *Journal of Economic Entomology*. 1983; 76 (3): 463-466.
- Reddy SP, Patil JV. *Genetic enhancement of rabi sorghum*. Academic Press, London. 2015; 1.



Sharma HC. Host-plant resistance to insects in sorghum and its role in integrated pest management. *Crop Protection*. 1993;**12** (1): 11-34.

Shylesha AN and Sravika A. Natural occurrence of predatory bugs, *Eocanthecona furcellata* (Wolff) and *Andrallus spinidens* (Fabr.) on *Spodoptera frugiperda* (Smith) (Hemiptera: Pentatomidae) in maize and their potential in management of fall armyworm. *Journal of Biological Control*. 2018; 32(3): 209-211.

Sidar YK, SonaliDeole, Nirmal A, Gajbhiye RK, Bisen MS. A study on the seasonal distribution of spider fauna in the maize field at Raipur, Chhattisgarh region. *Journal of Entomology and Zoology Studies*. 2017; 5 (2): 1105-1108.

Varaprasad PV, Staggenborg SA. Growth and Production of Sorghum and Millets. In Willy H. Verheye (ed.) *Soils, plant growth and crop production*. Encyclopedia of Life Support Systems (EOLSS), Oxford, United Kingdom, 2011; 2.

Wiseman BR, Painter RH, Wassom CE. Detecting corn seedling differences in the greenhouse by visual classification of damage by the fall armyworm. *Journal of Economic Entomology*. 1966; 59: 1211-1214.

Wyckhuys K, Neil RJ. Population dynamics of *Spodoptera frugiperda* Smith (Lepidoptera: Noctuidae) and associated arthropod natural enemies in Honduran subsistence maize. *Crop Protection*. 2006; 25: 1180-1190.

**Table 1. Seasonal incidence of fall armyworm and its natural enemies on sorghum during *kharif*, 2019**

SMW	Crop stage	Fall armyworm*				No. of natural enemies plant <sup>-1</sup> *			Weather parameters						
		No. of egg masses plant <sup>-1</sup>	No. of larvae plant <sup>-1</sup>	Plant infestation (%)	Leaf damage severity rating	Coccinellids	Spiders	Predatory bugs	Max. temp (°C)	Min. temp. (°C)	Relative humidity (%)		Rainfall (mm)	No. of rainy days	Wind speed (kmph)
											8:30 AM	5:30 PM			
34	10 DAS	0.03	0.00	0.00	0.00	0.00	0.00	0.00	33.75	25.67	82.57	67.42	5.20	1	1.86
35	17DAS	0.06	0.37	16.67	0.37	0.16	0.06	0.03	34.88	25.75	76.71	70.57	14.70	3	1.76
36	24DAS	0.10	0.67	26.67	0.90	0.20	0.10	0.03	32.52	25.94	81.14	64.00	2.50	0	1.93
37	31DAS	0.20	0.20	36.67	1.23	0.30	0.16	0.06	35.12	26.30	77.42	69.85	133.00	2	1.90
38	38DAS	0.13	0.60	43.33	1.53	0.06	0.00	0.00	31.64	25.40	85.85	82.57	28.60	3	1.79
39	45DAS	0.20	0.77	46.67	1.90	0.10	0.10	0.10	30.40	25.45	86.14	82.28	54.50	3	1.87
40	52DAS	0.30	1.33	53.33	2.67	0.20	0.43	0.13	32.35	25.70	86.00	77.00	0.50	0	2.10
41	59DAS	0.23	1.67	60.00	3.13	0.43	0.90	0.26	32.28	25.38	85.85	77.14	5.70	0	2.06
42	66DAS	0.16	0.30	56.67	2.90	0.26	0.53	0.16	30.07	24.45	86.14	84.71	114.60	6	1.93
43	73DAS	0.10	0.40	53.33	2.57	0.00	0.00	0.00	29.84	24.27	86.85	80.71	73.50	2	1.64
44	80DAS	0.06	0.80	53.33	2.00	0.06	0.10	0.06	30.28	24.22	85.42	80.57	78.90	3	1.76
45	87DAS	0.03	0.73	50.00	1.80	0.16	0.13	0.06	32.47	23.94	85.00	70.71	0.00	0	1.74
46	94DAS	0.03	0.57	53.33	1.30	0.13	0.13	0.03	31.32	23.48	86.28	75.71	0.20	0	2.00
47	101DAS	0.00	0.37	50.00	0.97	0.10	0.10	0.03	30.62	22.02	88.28	76.42	13.80	1	1.24
48	108DAS	0.00	0.10	46.67	0.60	0.06	0.06	0.00	30.18	22.01	86.85	75.57	0.20	0	1.86
49	115DAS	0.00	0.03	43.33	0.00	0.00	0.03	0.00	29.90	20.70	85.40	68.90	0.20	0	1.90

SMW: Standard Meteorological Week

DAS: Days After Sowing \*Mean of 30 observations

**Table 2. Correlation between the weather parameters and incidence of fall armyworm and its natural enemies on sorghum during *kharif*, 2019**

S. No.	Pest/ Natural enemies	Particulars	Correlation coefficient (r)						
			Max. temp. (°C)	Min. temp. (°C)	Relative humidity (%)		Rainfall (mm)	No. of rainy days	Wind speed (kmph)
					8:30 AM	5:30 PM			
1	Fall armyworm	Egg masses	0.219	0.668**	-0.081	0.331	0.334	0.198	0.529*
2		Larval population	0.029**	0.383	0.222	0.274	-0.192	-0.200	0.362
3		Plant infestation (%)	-0.633**	0.341	0.678**	0.664**	0.187	0.036	0.023
4		Leaf damage severity rating	-0.237	0.344	0.328	0.691**	0.429	0.325	0.214
5	Natural enemies	Coccinellids	0.405	0.453	-0.266	-0.008*	0.195	0.022*	0.417
6		Spiders	0.033*	0.230	0.157	0.262	0.063	0.021*	0.463
7		Predatory bugs	0.033*	0.329	0.130	0.336	0.162	0.135	0.439

\*Significant at 5 per cent level\*\* Significant at 1 per cent level

**Table 3 Multiple linear regression between the weather parameters and population of fall armyworm and its natural enemies in sorghum during *kharif*, 2019**

S. No.	Pest/ Natural enemies	Particulars	Multiple linear regression with abiotic factors	R <sup>2</sup>
1	Fall armyworm	Egg masses	$Y = -2.1107 + 0.0085X_1 + 0.0317X_2 + 0.0041X_3 + 0.0074X_4 + 0.0006X_5 - 0.0217X_6 + 0.1486X_7$	0.7577
2		Larval population	$Y = -5.9166 - 0.033X_1 + 0.1795X_2 - 0.017X_3 + 0.061X_4 - 0.0019X_5 - 0.1918X_6 + 0.1952X_7$	0.6591
3	Natural enemies	Coccinellids	$Y = -3.3657 + 0.0550X_1 - 0.0031X_2 + 0.0126X_3 + 0.0048X_4 + 0.0009X_5 - 0.0139X_6 + 0.2100X_7$	0.4186
4		Spiders	$Y = -10.254 + 0.119X_1 - 0.016X_2 + 0.063X_3 + 0.005X_4 + 0.001X_5 + 0.011X_6 + 0.643X_7$	0.4364

5		Predatory bugs	$Y = -2.7158 + 0.0272X_1 + 0.0032X_2 + 0.0159X_3 + 0.0023X_4 + 0.0003X_5 + 0.0027X_6 + 0.1635X_7$	0.4452
---	--	----------------	---	--------

$X_1$  - Maximum temperature ( $^{\circ}\text{C}$ ),  $X_2$  - Minimum temperature ( $^{\circ}\text{C}$ ),  $X_3$  - Morning relative humidity (%),  $X_4$  - Evening relative humidity (%),  $X_5$  - Rainfall (mm),  $X_6$  - No. of rainy days,  $X_7$  - Wind speed (kmph).