

Active and passive soil organic carbon pools as affected by different Land Use Patterns in red soils of Vikarabad district, Telangana, [India](#)

ABSTRACT

Carbon (C) fluxes are largely controlled by the small but highly bio-reactive, labile pools in terrestrial soils, while long-term C storage is determined by the long lived recalcitrant fractions. A survey was conducted in the years (2019-20 and 2020-21) in red soils of Vikarabad district covering eight *mandals* and soil samples were collected from four predominant land use patterns at two depths (0-15 and 15-30cm). Land use patterns studied included cultivated land with different cropping intensity *i.e.*, 100% cropping intensity (redgram-fallow), 200% cropping intensity (rice-rice), and two from natural conditions *i.e.*, forest land and fallow land to assess the impact of these land uses on various pools of SOC *viz.*, total organic carbon (TOC), oxidizable organic carbon and its pools. The SOC content of forest land recorded [the](#) highest mean [value](#) (7.07 g kg^{-1}) and that of fallow land recorded [the](#) lowest [value](#) (2.68 g kg^{-1}). The mean SOC stock was highest in forest land ($13.92 \text{ Mg C ha}^{-1}$) and that of fallow land was [the](#) lowest ($5.64 \text{ Mg C ha}^{-1}$). The highest accumulation of [mean](#) TOC ~~mean~~ was observed in forest (26.15 Mg ha^{-1}). About 60.00% and 53.16% of the TOC in forest land and 100% cropping intensity occurred as very active carbon pool suggesting that the accumulated C could easily be lost following the land use change. Contrarily, the majority of TOC of 200% cropping intensity (rice-rice) and fallow land (52.86% and 55.02% of TOC) was stabilized in the passive pool indicating more stable nature of the accumulated organic matter.

Keywords: Land Use Patterns, Soil Organic Carbon, ~~TOC~~ [Total Organic Carbon](#), Active Pool, Passive Pool, Forest

INTRODUCTION

Soil organic carbon (SOC) is reported as a sensitive indicator of soil quality and environmental sustainability [1]. To sustain fertility and productivity of soil, SOC is considered essential which can improve physical, chemical and biological properties of a soil [2] as well as it can be used for predicting climate change and effects on crop production [3]. The overall efficiency, resilience, adaptive capacity and mitigation potential of production or land use patterns towards the extremities of climate can be enhanced by increasing the quality as well as quantity of soil organic carbon pools in different land uses [4]. Global SOC pool in the top 1 m soil is approximately 1200 to 1600 Pg ($1 \text{ Pg} = 10^{15} \text{ g}$) and 695 to 930 Pg of inorganic [carbon](#). The amount of carbon (C) stored in soil is two times [that](#) of the global biotic C pool and three times of the global atmospheric C [pool](#). The concentration of atmospheric carbon dioxide (CO_2) may be greatly impacted with a small change in SOC pool, thus affecting the global carbon cycle [5]. It is therefore important to preserve, maintain and store SOC while addressing problems of climate change and food insecurity.

Land use/land cover change is one of the key factors which affect the soil organic carbon pool remarkably. The reason for this being the [difference in the](#) rates of input (e.g. plant litter) and ~~rate of~~ output (e.g. SOC mineralization) of soil organic matter (SOM) as a result of alterations in plant community and land management practices [6]. Land use changes and management are widely recognized as the most important driving forces of global carbon cycles [7], contributing [between e](#)

Comment [GK1]: Needs a reference

Comment [GK2]: Needs a reference

6%–39% of the ~~growth-increase~~ in CO₂ emissions and have profound impacts on SOC which are estimated to be one of the second largest source of human induced greenhouse gas emissions (1.5 Pg C/a) after fossil fuel combustion (5.3 Pg C/a) [8] because, soils may provide an immediate sink for atmospheric CO₂ with proper management. Tropical forests can act either as a carbon sink or source which contributes significantly in the modification of atmospheric C concentration. The conversion of tropical forests into other land uses such as plantations and croplands through anthropogenic activities may act as a carbon source leading to alterations of soil properties and processes. When natural forests are converted to croplands, the soil structure gets disrupted enhancing the mineralization of organic matter by microbes, subsequently leading to SOC loss.

Many attempts have been made to differentiate SOC into various pools of varying lability and more labile pools have been used as sensitive indicators of changes in soil quality in response to land use management [9] because these fractions may precede future changes of SOC [10]. SOC stocks in soil consist of labile and non labile pool with varying residence time. Labile pool (active pool or light fraction pool) are the source of nutrients with most rapid turnover rates that influences the quality and productivity of soil, it is also available in a relatively small proportion because it is easily decomposed and extremely sensitive to environmental fluctuation. Another type is the stable, resistant/recalcitrant pools with varying residence time (passive pool) which is slowly decomposed by microbial activity ~~semicroorganisms and so~~, it is not considered as a good parameter for soil quality but contributes in total organic carbon (TOC) stocks [11].

Although many studies have focused on SOC dynamics as consequence of land use change for last 20 years, SOC dynamics ~~was has been~~ poorly understood as they include climate, time, vegetation, topography and parent material. Thus, we studied on single soil type that holds many of these interactive controls as constant allowing the evaluation of SOC dynamics with better understanding in different land use patterns. Therefore, the objective of this research was to quantify various SOC ~~contents-pools~~ (very labile, labile, less labile and non-labile) and their relative proportions in total organic carbon (TOC) across different land use patterns.

2. MATERIALS AND METHODS

2.1 Study area

The ~~present~~ study was carried out in Vikarabad district, Telangana, India. The ~~geographical surface~~ area of the district is 3,386 ~~sq.km.km²~~ and is situated between ~~latitudes~~ 17°20'N to 11.15"N ~~latitude~~ and ~~longitudes~~ 77°54'N-54'E to 17.45"E ~~longitude~~. The net cropped area in the district is about 20.4 ~~lakh~~ ha and has a forest cover of 44,548 ha and fallow land to an extent of 20,769 ha. The agro climate of this sub region is characterized as hot, semiarid moist with dry summers and cold winters. The mean annual temperature varies from 22-~~°C-to~~ 32°C. The mean summer (April–June) temperature varies from 32-~~°C-to~~ 46°C, rising to a maximum of 48°C in May, and the mean winter (December–February) temperature varies from 12-~~°C-to~~ 26°C. The mean annual rainfall varies from 500 ~~-to~~ 1000 mm covering 45–47% of the mean annual potential evapotranspiration (PET) ranging between 1500-~~and~~ 1800 mm. To estimate the distribution of SOC stock and pools of different oxidisability among different land use patterns, we identified four land use patterns in Vikarabad district, viz.: 1) 100% cropping intensity (redgram-fallow), 2) 200% cropping intensity (Rice-rice), 3)

Comment [GK3]: Units are not coherent. You are mixing degrees decimal and degrees-minutes.

Formatted: Superscript

Comment [GK4]: Units are not coherent. You are mixing degrees decimal and degrees-minutes.

Comment [GK5]: Not clear

Forest land, and 4) Fallow land. A total of 320 sites were selected for the present study. Details of the site characteristics including age of the land uses, dominant species and management practices of the land use studied were recorded with the help of the villagers and are presented in Table 1.

Table 1. Site characteristics of different land use patterns

Land use	Age (year)	Dominant species	Management practices
100% cropping intensity (Redgram-fallow)	10	<i>Cajanus cajan</i>	Deep summer ploughing and plough planting, regular weeding and fertilizer application with supplemental irrigation as and when required by the crop.
200% cropping intensity (Rice-rice)	10	<i>Oryza sativa</i>	Summer ploughing, plough planting and transplanting of the crop, application of fertilizers.
Forest land	40	<i>Tectona grandis</i> , <i>Eucalyptus</i> , <i>Acacia nilotica</i> , <i>Tamarindus indica</i> , <i>Leucaena leucocephala</i> , <i>Dalbergia sissoo</i> , <i>Ficus benghalensis</i> , <i>Pongamia pinnata</i>	Mild anthropogenic disturbances for occasional tree felling, frequent collection of fuel wood and other non-timber forest products
Fallow land	25	<i>Eulalia trispicata</i> , <i>Imperata cylindrica</i> , <i>Cyrrondon dactylon</i> , <i>Cyperus rotundus</i>	Grazing by animals and erosional losses

2.2 Soil sampling and sample preparation

A survey was carried out in the years (2019-20 and 2020-21) and soil samples at two depths (0-15 and 15-30cm) were collected from eight mandals in Vikarabad district predominantly covered by red soils. Five villages in each mandal were selected and from each village, four land use pattern were selected out of which two were from agricultural land use with different cropping intensity i.e., 100% cropping intensity (redgram-fallow cropping system), 200% cropping intensity (rice-rice cropping system) and two from natural conditions i.e., forest and fallow land. Within each land use plot soils were collected from five points (4 in the corners and 1 in the center) at two depth classes i.e. 0–15 and 15–30 cm respectively. The five sub-samples at each location and depth class were pooled to get one composite sample for each depth class per plot. A total of 320 soil samples were collected at two depths were obtained for analysis. The soils were mixed thoroughly and large plant debris, roots and stones were removed manually by hand. Soil samples from the same depths were collected from undisturbed plots separately for bulk density by using a soil corer of known volume. In the laboratory, the soil samples were homogenized, air-dried, grounded and passed through a 2 mm sieve for further analysis.

Comment [GK6]: Could you provide an English translation for this word.

2.3 SOIL ANALYSIS

The soil samples were analysed for salient characteristics like pH, [Electrical Conductivity \(EC\)](#), bulk density, [and](#) soil organic carbon following standard procedures. The soil organic carbon stock was calculated by multiplying OC content (g g^{-1}) with bulk density (Mg m^{-3}) and depth (m) of surface soil [12].

$$\text{Soil organic carbon stock (Mg C ha}^{-1}\text{)} = \text{OC content (g g}^{-1}\text{)} \times \text{bulk density (Mg m}^{-3}\text{)} \times \text{depth (m)}$$

2.3.1 Different Pools of soil organic carbon

They were determined as described by [11] and [13]. One g of 0.5 mm sieved soil was taken in 500mL Erlenmeyer flask in three sets. For each set 10mL of 1N potassium dichromate was pipetted out and transferred to [the](#) Erlenmeyer flask. To the first set 5mL of concentrated H_2SO_4 was added, to the second set 10mL of concentrated H_2SO_4 and to the third set 20 mL of concentrated H_2SO_4 was added which corresponds to 12.0 N, 18.0 N and 24.0 N of H_2SO_4 . All the Erlenmeyer flasks were kept aside for 30 min to complete the reaction. Then 200 mL distilled water, 10 mL of 85% orthophosphoric acid, a pinch of NaF and 1 ml of diphenylamine indicator were added and titrated with 0.5N ferrous ammonium sulphate till color changed from violet blue to green. A blank was also titrated in similar manner to determine the quantity of ferrous ammonium sulfate consumed. The values of different organic C pools were expressed in Mg ha^{-1} .

The different pools of oxidisable organic carbon were determined as given below

1. Very labile (VL) Organic C oxidisable under 12.0 N H_2SO_4
2. Labile (L) The difference in SOC extracted between 18.0 N and 12.0N H_2SO_4
3. Less labile (LL) The difference in SOC extracted between 24.0 N and 18.0 N H_2SO_4
4. Non-labile (NL) Obtained by subtraction of VL, L and LL from TOC

2.3.2 Total Organic Carbon (TOC): Since access to TOC analyzer was not possible, the TOC was calculated by the formula given by [14].

$$\log_{10}(\text{TOC}) = 0.725 \times \log_{10}(\text{WBC}) + 0.198 \times \log_{10}(\text{SICL}) - 0.0759 \times \log_{10}(\text{MAR}) + 0.015$$

Where TOC is total soil organic C (%), WBC is the uncorrected Walkley and Black carbon (%) determined by wet digestion, SICL is the silt + clay content (%) and MAR is the [main-mean](#) annual rainfall (mm) of the region.

2.3.3 Active and Passive Pools of Soil Organic Carbon

The very labile and labile pool may be summed up and it may be designated as active pool. Similarly, less labile and non-labile pool may be summed up and designated as passive pool.

2.4 STATISTICAL ANALYSES

The data collected was statistically analyzed using SPSS statistical package version 18.0. All the soil parameters were analyzed by two-way ANOVA. The Duncan's multiple range test was used to segregate the significance of difference among the mean values obtained for soil parameters, in each land use pattern at $P=0.05$ which were considered to be statistically significant at 5% level of significance.

3. RESULTS AND DISCUSSION

The data on salient soil characteristics viz; bulk density, pH, EC, organic carbon and organic carbon stock are presented in (Table 2). The soil samples in general were sandy loam to sandy clay loam in texture with clay percent ranging from 35.96 to 40.28 %.

Comment [GK7]: In the materials and methods, include the methodology for soil texture.

Bulk density (Mg m^{-3})

The data pertaining to soil bulk density is presented in (Table 2). The values on an average, indicated that 200% cropping intensity (rice-rice) recorded the highest bulk density (1.45 Mg m^{-3}) followed by 100% cropping intensity (1.41 Mg m^{-3}), fallow land (1.39 Mg m^{-3}) and forest land recorded the lowest bulk density (1.33 Mg m^{-3}). The interaction effect between depth and land use pattern for soil bulk density was significant. Across the depth at 15- 30cm the highest BD was obtained by 200% cropping intensity (1.48 Mg m^{-3}) this was followed by 100% cropping intensity (1.44 Mg m^{-3}), fallow land (1.42 Mg m^{-3}) and the lowest were recorded under forest land use (1.36 Mg m^{-3}). It was found that there was an increase in bulk density from 0- 15cm (1.37 Mg m^{-3}) to 15-30cm (1.43 Mg m^{-3}). The increase in bulk density with increase in depth in all the land use system may be attributed to lower organic matter content and soil compaction from the pressure of upper soil layer [15].

Soil pH and EC

The soil pH under different land use pattern on an average, indicated that the soils under forest land (Table 2) recorded the lowest soil pH (6.68), this was followed by fallow land (7.43), 100% cropping intensity (redgram-fallow) (7.59) and the highest was recorded in 200% cropping intensity (rice-rice) (7.72). Among the depths there was increase in soil pH from (7.27) to (7.43). Interaction effect between soil pH and land use pattern was significant with forest soils recording the lowest soil pH (6.56) followed by fallow land (7.38), 100% cropping intensity (7.50), 200% cropping intensity (7.64) at 0-15cm. At 15-30cm soil depth the pH of the soil increased and same trend was followed among all the land use pattern.

The highest soil electrical conductivity was recorded under 200% cropping intensity (rice-rice) (0.28 dSm^{-1}), this was followed by 100% cropping intensity (redgram-fallow) cropping pattern (0.25 dSm^{-1}), fallow land (0.23 dSm^{-1}) and the lowest was recorded under forest land use (0.18 dSm^{-1}). With increase in soil depth the EC of the soil decreased in all the land use patterns. The interaction between depths and land use pattern for soil EC was significant with highest EC being recorded by 200% cropping intensity (rice-rice) (0.30 dSm^{-1}) at 0-15 depth. A close perusal of the data on soil pH and EC indicates that the soils sites having lower pH shows lower EC values especially in forest and fallow lands. Similar results were obtained by [16] and [17]. An increase in total soluble salt content has been reflected by an increase in EC under cultivated soils this which could be due to the addition of fertilizers and other amendments.

Formatted: Superscript

Soil organic carbon (g kg^{-1})

The data pertaining to soil organic carbon is presented in (Table 2). Irrespective of soil depth on an average, the highest soil organic carbon content was recorded under forest land (7.05 g kg^{-1}) which was followed by 200% cropping intensity (rice-rice) (4.14 g kg^{-1}), 100% cropping intensity (redgram-fallow) (3.42 g kg^{-1}) and lowest was seen in fallow land (2.68 g kg^{-1}). However with increase in depth the mean organic carbon decreased from (4.72 g kg^{-1}) to (3.87 g kg^{-1}). Interaction effect between soil depth and land use pattern was significant and it varied from (7.30 g kg^{-1}) in forest soils

at 0-15cm to (2.24 g kg⁻¹) in fallow land at 15-30cm. Higher amount of organic carbon under forest land could be attributed to leaf litter decomposition at the surface. It was observed that about 60-80% total carbon resources are in oxidisable form due to the presence of higher amount of soluble extractives like fat, waxes and alcohol soluble extractives in forest residues [18]. Among the cropping systems, 200% cropping intensity (rice-rice) ~~has shown~~ showed significantly higher SOC. This might be due to continuous submergence of soils for 8-9 months in ~~an a~~ year under rice-rice cropping system. ~~P~~rolonged water logging conditions may reduced the decomposing of added crop residues [11]. However the lower values of SOC under fallow lands could be attributed to the very low amount of addition of residues in the form of leaf litter to the soil though the soil was not disturbed for ~~a~~ longer period of time.

Soil organic carbon stock (Mg C ha⁻¹)

The values of soil organic carbon stock determined for different land use patterns are presented in (Table.2) and depicted in (Fig.1). The soil organic carbon stock varied across the land use patterns and soil depth. Among the land use patterns compared on an average, the highest soil organic carbon stock was recorded in forest land (13.92 Mg ha⁻¹) ~~,which was~~ followed by 200% cropping intensity (9.00 Mg ha⁻¹), 100% cropping intensity (7.28 Mg ha⁻¹) and ~~the~~ lowest was ~~seen observed~~ in fallow land (5.64 Mg ha⁻¹). Soil organic carbon stock ~~recorded showed~~ a significant decrease in soil along the depth at lower layers from (9.68 Mg ha⁻¹) to (8.24 Mg ha⁻¹). While considering the land use patterns across the depth the interaction effect for soil organic carbon stock was significant and the forest land use recorded the highest mean (14.33 Mg ha⁻¹) ~~,which was~~ followed by 200% cropping intensity (9.71 Mg ha⁻¹) and ~~the~~ lowest was recorded in fallow land (6.51 Mg ha⁻¹) at 0-15cm. At 15-30 cm soil depth the same trend was ~~followed observed~~ with the highest SOCS being recorded in forest land (13.50 Mg ha⁻¹), 200% cropping intensity (8.29 Mg ha⁻¹), 100% cropping intensity (6.39 Mg ha⁻¹) and ~~the~~ lowest was recorded in fallow land (4.77 Mg ha⁻¹).

Among the land use patterns the highest SOCS in the forest could be related to the presence of complete land cover generating more litter fall which are returned to the soil as organic matter. Hence there is a possibility that forest land use system can be considered as a stabilized system with very little disturbance. The lower content of soil organic carbon stock under 100% cropping intensity could be mainly due to no organic manure application, fallow without adequate crop residues cover after completion of crop cycle combined with poor soil structure [19]. The lowest values ~~were for soil organic carbon stock~~ obtained ~~for soil organic carbon stock~~ under fallow land could be due to the lack of shade under fallow land which facilitates the loss of carbon from soil [20]. The higher soil organic carbon stock in the surface layer could be due to the addition of cropping system organic matter, more root biomass and root exudates. The difference between surface and sub surface stock which is higher in forest land followed by 200% cropping intensity, 100% cropping intensity and lowest in fallow land ~~, could~~ be due to the addition of fresh organic matter through leaf fall in forest soils and addition of FYM in cropping system in the surface horizons.

Among the cropping systems ~~s~~, 200% cropping intensity (rice - rice) recorded highest SOC stock which can be attributed to anerobiosis and the associated chemical and biochemical changes that take place in the submerged soils following the prolonged flooding under water. ~~Further Also~~, the

decomposition of organic matter in the absence of oxygen is slow, incomplete and inefficient while under aerobic conditions oxygen acts as electron acceptor hastening the decomposition process. In addition, in submerged soils the formation of recalcitrant complexes with organic matter renders them less available for microbial attack. It was also noticed that higher productivity of wetlands and decreased humification of soil organic matter lead to the net accumulation of organic matter in wet lands soil.

Table 2. Effect of land use patterns on soil characteristics and soil organic carbon stock in Red soils of Vikarabad district

Depth	Bulk density (Mg m ⁻³)	pH	EC (dSm ⁻¹)	Organic carbon (g kg ⁻¹)	Soil organic carbon stock (Mg C ha ⁻¹)
D ₁ (0-15cm)	1.37 ^A	7.27 ^B	0.25 ^A	4.72 ^A	9.68 ^A
D ₂ (15-30cm)	1.43 ^B	7.43 ^A	0.23 ^B	3.87 ^B	8.24 ^B
Land use pattern					
L ₁ :100% cropping intensity (redgram-fallow)	1.41 ^b	7.59 ^b	0.25 ^{ab}	3.42 ^c	7.28 ^c
L ₂ :200% cropping intensity (rice-rice)	1.45 ^a	7.72 ^a	0.28 ^a	4.14 ^b	9.00 ^b
L ₃ :Forest land	1.33 ^d	6.68 ^d	0.18 ^b	7.05 ^a	13.92 ^a
L ₄ :Fallow land	1.39 ^c	7.43 ^c	0.23 ^c	2.68 ^d	5.64 ^d
Interaction					
D ₁ L ₁	1.39±0.01 ^d	7.50±0.05 ^c	0.27±0.04 ^{ab}	3.90±0.21 ^d	8.16 ± 0.53 ^d
D ₁ L ₂	1.43±0.02 ^{bc}	7.64±0.07 ^b	0.30±0.03 ^a	4.54±0.18 ^c	9.71 ± 0.38 ^c
D ₁ L ₃	1.30±0.02 ⁱ	6.56±0.09 ⁱ	0.19±0.01 ^{de}	7.30±0.14 ^a	14.33 ± 0.44 ^a
D ₁ L ₄	1.36±0.01 ^e	7.38±0.09 ^d	0.25±0.02 ^{bc}	3.12±0.16 ^e	6.51 ± 0.40 ^e
D ₂ L ₁	1.44±0.01 ^b	7.66±0.04 ^b	0.24±0.04 ^{bc}	2.94±0.26 ^e	6.39 ± 0.63 ^e
D ₂ L ₂	1.48 ± 0.02 ^a	7.79±0.05 ^a	0.26±0.03 ^{ab}	3.73 ± 0.24 ^d	8.29 ± 0.47 ^d
D ₂ L ₃	1.36±0.03 ^{ef}	6.80±0.06 ^e	0.17±0.01 ^e	6.80±0.15 ^b	13.50 ± 0.36 ^b
D ₂ L ₄	1.42 ± 0.01 ^b	7.48±0.03 ^c	0.22±0.02 ^{cd}	2.24 ± 0.16 ^f	4.77 ± 0.32 ^f

Mean values with different lower case superscript letters indicate significant difference between land use patterns for each soil depth and all land uses. Uppercase superscript letters indicate significant difference between depths for all land use system respectively at ($P=0.05$), ± indicates standard deviation of mean




Fig 1. Soil organic carbon stock (Mg C ha⁻¹) at two soil depths under different land use patterns in red soils

Active and Passive Soil Organic Carbon Pools

Very labile (VL) carbon pools varied across the land use patterns and soil depth and is present in (Table 3). It varied significantly among the land use patterns and soil depths. Among land use patterns compared on an average, the forest land recorded the highest content of very labile carbon (9.05 Mg ha⁻¹) which was followed by 200% cropping intensity (4.95 Mg ha⁻¹), 100% cropping intensity (3.80 Mg ha⁻¹) and the lowest was observed in fallow land (2.40 Mg ha⁻¹) irrespective of depths (2.40 Mg ha⁻¹).

A decrease in the content of this pool of SOC was noticed between 0-15 to 15-30 cm depth in all the land use patterns from (5.05 Mg ha⁻¹) to (4.50 Mg ha⁻¹). Interaction effect between depth and land use patterns was significant and among the land use patterns, the highest was recorded in forest land at 0-15 cm (9.60 Mg ha⁻¹) and the lowest VL pool of carbon was recorded in fallow land at 15-30 cm soil depth (1.80 Mg ha⁻¹). The decrease in very labile pools of soil carbon can be attributed to the lack of carbon supply from the top layers. Similar results were obtained by [20]. [21] reported higher content of very labile pool under forest land suggesting higher input of organic matter through plant residues. This increase is mainly related to free light fraction of soil organic carbon. Further Also, the higher levels of very labile pool in forest land can be attributed to no turning of soil and constant supply of plant residues [4, 22] [22] and [4]. In addition the higher amounts of very labile fraction which is sensitive to management practices [23] depends on largely on the amount of organic residue added to the soil [10] and [13]. This explains the reason for higher amounts of accumulation of very labile pool in surface layer as compared to subsurface layer. Percent contribution of VL pool to that of TOC (Fig2) was found to be highest in forest land (34.53 %) followed by 100% cropping intensity (28.75 %), 200% cropping intensity (26.97%) and the lowest was recorded in fallow land (25.88 %).

Labile pools of soil carbon under different land use patterns are presented in (Table 3). The amount of labile pool of SOC significantly varied among the land use patterns. On an average, its

amount was highest in forest land (5.25 Mg ha^{-1}) followed by 200% cropping intensity (3.70 Mg ha^{-1}) and 100% cropping intensity (3.22 Mg ha^{-1}). Both were ~~enpar-compared~~ with each other and ~~the~~ lowest was recorded in fallow land (1.77 Mg ha^{-1}). There were variations in the content of these pools of SOC along depth under different land ~~forms~~ ~~uses~~. A ~~d~~Decreasing trend in the content of labile pool of SOC with increasing depth was seen in all land use patterns from (3.48 Mg ha^{-1}) to (3.08 Mg ha^{-1}). Interaction effect between depth and land use pattern was significant with forest land recording the highest labile pool at 0-15cm (5.76 Mg ha^{-1}) and ~~the~~ lowest ~~was recorded~~ by fallow land at 15-30cm soil depth (1.44 Mg ha^{-1}).

Percent contribution of labile pool to that of TOC (Fig.2) was found highest in 100% cropping intensity (24.40%) followed by 200% cropping intensity (20.16%), forest land (22.67 %) and ~~the~~ lowest was recorded fallow land (19.09%). Less labile pool of carbon under different land use patterns are presented in (Table 2). It varied significantly among the land use patterns and depth. Among the land use patterns on an average, fallow land recorded the lowest less labile pool (2.20 Mg ha^{-1}) which was ~~enpar-compared~~ with 100% cropping intensity (2.46 Mg ha^{-1}) followed by 200% cropping intensity (3.30 Mg ha^{-1}) and ~~the~~ highest was recorded in forest land (3.50 Mg ha^{-1}). Whereas when the percent contribution of less labile pool to that of TOC was calculated (Fig.2) it was found that less labile pool percent was found highest in fallow land (23.73%) followed by 100% cropping intensity (18.61%), 200% cropping intensity (17.98 %) and lowest was recorded under forest land (13.35%).

The less labile carbon pools decreased with increase in soil depth. The clay content of soil could affect the changes and stabilization of carbon pools. ~~Due to The~~ natural and undisturbed conditions in forest land and ~~in-coupled with the~~ absence of tillage practices might have contributed to higher amount of less labile carbon. On the other hand lower values of less labile pool under the ~~agriculture-cropping~~ land use system could be attributed to lesser accumulation of organic carbon in conventional agricultural systems involving ploughing and other intercultural operations [24]. Similar results of less labile pools ~~were also have been~~ reported by [25] and [26].

The higher amount of less labile pool of carbon in 200% cropping intensity could be attributed to retarded rate of carbon oxidation in the soils ~~that~~ are flooded for almost 3 to ~~3.5~~ ~~4/2~~ years during rainy season [27]. This must have resulted in slower decomposition of carbon substances in soil as compared to upland conditions and the molecules like lignins, waxes may gradually accumulate in soil contributing to less labile pool. ~~Further-Also,~~ slower decomposition of submerged soils as compared to uplands soils lead to incorporation of phenolic substances in the soils organic matter fractions making them recalcitrant to decomposition [28]. Similar results ~~were also have been~~ reported by [29].

Land use patterns have significantly influenced the non labile pool of SOC and are presented in (Table.3). On ~~an~~ average, its amount was highest ~~with-in~~ forest land (8.40 Mg ha^{-1}) followed by 200% cropping intensity (6.40 Mg ha^{-1}), 100% cropping intensity (3.73 Mg ha^{-1}) and ~~the~~ lowest was recorded in fallow land (2.90 Mg ha^{-1}). With increase in depth the non labile pool on an average decreased from (5.35 Mg ha^{-1}) to (5.25 Mg ha^{-1}). Interaction effect between depth and land use patterns was significant and forest land recorded the highest non labile pool at both ~~the~~ soil depths.

Percent contribution of non labile pool of SOC to TOC (Fig.2) under different land use forms was opposite to that of very labile and labile pools, ~~and~~ the trend was ~~followed-asas follows~~: 200%

Formatted: Not Superscript/ Subscript

cropping intensity (34.87%) followed by forest land (32.05 %), fallow land (31.28%) and lowest was recorded under 100% cropping intensity (28.22%). With increase in ~~depth of soil~~ soil depth, the non labile pool of soil carbon decreased in all the land use patterns ~~except except the~~ 200% cropping intensity in which it showed an increase. Higher values of non labile pool under 200% cropping intensity can be attributed to fairly stable and more recalcitrant carbon content in soil which increases with increase in soil depth. ~~As almost 7-8 months the soil remains~~ Given that the soil remains for almost 7-8 months under submergence ~~and~~, this prolonged ~~submergence condition~~ might retard the oxidation and enhance the contribution of recalcitrant carbon to soil organic carbon. [30] also reported higher proportions of passive non labile pool in forest and it could be due to high input of plant litter in forest as compared to other land use patterns. They suggested that organic carbon in forest soils is more stable than the other land use types studied. The results of the present study indicate that appropriate management practices are needed to prevent the losses of soil organic carbon through decomposition in different land use patterns. further-Also, the land use systems s affect both the active and passive carbon pools. Our results are in accordance with findings of [31].

The amount of different pools of SOC at-in surface soil was as follows: very labile pool followed by non labile pool, labile pool and less labile pool, constituting about 27.77%, 29.46%, 19.18% and 15.57% of the TOC, respectively. At 15-30cm depth, the amount of different SOC pools-of SOC ~~followed the trend were~~ as follows: non labile ~~followed by~~, very labile, labile, and less labile pool constituting about 34.25%, 29.33%, 20.11% and 16.29 % of the TOC, respectively.

Total organic carbon

Total organic carbon (TOC) content ranged from 7.64 to 27.86 Mg ha⁻¹ under different land use patterns and depths (Table.3). On ~~an~~ average, the forest land had maintained significantly higher amount of mean TOC (26.15 Mg ha⁻¹) followed by 200% cropping intensity (18.35 Mg ha⁻¹), 100% cropping intensity (13.21 Mg ha⁻¹) and the lowest was recorded under fallow land (9.27 Mg ha⁻¹). With increase in soil depth the total organic carbon decreased from (18.18 Mg ha⁻¹) to (15.31 Mg ha⁻¹). Interaction effect between depth and land use pattern for TOC was significant with forest soils recording the highest TOC at 0-15cm (27.86 Mg ha⁻¹) and lowest was recorded at 15-30cm by fallow land (7.64 Mg ha⁻¹).

In our study, TOC content varied across different land use patterns and forest land had the highest TOC concentration. Total organic carbon content build up under any land use type is due to the balance between C inputs (litter fall deposits, crop residues, root exudates, root biomass and manure) and C losses (respiration by soil organisms) determined by the residue turnover, their quality and decomposition rate. A near-equilibrium between C inputs and C losses was found in undisturbed ecosystems. Higher SOC content in forest land use types in our ~~present~~ study could be due to high levels of inputs (leaf litter and root biomass) from the forest trees and its recalcitrant nature thus preventing microbial decomposition which in turn increased the TOC content in soil [29].

Table 3. Effect of land use patterns on soil carbon pools (Mg ha⁻¹) in red soils of Vikarabad district

	Very labile (Mg ha ⁻¹)	Labile (Mg ha ⁻¹)	Lesslabile (Mg ha ⁻¹)	Nonlabile (Mg ha ⁻¹)	Active pool (Mg ha ⁻¹)	Passive pool (Mg ha ⁻¹)	TOC (Mg ha ⁻¹)
Depth							
D ₁ (0-15cm)	5.05 ^A	3.48 ^A	2.85 ^A	5.35 ^A	9.49 ^A	8.69 ^A	18.18 ^A
D ₂ (15-30cm)	4.50 ^B	3.08 ^B	2.47 ^B	5.25 ^B	7.58 ^B	7.73 ^B	15.31 ^B
Land use patterns							
L ₁ : 100% cropping intensity (redgram-fallow)	3.80 ^b	3.22 ^b	2.46 ^c	3.73 ^c	7.02 ^c	6.19 ^{bc}	13.21 ^c
L ₂ : 200% cropping intensity (rice-rice)	4.95 ^c	3.70 ^b	3.30 ^b	6.40 ^b	8.65 ^b	9.70 ^b	18.35 ^b
L ₃ : Forest land	9.05 ^a	5.25 ^a	3.50 ^a	8.40 ^a	14.30 ^a	11.85 ^a	26.15 ^a
L ₄ : Fallow land	2.40 ^d	1.77 ^c	2.20 ^c	2.90 ^d	4.17 ^d	5.10 ^d	9.27 ^d
Interaction							
D ₁ L ₁	4.50 ± 0.32 ^d	3.70 ± 0.16 ^{cd}	2.82 ± 0.18 ^c	4.04 ± 0.29 ^d	8.20 ± 0.43 ^d	6.86 ± 1.51 ^d	15.06 ± 1.75 ^e
D ₁ L ₂	5.30 ± 0.35 ^c	4.00 ± 0.32 ^c	3.06 ± 0.17 ^b	6.00 ± 0.35 ^c	9.30 ± 0.59 ^c	9.60 ± 0.47 ^c	18.90 ± 0.40 ^c
D ₁ L ₃	9.60 ± 0.19 ^a	5.76 ± 0.36 ^a	3.90 ± 0.20 ^a	8.60 ± 0.69 ^a	15.36 ± 0.49 ^a	12.50 ± 0.43 ^a	27.86 ± 0.49 ^a
D ₁ L ₄	3.00 ± 0.49 ^e	2.10 ± 0.20 ^f	2.60 ± 0.30 ^{cd}	3.20 ± 0.30 ^{de}	5.10 ± 0.50 ^f	5.80 ± 0.52 ^e	10.90 ± 0.82 ^g
D ₂ L ₁	3.10 ± 0.37 ^e	2.75 ± 0.29 ^e	2.10 ± 0.33 ^d	3.42 ± 0.22 ^{de}	5.85 ± 0.45 ^e	5.52 ± 0.46 ^e	11.37 ± 0.81 ^f
D ₂ L ₂	4.60 ± 0.31 ^d	3.40 ± 0.26 ^d	3.00 ± 0.21 ^b	6.80 ± 0.41 ^b	8.00 ± 0.35 ^d	9.80 ± 0.39 ^c	17.80 ± 0.49 ^d
D ₂ L ₃	8.50 ± 0.18 ^b	4.75 ± 0.26 ^b	3.10 ± 0.23 ^b	8.20 ± 0.80 ^a	13.25 ± 0.39 ^b	11.20 ± 0.65 ^b	24.55 ± 0.62 ^b
D ₂ L ₄	1.80 ± 0.36 ^f	1.44 ± 0.20 ^g	1.80 ± 0.27 ^e	2.60 ± 0.26 ^f	3.24 ± 0.35 ^g	4.40 ± 0.61 ^f	7.64 ± 0.60 ^h

Mean values with different lower case superscript letters indicate significant difference between land use patterns for each soil depth and all land uses.

Uppercase superscript letters indicate significant difference between depths for all land use system respectively at (P<0.05). ± indicates standard deviation of mean.

The content of active SOC pool of SOC on an average (Table 3) was significantly higher under forest land (14.30 Mg ha^{-1}) which was followed by 200% cropping intensity (8.65 Mg ha^{-1}), 100% cropping intensity (7.02 Mg ha^{-1}) and the lowest was recorded in fallow land (4.17 Mg ha^{-1}). Interaction effect between land use patterns and soil depth for active pool was significant. With increase in depth active pool (AP) of soil organic carbon declined in all the land use patterns from (9.49 Mg ha^{-1}) to (7.58 Mg ha^{-1}). Interaction effect between depth and land use pattern was significant with forest soils having the highest active carbon pool at 0-15cm (12.50 Mg ha^{-1}).

The percent contribution of AP to TOC (Fig.3) was found highest in forest land (60.00%) followed by 100% cropping intensity (53.15 %), 200% cropping intensity (47.13%) and lowest in fallow land (44.98%).

Passive pool (PP) of SOC was also significantly influenced by land use patterns and is presented in (Table 3). On an average, the content of passive pool of SOC was significantly higher under forest land (11.85 Mg ha^{-1}), which was followed by 200% cropping intensity (9.70 Mg ha^{-1}), 100% cropping intensity (6.19 Mg ha^{-1}), and fallow land (5.10 Mg ha^{-1}). With increase in depth, on an average, passive carbon pools decreased from (8.69 Mg ha^{-1}) at 0-15cm to (7.73 Mg ha^{-1}) at 15-30cm soil depth. The percent contribution of passive pool to TOC (Fig.3) followed different trends where, fallow land contributed the highest percent (55.01%), which was followed by 200% cropping intensity (52.86%), forest land (41.00%) and the lowest was reported recorded by 100% cropping intensity (46.84%). The interaction effect between land use patterns and depth was significant with active and passive carbon pool decreasing with soil depth in all the land use patterns except in 200% cropping intensity where passive carbon pool increased with increase in depth of soil from 0-15 to 15-30cm.




Fig 2. Percent contribution of various pools of carbon of varying lability (0-30cm) towards total organic carbon (TOC) in different land use patterns




Fig 3. Distribution of active and passive pools up to 0-30cm depth (as % of TOC) under different land use patterns.

4. CONCLUSIONS

Within the 0–30 cm soil profile, TOC content and SOC stocks were higher-highest in forest land than the other land use patterns indicating the depletion of TOC with land use changes. The majority of soil organic carbon of forest land and 100% cropping intensity (redgram-fallow) was in an easily oxidisable active form, suggesting that the accumulated carbon could be easily lost following the land use changes. Conversely, preponderance of passive carbon pools in fallow land and 200% cropping intensity (rice-rice system) indicated a more stable nature of the accumulated soil organic carbon. The data acquired revealed that the applied soil management regimes had significant effects of varying magnitude on TOC and its fractions. Thus, size and distribution of active and passive carbon pools can be used as indicators for change in soil quality.

REFERENCES

1. Lal R. Soil carbon dynamics in cropland and rangeland. *Environmental Pollution*. 2002;116(3): 353–362.
2. Pan G, Zhou P, Li Z, Pete S, Li L, Qiu D, Zhang X, Xu X, Shen S, Chen X. Combined inorganic/organic fertilization enhances N efficiency and increases rice productivity through organic carbon accumulation in a rice paddy from the Tai Lake region, China. *Agr.* 2009;52:65.
3. Kirschbaum MUF. Will changes in soil organic carbon act as a positive or negative feed-back on global warming? *Biogeochemistry*. 2000;48: 21–51.
4. Srinivasarao Ch, Venkateswarlu B, Lal R, Singh A K and Kundu S. Sustainable management of soils of Dryland ecosystems of India for enhancing agronomic productivity and sequestering carbon. *Advances in Agronomy*. 2013;121:253-329.
5. Tedla A, Zhou X, Su B, Wan S, Luo Y. Labile, recalcitrant, and microbial carbon and nitrogen pools of a tallgrass prairie soil in the US Great Plains subjected to experimental warming and clipping. *Soil Biol Biochem.* 2009;41:110–116.
6. Poeplau C, Don A. Sensitivity of soil organic carbon stocks and fractions to different land-use changes across Europe. *Geoderma* 2013;192:189–201.
7. Zhang L, Xie Z, Zhao RF, Wang YJ. The impact of land use change on soil organic carbon and labile organic carbon stocks in the Longzhong region of Loess Plateau. *Journal of Arid Land* 2012; 4(3):241–250

8. IPCC. ClimateChange.Mitigation of Climate Change.2007. Cambridge University Press, Cambridge.
9. Blair GJ, Lefroy RDB and Lisle L. Soil carbon fractions based on their degree of oxidation and the development of a carbon management index for agricultural systems. Australian Journal of Soil Research.1995;46:1459-1466.
10. Chen CR, Xu ZH and Mathers NJ. Soil carbon pools in adjacent natural and plantation forests of subtropical Australia.Soil Science Society of America Journal.2004;68:282-291.
11. Mandal B, Majumder B, Adhya TK, Bandyopadhyay PK, Gangopadhyay A, Sarkar D, Kundu MC, Choudhury SG, Hazra G C, Kundu S, Samantaray RN and Mishra AK. The potential of double-cropped rice ecology to conserve organic carbon under subtropical climate. Global Change Biology.2008;14: 2139-2151.
12. Bhattacharya R, Tuti MD, Kundu S, Bisht J K and Bhatt J C. Conservation tillage impacts on soil aggregation and carbon pools in a sandy clay loam soil of the Indian Himalayas. Soil Science Society of America Journal.2012;76:1-11.
13. Chan KY, Bowman A, Oates AL.2001.Oxidizable organic carbon fractions and soil quality changes oxic paleustaff under different pastures. Soil science.2001;166:61-67.
14. Jha P, Lakaria BL, Biswas A K, Srivastava S, Neenu R and Subba Rao. A manual on soil carbon analysis. Indian institute of soilscience, Bhopal.2003;41.
15. Devi B, Bhardwaj D R, Panwar P, Pal S, Gupta N K and Thakur C L. Long term effects of natural and plantation forests on carbon sequestration and soil properties in mid-hill subhumid condition of Himachal Pradesh, India. Soil Biology and Biochemistry.2013;34(1):19-2.
16. Barros, J.D.S and Chaves, L.H.G.2014.Changes in soil chemical properties under different farming systems exploration in semiarid region of Paraiba. 9(31):2436-2442
17. Shah S, Sharma D P, Pala N A, Tripathi P and Dar A. Carbon stock and density of soils under pine (Sargent) forests of Solan forest division, Himachal Pradesh. Soil Biology and Biochemistry.2013;41(3): 279-286.
18. Lorenz K, Lal R and Shipitalo MJ. Stabilization organic carbon in chemically separated pools in no-till and meadows soils in Northern Appalachia.Geoderma.2006;52-56.
19. Meena VS, Mondal T, Pandey BM, Mukherjee A, Yadav RP, Choudhary M, Singh S, Bisht JK and Pattanayak A. Land use changes: strategies to improve soil carbon and nitrogen storage pattern in the mid-Himalaya ecosystem, India.Geoderma.2018; 321:69–78.
20. Gupta M K and Sharma S D.Sequestered Organic Carbon Stock in the Soils under Different Land Uses in Uttarakhand State of India.Journal of Life Sciences Research.2014;1(1):5-9.
21. Rangel OJP, Silva CA, Guimaraes PTG and Guilherme L R G.Fractions of oxidizable organic carbon *Oxisol* cultivated with coffee at different planting spacings. *Agrotechnology and Science*.2008;32: 429-437.
22. Lal R. Sequestering carbon in soils of agro-ecosystems. Food Policy.2011;36:33-39.
23. Guareschi RF, Pereira, MG and Perinee A. Oxidizable carbon fractions in red *Latosol* under different management systems. Revista Ciencia Agronomica.2013;44:242-250.
24. Bayer C, MartinNeto L, Mielniczuk J, Pavinato A, Dieckow J. Carbon sequestration in two Brazilian Cerrado soils under no-till. Soil and Tillage Research.2006;86:237-245.
25. Zhang L, Xie Z, Zhao R F and Wang Y J. The impact of land use change on soil organic carbon and labile organic carbon stocks in the Longzhong region of Loess Plateau. Journal of Arid Land.2005;4(3):241-250.
26. Benbi D K, Brar K , Toor A S, Singh P and Singh P. Soil carbon pools under poplar-based agroforestry, rice-wheat, and maize-wheat cropping systems in semi-arid India. Nutrient Cycling in Agroecosystem.2012;92:107-118.
27. Jarecki M K and Lal R.Crop management for soil carbon sequestration. Critical Reviews in Plant Sciences.2003;22:471-502.
28. Somasundarama RS, Chaudhary D, Awanish Kumar A, Biswas A K, Sinha N, Mohantya M, Hati K M, Jha P, Sankar M, Patra AK, Dalal R and Chaudhari S K. Effect of contrasting tillage and

cropping systems on soil aggregation, carbon pools and aggregate-associated carbon in rainfed Vertisols. *European Journal of Soil Science*.2013;69:879-891.

29. Sahoo UK, Singh SL, Gogoi A and Kenye A. Active and passive soil organic carbon pools as affected by different land use types in Mizoram, Northeast India. *Plos One*.2019;1-16.
30. Kalambukattu JG, Singh R and Arunkumar K. Soil carbon pools and carbon management index under different land use systems in the Central Himalayan region. *Soil and Plant Science*.2013;63(3):200-205.
31. Datta A, Basak N, Chaudhari S K and Sharma D K. Soil properties and organic carbon distribution under different land uses in reclaimed sodic soils of North-West India. *Geoderma Regional*.2015;4:134-146.

UNDER PEER REVIEW

UNDER PEER REVIEW