

## **Influence /Effect of Dates of Sowing and Planting Geometry on Yield, Nutrient Uptake and Economics of Baby corn**

**ABSTRACT:** During the *summer* season of 2018, a field experiment was conducted at Assam Agricultural University, Jorhat. The treatments consisted of four date of sowing viz., 20<sup>th</sup> February (D<sub>1</sub>), 2<sup>nd</sup> March (D<sub>2</sub>), 12<sup>th</sup> March (D<sub>3</sub>) and 22<sup>nd</sup> March (D<sub>4</sub>) in main plot and four planting geometry practices viz., 40 cm x 20 cm (S<sub>1</sub>), 40 cm x 25 cm (S<sub>2</sub>), 45 cm x 20 cm (S<sub>3</sub>) and 45 cm x 25 cm (S<sub>4</sub>), in sub-plot with three replications. The results revealed that 2<sup>nd</sup> March sowing recorded *higher* plant height, number of cobs per plant, weight of cob, cob yield and fodder yield which was at par with 12<sup>th</sup> March sowing and significantly higher than other sowing dates. Similar effects of these treatments were also observed in respect to N, P, K uptake and economics. Among the different planting geometry, yield of cob without husk and fodder yield, gross return and net return were found to be higher under spacing 45 cm x 20 cm as compared to others. Total N, P and K uptake were also found to be higher under 45 cm x 20 cm spacing than rest of the planting geometry. Planting geometry had non-significant effect on available N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O status of soil after harvest of the crop.

**Key words:** Baby corn, Planting geometry, Dates of sowing, Yield, Nutrient uptake.

**INTRODUCTION :** Baby corn is a very high value crop, used as vegetable, which can boost the economy of poor farmers by diversification in their agriculture. It forms a major use in various salads and soups. The waste portion which is not edible like stem, leaves, fiber *etc.*, can be fed to cattle as green fodder. Baby corn as a cash crop for intensive agro ecosystems where small farmers grow three or more crops in highly diverse cropping system. Baby corn as the name implies, is not genetically dwarf maize but it is the immature ear of normally grown maize harvested within 2-3 days of silking. Although agronomic requirements of baby corn are similar to normal maize except for the suitable variety, high plant population per unit area, nitrogen requirements and suitable date of sowing which need to be studied for local agro-climatic conditions. Other than different components, date of sowing and plant population per unit area are critical elements deciding yield and quality of baby corn. Date of sowing is a non-monetary input which plays significant role in production and productivity of any crop.

Optimum crop geometry is one of the important factors for higher production leading to efficient utilization of resources and also harvesting as much as solar radiation and in turn better photosynthesis. Above red mark sentence is not required. Optimum crop geometry is one of the important factors for higher productivity, by virtue of which there is efficient utilization of underground resources and also harvesting maximum solar radiation which in turn results in better photosynthesis as reported by Monneveux *et al.* [3]

**Materials and Methods:** The field experiment was carried out at Assam Agricultural University, Jorhat during the *summer* season, 2018. The site is situated at 26°47'N latitude and 94°12'E longitude with an altitude of 86.56 meter above the mean sea level. Check the MSL.

The experiment was laid out in a split-plot design with three replications. The treatments consisted of four dates of sowing viz., 20<sup>th</sup> February (D<sub>1</sub>), 2<sup>nd</sup> March (D<sub>2</sub>), 12<sup>th</sup> March (D<sub>3</sub>), 22<sup>nd</sup> March (D<sub>4</sub>) in main plot and four planting geometry practices viz., 40 cm x 20 cm (S<sub>1</sub>), 40 cm x 25 cm (S<sub>2</sub>), 45 cm x 20 cm (S<sub>3</sub>), 45 cm x 25 cm (S<sub>4</sub>), in sub-plot. The soil

of the experimental site was sandy loam in texture, acidic in reaction (pH 5.06), medium in organic carbon (0.74%), available N (232.21 kg/ha), available P<sub>2</sub>O<sub>5</sub> (25.36 kg/ha) and available K<sub>2</sub>O (168.72 kg/ha).

Jorhat district is characterized with sub-tropical and humid climate having warm summers and cold winters. This region receives an average annual rainfall of 2500 mm. The monsoon starts from the month of June and continues up to the month of September with the pre-monsoon shower starting from mid-March, most of which is received during the monsoon period. During the crop growth period, the weekly mean maximum temperature ranged from 23.7°C to 34.5°C and the weekly mean minimum temperature ranged from 14.5°C to 25.1°C. The weekly average relative humidity during the morning hours ranged from 85 per cent to 97 per cent while mean evening relative humidity ranged from 56 per cent to 82 per cent.

The total precipitation received was 617.5 mm during the crop season from 19<sup>th</sup> Feb to 15<sup>th</sup> June, 2018, out of which maximum amount (86.3 mm and 99.4 mm) was received in the last week of May and second week of June. The bright sunshine hours during the crop growing season ranged from 1.0 to 7.1 hours/day.

## RESULTS AND DISCUSSION

### Growth, yield attributes and baby corn yield without husk (q/ha)

Data in Table 1 revealed that sowing on 2<sup>nd</sup> March recorded highest plant height which was at par with 12<sup>th</sup> March sowing. Sowing dates had a significant effect on number of cobs per plant and cob weight without husk. Maximum number of cobs per plant and weight of cobs without husk were recorded in 2<sup>nd</sup> March sowing which was at par with 12<sup>th</sup> March sowing and significantly higher than the other dates of sowing. The maximum baby corn yield without husk was found (18.08 q/ha) on 2<sup>nd</sup> March sowing which was statistically comparable with the yield obtained in 12<sup>th</sup> March and both were significantly higher than the sowing dates 20<sup>th</sup> Feb and 22<sup>nd</sup> March. This might be due to the fact that late sowing dates were badly affected by higher rainfall before harvesting. Another reason may be that 2<sup>nd</sup> March sowing of baby corn recorded highest value of growth and yield attributes which finally increased the baby corn yield. This might be due to the fact that timely sowing of baby corn resulted in greater partitioning of photosynthates from source to sink resulting in higher baby corn and fodder yield. Similar findings was reported by Tamadon [7]

Among the different planting geometry, highest plant height was recorded in 40 cm x 20 cm spacing which was at par with 45 cm x 20 cm. The higher plant height in closer spacing might be due to increase in competition for sunlight, nutrients, space and water by the plants. The results are in conformity with the findings of Verma *et al* [10]. Number of cobs per plant and weight of cobs per plant were highest in 45 cm x 25 cm. But baby corn yield without husk was the highest with spacing of 45 cm x 20 cm (16.96 q/ha) which was at par with 45 cm x 25 cm but significantly higher than 40 cm x 20 cm and 40 cm x 25 cm. The lowest yield of cob without husk was noticed with 40 cm x 20 cm (13.41 q/ha). Yield of the crop is function of several yield components which are dependent on complementary interaction between vegetative and reproductive growth of the crop. The highest yield of baby corn without husk was recorded with spacing 45 cm x 20 cm which might be due to optimum plant population (1, 11,000 plants/ha). The yield obtained under spacing 45 cm x 20 cm was statistically at par with 45 cm x 25 cm (89,000 plants/ha) but significantly higher than 40 cm x 25 cm (1, 00,000 plants/ha) and 40 cm x 20 cm (1, 25,000 plants /ha). The lowest yield was associated with spacing 40 cm x 20 cm. Under wider spacing of 45 cm x 25 cm, all the yield attributing characters were at their best but due of lesser plant population per unit area, it

could not compensate the baby corn yield obtained under spacing 45 cm x 20 cm. These results are in conformity with the findings of several earlier researchers Arvadiya *et al.* [1], Thavaprakaash and Velayudham [8], Ghosh *et al.* [2], Neelam and Dutta [4].

#### **Dry yield of baby corn (q/ha)**

Result presented on Table 1 indicated that the date of sowing had a significant effect on the dry yield of baby corn without husk. The highest dry yield of baby corn was recorded on sowing date 2<sup>nd</sup> March (3.05 q/ha), which was significantly higher than other date of sowing.

Dry yield of baby corn without husk was the highest with spacing of 45 cm x 20 cm (2.92 q/ha) which was at par with 45 cm x 25 cm but significantly higher than 40 cm x 20 cm and 40 cm x 25 cm.

#### **Fodder yield (q/ha)**

Table 1 indicates that the green fodder yield of baby corn was significantly influenced by date of sowing. The highest green fodder yield (276.55 q/ha) was recorded by sowing baby corn on 2<sup>nd</sup> March followed by 12<sup>th</sup> March (266.45 q/ha) which were at par with each other. The lowest green fodder yield was obtained in 22<sup>nd</sup> March (245.84 q/ha). It might be due to highest plant height and better growth in these two dates of sowing.

The effect of planting geometry on fodder yield was found to be significant. The maximum fodder yield was obtained in 45 cm x 20 cm spacing which was statistically at par with yield obtained in 40 cm x 20 cm but significantly higher than 40 cm x 25 cm and 45 cm x 25 cm spacing. The possible reason for increased yield under spacing 45 cm x 20 cm might be due to optimum number of plants per unit area resulting in higher green fodder yield of baby corn which is comparable with spacing 40 cm x 20 cm.

#### **Dry fodder yield (q/ha)**

The effect of date of sowing was found to be significant on dry fodder yield. The highest yield was recorded on 2<sup>nd</sup> March which was at par with 12<sup>th</sup> March. On the other hand the lowest yield was obtained in the sowing date 22<sup>nd</sup> March because late sown crop was badly affected by rainfall before harvest.

The effect of planting geometry on dry fodder yield was found to be significant. The maximum dry fodder yield was obtained in 45 cm x 20 cm spacing which was statistically at par with yield obtained in 40 cm x 20 cm but significantly higher than 40 cm x 25 cm and 45 cm x 25 cm. However, the yield obtained in spacing 40 cm x 25 cm was at par with 45 cm x 25 cm.

The green fodder yield obtained under closer spacing of 40 cm x 20 cm was lower as compared to 45 cm x 20 cm but was statistically at par with 45 cm x 20 cm. It might be because of the increased barrenness under 40 cm x 20 cm spacing where supply of growth factors such as light, water and nutrients to plants is affected by interaction between the plants and by the efficiency of use of limiting resources. The possible reason for increased dry fodder yield under spacing 45 cm x 20 cm might be due to optimum number of plants per unit area resulting in higher green fodder yield of baby corn which is comparable with spacing 40 cm x 20 cm. The lower green fodder yield was noticed under 45 cm x 25 cm spacing which was due to lesser number of plants per unit area under this planting geometry as compared to rest of the treatments. Similar results regarding the green fodder yield of baby corn were reported by Sobhana *et al.* [6], Thakur and Sharma [9] and Neelam and Dutta [4] who also envisaged that green fodder yield increased significantly with decrease in spacing *i.e.* with increase in plant population

### **Nitrogen content and uptake**

There was no significant difference was observed on nitrogen content in both baby corn and fodder (Table 2).

The effect of date of sowing was found to be statistically significant in case of nitrogen uptake by baby corn and fodder. The highest uptake was recorded under 2<sup>nd</sup> March sowing which was at par with 12<sup>th</sup> March sowing. This might be due to higher dry matter production in these dates. Similar results were observed in total N uptake.

The effect of planting geometry on nitrogen uptake by cob and fodder were found to be significant. The N uptake was the highest in 45 cm x 20 cm spacing which was comparable to 45 cm x 25 cm and 40 cm x 25 cm but all these were significantly higher than 40 cm x 20 cm. Total uptake also follows the similar trend. It might be because of the availability of growth resources (light, nutrient, moisture *etc.*) in optimum quantity.

### **Phosphorus content and uptake**

The effect of dates of sowing could not produce any significant difference in case of P content by cob and fodder. But phosphorus uptake by cob and fodder were significantly affected and highest was recorded on 2<sup>nd</sup> March sowing which was at par with 12<sup>th</sup> March sowing. Same trend was observed in total P uptake.

But planting geometry significantly affected the phosphorus uptake. Total P uptake by baby corn was higher in 45 cm x 25 cm which was statistically at par with spacing 45 cm x 20 cm but both were significantly higher than 40 cm x 20 cm and 40 cm x 25 cm. Significantly higher uptake by fodder was noticed in spacing 45 cm x 20 cm as compared to 40 cm x 20 cm but was at par with 45 cm x 25 cm and 40 cm x 25 cm.

There was significant effect of dates of sowing on total uptake of P by baby corn (Table 3). The significantly higher total P uptake was noticed in spacing 45 cm x 20 cm which was at par with 45 cm x 25 cm but both were significantly higher than the spacing 40 cm x 20 cm and 40 cm x 25 cm. This might be due to higher dry matter production in 45 cm x 20 cm which leads to higher uptake.

### **Potassium content and uptake**

The effect of date of sowing and planting geometry were found to be statistically non-significant on K content in cob and fodder.

Data presented in Table 4 showed that K uptake by the baby corn, fodder and total uptake were significantly influenced by the sowing dates and planting geometry. The highest K uptake was recorded in crop sown on 2<sup>nd</sup> March which was at par with 12<sup>th</sup> March and significantly higher than 22<sup>nd</sup> March sowing. Similar trend was observed in total K uptake also.

Among the different planting geometry, the highest uptake was noticed in spacing 45 cm x 20 cm which was at par with 45 cm x 25 cm but significantly higher than 40 cm x 25 cm and 40 cm x 20 cm. The lowest uptake of K was recorded under 40 cm x 20 cm spacing. Total uptake also followed the similar trend.

At wider spacing (45 cm x 25 cm), the rhizosphere will develop properly, without any mutual competition among the plants and thereby creating favourable environment for higher uptake of nutrients. N, P and K uptake in 40 cm x 20 cm spacing was significantly lesser than 45 cm x 20 cm and 45 cm x 25 cm because less amount of nutrients were available per plant due to different types of competition during different crop growth stages and thereby plants failed to extract desired quantity of nutrients from the soil. Higher N, P and K content by

fodder under the wider spacing of 45 cm x 25 cm might due to better vegetative growth of the plants due to more availability of sunlight, water and space, which favoured the growth of plants. These results are in close conformity with Thavaprakash and Velayudham [7], Sobhana *et al.* [5], Neelam and Dutta [3].

#### **Available nutrient content in soil (kg/ha)**

Different treatments failed to produce any significant differences in case of available nutrient status in soil after harvest.

#### **Economics**

The date of sowing 2<sup>nd</sup> March was found most remunerative and gave maximum gross returns (₹1,72,327.69/ha) and net returns (₹1,38,831.44 /ha) and benefit-cost ratio (4.15), followed by 12<sup>th</sup> March given gross returns (₹1,57,325.23/ha) and net returns (₹1,23,828.98/ha) and benefit-cost ratio (3.71). The lowest economic return was recorded during the sowing date 22<sup>nd</sup> March which gave gross returns (₹1, 22,930.60/ha) and net returns of (₹89,434.35/ha) and benefit-cost ratio (2.67).

In the present study, the spacing 45 cm x 20 cm recorded the higher gross return (₹1,62,763.61/ha), net return (₹1,29,209.61/ha) and benefit-cost ratio (3.85) than the other spacing treatments of 40 cm x 20 cm, 40 cm x 25 cm and 45 cm x 25 cm (Table 4). Spacing 40 cm x 20 cm recorded the lowest gross return (₹1,33,569.97/ha), net return (₹99,112.97/ha) and benefit-cost ratio (2.88).

**CONCLUSION:** Sowing of baby corn on 2<sup>nd</sup> March had a significant effect on the growth, yield attributes, cob yield and fodder yield of baby corn. Similarly planting geometry of 45 cm x 20 cm resulted in higher baby corn as well as green fodder yield. The best performance of baby corn during the *summer* season could be achieved by sowing the seed on 2<sup>nd</sup> March with the spacing of 45 cm x 20 cm with higher productivity as well as remunerative economic return in Assam.

#### **REFERENCES**

1. Arvadiya, L.K.; Raj, V.C.; Patel, T.U. and Arvadiya, M.K. . Influence of plant population and weed management on weed flora and productivity of sweet corn. *Indian Journal of Agronomy*. 2012; 57(2): 162-167.
2. Ghosh, M.; Maity, S.K.; Gupta, S.K. and Chowdhury, A.R. (2017). Performance of baby corn under different plant densities and fertility levels in lateritic soils of Eastern India. *Intern. J. Pure App. Biosci.* 2017; 5(3): 696-702.
3. Monneveux, P., Zaidi, P.H. and Sanchez, C. Population density and low nitrogen affects yield. *Associated Traits in Tropical Maize. Crop Science*. 2005; **45**(2): 103-106.
4. Neelam and Dutta, R. Production of baby corn as influenced by spacing and nutrient management. *Int. J. Curr. Microbiol. App. Sci.* 2018;7(12): 1332-1339.
5. Ramachadrappa, B.K.; Nanjappa, H.V.; Thimmegowda, M.N. and Soumya, T.M. Production management of profitable baby corn cultivation. *Indian Farming*. 2004; 39: 1-2.
6. Sobhana, V.; Kumar, A.; Idnani, L.K.; Singh, I. and Shivadhar. Plant population and nutrient requirement for baby corn hybrids (*Zea mays*). *Indian Journal of Agronomy*. 2012; 57(3): 294-296.
7. Tamadon, R.M. (2000). Effects of sowing date and plant density, yield and yield components of sweet corn var. KSC403 su in Mazandaran conditions. M.Sc. Thesis of Agronomy, University of Mazandaran, Iran. 2000; 104.

8. Thavaprakash, N. and Velayudham, K. Influence of crop geometry, intercropping system and INM practices on cob yield and nutrient uptake of baby corn. *Mysore Journal of Agricultural Science*.2009; 43: 686-695.
9. Thakur, D.R. and Sharma, V. Effect of planting geometry on baby corn yield in hybrid and composite cultivars of maize. *Journal of Agricultural Science*.70(4): 246-247.
10. Verma, A.K.; Harika, A.S.; Singh, P.K.; Kaur, K. and Yadav, A. International Conference on Sustainable Agriculture for Food and Livelihood Security. 2012: (701-702), Punjab Agricultural University, Ludhiana.

UNDER PEER REVIEW

**Table 1: Effect of dates of sowing and planting geometry on yield of baby corn**

Treatments	Plant height at harvest(cm)	No. of baby corn per plant	Weight of baby corn without husk (g)	Baby corn yield without husk (q/ha)	Dry yield of baby corn without husk (q/ha)	Green fodder yield (q/ha)	Dry fodder Yield (q/ha)
<b>Dates of sowing</b>							
20 <sup>th</sup> February	168.11	1.45	9.80	12.96	2.15	250.52	92.51
2 <sup>nd</sup> March	185.03	2.34	11.40	18.08	3.05	276.55	102.01
12 <sup>th</sup> March	177.56	2.20	11.16	16.34	2.72	266.45	98.36
22 <sup>nd</sup> March	166.28	1.18	8.80	12.29	2.07	245.84	90.77
S.Ed(±)	3.47	0.29	0.53	1.10	0.13	6.50	2.31
CD(0.05)	8.50	0.71	1.30	2.69	0.31	15.91	5.66
<b>Planting geometry</b>							
40 cm x 20 cm	179.51	1.22	8.75	13.41	2.14	262.95	97.30
40 cm x 25 cm	170.87	1.92	10.99	13.70	2.20	256.72	94.96
45 cm x 20 cm	177.06	1.56	9.86	16.96	2.92	270.47	100.09
45 cm x 25 cm	169.54	2.46	11.60	15.60	2.72	249.21	91.30
S.Ed(±)	2.91	0.20	0.41	1.22	0.25	6.64	2.43
CD (0.05)	6.00	0.42	0.85	2.52	0.51	13.70	5.02

**Table 2: Effect of date of sowing and planting geometry on nitrogen content and nitrogen uptake by baby corn**

Treatments	N content (%)		N uptake (kg/ha)		Total N uptake (kg/ha)
Dates of sowing	Baby corn	Fodder	Baby corn	Fodder	
20 <sup>th</sup> February	1.97	0.63	4.13	58.00	62.13
2 <sup>nd</sup> March	1.79	0.69	5.42	69.64	75.07
12 <sup>th</sup> March	1.74	0.67	4.76	65.10	69.86
22 <sup>nd</sup> March	2.05	0.57	4.06	55.86	59.92
S.Ed(±)	0.22	0.07	0.47	3.88	4.10
CD(0.05)	NS	NS	1.27	8.00	8.45
<b>Planting geometry</b>					
40 cm x 20 cm	1.89	0.58	3.70	57.21	60.91
40 cm x 25 cm	2.07	0.61	4.59	59.21	63.81
45 cm x 20 cm	1.77	0.65	5.14	66.68	71.82
45 cm x 25 cm	1.83	0.71	4.94	65.51	70.44
S.Ed(±)	0.15	0.05	0.29	3.61	3.72
CD(0.05)	NS	NS	0.61	7.46	7.68

**Table 3: Effect of dates of sowing and planting geometry on phosphorus content and phosphorus uptake by baby corn**

Treatments	P content (%)		P uptake (kg/ha)		Total P uptake (kg/ha)
	Baby corn	Fodder	Baby corn	Fodder	
<b>Dates of sowing</b>					
20 <sup>th</sup> February	0.49	0.15	1.04	13.71	14.74
2 <sup>nd</sup> March	0.45	0.16	1.30	15.91	17.20
12 <sup>th</sup> March	0.43	0.15	1.11	15.00	16.11
22 <sup>nd</sup> March	0.52	0.14	1.04	13.48	14.51
S.Ed(±)	0.04	0.01	0.07	0.62	0.83
CD (0.05)	NS	NS	0.14	1.27	1.71
<b>Planting geometry</b>					
40 cm x 20 cm	0.46	0.13	0.92	13.35	14.28
40 cm x 25 cm	0.48	0.14	1.03	13.83	14.86
45 cm x 20 cm	0.46	0.15	1.27	15.48	16.74
45 cm x 25 cm	0.49	0.17	1.28	15.44	16.69
S.Ed(±)	0.03	0.01	0.11	0.87	0.87
CD (0.05)	NS	NS	0.22	1.80	1.79

**Table 4. Effect of dates of sowing and planting geometry on potassium content and potassium uptake by baby corn**

Treatments	K content (%)		K uptake (kg/ha)		Total K uptake (kg/ha)
	Baby corn	Fodder	Baby corn	Fodder	
<b>Dates of sowing</b>					
20 <sup>th</sup> February	0.56	0.84	1.34	77.57	78.91
2 <sup>nd</sup> March	0.54	0.87	1.48	85.20	86.68
12 <sup>th</sup> March	0.55	0.85	1.40	83.43	84.83
22 <sup>nd</sup> March	0.57	0.82	1.30	76.39	77.69
S.Ed(±)	0.01	0.03	0.07	3.45	3.71
CD (0.05)	NS	NS	0.14	7.10	7.64
<b>Planting geometry</b>					
40 cm x 20 cm	0.56	0.79	1.18	76.35	77.53
40 cm x 25 cm	0.57	0.81	1.25	76.87	78.12
45 cm x 20 cm	0.53	0.86	1.57	86.39	87.95
45 cm x 25 cm	0.56	0.92	1.51	82.98	84.50
S.Ed(±)	0.02	0.05	0.12	3.34	3.37
CD (0.05)	NS	NS	0.25	6.89	6.95



**Table 5: Effect of date of sowing and planting geometry on available N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O content of soil after harvest**

Treatments	N (kg/ha)	P <sub>2</sub> O <sub>5</sub> (kg/ha)	K <sub>2</sub> O (kg/ha)
<b>Dates of sowing</b>			
20 <sup>th</sup> February	237.22	23.97	133.71
2 <sup>nd</sup> March	230.21	20.62	131.39
12 <sup>th</sup> March	233.23	22.98	132.59
22 <sup>nd</sup> March	241.62	25.81	135.29
S.Ed(±)	33.63	3.91	28.79
CD (0.05)	NS	NS	NS
<b>Planting geometry</b>			
40 cm x 20 cm	240.24	24.21	135.14
40 cm x 25 cm	237.74	23.33	134.41
45 cm x 20 cm	230.72	22.79	130.71
45 cm x 25 cm	233.58	23.06	132.73
S.Ed(±)	13.42	3.03	30.95
CD (0.05)	NS	NS	NS

**Table 6: Effect of date of sowing and planting geometry on economics of baby corn**

Treatments	Cost of cultivation (₹/ha)	Gross return (₹/ha)	Net return (₹/ha)	B:C ratio
<b>Dates of sowing</b>				
20 <sup>th</sup> February	33496.25	128726.42	95230.17	2.85
2 <sup>nd</sup> March	33496.25	172327.69	138831.44	4.15
12 <sup>th</sup> March	33496.25	157325.23	123828.98	3.71
22 <sup>nd</sup> March	33496.25	122930.60	89434.35	2.67
<b>Planting geometry</b>				
40 cm x 20 cm	34457.00	133569.97	99112.97	2.88
40 cm x 25 cm	33857.00	135248.83	101391.83	2.99
45 cm x 20 cm	33554.00	162763.61	129209.61	3.85
45 cm x 25 cm	32117.00	149727.51	117610.51	3.66