

## Original Research Article

### Competitive behaviour of new cultivars of Indian mustard on weeds in lower Siwaliks of North-West Himalayas

**Guide on the colours used:**

**The red colours:** These are correction which have been made and they are to be retained;

**The words in green colour:** are my comments which are to be responded to and deleted eventually.

**The words in blue colour:** These are words that require recasting for clarity.

#### ABSTRACT

**Aims:** To study the competitive ability of new varieties of Indian mustard with weeds during rabi season at Jammu

**Study design:** Split Plot Design

**Place and Duration of Study:** Oilseed Experimental Area, Research Farm, SKUAST-Jammu, Chatha, Jammu & Kashmir (UT)

**Methodology:** A field experiment was conducted at the Research farm, Chatha, Jammu during the *rabi* season of 2010-11 and 2011-12 to study the effect of various cultivars of Indian Mustard on weed density and dry weight in Indian mustard (*Brassica juncea* L.) under weedy check and weed free conditions. Among the different treatments arranged in split plot design the cultivars of Indian mustard were kept in main plots and two treatments viz. Weed free and weedy check plots were kept in sub-plots.

**Results:** The varieties NRCDR 2 and Navgold were found to be the most competitive cultivars of Indian mustard crop in suppressing the weeds at different crop growth stage at 60 DAS and exhibited lowest weed Index during both the years of experimentation.

**Conclusion:** Based on two years of experimentation, it may be concluded that maximum B:C ratio (give full meaning of B:C ratio here) was observed with cultivar NRCDR 2 (1.51 and 1.03) during both the years of experimentation and was found to be the most competitive cultivar of Indian mustard in suppression of weeds during the early stages of crop growth.

**Key Words:** Weeds, Weed population, Competitive cultivars, Indian mustard

#### 1. INTRODUCTION

Mustard is an important *Rabi* oilseed crop (apart from being important oilseed crop, it has some other useful qualities which can be added) of India and is grown on an area of 6.12 million hectares (approx.) with an average productivity of 1511 kg/ha during the year 2018-19 [1]. India ranks third among the major rapeseed mustard growing countries of the world with

9.98 % of the world's area under Rapeseed mustard cultivation. Among the nine oilseed crops grown in the country, it occupies about 24.70 per cent of area and 48.28 percent of production of the total oilseed production in India. Rapeseed and mustard are the major group of oilseed crops grown in Jammu region on a sizeable acreage in Jammu and Kashmir on an area of more than 55000 hectares with an average productivity of 1149 kg/ha which is far below the national average. Weeds ~~are one of the major biotic stress which~~ cause considerable reduction in the yield of Indian mustard which ranges ~~to an extent of~~ 30-45 per cent ~~and more~~ (if it is more than what is in the range, then include it in that range) and weed management during the critical crop weed competition period result in minimum economic losses [2]. Under irrigated conditions, Indian mustard is infested predominantly with broad leaved as well as grassy weeds. The critical period of crop weed competition in Indian mustard is between 30-45 days after sowing [3]. Different weed management strategies viz. herbicides, cultural and biological etc. are used to manage the weeds below economic threshold [4]. Recently some mustard genotypes have been found to have vigorous growth (what?) do you mean growth characteristics? during the initial crop growth period up to 45 days after sowing thereby suppressing the weeds thus benefiting the crop during initial days of crop-weed competition period. Therefore, growing of recently developed competitive cultivars of Indian mustard for managing weeds holds promise. ~~Therefore.~~ **Consequently**, the present study was undertaken with an objective to study the competitive behaviour of newly released varieties of Indian mustard against weeds under Jammu conditions.

It is good information to know the world first two leading countries in rapeseed mustard growing.

Characteristic	Rapeseed production in million metric tons
Canada	19.49
European Union	16.29
China	14.05
India	8.5

<https://www.google.com/search?q=countries+that+produce+rapeseed+mustard&oq=countries+that+produce+rapeseed+mustard+&aqs=chrome..69i57.21656j0j15&sourceid=chrome&ie=UTF-8>

downloaded 29/03/2022

for now India is number four and not three.

## 2. MATERIALS AND METHODS

A two year experiment to study the effects of different cultivars of Indian Mustard on weeds and its suppression was laid out in RBD (**Write out the full meaning of RBD**) with ~~3~~ **three** replications under AICRP (Rapeseed Mustard) programme ~~at~~ during the *rabi* season of 2010-11 and 2011-12 at the Research Farm, Chatha of the Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu, Jammu and Kashmir, India, which is situated at 32° 40' N latitude and 74° 58' E longitude with an altitude of 332 m above mean sea level. The soil of the experimental field was low in organic carbon (0.37%) and nitrogen (208 kg/ha), medium in available phosphorus (15.3 kg/ha) and potassium (128 kg/ha) and slightly alkaline in pH (7.1). Indian Mustard crop was sown in second fortnight of October in rows 30 cm apart and 10±15 cm (**it is desirable to use a specific spacing so that the outcome of the study can be the product of a particular plant population**) plant to plant distance using 5 kg seeds/ha during both the years of experimentation. Recommended dose of 60:30:15:20 kg/ha of N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O:S was uniformly applied to all the treatments using urea, DAP, MOP and Gypsum as fertilizers. **Full dose of P, K and S besides half dose of N were applied as basal dose at the time of sowing whereas rest of the N was given (recast this statement in blue)** as 2 split doses during both ~~the~~ years of experimentation. Crop was raised as per standard package and practices during both the years of experimentation and was harvested during the last week of March during both the years of experimentation. The experiment consisted of twelve treatments which were laid down in factorial RBD with 3 replications and comprised of two factors namely six Indian mustard (*Brassica juncea* L.) varieties (kranti, RL 1359, Navgold, NRCDR 2, CS-56/NRCDR 601 and RSPR 01.) as Factor A and two levels of weed management as Factor B (weed free and weedy check). In the weed free plots ~~3~~ **three** hand weeding at 25-30 days interval were done to keep the field weed free throughout the crop season. The variety CS-56 ~~which was involved was taken in experiment~~ in the first year of experimentation ~~and~~ was replaced by NRCDR 601 during the second year of experimentation. **You may need to support your action here with reasons for taking such an action.** ~~All~~ The standard package and practices were followed for raising the crop. ~~The Data from the study were was analyzed and presented using standard formulas and statistical tools. All the standard package and practices were followed and data was analysed~~ using standard statistical procedures (**it may improve the work if one is specific on the procedure of statistical analysis used**).

**This section is expected to contain description on how Net Returns (Rs./ ha) and B:C ratio were obtained (calculated).**

### 3. RESULTS AND DISCUSSION

The experimental field was infested mainly with broad leaved weeds (*Trachyspermum* sp., *Anagallis arvensis*, *Euphorbia helioscopia*, *Medicago denticulata*, *Vicia sativa*, *Chenopodium album*, *Rumex maritimus*, *Fumaria parviflora* and *Cirsium arvense*), grassy

The information above is from the source below.

Bio-efficacy of different herbicides on weeds and yield of ...

The experimental field was infested mainly with broad leaved weeds (*Trachyspermum* sp., *Anagallis arvensis*, *Euphorbia helioscopia*, *Medicago denticulata*, *Vicia sativa*, *Chenopodium album*, *Rumex maritimus*, *Fumaria parviflora* and *Cirsium arvense*), grassy weeds (*Poa annua*, *Phalaris minor* and *Avena ludoviciana*) and sedges (*Cyperus rotundus*) during both the years of ...

<https://www.thepharmajournal.com/archives/2022/vol11issue2/PartK/11-1-389-195.pdf>

Even if you have similar outcome, please state the outcome in your own way, in order to minimize similarity with other research reports.

weeds (*Poa annua*, *Phalaris minor* and *Avena ludoviciana* (this second name must be in small letters all through) and sedges (*Cyperus rotundus*) during both the years of experimentation. ~~Weedy check caused an average reduction in seed yield of Indian mustard (27.84%) of 27.84% which was significantly lower than weed free situation in comparison.~~ Uncontrolled weed growth in the weedy check caused a significant yield loss of 27.84% relative to weed free. (It would have been interesting to have the % yield loss caused by unchecked weed growth in the two seasons before arising at the average). Maximum weed population and dry weed biomass was recorded in weedy check (control) plots (it will add flesh to the work if some of the figures of the weed biomass are given), whereas no weed population and weed dry weight was recorded in weed free plots. Since this Section was subtitled: **RESULTS AND DISCUSSION**, let us see the author(s) reflect it by doing some discussions of the results by relating the outcome with what has been reported in the past.

Among the different Indian mustard cultivars NRCDR 2 (1398 kg/ha) though at par with Nov gold (1316 kg/ha) resulted in significant increase in the seed yield and other yield attributes viz. Siliquae per plant, Seeds per Siliquae, 1000- seed weight and—of Indian mustard than other cultivars in comparison during the ~~second?~~ first year of experiment. However, during the second year of experimentation, Indian mustard cultivar NRCDR 601(1298 kg/ha) proved to be equally competitive along with cultivars NRCDR 2 (1368 kg/ha) and Navgold (1274 kg/ha) in increasing the seed yield of Indian mustard than other cultivars in comparison (Table 2). Also, lowest weed density and weed dry weight was recorded in plots planted with Indian Mustard cultivars viz. NRCDR 2 and Navgold

during both ~~the~~ years of experimentation. However, variety NRCDR 601 proved to be equally effective in suppressing the grassy and broad leaved weeds during the second year of experimentation (Table 1). This suppression of weeds during the initial period of crop growth may have been resulted due to better utilization of nutrients, moisture, space and light thereby resulting in higher LAI of the said genotypes thereby increasing their ability over other genotypes in comparison [5,6]. The lowest weed Index was recorded with cultivar NRCDR 2 (4.77 and 7.88 % respectively) during both ~~the~~ years of experimentation. Among the sub plots lowest weed population was recorded in weed free plots in comparison to weedy check plots.

## CONCLUSION

Based on average of two years of experimentation in 2010-11 and 2011-12 on competitive behavior of Indian mustard (*Brassica juncea* L.) varieties against weeds vis-à-vis weed smothering potential of various newly developed genotypes at SKUAST-J, Chatha centre it was concluded that new high yielding variety NRCDR 2 gave highest seed yield of mustard and was found to be most competitive cultivars of Indian mustard for suppression of weeds in Indian mustard crop in the Jammu region.

## REFERENCES

1. Anonymous. Agriculture Statistics at a glance, Directorate of Economics & Statistics. Department of Agriculture and Cooperation & Family Welfare, GoI, 4<sup>th</sup> Advance estimates, Ministry of Agriculture, Govt. of India, Krishi Bhawan, New Delhi, 2020. Pp. 72-109
2. Bharat R, Gupta M, Bazaya BR and Gupta R Bio-efficacy of different herbicides on weeds and yield of Indian mustard (*Brassica juncea* L.) in Jammu, The Pharma Innovation Journal. 2022; **11** (2) : 725-727
3. Bharat, R and Kachroo, D. Bio-efficacy of various herbicides and their mixtures on weeds and yield of wheat (*Triticum aestivum*) under subtropical agro ecosystem, *Indian Journal of Agronomy*. 2007; **52** (1): 53-59.
4. Chauhan YS, Bhargava MK, Jain VK Weed management in Indian mustard. *Indian Journal of Agronomy*. 2005; **50** (2): Pp: 149-151
5. Raj P, Singh RP, Pal R, Rajput P, Rana SS Integrated weed management in Indian mustard (*Brassica juncea* L.). *Int. J. Curr. Microbiol. App. Sci.*, 2020; Spl. Issue (10): 271-276
6. Singh RK, Singh RP, Singh MK Weed management in Indian mustard- A review. *Agricultural Reviews*. 2013; **34** (1): Pp. 36-49

UNDER PEER REVIEW

**Table 1. Effect of different cultivars on weed count and weed dry weight in Indian mustard at 60 days after sowing of Indian mustard**

Treatments	Weed Population (no./m <sup>2</sup> )						Weed dry weight (g/m <sup>2</sup> )					
	2010-11			2011-12			2010-11			2011-12		
	Broad leaved	Grasses	Total	BLW	Grasses	Total	BLW	Grasses	Total	BLW	Grasses	Total
<b>Varieties</b>												
Kranti	33.64 ( 5.80)	9.67 ( 3.11)	43.31 ( 6.58)	42.33 (6.51)	38.00 (6.16)	80.33 (8.96)	23.61 ( 4.86)	7.40 ( 2.72)	31.01 (5.57)	39.77 (6.31)	23.98 (4.90)	63.75 (7.98)
RL 1359	35.88 ( 5.99)	13.69 ( 3.70)	49.57 (7.05)	42.67 (6.53)	40.00 (6.32)	82.67 (9.09)	24.90 ( 4.99)	8.29 ( 2.88)	33.19 (5.76)	40.19 (6.34)	24.26 (4.93)	64.45 (8.03)
NRCDR 2	29.60 (5.44)	9.12 ( 3.02)	38.72 (6.22)	36.67 (6.06)	33.33 (5.77)	70.0 (8.37)	22.84 ( 4.78)	6.96 ( 2.64)	29.8 (5.46)	36.48 (6.04)	21.67 (4.66)	58.15 (7.63)
Novgold	33.99 ( 5.83)	11.56 ( 3.40)	45.55 (6.75)	39.33 (6.27)	36.67 (6.06)	76.0 (8.72)	23.42 (4.84)	7.95 ( 2.82)	31.37 (5.60)	39.85 (5.69)	22.55 (4.75)	62.40 (10.08)
NRCDR 601	32.72 ( 5.72)	12.74 ( 3.57)	45.46 (6.64)	40.33 (6.35)	35.33 (5.94)	75.66 (8.69)	24.20 ( 4.92)	8.53 ( 2.92)	32.73 (5.72)	37.50 (6.31)	22.09 (4.70)	59.59 (7.89)
RSPR 01	39.56 ( 6.29)	14.51 ( 3.81)	54.07 (7.35)	43.33 (6.58)	41.33 (6.43)	84.66 (9.20)	24.41 ( 4.94)	8.64 ( 2.94)	33.05 (5.75)	40.92 (6.40)	24.87 (4.99)	65.79 (8.11)
CD (p=0.05)	NS	NS		NS	NS		NS	NS		NS	NS	
<b>Weed Management</b>												
Weedy check	85.56 ( 9.25)	23.23 ( 4.82)	108.80 (10.43)	72.95 (8.54)	63.81 (7.99)	136.76 (11.69)	64.96 (8.06)	17.55 ( 4.19)	82.51 (9.08)	71.04 (8.43)	42.27 (6.50)	113.31 (10.64)
Weed free	5.80 ( 2.41)	4.20 ( 2.05)	10.00 (3.16)	8.95 (2.99)	11.81 (3.44)	20.76 (4.56)	2.92 (1.71)	2.13 (1.46)	5.05 (2.25)	7.33 (2.71)	4.65 (2.16)	11.98 (3.46)
CD (p=0.05)	0.32	0.35	0.68	0.29	0.36	0.65	0.12	0.06	0.09	0.20	0.23	0.22

**BLW =? (Write out the full meaning of BLW here).**

**Table 2. Effects of different cultivars on yield, yield attributes, net returns and economics of Indian Mustard**

**Close the gap above with green line**

Treatments	Seed yield (q/ha)		Siliquae/plant		Seed/Siliquae		1000-seed weight		Net Returns (Rs./ ha)		B:C ratio	
	2010-11	2011-12	2010-11	2011-12	2010-11	2011-12	2010-11	2011-12	2010-11	2011-12	2010-11	2011-12
<b>Varieties</b>												
Kranti	12.84	12.52	397.16	424.83	15.16	12.83	3.70	3.59	13283	10741	1.26	0.85
RL 1359	12.66	12.09	376.13	409.17	15.10	12.53	3.63	3.55	12954	9936	1.23	0.78
NRCDR 2	13.98	13.68	414.53	443.33	15.90	14.93	3.8	3.79	15912	12893	1.51	1.03
Novgold	13.18	12.74	412	433.67	15.36	14.33	3.74	3.68	13907	11148	1.32	0.88
CS 56/NRCDR 601	12.97	12.97	408.06	439.33	16.20	14.67	3.82	3.77	13500	11579	1.28	0.91
RSPR 01	11.99	11.74	365.43	397.33	14.26	12.07	3.46	3.38	11705	9289	1.11	0.73
CD (p=0.05)	0.82	1.09	26.64	11.78	NS	0.83	0.16	0.13	-	-	-	-
<b>Weed Management</b>												
Weedy check	11.17	10.18	346.78	389.24	14.70	12.00	3.39	3.42	11436	7668	1.20	0.69
Weed free	14.68	14.85	444.83	453.62	15.96	14.82	3.76	3.79	15652	13973	1.36	1.01
<b>CD (p=0.05)</b>	NS	0.58	15.38	6.30	1.0	0.45	0.09	0.07	-	-	-	-

**BLW =? (Write out the full meaning of BLW here).**