

# Original Research Article

## Interaction effect of Organic Manures and Alkali Water on Yield of Wheat

### Abstract

Field experiment were carried out to investigate "Interactive effect of organic manures and RSC water on yield of wheat" in *rabi* 2019-20 and 2020-21. Result found that the irrigation with 10 mmol L<sup>-1</sup> RSC water significantly decreased grain (26.02 q ha<sup>-1</sup>) and straw yield (38.15 q ha<sup>-1</sup>) of wheat in comparison to lower levels of RSC in water in wheat crop. Application of organic manures significantly increased grain yield of wheat during both the years and in pooled analysis. The highest grain (34.06 q ha<sup>-1</sup>) and straw yield (51.60 q ha<sup>-1</sup>) were obtained under vermicompost application and the effect of vermicompost (M<sub>2</sub>) and FYM (M<sub>1</sub>) application remained statistically at par with each other. Interactive effect of RSC water and organic manures on grain and straw yield was found significant during both the years and in pooled.

**Keywords:-** FYM, RSC water, Vermicompost, Wheat

### Introduction

The usage of poor-quality water for irrigation of agricultural crops would inevitably aggravate India sodicity problem. Soil sodicity influence physiological, biochemical, and morphological changes in plants, which represent the plant's overall performance. However, if inadequate minerals are made available to specific plant species, such as wheat, the most important *rabi* crop of salty regions, they may grow and yield well. Most saline sodic soils are generally poor in Ca, N, Zn, Mn, Cu, Fe and organic matter. (Singh *et al.*, 2013, Kumawat and Yadav 2013). Organic matter importance has been proven by its ability to improve the physical conditions of soils for crop development in addition to its role as fertilizer. The efficacy of various organic additions such as manure and compost in the restoration of salty sodic soils has been studied (Diez & Krauss 1997). Organic additions to soil reduce bulk density while increasing CEC and water holding capacity, resulting in a diluting impact on alkali soil. Organic matter works to prevent salt from building up in soils (Gupta *et al.*, 1984). The microbial processes of nutrient immobilization and mineralization are strongly linked to organic matter dynamics and nutrient cycle.

### Material and methods

The experiments were conducted at Agronomy Farm, S.K.N. College of Agriculture, Jobner (Rajasthan), Geographically Jobner is situated at 26°05' North latitude and 75°28' East longitude at an altitude of 427 meter above mean sea level (MSL). Maximum temperature in summer ranges between 28 to 44°C, whereas temperature falls down to as low as 3°C during winter. The average annual rainfall varies between 400 to 500 mm. The experiment comprising of 36 treatment combinations were replicated three time was laid out in split -plot design with three levels of alkali water (2, 6, 10 RSC mmol L<sup>-1</sup>), three organic manures (Control, FYM @ 15 t ha<sup>-1</sup> and vermicompost @ 5 t ha<sup>-1</sup>) in main plot and four levels of iron (Control, 50 kg, 100 kg and 150 kg FeSO<sub>4</sub>.7H<sub>2</sub>O ha<sup>-1</sup>) in sub plot were tested. The experiment soil is low in organic carbon, available nitrogen, iron and medium in available phosphorous, potassium content in soil. Organic manures and iron were applied prior to sowing through FeSO<sub>4</sub>.7H<sub>2</sub>O respectively as per treatments.

The experiment crop was irrigated by different RSC waters. The different levels of RSC in irrigation water were prepared by dissolving required amount of NaCl, Na<sub>2</sub>SO<sub>4</sub>, NaHCO<sub>3</sub>, CaCl<sub>2</sub> and MgCl<sub>2</sub> in base water (control) (Table 1). The normal tube well water (base water) was used for first irrigation in all plots at CRI (Crown Root Initiation) stage and later on crop was irrigated five times with varying alkali water during whole growth period as per plan of experiment. To check the lateral movement of water and salts buffer strips around each irrigation channel were kept.

The composition of the prepared waters is given in Table 1.

**Comment [I1]:** State the type of experimental design used in the study and the respective treatments before the result.

**Comment [I2]:** State What you are recommending to the farmers after the pooled.

**Comment [I3]:** Give the full meaning

**Comment [I4]:** Write on Alkali water and its impact on yield of wheat

**Comment [I5]:** Are there no recent literature?

**Comment [I6]:** Old

**Comment [I7]:** at

**Comment [I8]:** at

**Comment [I9]:** check spelling

**Comment [I10]:** case sensitive

**Comment [I11]:** Give full meaning

**Table 1 Composition of synthetic irrigation water**

RSC (mmol L <sup>-1</sup> )	EC (dSm <sup>-1</sup> )	SAR	Ionic composition (mmol L <sup>-1</sup> )						
			Na <sup>+</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	HCO <sub>3</sub> <sup>-</sup>	CO <sub>3</sub> <sup>2-</sup>	Cl <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>
2.0 (Base water)	1.3	8.2	10	1.5	1.5	4.5	0.5	6.5	1.5
6.0	3.2	16.7	26.4	2.5	2.5	10.0	1.0	10.5	10.5
10.0	3.2	16.7	26.4	2.5	2.5	12.0	3.0	8.5	8.5

## Result and Discussion

### Yield

The results presented in Table 2 and Figure 1 for grain and straw yield of wheat decreased significantly with increase in level of RSC in irrigation water during both the years of study and also when data were pooled. Decline in crop yields due to high RSC water irrigations could be ascribed to the deteriorating effect of high pH and RSC on chemical, physical and biological soil fertility parameters along with Ca deficit in the soil solution (Choudhary *et al.* 2019). This may be explained on the basis that increasing RSC in irrigation water increased the ESP and pHs of soil resulting into decreased availability of P, K, Ca and Mg but increased the uptake of Na which is toxic to plant. The higher amount of Na may adversely affect the physiological, metabolic and enzymatic activities and utilization of photosynthates in plant. The cell elongation and cell division may also be adversely affected due to higher accumulation of Na. This causes a reduction in number of tillers and plant height. In this study, also the number of tillers and plant height decreased drastically with increasing level of RSC in irrigation water. This may be a reason for decrease in grain and straw yield of wheat. Further, the reduction in grain and straw yield may also be due to the poor physical condition of soil caused by accumulation of higher amount of Na in soil. Under high RSC water increased exchangeable Na has detrimental effect on physical condition and nutritional properties, resulting in poor root development and plant growth. This resulted in decrease in yield of wheat. Choudhary *et al.* (2019) and Sheoran, *et al.* (2021) also expressed similar views and observed reduction in the grain and straw yield with an increase in levels of RSC in water.

The significantly maximum increased in grain and straw yield of wheat was observed with the application of vermicompost @ 5 t ha<sup>-1</sup> (34.62 %) and FYM @ 15 t ha<sup>-1</sup> (30.71%) over control during both the years and during pooled analysis (Table 2) but the effect of vermicompost and FYM was statistically at par with each other. The improvement in yield might be due to the result of overall improvement in soil environment due to decrease in pHs and ESP under the influence of applied organic manures. These beneficial effects favoured greater availability of plant nutrients and their steady supply throughout growth for optimum development as explained earlier in preceding paragraphs and observed in the present study. The higher nutrients availability and congenial environment for their uptake, favored greater synthesis of carbohydrates and their efficient partitioning into different sinks including reproductive structures which ultimately brought significant improvement in effective tillers and grain yield. The straw yield increased due to improvement in vegetative growth of wheat. These results are in conformity with those of Bhakher *et al.*, (1997) and (Kamaleshwaran and Elayaraja, 2021) who have reported increased in yield attributes and yield of rice and wheat due to application of vermicompost and farmyard manure might have supplied the essential minerals and worked as catalyst for efficient use of applied nutrients for increasing the yield attribute. Similar results were also reported by Sharma *et al.* (2009), Abbas *et al.* (2009), Thakur *et al.* (2011), Prativa and Bhattarai (2011) and Anand *et al.* (2014), Vishram *et al.* (2014) and Choudhary *et al.* (2020) also studied on response of FYM on seed yield and quality of crop under high RSC water.

The grain and straw yield also increased significantly with increase in levels of Fe during both the years and in pooled analysis (Table 2). The favourable influenced of applied Fe on these

Comment [I12]: at

Comment [I13]: at

characters might be due to its catalytic effect on most on the physiological and metabolic process of plant. Fe is involved in the formation of chlorophyll and plays a somewhat similar role to Mg in the porphyrin structure of chlorophyll. Fe is also a constitute of large no. of metabolically active compound like cytochromes, haeme and non-haeme enzymes and other functional metallo proteins such as ferredoxin and haemoglobin. Fe plays an important role in Zn, Cu, Mg and Mn metabolism. It is widely agreed that Fe is directly involved in protein synthesis. The application of Fe in a soil deficient in its status, improved overall growth and development of plants and ultimately the effective tillers, plant height, grain and straw yield increased under irrigation with high RSC waters. The findings of present investigation are supported by Radder and Husen (2017) and Janmohammadi *et al.* (2018).

UNDER PEER REVIEW


**Table 2 Effect of alkali water, organic manures and iron application on grain and straw yield (q ha<sup>-1</sup>) of wheat**

Treatments	Grain yield (q ha <sup>-1</sup> )			Straw yield (q ha <sup>-1</sup> )		
	2019-20	2020-21	Pooled	2019-20	2020-21	Pooled
<b>Alkali water (RSC mmol L<sup>-1</sup>)</b>						
W <sub>2</sub> (2 Base water)	35.56	34.10	34.83	56.32	54.32	55.32
W <sub>6</sub> (6)	32.13	31.03	31.58	48.33	46.54	47.43
W <sub>10</sub> (10)	26.72	25.31	26.02	38.65	37.66	38.15
SE <sub>m</sub> ±	0.55	0.47	0.36	0.84	0.63	0.53
CD (P=0.05)	1.66	1.41	1.05	2.52	1.90	1.52
<b>Organic manures</b>						
M <sub>0</sub> (Control)	26.13	24.47	25.30	39.36	37.92	38.64
M <sub>1</sub> (FYM @ 15 t ha <sup>-1</sup> )	33.69	32.44	33.07	51.67	49.65	50.66
M <sub>2</sub> (Vermicompost @ 5 t ha <sup>-1</sup> )	34.58	33.53	34.06	52.26	50.94	51.60
SE <sub>m</sub> ±	0.55	0.47	0.36	0.84	0.63	0.53
CD (P=0.05)	1.66	1.41	1.05	2.52	1.90	1.52
<b>Iron levels (FeSO<sub>4</sub> .7H<sub>2</sub>O)</b>						
F <sub>0</sub> (Control)	27.10	25.21	26.16	39.61	37.05	38.33
F <sub>50</sub> (50 kg ha <sup>-1</sup> )	31.22	30.46	30.84	47.11	45.85	46.48
F <sub>100</sub> (100 kg ha <sup>-1</sup> )	34.37	33.29	33.83	52.58	51.57	52.07
F <sub>150</sub> (150 kg ha <sup>-1</sup> )	33.17	31.63	32.40	51.76	50.23	50.99
SE <sub>m</sub> ±	0.72	0.71	0.51	0.80	0.82	0.57
CD (P=0.05)	2.05	2.02	1.42	2.28	2.32	1.61

Comment [I14]: capitalize

Comment [I15]: capitalize

Comment [I16]: capitalize



**Fig. 1 Effect of alkali water, organic manures and iron on grain and straw yields of wheat**

### Interactive effect of RSC water and organic manures on grain and straw yield

The interactive effect of RSC water and organic manures on grain and straw yield was found significant during both the years and in pooled analysis (Table 3 & 4 and Fig. 2 & 7) and data reveal that grain and straw yield increased significantly with organic manures treatment under all the levels of RSC in water. Grain and straw yield obtained with RSC water ( $W_2$ ) and  $5 \text{ t ha}^{-1}$  vermicompost ( $M_2$ ) was significantly superior to the yield obtained over control and statically at par with  $15 \text{ t ha}^{-1}$  FYM along with other level of RSC water  $W_6$  and  $W_{10}$ . The maximum grain yield was recorded under  $W_2$  ( $34.83 \text{ q ha}^{-1}$ ) and minimum under  $W_{10}$  ( $26.02 \text{ q ha}^{-1}$ ). This indicates that harmful effect of high RSC water can be mitigated by applying organic manures resulting in overall improvement in soil physical environment due to decrease in pH, E<sub>c</sub>, ESP and increase CEC and reducing the accumulation of Na in soil resulting in higher grain and straw yield under  $W_2 M_2$  over rest of the treatment combinations. The results are in close conformity with Pareek and Yadav (2011) and Rathi *et al.* (2020).

**Table 3 Interactive effect of application of alkali water and organic manures on grain yield (q ha<sup>-1</sup>) of wheat**

Treatments	2019-20			Mean
	M <sub>0</sub> (Control)	M <sub>1</sub> (FYM @ 15 t ha <sup>-1</sup> )	M <sub>2</sub> (Vermicompost @ 5 t ha <sup>-1</sup> )	
W <sub>2</sub> (2 Base water)	31.31	37.29	38.07	35.56
W <sub>6</sub> (6)	27.91	34.13	34.36	32.13
W <sub>10</sub> (10)	19.19	29.66	31.31	26.72
Mean	26.13	33.69	34.58	
SE <sub>m±</sub>	0.96			
CD (P=0.05)	2.88			
Treatments	2020-21			Mean
	M <sub>0</sub> (Control)	M <sub>1</sub> (FYM @ 15 t ha <sup>-1</sup> )	M <sub>2</sub> (Vermicompost @ 5 t ha <sup>-1</sup> )	
W <sub>2</sub> (2 Base water)	29.64	36.05	36.62	34.10
W <sub>6</sub> (6)	26.24	33.03	33.81	31.03
W <sub>10</sub> (10)	17.52	28.25	30.17	25.31
Mean	24.47	32.44	33.53	
SE <sub>m±</sub>	0.82			
CD (P=0.05)	2.44			
Treatments	Pooled			Mean
	M <sub>0</sub> (Control)	M <sub>1</sub> (FYM @ 15 t ha <sup>-1</sup> )	M <sub>2</sub> (Vermicompost @ 5 t ha <sup>-1</sup> )	
W <sub>2</sub> (2 Base water)	30.47	36.67	37.35	33.909
W <sub>6</sub> (6)	27.07	33.58	34.08	30.577
W <sub>10</sub> (10)	18.36	28.95	30.74	24.549
Mean	24.41	32.81	34.04	
SE <sub>m±</sub>	0.63			
CD (P=0.05)	1.82			

Comment [I17]: capitalize

Comment [I18]: capitalize

Comment [I19]: capitalize

**Table 4** Interactive effect of application of alkali water and organic manures on straw yield (q ha<sup>-1</sup>) of wheat

Treatments	2019-20			
	M <sub>0</sub> (Control)	M <sub>1</sub> (FYM @ 15 t ha <sup>-1</sup> )	M <sub>2</sub> (Vermicompost @ 5 t ha <sup>-1</sup> )	Mean
W <sub>2</sub> (2 Base water)	50.61	58.93	59.41	56.32
W <sub>6</sub> (6)	36.59	54.14	54.26	48.33
W <sub>10</sub> (10)	30.89	41.95	43.10	38.65
Mean	39.36	51.67	52.26	
SEm±	1.46			
CD (P=0.05)	4.37			
Treatments	2020-21			
	M <sub>0</sub> (Control)	M <sub>1</sub> (FYM @ 15 t ha <sup>-1</sup> )	M <sub>2</sub> (Vermicompost @ 5 t ha <sup>-1</sup> )	Mean
W <sub>2</sub> (2 Base water)	48.87	56.53	57.55	54.32
W <sub>6</sub> (6)	34.86	52.22	52.55	46.54
W <sub>10</sub> (10)	30.04	40.21	42.73	37.66
Mean	37.92	49.65	50.94	
SEm±	1.10			
CD (P=0.05)	3.29			
Treatments	Pooled			
	M <sub>0</sub> (Control)	M <sub>1</sub> (FYM @ 15 t ha <sup>-1</sup> )	M <sub>2</sub> (Vermicompost @ 5 t ha <sup>-1</sup> )	Mean
W <sub>2</sub> (2 Base water)	49.74	57.73	58.48	54.11
W <sub>6</sub> (6)	35.72	53.18	53.40	44.56
W <sub>10</sub> (10)	30.47	41.08	42.91	36.69
Mean	40.10	49.41	50.70	
SEm±	0.91			
CD (P=0.05)	2.64			

Comment [I20]: capitalize

Comment [I21]: capitalize

Comment [I22]: capitalize






Fig. 2 Interactive effect of application of alkali water and organic manures on grain yield of wheat (*Rabi* 2019-20)




Fig. 3 Interactive effect of application of alkali water and organic manures on grain yield of wheat (*Rabi* 2020-21)




Fig. 4 Interactive effect of application of alkali water and organic manures on grain yield of wheat (Pooled)




Fig. 5 Interactive effect of application of alkali water and organic manures on straw yield of wheat (*Rabi* 2019-20)




Fig. 6 Interactive effect of application of alkali water and organic manures on straw yield of wheat (*Rabi* 2020-21)




Fig. 7 Interactive effect of application of alkali water and organic manures on straw yield of wheat (Pooled)

## Conclusion

Our results showed that effect of RSC water and organic manure application on grain and straw yields of wheat was found significant and maximum being with 2 mmol L<sup>-1</sup> RSC water and vermicompost @ 5 t ha<sup>-1</sup> treatment combination indicating the magnitude of reduction in yields (grain and straw) being less with vermicompost and FYM at all the levels of RSC water.

Comment [I23]: at

## References

- Abbas, G., Khan, M.Q., Khan, M.J., Hussain, F. and Hussain, I. 2009. Effect of iron on the growth and yield contributing parameters of wheat (*Triticum aestivum* L.). *Journal of Animal Plant Science*, **19**,135-139.
- Anand, S.R., Vishwanatha, J., Ravishankar, G., Karegoudar, A.V. and Rajkumar, R.F. 2014. Performance of chemical and organic amendments on cotton in alkali Vertisols of northern Karnataka. *Environment and Ecology*, **32**, 1657-1660.
- Bhakher, J.R., Sharma, O.P. and Jat, B.C. 1997. Effect of nitrogen and FYM on yield and yield attributes of barley in a loamy sand soil. *Annals of Agricultural Research*, **18**, 244-245.
- Choudhary, D.R., Singh, A., Panghal, V.P.S. and Bhuker, A. 2020. Response of FYM and gypsum on seed yield and quality of radish under high RSC water. *International Journal of Chemical Studies*, **8**,1544-1548.
- Choudhary, O.P., Bhalla, M., Sharma, S., Sharda, R. and Mavi, M.S. 2019. Long-term impact of cyclic use of sodic and canal water for irrigation on soil quality and wheat yield in cotton-wheat cropping system. *Journal of the Indian Society of Soil Science*, **67**, 34-43.
- Choudhary, O.P., Ghuman, B.S., Thuy, N. and Buresh, R.J. 2011. Effects of long-term use of sodic water irrigation, amendments and crop residues on soil properties and crop yields in rice-wheat cropping system in a calcareous soil. *Field Crops Research*, **121**, 363-372.
- Diez, T. and M. Krauss. 1997. Effect of long-term compost application on yield and soil fertility. *Agrobiological Research-Zeitschrift Fur Agrarbiologie-Agricuturechemie Okologie*, 50: 78-84. Germany
- Gupta, R.K., Bhumbra, D.R. and Abrol, I.P. 1984. Effect of soil pH, organic matter and calcium carbonate on dispersion behavior of alkali soils. *Soil Science*, 137: 145-251.
- Janmohammadi, M., Abdoli, H., Sabaghnia, N., Esmailpour, M. and Aghaei, A. 2018. The effect of iron, zinc and organic fertilizer on yield of chickpea (*Cicer artietinum* L.) in Mediterranean climate. *Acta Universitatis Agriculturae Et Silviculturae Mendelianae Brunensis*, **66**, 49-60.
- Kamaleshwaran, R. and Elayaraja, D. 2021. Influence of vermicompost and FYM on soil fertility, rice productivity and its nutrient uptake. *International Journal of Agriculture and Environmental Research*, **7**, 575-583.
- Kumawat, S.R. and Yadav, B.L. 2013. Sodicity tolerance of fenugreek (*Trigonella foenum-graecum* L.) as influenced by application of zinc and vermicompost. *Legume Research*, **36**, 312-317.
- Pareek, N. and Yadav, B.L. 2011. Effect of organic manure on soil physico-chemical properties, soil microbial biomass and yield of mustard under irrigation of different residual sodium carbonate water. *Journal of the Indian Society of Soil Science*, **59**, 336-342.
- Prativa, K.C. and Bhattarai, B.P. 2011. Effect of integrated nutrient management on the growth, yield and soil nutrient status in tomato. *Nepal Journal of Science and Technology*, **12**, 23-28.
- Radder, B.M. and Husen, S.P. 2017. Effect of iron sulfate application on yield, nutrient uptake and available nutrient status of soybean (*Glycine max* L.) at harvest in vertisols of Karnataka, India. *Environment and Ecology*, **35**, 1336-1340.
- Rathi, D., Antil, R.S., Sharma, M.K. and Sheoran, S. 2020. Effect of FYM and gypsum on distribution of micronutrient in soil under sodic water irrigation, A long-term study. *Journal of the Indian Society of Soil Science*, **68**,100-106.
- Sharma, O.P., Datt, N. and Chander, G. 2009. Effect of vermicompost, farmyard manure and chemical fertilizers on yield, nutrient uptake and soil fertility in okra (*Abelmoschus*

*esculentus*) - onion (*Allium cepa*) sequence in Wet Temperate Zone of Himachal Pradesh. *Journal of the Indian Society of Soil Science*, **3**, 357-361.

Sheoran, P., Basak, N., Kumar, A., Yadav, R.K., Singh, R., Sharma, R., Kumar, S., Singh, R.K. and Sharma, P.C. 2021. Ameliorants and salt tolerant varieties improve rice-wheat production in soils undergoing sodification with alkali water irrigation in Indo-Gangetic Plains of India. *Agricultural Water Management*, **243**, 106492.

Singh, K., Singh, B. and Singh, R.R. 2013. Effect of land rehabilitation on physicochemical and microbial properties of a sodic soil. *Catena*, **109**, 49-57.

Thakur, R., Sawarkar, S.D., Vaishya, U.K. and Singh, M. 2011. Impact of continuous use of inorganic fertilizers and organic manure on soil properties and productivity under soybean-wheat intensive cropping of a vertisol. *Journal of the Indian Society of Soil Science*, **59**, 74-81.

Vishram, S., Sanjai, C., Verma, V.K. and Srivastava, A.K., 2014. Studies on integrated nutrient management in mustard [*Brassica juncea* (L.)]. *International Journal of Agricultural Sciences*, **10**, 667-670.