

Influence of Sulphur and Zinc levels on growth and yield of Cowpea

(*Vigna unguiculata* L.)

ABSTRACT: A field experiment entitled “Influence of Sulphur and Zinc levels on growth and yield of Cowpea (*Vigna unguiculata* L.)”[†] was carried out during *kharif* season of 2021 at experimental field of the Crop Research Farm, **Department of Agronomy**, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj, and Uttar Pradesh, India. The soil of experimental field is Sandy loam in texture, nearly neutral in soil reaction (pH 7.0). To determine the treatments which are T₁ 20 kg/ha S + 10 kg/ha Zn, T₂ 20 kg/ha S + 15 kg/ha Zn, T₃ 20 kg/ha S + 20 kg/ha Zn, T₄ 30 kg/ha S + 10 kg/ha Zn, T₅ 30 kg/ha S + 15 kg/ha Zn, T₆ 30 kg/ha S + 20 kg/ha Zn, T₇ 45 kg/ha S + 10 kg/ha Zn, T₈ 45 kg/ha S + 15 kg/ha Zn, T₉ 45 kg/ha S + 20 kg/ha Zn, T₁₀. Control was used. The Experiment was laid out in Randomized Block Design with Ten treatments which was replicated thrice.

Background: Cowpea is known as drought hardy nature, its wide and droopy leaves keeps soils and soil moisture conserved due to shading effect. It is also known as black-eyed pea or southern pea etc. and has multiple uses like food, feed, forage, fodder, green manuring and vegetable.

Objectives: To study the retaliation performances of Sulphur and zinc on growth, yield and yield attributes of cowpea (*Vigna unguiculata* L.)

Conclusion: It was determined that the application of Sulphur – 45 kg/ha + Zinc – 20 kg/ha resulted in a higher grain yield (13.67 q/ha), which was significantly superior to the other treatments.

Key words: cowpea, growth, sulphur, yield and zinc.

Comment [h1]: Remove

Comment [h2]: remove

Comment [h3]: 10 treatments were used for conducting the experiments namely

Comment [h4]: Insert = or : after T1, T2, T10

Comment [h5]: remove

Comment [h6]: remove

Comment [h7]: 45 kg/ha sulphur + 20 kg/ha zinc

INTRODUCTION

Cowpea (*Vigna unguiculata* L.) is one of the most important vegetable crops grown as pulse, vegetable and fodder. It is poor man's protein source and considered one of the most ancient human food sources and has probably been used as a crop plant since Neolithic times

(Ng and Marechal 1985) (7). Cowpea is a vital multipurpose grain legume extensively cultivated in arid and semiarid tropics. It is an important source of nutrients and provides high quality, inexpensive protein diet based on cereal grains and starch foods. Cowpea is a good source of food, fodder, vegetables and certain snakes (Singh *et al.*, 2012)(11). In India pulses are grown nearly in 25.43 m ha with an annual production of 17.28 m t and a median productivity of 679 kg/ha (Anonymous *et al.*, 2012)(2). In Rajasthan the realm under pulses is 47.54 lakh ha with an annual production of 32.54 lakh m t and an average productivity of 684 kg/ha and also the area under Cowpea is 1.02 lakh ha with the production of 0.64 lakh tonnes and productivity of 529 kg/ha. The per capita availability of pulses in India is 35.5 g/day as against the minimum requirement of 70 g/day/capita.

Comment [h8]: Add recent area and production of pulses

Comment [h9]: Add recent data

Sulphur is the fourth major nutrient next to N, P and K and an essential element for plant growth particularly for legumes crops which play an important role in plant metabolism system. Sulphur (S) is a constituent of some important amino acids namely cystine, cysteine and methionine. Sulphur is essential for the growth and development, plays a key role in plant metabolism. S containing amino acids (cystine, cysteine and methionine) and promotes nodulation in legumes. It helps in chlorophyll formation and encourages vegetative plant growth (Karche and co-workers 2012) (4).

Comment [h10]: remove

Comment [h11]: provide space instead of full stop

Zinc, which is important for growth and reproduction in plants, animals and humans, is one of the seventh essential micronutrients. In plants, it plays a key role during physiological growth, DNA stabilization, gene expression, enzyme activity, protein synthesis and improved chlorophyll function (Prasad, A.S. 2003)(8). Zinc deficiency is a major limiting factor in several Asian countries. It is now being recorded as third most deficient nutrient in crop production after nitrogen and phosphorus. In India, zinc deficient soils occupy almost 50% of the agricultural area and are a critical constraint on yield.

Comment [h12]: most important essential

Comment [h13]: By keeping these facts in mind the present study was executed for find out the effect of sulphur and zinc on performance of cowpea.

MATERIALS AND METHODS

The experiment was carried out during *kharif* season of 2021 at the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, Uttar Pradesh which is located at 25°24'41.27" N latitude, 81°50'56" E longitude (2021) and 98 m altitude above the mean sea level. This area is situated on the right side of the river *Yamuna* and south east side of Prayagraj city. Having nearly neutral in soil reaction (pH 7.0), organic carbon (0.375 %), available nitrogen (168.75 kg/ha), available phosphorus (17.4 kg/ha) and available potassium (231.7 kg/ha). The climate of the region is semi-arid subtropical. In this experiment a total of ten

Comment [h14]: Mention the range of availability low/medium/high

Comment [h15]: Please check this sentence

treatments has been developed and were tested. Treatment composed of T₁- Sulphur – 20 kg/ha + Zinc – 10 kg/ha, T₂- Sulphur – 20 kg/ha + Zinc – 15 kg/ha, T₃- Sulphur – 20 kg/ha + Zinc – 20 kg/ha, T₄- Sulphur – 30 kg/ha + Zinc – 10 kg/ha, T₅- Sulphur – 30 kg/ha + Zinc – 15 kg/ha, T₆- Sulphur – 30 kg/ha + Zinc – 20 kg/ha, T₇- Sulphur – 45 kg/ha + Zinc – 10 kg/ha, T₈- Sulphur – 45 kg/ha + Zinc – 15 kg/ha, T₉- Sulphur – 45 kg/ha + Zinc – 20 kg/ha, T₁₀- Control (20:50:20 NPK) kg/ha. Ten treatments were replicated thrice in Randomized Complete Block Design. The recommended dose of fertilizer (N:P:K) is 20:50:20 kg/ha. The statistics were calculated and analyzed by using the Ali, J., Singh, S.P. and Singh, S. (2013)(1). statistical approach.

Comment [h16]: Briefly write the methodology of taking the observations

RESULT AND DISCUSSION

Plant height (cm)

Observations recorded in respect to the plant height of cowpea were represented in Table 1 there was an increase in crop age and plant height was progressively noticed with the advancement during the experimentation period. The analysis on plant height was significantly higher in all the different growth intervals with the effect of different levels Sulphur and zinc. At harvest maximum plant height was recorded (75.64 cm) with the application of Sulphur – 45 kg/ha + Zinc – 20 kg/ha which was significantly superior over all other treatments except with the application of Sulphur - 45 kg/ha + Zinc - 15 kg/ha (73.27 cm). The probable reason for increasing plant height might be due to presence of sulphur in the application plays important role in several physiological and biochemical processes which are of vital importance for growth and development of plant. Similar results were earlier reported by Raiger *et al.*, (2017) (9). Application of zinc increase in plant height might be due to its role in biosynthesis of indole acetic acid (IAA) and especially due to its role in initiation of primordial for promoting of photosynthesis which resulted in better plant growth and yield Ali *et al.* (2013)(1).

Number of Branches per plant

Comment [h17]: Remove

The optimum result observations were recorded in no. of branches per plant were depicted in Table 1 and was increased with advancement of the crop during the crop growth period. At harvest maximum no of branches per plant (9.66) recorded with the application of Sulphur – 45 kg/ha + Zinc – 20 kg/ha which was significantly superior over all other treatments except with the application of Sulphur - 45 kg/ha + Zinc - 15 kg/ha (8.89). Maximum no. of branches was observed with application of sulphur 40 kg/ha zinc 20 kg/ha availability of zinc might have stimulated the metabolic and enzymic activity and there by increases the plant

growth attributes which increases the number of branches/plant similar results have also reported by Kasturi krishna and Ahlawat (2000)(5).

Plant dry weight (g/plant)

Observations to be recorded in the dry weight of cowpea were represented in Table 1 the maximum dry weight had given at harvest. At harvest the maximum dry weight (67.90 g/plant) was found in the treatment with application of Sulphur – 45 kg/ha + Zinc – 20 kg/ha which was significantly superior over all other treatments except with the application of Sulphur - 45 kg/ha + Zinc - 15 kg/ha where it was recorded (64.98 g/plant). Zn and S application created a balanced nutritional environment which enhanced metabolic activities and photosynthetic rate, resulting in improvement in plant height and ultimately increases plant dry weight. Similar results were reported by Meena *et al.* (2006)(6), Ramawtar *et al.* (2013)(10).

Yield, Straw yield and Harvest Index of Cowpea

Observation regarding yield are given in Table 2. Grain yield is an important and considerable trait at all the time. Maximum seed yield (13.67 q/ha) and Straw yield (15.26 q/ha) were recorded with application Sulphur – 45 kg/ha + Zinc – 20 kg/ha which was significantly over all the treatments except with the application of Sulphur - 45 kg/ha + Zinc - 15 kg/ha in grain yield (12.03 q/ha) and straw yield (14.27 q/ha) which was statistically at par with the applications of Sulphur – 45 kg/ha + Zinc – 20 kg/ha. The grain yield being the function of cumulative effect of yield attributes, increased significantly due to addition of Sulphur. Sulphur of chloroplast protein resulted in greater photosynthetic efficiency which in turn translated in terms of increase in yield Karche *et al.* (2012)(4). Similar results were also reported by Singh *et al.* (2006)(12), Singh and Mann (2007)(13). Zinc play an important role in biosynthesis of indole acetic acid which is responsible for initiation of primordial for reproductive parts and partitioning of photosynthesis towards them which resulted in better yield. While maximum Harvest index was recorded in the application of Sulphur - 45 kg/ha + Zinc - 10 kg/ha (47.93 %) and found non significant. Similar finding is found due to the increase in grain yield and straw yield the harvest index increases Baviskar *et al.* (2010)(3).

CONCLUSION

It is concluded that application of Sulphur – 45 kg/ha + Zinc – 20 kg/ha was found more productive in grain yield (13.67 q/ha) which was significantly superior over rest of the treatments.

Comment [h18]: 45 kg/ha sulphur + 20 kg/ha zinc

References

1. Ali, J., Singh, S.P. and Singh, S. (2013). Response of fababean to boron, zinc and sulphur application in alluvial soil. *Journal of the Indian Society of Soil Science*. 61(3):202-206.
2. Anonymous. Economic survey of India, ministry of finance (economic division) GOI, New Delhi 2012, 17-22.
3. Baviskar, V.S., Shete, P.G. and Daspute, R.A. (2010). Response of summer cluster bean [*Cyamopsis tetragonoloba* (L.) Taub] to organic fertilizers and different levels of sulphur for vegetable purpose. *International Journal of Agricultural Science*. 6(2):456-458.
4. Karche, R.P., Dalwadi, M.R., Patel, J.C., Gaikwad, V.P. and Panchal, D.B. (2012). Influence of phosphorus and sulphur on yield and nutrient uptake by summer cluster bean grown on typicustochrept of Anand. *Asian Journal of Soil Science*, 7(2): 239-241.
5. Kasturi Krishna, S. and Ahlawat, I.P.S. (2000). Effect of moisture stress, phosphorus, sulphur and zinc fertilizers on growth and development of pea (*Pisum sativum*). *Indian Journal of Agronomy*, 45(2): 353-356.
6. Meena, K.R., Dahama, A.K. and Reager, M.L. (2006). Effect of phosphorus and zinc fertilization on growth and quality of cluster bean (*Cyamopsis tetragonoloba* (L.) Taub.). *Annual Agricultural Research*, 27(3): 224-226.
7. Ng, N.Q. and Marechal, R (1985) Cowpea taxonomy, origin, germplasm. In *Cowpea Research Production and Utilization* (R.S. Singh and K.O. Rachie, Eds.), John Wiley and Sons, NewYork, 11-22.
8. Prasad, A.S. 2003. Zinc deficiency Has been known for 40 years but ignored by global health organizations. *British Medical Journal*, 326:409-410.
9. Raiger, R., Kumawat, B.L., Sanwal, R.C., Kumawat, S.R. and Kumawat, S.R. (2017). Response of Cluster Bean (*Cyamopsis tetragonoloba* L.) to Phosphorus and Sulphur in Torripsamments of Rajasthan, India. *International Journal of Current Microbiology and Applied Sciences*. 6(6): 2003-2008.

10. Ramawtar, A.C. Shivran and Yadav, B.L. (2013). Effect of NP fertilizers, vermicompost and sulphur on growth, yield and quality of cluster bean [*Cymopsis tetragonoloba* (L.)] and their residual effect on grain yield of succeeding wheat [*Triticum aestivum* (L.)]. *Legume Research*, 36 (1): 74-78.
11. Singh, A.K., Bhatt, B.P., Sundaram, P.K., Kumar, S., Bahrati, R.C., Chandra, N. and Rai, M. (2012). Study of site-specific nutrient management of cowpea seed production and their effect on soil nutrient. *Journal of Agriculture Sciences* 4, 191-198.
12. Singh, R.N. Singh, S. and Singh, B. (2006). Interaction effect of sulphur and boron on yield, nutrient uptake and quality characters of soybean [*Glycine Max* (L.) Meril] grown in acidic upland soil. *Journal of Indian Society of Soil Sciences*, 54(4): 516-518.
13. Singh, Y.P. and Mann, J.S. (2007). Interaction effect of sulphur and zinc in groundnut and their availability in Tonk district of Rajasthan. *Indian Journal Agronomy*, 52 (1): 70-73.

Table 1: Influence of Sulphur and Zinc growth attributes of Cowpea

| | Treatments | Plant height (cm) | No. of Branches per plant | Dry weight (g/plant) |
|-----|--------------------------------------|-------------------|---------------------------|----------------------|
| | | At harvest | At harvest | At harvest |
| T1 | Sulphur - 20 kg/ha + Zinc - 10 kg/ha | 65.85 | 8.40 | 58.20 |
| T2 | Sulphur - 20 kg/ha + Zinc - 15 kg/ha | 66.84 | 8.43 | 58.63 |
| T3 | Sulphur - 20 kg/ha + Zinc - 20 kg/ha | 67.94 | 8.46 | 59.01 |
| T4 | Sulphur - 30 kg/ha + Zinc - 10 kg/ha | 68.38 | 8.53 | 61.2 |
| T5 | Sulphur - 30 kg/ha + Zinc - 15 kg/ha | 69.97 | 8.60 | 61.42 |
| T6 | Sulphur - 30 kg/ha + Zinc - 20 kg/ha | 71.33 | 8.70 | 63.70 |
| T7 | Sulphur - 45 kg/ha + Zinc - 10 kg/ha | 70.46 | 8.63 | 63.02 |
| T8 | Sulphur - 45 kg/ha + Zinc - 15 kg/ha | 73.27 | 8.93 | 64.78 |
| T9 | Sulphur - 45 kg/ha + Zinc - 20 kg/ha | 75.64 | 9.66 | 67.90 |
| T10 | Control (NPK 20: 50: 20) kg/ha | 63.33 | 7.86 | 56.58 |
| | SEm (\pm) | 0.845 | 0.264 | 1.124 |
| | CD (0.05%) | 2.51 | 0.784 | 3.342 |

Comment [h19]: $p=0.05\%$

Table 2: Influence of Sulphur and Zinc on Yield and Economics of Cowpea

| | Treatments | Yield | | |
|-----|--------------------------------------|--------------------|---------------------|-------------------|
| | | Grain yield (q/ha) | Stover yield (q/ha) | Harvest index (%) |
| T1 | Sulphur - 20 kg/ha + Zinc - 10 kg/ha | 7.98 | 9.44 | 45.8 |
| T2 | Sulphur - 20 kg/ha + Zinc - 15 kg/ha | 8.91 | 10.30 | 46.37 |
| T3 | Sulphur - 20 kg/ha + Zinc - 20 kg/ha | 9.00 | 10.46 | 46.23 |
| T4 | Sulphur - 30 kg/ha + Zinc - 10 kg/ha | 9.62 | 11.08 | 46.52 |
| T5 | Sulphur - 30 kg/ha + Zinc - 15 kg/ha | 10.83 | 12.45 | 46.51 |
| T6 | Sulphur - 30 kg/ha + Zinc - 20 kg/ha | 11.53 | 13.17 | 46.66 |
| T7 | Sulphur - 45 kg/ha + Zinc - 10 kg/ha | 11.27 | 12.29 | 47.93 |
| T8 | Sulphur - 45 kg/ha + Zinc - 15 kg/ha | 12.03 | 14.27 | 45.78 |
| T9 | Sulphur - 45 kg/ha + Zinc - 20 kg/ha | 13.67 | 15.26 | 47.30 |
| T10 | Control (NPK 20: 50: 20) | 7.90 | 8.70 | 47.45 |
| | S.Em(±) | 0.345 | 0.340 | 1.018 |
| | CD (p=0.05) | 1.02 | 1.010 | - |