

Original Research Article

Variability, correlation, path coefficient and genetic diversity analysis in Indian mustard

ABSTRACT

Phenotypic and genotypic coefficient of variation (PCV & GCV), heritability, genetic advance (GA), correlation, path and diversity analysis for fourteen characters in twenty-five genotypes of Indian mustard (*Brassica juncea* L.) were evaluated for seed yield and thirteen other yield contributing traits. Genetic variability indicated that, the PCV was greater than GCV for all the traits studied was majorly due to the influence of environment. High heritability along with high genetic advance as ~~per cent~~ percent of mean were recorded for plant height, number of primary branches per plant, number of secondary branches per plant, length of main raceme, number of siliquae on main raceme, number of siliquae per plant, 1000 seed weight, biological yield per plant, seed yield per plant and harvest index, Indicating the breeding improvement through direct selection. In correlation analysis, seed yield per plant had significant and positive correlation with number of primary branches per plant, number of secondary branches, number of siliquae on main raceme, number of siliquae per plant, length of siliqua, number of seeds per siliqua, 1000 seed weight, biological yield per plant and harvest index at both genotypic and phenotypic levels. Path coefficient analysis indicated that, the highest positive direct effects was noted for days to 50% flowering, plant height, number of siliquae per plant, number of seeds per siliqua, length of main raceme, number of primary branches per plant, 1000 seed weight, biological yield per plant, harvest index. Therefore, these traits should be given more priority for selection in breeding programme. Genetic divergence assessed using D^2 statistics for characters enabled grouping of all the genotypes in ~~Six~~ six clusters. Diversity among the clusters varied from 9.67 to 24.03 inter-cluster distances, cluster II and V showed maximum inter cluster distance followed by that between cluster V and VI. The genotypes falling in these clusters could be utilized for hybridization programme in Indian Mustard.

Keywords: Indian mustard, variability, genetic advance, heritability, correlation, path analysis and diversity.

INTRODUCTION

Oilseed Brassicas, often known as rapeseed-mustard, are a significant category of oilseed crops worldwide, consisting of eight cultivated crops from the Brassiceae tribe of the Brassicaceae family (Cruciferae). The term "mustard" derives from the European practice of preparing a heated paste by combining the sweet 'must' of old wine with crushed seeds of black mustard (*Brassica nigra*) [1]. In India, Indian mustard (*Brassica juncea* (L.)2n=36) is a major oilseed crop. It is grown for oil, condiments, and therapeutic purposes. However, the crop is mostly grown for the manufacture of edible vegetable oil. Despite belonging to the same family and genus, rapeseed and mustard exhibit unique botanical

characteristics. Rapeseed is an annual herb that grows to a height of 45 to 150 cm. In most situations, a waxy covering covers the stems. Plant leaves are easily distinguished from mustard (rai) plants. Rapeseed leaves are sessile, glabrous, and hairy. The stalk is partially or completely grasped by the lowest segment of the blade (lamina). The fruits are thicker than mustard (rai) and feature a beak that is one-third to half the fruit's length. The seeds are golden or brown in colour, with a smooth seed coat. Mustard plants are tall (90-200 cm), erect, and highly branched. Instead of being dilated at the base and clasping like rapeseed, the leaves are stalked, broad, and pinnatifid. The fruits are narrow, ranging in length from 2 to 6.5 cm, with strong ascending or erect stems and short, thick beaks. The seed has a rough seed coat and is brown to dark brown in colour. Mustard is the world's third most significant oilseed crop, behind soybeans and palm oil. Among the several oilseed crops produced globally, the estimated area, production, and yield of mustard were 34.88 mha, 69.22 mt, and 1.98 Mt/ha, respectively [2]. Rapeseed production, area, and yield in India are 8.30 million hectares, 1.3 million tonnes, and 1.1 million tonnes per acre, respectively. Where rapeseed yield is lower than that of other crops.

Comment [a1]: ?

Genetic variety and diversity are crucial requirements for crop development because they allow for more selective breeding. Because seed yield is a polygenic characteristic that frequently leads to changes in other characters, the link between seed yield and other traits is useful for selecting the proper selection in breeding programmes. During the selection process, correlation analysis measures the degree, direction, and strength of the association between two or more variables. Path coefficient analysis quantifies the direct and indirect impacts of numerous independent variables on a dependent variable [3]. As a result, the current study is being carried out to estimate the heritability and diversity of genotypes for yield and yield contributing variables, as well as to assess the amount of direct and indirect reasons of linkage among various characters through path analysis in mustard.

MATERIALS AND METHODS

The present investigation entitled "Genetic Diversity and Characters Association in Mustard" was carried out under, Department of Plant Breeding & Genetics at Agriculture Farm, School of Agriculture, Lovely Professional University, Jalandhar (Punjab) during the *rabi* 2021-22. The experimental design comprised 25 different genotypes of Mustard (Table 1) were grown in a RBD with three replications during *rabi* 2021-22. Each plot consists of four rows of 5.0 m length. Five competitive plants were selected from each replication for 14 quantitative traits *viz*: days to 50% flowering, days to maturity,

plant height, number of primary branches per plant, number of secondary branches per plant, number of siliqua per plant, number of seed per siliqua, siliqua length of siliqua, number of siliqua on main raceme, 1000- seed weight, biological yield per plant, harvest index, seed yield per plant. The mean values of each genotype were computed for statistical analysis. The standard method of analysis of variance was given [4], phenotypic and genotypic coefficient of variation, heritability (Broad Sense) and genetic advance as percent of mean were estimated by the formula as suggested by [5] and [6]. The formula of genotypic correlation coefficients was estimated by [7]. Path analysis along with genotypic correlation coefficient is applied to know the direct and indirect effects of the components on yield as suggested by [8] and illustrated by [3]. The replicated data were subjected to genetic divergence analysis using Mahalanobis's D² - statistic [9].

RESULTS AND DISCUSSION

The analysis of variance indicated that the mean sum of squares due to genotypes were highly significant at 1% and 5% levels for all the traits except for four traits number of primary branches, number of secondary branches, length of siliqua and 1000 seeds weight (g) (Table 2). Similar results were also obtained by [10],[11],[12] and [13] in which analysis of variance showed significant differences for all the characters under their study.

In the present study, the phenotypic coefficient of variation (PCV) was greater than genotypic coefficient of variation (GCV) for all the traits. The high magnitude of both coefficients was recorded for traits; number of siliqua per plant recorded the highest PCV and GCV followed by seed yield per plant, number of siliqua on main raceme, harvest index, biological yield per plant and plant height (Table 3). The estimates of phenotypic and genotypic coefficient of variability for the characters under study are in accordance with earlier reports, [14],[15],[16],[17],[18],[19],[11] and [20].

High heritability with high genetic advance was recorded for days plant height, number of primary branches per plant, number of secondary branches per plant, length of main raceme, number of siliquae on main raceme, number of siliquae per plant, 1000 seed weight, biological yield per plant, seed yield per plant and harvest index (Table 3). This might be attributed to additive gene action responsible for their expression and hence, phenotypic selection for their amenability, can be brought about. Similar result found, [19] for 1000 seed weight, [21] for number of Siliqua per plant, [22] for 50% flowering, [23],[24] for 50% flowering, plant height, seeds per Siliqua, 1000 seed weight, [25] for seed yield per plant, number of secondary branches, 1000 seed weight, number of seeds per Siliqua, primary branches per plant and Siliqua length, [26],[27],[28] and [29].

The significant phenotypic correlation of seed yield per plant was found positive for the characters namely, number of primary branches per plant, number of secondary branches, number of siliquae on main raceme, number of siliquae per plant, length of siliqua, number of seeds per siliqua, 1000 seed weight, biological yield per plant and harvest index (Table 4). This suggests that these characters should be considered while selecting for improvement in seed yield per plant provided the character should show high variability, which is the basis for selection. However, days to maturity and plant height

showed negative correlation with seed yield per plant. In the present findings, the magnitudes of genotypic correlation coefficient were greater than corresponding phenotypic correlation coefficient (in general also) similar findings by [11],[30],[20],[13] and [19].

The path coefficient analysis of different characters revealed that positive direct effect on seed yield per plant per plant was observed by days to 50% flowering, plant height, number of siliquae per plant, number of seeds per silique, length of main raceme, number of primary branches per plant, 1000 seed weight, biological yield per plant, harvest index (Table 5). We can conclude based on present finding that those all character which had positive direct effect on seed yield per plant that universal traits for overall improvement. Negative direct effect on seed yield per plant was exerted by days to 2 maturity, number of secondary branches per plant, number of silique on main raceme and length of silique. In the present findings, the similar results are also reported by Direct and positive effect by [30],[31],[13],[19],[20],[32] and [31] viz seed yield per plant.

Genetic divergence analysis based on Mahalanobis D² statistics, twenty-five genotypes were grouped into six clusters in mustard (Table 6). The maximum differences among the genotypes within the same cluster (intra-cluster) were shown by cluster I followed by cluster II. Clusters III, IV & V showed zero intra-cluster distances (Table 7). The genotypes showing maximum diversity from clusters I & II could be utilized directly for future hybridization programs. In conformity to the present investigation, similar findings were found by [33],[34] and [35].

It was observed that biological yield per plant was the highest contributor towards divergence followed by all except Length of main raceme (Table 8) don't contribute significantly to the total divergence. Similarly, Devi *et al.*, (2017) was observed, biological yield, Harvest index 1000 seed weight and seed yield per plant were the major contributors for genetic diversity among the genotypes. The cluster mean values for 14 characters of 25 genotypes have been represented in (Table 8).

Table 1: Detail of the Genotypes given below.

Sr. No.	Entries	Pedigree/Source (DRMR)
1	BHAGIRATHI	SELECTION FROM PUSA JAIKISHAN
2	NRCHB101	BL-4X PUSA BOLD
3	VAIBHAV	DRIVED THOUGHT BIPARENTAL MATTING INVOLVING VARUNA, KESHARI, CSU10 AND B1775, B1786, B1866
4	TM4	VARUNAXTM-1

5	KRANTHI	SELECTIONFORVARUNA
6	RH30	SELECTIONFROMP26/3-1
7	DMH1	DEVELOPEDBYCMS
8	RH119	PUSA BOLD XRAJAT(PCR-7)
9	KBS3	PUSAKALYANIX YUKINA
10	ROHINI	SELECTIONFROMNATURALPOPULATIONOF VARUNA
11	URVASHI-18	VARUNAXKRANTI
12	PUSAMUSTARD 24	PUSABOLDXLEB) XLES-29
13	PUSAJAIKISHAN	SOMACLONEOFVARUNA
14	GUJARATH MUSTARD2	SELECTIONFROMMATERIALCOLLECTED FROMVENDACHU,GUJRAT
15	GUJARATH MUSTARD1	MR71-3-2XTM-4
16	PUSABOLD	VARUNAXBIC1780
17	SMR9	-
18	RH749	DRMR
19	BR 23	SELECTIONFROMLOCALGERMPLASMOF PURNEA,BHIHAR
20	TL 17	-
21	JD6	PUSABOLD XGLOSSY
22	PUSAMUSTARD 28	SEJ8XPUSA JAGANNATH
23	RH701	-
24	PARWATHI MUSTARD	SELECTIONFROMGAMMAIRRIGATEDBT4
25	PUSAMUSTARD 27	DIVYA/PUSABOLD/PR666EPS/PR704EPS2

Table 2: Analysis of variance for fourteen characters in Mustard.

Sr. No.	Characters	Sourceofvariation		
		Replications	Genotypes	Error
	Degreeoffreedom	2	24	48
1	Daysto 50%flowering	1.33	27.18**	3.20
2	Daystomaturity	10.09	20.72**	3.88

3	Plantheight(cm)	59.43	1597.16**	180.49
4	Numberofprimarybranches	0.23	0.72	0.15
5	Numberofsecondarybranches	0.06	2.92	0.32
6	Lengthofmain raceme	72.74	178.52**	26.10
7	Numberofsiliquaein main raceme	28.40	96.77**	9.30
8	Number ofsiliquaeper plant	36.30	11410.59**	846.89
9	Lengthofsiliquae	0.01	0.68	0.03
10	Numberofseeds per siliquae	1.01	4.40**	0.59
11	1000seedweight	0.03	2.71	0.07
12	BiologicalYield	271.31	2105.68**	92.50
13	Seedyieldperplant	20.20	387.61**	16.91
14	Harvestingindex	21.04	80.52**	13.94

Table 3.: Estimate of coefficient of variation, h₂ (broad sense) and genetic advance in per cent of mean in Mustard.

Sr. No.	Characters	Range	Mean	Coefficient of variation		Heritability in broad sense (%)	Genetic advance in % of mean
				PCV	GCV		
1	Day to 50% flowering	43.33-52.00	48.23	6.24 %	5.86%	88.20	11.34%
2	Day to maturity	116.00-128	119.33	2.20%	1.98%	81.25	3.68%

3	Plant height(cm)	103.33-202.67	144.08	16.01%	15.08%	88.70	29.26%
4	Number of primary branches	3.06-5.46	3.73	13.19%	11.67%	78.19	21.25%
5	Number of secondary branches	5.20-9.60	6.85	14.43%	13.50%	87.64	26.05%
6	Length of main raceme	51.47-84.40	62.25	12.39%	11.44%	85.38	21.79%
7	Number of siliques in main raceme	23.47-54.13	30.98	18.33%	17.42%	90.39	34.13%
8	Number of siliques per plant	193.27-414.27	277.40	22.23%	21.39%	92.58	42.39%
9	Length of siliques	4.60-6.40	5.73	8.35%	8.13%	94.77	16.31%
10	Number of seeds per siliques	16.60-21.40	19.33	6.26%	5.82%	86.55	11.17%
11	1000 seed weight	3.27-5.03	4.13	13.87%	13.77%	98.56	28.17%
12	Biological Yield	120.30-213.60	162.50	16.29%	15.93%	95.61	32.09%
13	Seed yield per plant	23.10-68.70	51.20	22.21%	21.7253 %	95.64	43.76%
14	Harvesting index	18.90-41.90	31.60	16.39%	14.9109 %	82.68	27.92%

Table 4:
Summary of phenotypic correlation (above diagonal) coefficient & genotypic correlation (below diagonal) coefficient for yield and yield contributing traits in mustard under study.

S.N.	Characters	DFF	DM	PH	NPBP	NSBP	LMR	NSMR	NSPP	LS	NSPS	BYPP	TSW	HI	SYPP
1	DFF	1.0000	+	++	+	+++	+	+	-**	+++	+	-	-	-	-
2	DM	+	1.0000	+	+	+	+	+	+	+	+	-	-	+	-
3	PH	++	++	1.0000	++	++	++	+	-*	++	-	-	-	+	-
4	NPBP	+	+	++	1.0000	++	+	++	-	+	-	-	-	+	+
5	NSBP	++	+	++	++	1.0000	+	+	-*	+	+	-	-	+	+
6	LMR	++	+	++	+	++	1.0000	++	-	+	+	+	+	+	+
7	NSMR	+	++	+	++	+	++	1.0000	+	-	-	-	-	+	+
8	NSPP	-**	+	-*	-	-*	-*	+	1.0000	-**	+	+	+	-	+
9	LS	+	+	+	+	+	+	-	-**	1.0000	+	-	-	+	+
10	NSPS	+	+	-	-	+	+	-	+	+	1.0000	++	+	-	+
11	BYPP	-	-	-	-	-	+	-	+	-	+	1.0000	+	++	++
12	TSW	-	-	-	-	-	+	-	+	-	+	++	1.0000	-	++
13	HI	+	+	+	+	+	+	+	-	+	-	+	+	1.0000	++
14	SYPP	-	-	-	+	+	+	+	+	+	+	++	++	++	1.0000

**indicates significant at 5% and 1% probability level, respectively. DF = Days to 50% flowering, DM = Days to maturity, PH = Plant height, NPBP = Number of primary branches, NSBP = Number of secondary branches, LMR = Length of main raceme, NSMR = Number of siliqua on main raceme, NSP = Number of siliqua/plants, LS = Length of siliqua, NSS = Number of seeds/siliqua, TSW = 1000 seed weight, BY = Biological yield/plant, SY = Seed yield/plant and HI = Harvest index.

Table 5: Genotypic direct and indirect effect of different characters on seed yield per plant in Mustard

S.N.	Characters	DFF	DM	PH	NPBP	NSBP	LMR	NSMR	NSPP	LS	NSPS	BYPP	TSW	HI
1	DFF	0.0142	0.0042	0.0108	0.0069	0.0105	0.0079	0.0021	-0.0085	0.0101	0.0022	-0.0015	-0.0028	0.0024
2	DM	-0.0303	-0.1025	-0.0520	-0.0486	-0.0250	-0.0504	-0.0677	-0.0051	-0.0107	-0.0059	0.0078	0.0205	-0.0217
3	PH	0.1887	0.1263	0.2489	0.1396	0.1807	0.1817	0.1124	-0.1100	0.1443	-0.0423	-0.0608	-0.0483	0.0283
4	NPBP	0.0241	0.0235	0.0278	0.0496	0.0402	0.0217	0.0407	-0.0136	0.0055	-0.0127	-0.0024	-0.0178	0.0296
5	NSBP	-0.0312	-0.0102	-0.0305	-0.0340	-0.0421	-0.0238	-0.0183	0.0203	-0.0171	-0.0018	0.0018	0.0050	-0.0179
6	LMR	0.0622	0.0549	0.0815	0.0489	0.0633	0.1117	0.0652	-0.0483	0.0462	0.0069	0.0071	0.0055	0.0418
7	NSMR	-0.0400	-0.1761	-0.1205	-0.2187	-0.1162	-0.1558	-0.2669	-0.0483	0.0369	0.0273	0.0207	0.0585	-0.1331
8	NSPP	-0.1202	0.0100	-0.0886	-0.0551	-0.0969	-0.0866	0.0363	0.2005	-0.1132	0.0802	0.0173	0.0484	-0.0028
9	LS	-0.0802	-0.0119	-0.0656	-0.0126	-0.0461	-0.0469	0.0156	0.0639	-0.1132	-0.0161	0.0009	0.0095	-0.0097
10	NSPS	0.0009	0.0003	-0.0010	-0.0015	0.0002	0.0004	-0.0006	0.0023	0.0008	0.0058	0.0027	0.0032	-0.0004
11	BYPP	-0.0294	-0.0214	-0.0685	-0.0137	-0.0122	0.0177	-0.0218	0.0242	-0.0023	0.1316	0.2805	0.2247	0.1057
12	TSW	-0.0783	-0.0806	-0.0780	-0.1445	-0.0477	0.0198	-0.0881	0.0971	-0.0336	0.2223	0.3221	0.4020	0.0176
13	HI	0.1115	0.1393	0.0747	0.3915	0.2804	0.2457	0.3279	-0.0092	0.0562	-0.0459	0.2477	0.0287	0.6573
14	SYPP	-0.0081	-0.0442	-0.0610	0.1076	0.1892	0.2430	0.1368	0.1653	0.0098	0.3517	0.8441	0.7372	0.6969
15	Partial R ²	-0.0001	0.0045	-0.0152	0.0053	-0.0080	0.0271	-0.0365	0.0331	-0.0011	0.0020	0.2367	0.2964	0.4581

RSQUARE=1.0025 RESIDUAL EFFECT=SQRT(1-1.0025), Bold figures indicated direct effect. DF = Days to 50% flowering, DM = Days to maturity, PH = Plant height, NPBP = Number of primary branches, NSBP = Number of secondary branches, LMR = Length of main raceme, NSMR = Number of siliqua on main raceme, NSP = Number of siliqua/plants, LS = Length of siliqua, NSPS = Number of seeds/siliqua, TSW = 1000 seed weight, BY = Biological yield/plant, SY = Seed yield/plant and HI = Harvest index.

Table6:Clusteringpatternof25Mustardgenotypes&basedonMahalanobis'sD²statistic.

ClusterNo.	No. ofGenotype s	Genotypes
Cluster1	15	URVASHI-18, KBS-03, TL-17, TM-04, PUSAJAI KISHAN, RH-749, ROHINI, NRCHB-101,VAIBHAV, GUJRAT MUSTARD 1, PUSAMUSTARD28,PUSAMUSTARD24,RH119, JD-06,PUSAMUSTARD27.
Cluster2	6	RH-30, KRANTHI, RH-701, BR-23,PARWATHIMUSTARD,BHAGIRADHI
Cluster3	1	PUSABOLD
Cluster4	1	SMR-09
Cluster5	1	GUJARATMUSTARD2
Cluster6	1	DMH-01

Table 7:Estimatesofaverageintraandinter-clusterdistancesforthe fiveclustersinMustard

Clusters	Cluster1	Cluster2	Cluster3	Cluster4	Cluster5	Cluster6
Cluster1	9.71	14.20	12.72	13.35	15.25	13.96
Cluster2		10.58	19.09	18.92	24.03	16.40
Cluster3			0.00	9.67	14.37	17.03
Cluster4				0.00	15.08	17.22
Cluster5					0.00	20.64
Cluster6						0.00

Table 8: Cluster means and percentage contribution for different characters in mustard

Characters	DFF	DM	PH	NPBP	NSBP	LMR	NSMR	NSPP	LS	NSPS	BYPP	TSW	HI	SYP
Cluster1	48.36	118.87	142.64	3.70	6.83	62.10	29.43	251.09	5.80	18.83	4.03	154.38	32.52	49.96
Cluster2	47.89	118.78	134.30	3.64	6.66	60.71	31.11	321.97	5.66	20.53	4.81	194.77	32.40	62.74
Cluster3	45.67	120.67	136.53	3.87	6.33	51.47	32.73	319.00	4.59	18.40	3.49	138.13	27.55	38.03
Cluster4	43.33	119.33	142.13	3.27	6.20	60.99	30.27	394.53	5.27	20.00	3.29	185.67	18.92	34.93
Cluster5	52.00	119.00	175.40	3.47	6.73	63.73	29.33	252.33	6.42	19.53	3.27	120.27	19.20	23.08
Cluster6	52.00	128.00	202.67	5.47	9.60	84.40	54.13	270.93	6.09	19.67	3.98	136.53	41.86	57.30
MEAN	48.2	120.7	155.6	3.9	7.1	63.9	34.5	301.7	5.7	19.5	3.8	154.9	28.7	44.3
CV%	3.71	1.65	9.32	10.68	8.79	8.21	9.85	10.49	3.31	3.98	5.92	2.88	11.82	8.04
Time ranked 1 st	8	2	2	2	2	0	5	20	38	6	144	10	55	6
% Contribution	2.67%	0.67%	0.67%	0.67%	0.67%	0.00%	1.67%	6.67%	12.67%	2.00%	48%	3.33%	18.33%	2.00%

DF = Days to 50% flowering, DM = Days to maturity, PH = Plant height, NPBP = Number of primary branches, NSBP = Number of secondary branches, LMR = Length of main raceme, NSMR = Number of siliqua on main raceme, NSP = Number of siliqua/plants, LS = Length of siliqua, NSS = Number of seeds/siliqua, TSW = 1000 seed weight, BY = Biological yield/plant, SY = Seed yield/plant and HI = Harvest index.

CONCLUSION

The conclusion that can be reached from variability, correlations, pathcoefficient analysis and genetic divergence is that plant height, number of primary branches, number of secondary branches, number of siliqua on main raceme, 1000 seeds weight, biological yield per plant, seed yield per plant and harvesting index found the most important component characters. Hence, these traits should be considered as selection criteria for yield improvement in mustard.

REFERENCES

- [1] Hemingway J. S. (1976). Mustard: *Brassica spp.* and *Sinapis alba* (*Cruciferae*). pp56-9. (in) *Evolution of Crop Plants*. Simmonds (Ed.). Longmans, London.
- [2] USDA, N. (1999). United States department of agriculture. Natural Resources Conservation Service. *Plants Database*. <http://plants.usda.gov> (accessed in 2000).
- [3] Dewey, D. R. and Lu, K. H. (1959). Correlation and path coefficient analysis of seed production, *Agronomy J.*, 51: 515-518.
- [4] Panse V. G. and Sukhatme P. V. (1961). Statistical methods for agricultural workers, ICAR, New Delhi.
- [5] Burton, G. W. and De Vane, E. H. (1953). Estimating Heritability in Tall Fescue (*Festuca Arundinacea*) from Replicated Clonal Material. *Agronomy Journal*, 45(10), 478-481.
- [6] Johanson, H. W., Robinson, H. F. and Comstock, R. E. 1955. Estimates of genetic and environmental variability in Soyabean. *Agronomy J.*, 47(7): 314- 315.
- [7] Al-Jibouri, H. and Miller, P. and Robinson, H. (1958). Genotypic and Environmental Variances and Covariances in an Upland Cotton Cross of Interspecific Origin 1. *Agronomy Journal - AGRON J.*, 50. 10.
- [8] Wright S. (1921). Correlation and causation. *J. agric. Res.*, 20: 557-587.
- [9] Mahalanobis, P. C. (1936). On the generalized distance in statistics. National Institute of Science of India.
- [10] Sohan, R., and Verma, N. (2010). Genetic Variability and yield components in Indian mustard, *Journal of Oilseeds Research*, 27(2): 170-171.
- [11] Akabari, V. R. and Niranjana, M. (2015). Genetic variability and trait association studies in Indian mustard (*Brassica juncea* L.). *International Journal of Agricultural Sciences*, 11. 35-39.

- [12] Devi, T. R., Devi, N. D., Vivekananda, Y. and Sharma, P. R. (2017). Genetic diversity analysis in Indian mustard (*Brassica juncea* L. Czern and Coss) genotypes using agromorphological parameters. *Electronic Journal of Plant Breeding*, 8(3), 749-753.
- [13] Kumar, R., Kaur, S., Bala, K., Kaur, S. and Sharma, L., (2019). Assessment of Genetic Variability, Correlation and Path Analysis for Yield Traits in F1 Hybrids of Indian Mustard [*Brassica juncea* (L.) Czern & Coss.]. *Agriways*, 7(1):1-7.
- [14] Chowdhury, P. R and Goswami, G. D. (1991). Genetic variability studies in Indian mustard (*Brassica juncea* L. Czern. & Coss.). *Environ. and Ecol.*, 9(4):1003-1006.
- [15] Pant, S. C. and Singh, P. (2001). Genetic variability in Indian mustard. *Agricultural Science Digest*, 21(1):28-30.
- [16] Singh, Poonam, Singh, D. N., and Chakraborty, M. (2003). Variability, heritability and genetic advance in Indian mustard (*Brassica juncea* L.).
- [17] Rai, S. K., Verma, A. and Pandey, D. D. (2005). Genetic variability and character association analysis in Indian mustard (*Brassica juncea* (L.) Czern and Coss.). *Annals of Agri. Bio. Res.*, 10(1):29-34.
- [18] Singh, S. K., Singh, A. K. and Kumar, K. (2007). Genetic variability for yield and its components in Indian mustard (*Brassica juncea* L. Czern & Coss.). *New Botanist*, 34(1/4):147-150.
- [19] Yadav, V. K., Srivastava, K. K., Mishra, V. K. and Negi, S. (2021). Studies on Genetic Variability, Character Association and Genetic Divergence in Indian Mustard (*Brassica juncea* L. Czern and Coss.). *Int. J. Curr. Microbiol. App. Sci.*, 10: 132-143.
- [20] Devi, B. (2018). Correlation and path analysis in Indian mustard [*Brassica juncea* (L.) In agro-climatic conditions of Jhansi (U.P.)]. *Journal of Pharmacognosy and Phytochemistry*, 7(1): 1678-1681.
- [21] Gangapur, D. R., Prakash, B. G. and Hiremath, C. P. (2011). Genetic variability studies of germplasm accessions in Indian mustard under protected and unprotected conditions at Bijapur, Karnataka. *International Journal of Agricultural Sciences*, 7(1), 54-57.
- [22] Singh, M., Tomar, A., Mishra, C. N. and Srivastava, S. B. L. (2011). Genetic parameters and character association studies in Indian Mustard. *Journal of Oilseed Brassica*, 2(1): 35-38.
- [23] Kumar, H., Srivastava, Ayushi, Vishwakarma, M. K. and Lal, J. P. (2012). Genetic Enhancement of Variability through Induced Mutagenesis in Two Genotypes of (*Brassica napus* L.). *Madras Agric. J.*, 99 (4-6):228-231.
- [24] Nasim, A., Farhatullah, Iqbal, S., Shah, S., and Azam, S. M. (2013). Genetic variability and correlation studies for morphological traits in (*Brassica napus* L.). *Pak. J. Bot.* 45(4): 1229-1234.
- [25] Singh, Maharaj, and Chauhan. (2013). Genetic variation and correlation for some physiological character and seed yield in Indian Mustard (*Brassica juncea* L.) Under Rainfed Condition. *J. Oilseeds Res.* 30(2):167-170.

- [26] Tripathi, N., Kumar, K. and Verma, O. P. (2013). Genetic variability, heritability and genetic advance in Indian Mustard (*Brassica juncea* L. Czern and Coss.) for seed yield and its contributing attributes under normal and saline/alkaline condition. *International Journal of Science and Research*, 983-985.
- [27] Meena, P., Chauhan, J. S., Singh, M., Singh, K. H., Rathore, S. S. and Meena, M. L. (2014). Genetic parameters and correlations for seed yield and morphological characters in Indian Mustard [*Brassica juncea* (L.) Czern. & Coss.]. *J. Oilseeds Res.*, 31(2): 114-117.
- [28] Bind, D., Singh, D. and Dwivedi, V. K. (2014). Genetic variability and character association in Indian mustard [*Brassica Juncea* (L.) Czern & Coss.]. *Agric. Sci. Digest.*, 34 (3): 183– 188.
- [29] Synrem, G. I., Rangare, N. R., Myrthong, I. and Bahadure, D. M. (2014). Variability studies in intra specific crosses of Indian mustard [*Brassica juncea* (L.) Czern and Coss.] Genotypes. *IOSR Journal of Agriculture and Veterinary Science* (IOSR-JAVS), 7 (9): 29-32.
- [30] Rauf, M. A., and Rahim, M. A. (2018). Genetic variability studies among yield and its contributing traits in mustard (*Brassica napus* L.). *Adv. Zool. Bot.*, 6, 101–108.
- [31] Dawar, S., Kumar, N., and Mishra, S. P. (2018). Genetic Variability, Correlation and Path Coefficient Analysis in the Indian Mustard (*Brassica juncea* L. Czern and Coss) Varieties Grown in Chitrakoot, India. *Int. J. Curr. Microbiol. App. Sci.*, 7(3): 883-890.
- [32] Nagda, R., Dubey, D., Avinash, H. and Tamatam, D. (2018). Assessment of Genetic Diversity in Mustard Genotypes. *Plant Archives*, Vol. 18 No. 2, 2018 pp. 2091-2096.
- [33] Doddabhimappa, R., Gangapur, B., Prakash, G. and Hiremath, C. P. (2010). Genetic diversity analysis of Indian mustard (*Brassica juncea* L.). *Electronic Journal of Plant Breeding*, 1(4), 407-413.
- [34] Chaurasia, R. K., Ram, B. and Gr Chougule. (2014). Genetic divergence for seed yield and component traits in Indian mustard [*Brassica juncea* (L.) Czern & Coss.]. G. B. Pant University of Agriculture & Technology, Pantnagar (Uttarakhand). *AG RES-An internationale-journal*, 3(3): 260-269.
- [35] Gupta, A., Pant, N. C., Dwivedi, U., Tiwari, S., Pandey, C. S., Dhoundiyal, R. and Verma, O. P. (2018). Studies on correlation and path coefficient analysis for yield and yield related traits in Indian mustard (*Brassica juncea* L. Czern & Coss.) under timely and late sown conditions. *Journal of Pharmacognosy and Phytochemistry*, 7(2), 2545-2551.