

## **Original Research Article**

**Correlation and path coefficient analysis in-of some rice varieties (*Oryza sativa* L.) and rice  
landraces**

**Comment [S1]:** genotypes

### **Abstract**

The present research was carried out to study the correlation and path coefficient analysis in twenty six rice (*Oryza sativa* L.) varieties, among 26 rice genotypes, 15 rice varieties and 11 landraces collections from hilly region of Mirzapur in eastern Uttar Pradesh. At phenotypic and genotypic level, GPP (0.603), Chl (0.326), EBT(0.356) showed highly positive significant correlation with YPP to emerge as most important associates of grain yield in rice. Path analysis identified the highest positive direct effect on grain yield per plant was exhibited by GPP(0.485), DFF(0.414), PH @at the time of 45 DAT(0.255), Chl(0.255), EBT(0.173), PH @at the time 30 DAT (0.042), NOT(0.029), PL(0.024) are the most significant direct as well as indirect effect. Yield contributing components which under consideration at time of devising selection strategy aimed at developing varieties having higher yield. In reference to most of the previous reports on rice, comparatively small proportion of direct and indirect effects of different components attained high order value in the present research.

**Keywords:** Correlation, Path analysis, landraces, Rice

### **Introduction**

Rice (*Oryza sativa* L.) occupies a pivotal place in Indian agriculture, as it forms the staple food for two-thirds of the population and provides 43 per cent calories requirement and 20-25% agriculture income. More than 90 percent of the world's rice is grown and consumed in Asia, where 60 per cent of the earth's people and two third of world's poor live (Khush and Virk, 2000). Rice farming is about 10,000 year old and largest single use of land for producing food. About 11% of total Earth's arable land was covered by rice fields. The frequent occurrence of drought as well as other abiotic stresses has been identified as the major issue to the low productivity of rice in rainfed ecosystems, particularly in eastern region of India. Most of agronomical traits are quantitative traits showing normal distributions in phenotype of the traits. Information on association of characters, direct and indirect effects contributed by each character towards yield will be an added advantage in helping the selection process. (Singh, et. al., 2018)—Correlation and path analysis establish the extent of association between yield and yield components and also bring out relative importance of their direct and indirect effects, thus giving an obvious understanding of their association with grain yield (Singh, et. al., 2018). Ultimately, this kind of analysis could help the breeder to design his selection strategies to improve grain yield. In the light of the above scenario, the present investigation is carried out with the objective of studying the character associations in rice for yield improvement.

**Formatted:** Font: Italic

### **Materials and methods**

The present experiment was carried out at Student's Instructional Farm (SIF), Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya, India. Seeds of the 26 genotypes were sown in raised nursery bed. The seedlings were transplanted to the main field at the rate of one seedling per hill, after 21 days, with a spacing of 20cm x 15cm. The experiment was arranged in a randomized block design (RBD) with thrice replications. The recommended agronomical practices and plant protection measures were followed to ensure a normal crop growth and development. Observations

were recorded on five randomly selected plants in each replication from the two centre rows. 14 traits *viz.* Days to flowering Initiation (DFIT), days to 50% flowering (DFF), days to 100% flowering (DHF), Plant height @at the time of 15, 30, 45 DAT, Maturity (PH), no. of tillers (NOT), panicle length (PL), Spiklete's per panicle (SPP), ear bearing tillers (EBT), grain per panicle (GPP), Chlorophyll (Chl), Grain yield per plant (GYP). Correlation coefficient at the genotypes and phenotypic levels was computed by Singh and Chaudhary (1995) and Dewey and Lu (1959) for path analysis.

UNDER PEER REVIEW

**Table 1. Detailed list of selected rice genotypes and their origin:**

| Name of variety             | Parentage                          | Year of release | Duration (in days) | Eco-System                   | Salient Features   | Recommended for cultivation       |
|-----------------------------|------------------------------------|-----------------|--------------------|------------------------------|--|-----------------------------------|
| Pusa Basmati-1              | Pusa-150 x Karnal Local            | 1989            | 135                | Irrigated Areas              | Semi dwarf (85-95 cm), grains: super fine aromatic, Yield: 45 Q/ha.  | Goa, Mizoram and Uttar Pradesh    |
| NDR-359                     | BG-90-2-4 x 08677                  | 1994            | 115-125            | Irrigated Areas              | Semi dwarf (90-95 cm), grains: short tipped, Yield: 50 Q/ha.   | Uttar Pradesh, Bihar and Orissa   |
| Pusa 1121 (Pusa Sugandha-5) |                                    | 2005            | 135-140            | Irrigated saline soils       | Medium (97.3 cm); Grain-slender, mod. resit. to RTV, sheath rot, & BLB; Yield: 55-65 q/ha.                             | AP and Kerala                     |
| DRR Dhan-44                 | IR93376-B-B-130                    | 2014            | 120                | Upland and drought prone     | Drought tolerant, HYV, Semi tall   | Uttrakhand, Haryana and Bihar     |
| Sahbhagi dhan               |                                    | 2009            | 105                | Rainfed upland/lowland       | LB grain, tolerant to drought. Res to leaf blast, mod. res to brown spot, sheath rot, SB and leaf folder, 3.5-4 t/ha   | Orissa and Jharkhand              |
| Swarna Sub-1                | Swarna 3/IR 49830-7-1-2-3          | 2009            | 145                | Flood prone shallow lowlands | Plant height- 83.3cm<br>Tolerant to complete submergence between 15-17 days, 5.2 t/ha yield                            | UP, Uttrakhand, Haryana and Bihar |
| NDR 2064                    | Pant Dhan 4/Saket4 // NDR 2017     | 2007            | 115                | irrigated areas              | 50-55 Q/ha yield grains are (M.S)Medium Size, High yielding  | UP, Orissa and West Bengal        |
| NDR 2065                    | Pant Dhan 4/Saket 4// NDR 2018     | 2011            | 120-125            | irrigated areas              | 50-55Q/ha grains are LB(Long Bold), High yielding  | UP, Orissa and West Bengal        |
| NDR 97                      | Nanina-22 x Ratna                  | 1992            | 90-95              | Rain fed Uplands             | Dwarf (75-80 cm), Yield: 25-30 Q/ha.   | UP, Orissa and West Bengal        |
| IR-64                       | IR-5857-33-2-1 x IR-2061-465-1-5-5 | 1991            | 115-120            | Irrigated Areas              | Semi dwarf (100 cm), grains: Yield: 58 Q/ha.   | All India                         |
| Sarjoo-52                   | T(N)1 x Kashi                      | 1982            | 130-133            | Irrigated                    | Semi dwarf (98 cm), erect, grains: long bold, white, moderately resistant to Bacterial Leaf Blight, Yield: 50-60 Q/ha. | Uttar Pradesh.                    |

**Table 2: Details of Rice landraces and their area of collection:**

| S No. | Local Name                 | Village     | District                 | Characteristics   |
|-------|----------------------------|-------------|--------------------------|---|
| 1.    | Local Selection 1( LS 1)   | Tisuhi      | Mirzapur, Uttar Pradesh  | Brown grain, Semi long, Drought tolerant, short stature, early maturity             |
| 2.    | Local Selection 3( LS 3)   | Bharko      | Mirzapur, Uttar Pradesh  | Brown grain, elongated, Drought tolerant, medium stature                            |
| 3.    | Local Selection 4( LS 4)   | Pochkhora   | Mirzapur, Uttar Pradesh  | Red grain, late maturity  |
| 4.    | Local Selection 5( LS 5)   | Jamunia     | Mirzapur, Uttar Pradesh  | Brown grain, Semi long, short stature, late maturity                                |
| 5.    | Local Selection 6( LS 6)   | Bharko      | Mirzapur, Uttar Pradesh  | Brown grain, semi spherical, medium stature   |
| 6.    | Local Selection 7( LS 7)   | Pochkhora   | Mirzapur, Uttar Pradesh  | Brown grain, bold grain, Semi long, Drought tolerant, short stature, early maturity |
| 7.    | Local Selection 8( LS 8)   | Manihan     | Mirzapur, Uttar Pradesh  | Brown grain, semi long, medium stature  |
| 8.    | Local Selection 9( LS 9)   | Deep Nagar  | Mirzapur, Uttar Pradesh  | Brown grain, Semi long, Drought tolerant, short stature, early maturity             |
| 9.    | Local Selection 10( LS 10) | Jamunia     | Mirzapur, Uttar Pradesh  | Brown grain, semi long, medium stature, early maturity                              |
| 10.   | Local Selection 11( LS 11) | Tisuhi      | Mirzapur, Uttar Pradesh  | Brown grain, elongated, long stature, late maturity                                 |
| 11.   | Local Selection 12( LS 12) | Mugal sarai | Chaundali, Uttar Pradesh | Black grain, elongated, Drought tolerant, long stature, late maturity               |

**Comment [S2]:** Write the local name of the land races

## Results and Discussion

### Phenotypic correlation coefficients

In the present investigation, the genotypic correlation coefficients were generally higher than their respective phenotypic correlation coefficients (Table 3). At phenotypic level, YPP showed highly positive significant correlation by GPP (0.603), Chl (0.326). Similar trends of results were also reported by Kishore *et al.* 2007. Positive significant correlation was obtained in EBT (0.249). Highly negative significant correlation was showed with PH@at the time of maturity (0.341). Negative significant correlation was showed by PH @at the time of 45DAT (0.223). Positive non-significant correlation was showed with DFF (0.149), DHF (0.102), DTFI (0.095), PL (0.088), SPP (0.072), PH @at the time of 30DAS (0.006). Negative non-significant correlation was with PH@at the time of 15DAS (0.012), NOT (0.088). Chl content showed highly negative significant correlation by PH@at the time of 15DAS, Positive significant correlation with GPP. Positive non-significant correlation with PL, SPP and PH@at the time of maturity, PH @at the time of 30 DAS, PH @at the time of 45 DAT show with Negative non-significant correlation. GPP showed highly negative significant correlation with PH@at the time of maturity, NOT. Positive non-significant correlation was obtained in GPP, PH@at the time of 15DAS, PH @at the time of 45DAT. Negative non-significant correlation in DHF, followed by EBT, PH @at the time of 30DAS and EBT showed positive significant correlation in DHF. Positive non-significant correlation with DFF, PH@at the time of 15DAS, DTFI, SPP, PH @at the time of 45DAT, PH @at the time of 30DAS, PH@at the time of maturity, PL. SPP showed highly negative significant correlation in PH@at the time of maturity and Negative significant correlation was showed with PH @at the time of 45DAT. Positive non-significant correlation was obtained in NOT, DHF, DFF, DTFI, PL. Negative non-significant was obtained in PH @at the time of 30DAS, PH@at the time of 15DAS. PL showed positive significant correlation in PH @at the time of 45DAS. Positive non-significant correlation was obtained in NOT, DFF, PH @at the time of 30DAS, DTFI, PH@at the time of maturity. Negative non-significant correlation was obtained in PH@at the time of 15DAS, DHF. Similar results were reported by Lalitha and Shreedhar (1996) .NOT showed positive non-significant correlation in PH@at the time of maturity. Negative non-significant correlation was obtained in PH @at the time of 30DAS, PH @at the time of 45DAT, DTFI, DHF, DFF, PH@at the time of 15DAS. PH@at the time of maturity showed highly positive significant correlation in DTFI, PH @at the time of 45DAT, DFF. Positive significant correlation was obtained in PH@at the time of 15DAS, DHF. Positive non-significant correlation was obtained in PH @at the time of 30DAS. PH@at the time of 45DAT showed highly positive correlation in DHF. Positive significant correlation was showed in PH@at the time of 15DAS. Positive non-significant correlation was obtained in DTFI, DFF. PH@at the time of 30DAS showed highly positive significant correlation in DHF. Positive significant correlation was obtained in PH@at the time of 15DAS. Positive non-significant correlation was obtained in DTFI, DFF(0.054). PH @at the time of 15DAS showed highly positive significant correlation with DFF, DHF. DTFI showed positive significant correlation. DHF showed highly positive significant correlation with DTFI, DFF. DFF showed highly positive significant correlation with DTFI. These positive association between these characters have also been reported by Chand *et. al.*, 2007, Borbora *et. al.*, 2005.

### Genotypic correlation coefficient

Yield per plant showed highly positive significant correlation with GPP(0.637), Chl(0.417), EBT(0.356). Highly negative significant correlation was obtained in PH@at the time of maturity(0.375). Negative non-significant correlation was obtained with PH @at the time of 45DAT(0.251), NOT (0.246).

Positive non-significant correlation was showed by PL (0.199), DFF(0.154), DHF(0.122), SPP (0.117), DTFI (0.100). Negative non-significant correlation was obtained in PH@at the time of15DAS (0.002), PH @at the time of30DAS (0.070). Chl showed highly positive significant correlation by GPP (0.280). Positive significant correlation with PH@at the time of15DAS, PH @at the time of30DAS, DHF, DTFI(0.339), DFF, NOT and positive non- significant correlation was showed by EBT, PL, SPP. Negative non-significant correlation was obtained with PH@at the time of maturity, PH @at the time of45DAT. Characters mentioned above has also being reported in rice by earlier workers (Qamar *et al.* 2005; Ram Krishan *et al.* 2006) GPP showed highly positive significant correlation was obtained in EBT and negative highly significant correlation was obtained in NOT, PH @at the time of45DAT, PH@at the time of maturity. Positive non-significant correlation was obtained in PH @at the time of30DAS, SPP, DHF. Negative non-significant correlation was found in DFF, DTFI, PH@at the time of15DAS, PL. EBT showed highly positive significant correlation in PH @at the time of30DAS, PH@at the time of15DAS, DHF, DTFI, DFF, SPP, PH @at the time of45DAT. Negative non-significant correlation was obtained in NOT, PL. Positive non-significant correlation was obtained in PH@at the time of maturity(0.082). SPP showed highly positive significant correlation in NOT. Highly negative significant correlation was found in PH @at the time of30DAS, PH@at the time of maturity, PH @at the time of45DAT. Positive non-significant correlation was obtained in PL, DFF, DTFI, DHF. Negative significant correlation was obtained in PH@at the time of15DAS and PL showed highly significant correlation was obtained in PH @at the time of45DAT. Highly negative significant correlation was obtained in PH @at the time of30DAS. Negative significant correlation was obtained in NOT. Positive non-significant correlation was obtained in DFF, DTFI, PH@at the time of maturity. Negative non-significant correlation was obtained in PH@at the time of15DAS, DHF. NOT showed highly negative significant correlation was obtained in PH @at the time of30DAS(1.088), PH@at the time of15DAS, DFF, DHF. Positive non-significant correlation was obtained with PH @at the time ofmaturity. Negative non-significant correlation was found in PH @at the time of45DAT, DTFI. PH@at the time ofmaturity showed highly positive significant correlation in PH @at the time of45DAT, PH@at the time of15DAS, DTFI, DFF. Positive significant correlation was showed in DHF, PH @at the time of30DAS. PH@at the time of 45DAT showed highly positive significant correlation was obtained in PH @at the time of30DAS, PH@at the time of15DAS. Positive significant correlation was obtained in DFF. Positive non-significant correlation was obtained in DHF, DTFI. PH@at the time of 30DAS showed highly positive significant correlation in DHF, PH@at the time of15DAS. Positive significant correlation was obtained in DTFI, DFF. In PH @at the time of15DAS showed highly positive significant correlation with DTFI, DFF, DHF. DHF showed highly positive significant correlation with DTFI, DFF. DFF showed highly positive significant correlation with DTFI. (Mahto *et al.* (2003), Chand *et al.* 2007)

#### **Path-coefficient Analysis**

##### **Phenotypic path coefficients**

The direct and indirect effect of different characters on grain yield/plant computed by using phenotypic correlations are presented in Table-4. The highest positive direct effect on grain yield per plant was exhibited by GPP(0.485), DFF(0.414), PH @at the time of45DAT(0.255), Chl(0.255), EBT(0.173), PH @at the time of30DAS(0.042), NOT(0.029), PL(0.024) the direct effects of remaining characters were too low to be considered important. GPP exhibited indirect positive effect on grain yield per plant via PH@at the time of maturity(0.161), Chl(0.064), EBT(0.02), DTFI(0.002), PH@at the time of 15 DAS(0.002), PH @at the time of30 DAS(0.001), SPP(0.001), High direct effect of filled

spikelets/panicle on single plant yield was reported by Eidi kohnaki *et al.*, (2013), Kiani and Nematzadeh (2012), Seyoum *et al.*, (2012), Bagheri *et al.*, (2011), Bhadru *et al.*, (2011) and Chandra *et al.*, (2009).

#### **Genotypic path coefficients**

The highest positive direct effect on grain yield per plant was exhibited by DTFI (1.034), GPP(0.372), SPP(0.358), PL(0.295), PH @at the time ofmaturity(0.187), EBT(0.135), PH @at the time of30DAS (0.100), Chl (0.046). DTFI exhibited indirect positive effect on grain yield per plant *via* EBT, PH @at the time of maturity , NOT, SPP, PL, PH @at the time of 30 DAS, similar result supported that Bhadru *et al.*, (2011) and Chandra *et al.*, (2009) reported positive direct effect of days to 50% flowering and Eidi kohnaki *et al.*, (2013) and Nematzadeh (2012) found the positive direct effect.

**Table 3: Estimation of phenotypic correlation coefficient for 14 characters in selected rice germplasm**

|  | Traits                            | DTF<br>I | DFF         | DHF         | PH<br>@at<br>the<br>time<br>of15<br>DAS | PH<br>@at<br>the<br>time<br>of30<br>DAS | PH<br>@at<br>the<br>time<br>of45<br>DAT | PH<br>@at<br>the<br>time of<br>Maturi<br>ty | NOT    | PL     | SPP          | EBT    | GPP          | Chl          | GYP          |
|--|-----------------------------------|----------|-------------|-------------|---|---|---|---|--------|--------|--------------|--------|--------------|--------------|--------------|
|  | DTFI                              | 1.000    | 0.968<br>** | 0.865<br>** | 0.268<br>*                              | 0.068                                   | 0.187                                   | 0.315<br>**                                 | -0.041 | 0.051  | 0.079        | 0.172  | -0.039       | -0.307<br>** | 0.095        |
|  | DFF                               |          | 1.000       | 0.832<br>** | 0.322<br>**                             | 0.054                                   | 0.224<br>*                              | 0.305<br>**                                 | -0.090 | 0.136  | 0.112        | 0.184  | -0.010       | -0.275<br>*  | 0.149        |
|  | DHF                               |          |             | 1.000       | 0.300<br>**                             | 0.311<br>**                             | 0.206                                   | 0.255<br>*                                  | -0.072 | -0.034 | 0.114        | 0.246* | 0.012        | -0.301<br>** | 0.102        |
|  | PH @at the<br>time of15 DAS       |          |             |             | 1.000                                   | 0.262<br>*                              | 0.484<br>**                             | 0.288<br>*                                  | -0.171 | -0.018 | -0.188       | 0.177  | -0.033       | -0.342<br>** | -0.012       |
|  | PH @at the<br>time of30 DAS       |          |             |             |   | 1.000                                   | 0.222<br>*                              | 0.107                                       | -0.037 | 0.057  | -0.071       | 0.061  | 0.012        | -0.169       | 0.006        |
|  | PH @at the<br>time of 45 DAS      |          |             |             |   |   | 1.000                                   | 0.740<br>**                                 | -0.039 | 0.237* | -0.283<br>*  | 0.086  | -0.480<br>** | -0.189       | -0.223<br>*  |
|  | PH @at the<br>time of<br>Maturity |          |             |             |   |   |   | 1.000                                       | 0.002  | 0.024  | -0.347<br>** | 0.022  | -0.439<br>** | -0.160       | -0.341<br>** |
|  | NOT                               |          |             |             |   |   |   |   | 1.000  | 0.201  | 0.193        | -0.124 | -0.120       | -0.009       | -0.088       |
|  | PL                                |          |             |             |   |   |   |   |        | 1.000  | 0.046        | 0.001  | -0.133       | 0.051        | 0.088        |
|  | SPP                               |          |             |             |   |   |   |   |        |        | 1.000        | 0.170  | -0.023       | 0.045        | 0.072        |
|  | EBT                               |          |             |             |   |   |   |   |        |        |              | 1.000  | 0.117        | -0.051       | 0.249<br>*   |
|  | GPP                               |          |             |             |   |   |   |   |        |        |              |        | 1.000        | 0.251<br>*   | 0.603<br>**  |
|  | Chl                               |          |             |             |   |   |   |   |        |        |              |        |              | 1.000        | 0.326<br>**  |



Table 4: Estimation of genotypic correlation coefficient for 14 characters in selected rice germplasm

| Traits                      | DTFI  | DFF     | DHF     | PH @at the time of 15 DAS | PH @at the time of 30 DAS | PH @at the time of 45 DAT | PH @at the time of Maturity | NOT      | PL       | SPP      | EBT      | GPP      | Chl      | GYP      |
|-----------------------------|-------|---------|---------|---------------------------|---------------------------|---------------------------|-----------------------------|----------|----------|----------|----------|----------|----------|----------|
| DTFI                        | 1.000 | 0.972** | 0.875** | 0.307**                   | 0.171                     | 0.188                     | 0.319**                     | -0.185   | 0.062    | 0.123    | 1.130**  | -0.043   | -0.339** | 0.100    |
| DFF                         |       | 1.000   | 0.845** | 0.379**                   | 0.088                     | 0.230*                    | 0.309**                     | -0.367** | 0.202    | 0.127    | 1.121**  | -0.009   | -0.306** | 0.154    |
| DHF                         |       |         | 1.000   | 0.353**                   | 0.755**                   | 0.205                     | 0.262*                      | -0.300** | -0.014   | 0.122    | 1.752**  | 0.010    | -0.340** | 0.122    |
| PH @at the time of 15 DAS   |       |         |         | 1.000                     | 0.481**                   | 0.561**                   | 0.343**                     | -0.455** | -0.004   | -0.166   | 1.986**  | -0.080   | -0.485** | -0.002   |
| PH @at the time of 30 DAS   |       |         |         |                           | 1.000                     | 0.592**                   | 0.229*                      | -1.088** | -0.423** | -0.589** | 2.599**  | 0.018    | -0.390** | -0.070   |
| PH @at the time of 45 DAS   |       |         |         |                           |                           | 1.000                     | 0.757**                     | -0.173   | 0.326**  | -0.374** | 0.561**  | -0.509** | -0.203   | -0.251*  |
| PH @at the time of Maturity |       |         |         |                           |                           |                           | 1.000                       | 0.123    | 0.036    | -0.485** | 0.082    | -0.457** | -0.183   | -0.375** |
| NOT                         |       |         |         |                           |                           |                           |                             | 1.000    | -0.284*  | 0.978**  | -1.707** | -0.570** | -0.280** | -0.246*  |
| PL                          |       |         |         |                           |                           |                           |                             |          | 1.000    | 0.204    | -0.429** | -0.169   | 0.138    | 0.199    |

|            |  |  |  |  |  |  |  |  |  |              |              |              |              |         |
|------------|--|--|--|--|--|--|--|--|--|--------------|--------------|--------------|--------------|---------|
| <b>SPP</b> |  |  |  |  |  |  |  |  |  | <b>1.000</b> | 0.592**      | 0.014        | 0.042        | 0.117   |
| <b>EBT</b> |  |  |  |  |  |  |  |  |  |              | <b>1.000</b> | 1.121**      | 0.177        | 0.356** |
| <b>GPP</b> |  |  |  |  |  |  |  |  |  |              |              | <b>1.000</b> | 0.280**      | 0.637** |
| <b>Chl</b> |  |  |  |  |  |  |  |  |  |              |              |              | <b>1.000</b> | 0.417** |

**Table 5. Phenotypic Path Coefficient for 14 character in selected rice germplasm**

| Trait                       | DTFI          | DHF          | DHF           | PH@at the time of 15 DAS | PH @at the time of 30 DAS | PH @at the time of 45 DAS | PH@at the time of maturity | NOT          | PL           | SPP           | EBT          | GPP          | Chl          | GYP      |
|-----------------------------|---------------|--------------|---------------|--------------------------|---------------------------|---------------------------|----------------------------|--------------|--------------|---------------|--------------|--------------|--------------|----------|
| DTFI                        | <b>-0.055</b> | 0.343        | -0.095        | -0.015                   | 0.003                     | 0.047                     | -0.116                     | -0.001       | 0.001        | -0.005        | 0.030        | -0.019       | -0.078       | 0.095    |
| DFF                         | -0.053        | <b>0.414</b> | -0.092        | -0.019                   | 0.002                     | 0.057                     | -0.112                     | -0.003       | 0.003        | -0.007        | 0.032        | -0.005       | -0.070       | 0.149    |
| DHF                         | -0.048        | 0.343        | <b>-0.110</b> | -0.017                   | 0.013                     | 0.052                     | -0.094                     | -0.002       | -0.001       | -0.007        | 0.042        | 0.006        | -0.077       | 0.102    |
| PH @at the time of 15 DAS   | -0.015        | 0.133        | -0.034        | <b>-0.058</b>            | 0.011                     | 0.122                     | -0.106                     | -0.005       | 0.000        | 0.011         | 0.031        | -0.016       | -0.087       | -0.012   |
| PH @at the time of 30 DAS   | -0.004        | 0.022        | -0.034        | -0.015                   | <b>0.042</b>              | 0.056                     | -0.039                     | -0.001       | 0.001        | 0.004         | 0.010        | 0.006        | -0.043       | 0.006    |
| PH @at the time of 45 DAT   | -0.010        | 0.088        | -0.024        | -0.028                   | 0.009                     | <b>0.252</b>              | -0.272                     | -0.001       | 0.006        | 0.017         | 0.015        | -0.233       | -0.048       | -0.223*  |
| PH @at the time of maturity | -0.012        | 0.129        | -0.031        | -0.017                   | 0.005                     | 0.187                     | <b>-0.367</b>              | 0.000        | 0.001        | 0.020         | 0.004        | -0.213       | -0.041       | -0.341** |
| NOT                         | 0.002         | -0.037       | 0.008         | 0.010                    | -0.002                    | -0.010                    | -0.001                     | <b>0.029</b> | 0.005        | -0.011        | -0.021       | -0.058       | -0.002       | -0.088   |
| PL                          | -0.003        | 0.056        | 0.005         | 0.001                    | 0.002                     | 0.060                     | -0.009                     | 0.006        | <b>0.024</b> | -0.003        | 0.000        | -0.064       | 0.013        | 0.088    |
| SPP                         | -0.005        | 0.043        | -0.013        | 0.011                    | -0.003                    | -0.071                    | 0.127                      | 0.006        | 0.001        | <b>-0.058</b> | 0.029        | -0.011       | 0.012        | 0.072    |
| EBT                         | -0.009        | 0.087        | -0.027        | -0.010                   | 0.003                     | 0.022                     | -0.008                     | -0.004       | 0.000        | -0.010        | <b>0.173</b> | 0.057        | -0.013       | 0.249*   |
| GPP                         | 0.002         | -0.003       | -0.001        | 0.002                    | 0.001                     | -0.121                    | 0.161                      | -0.004       | -0.003       | 0.001         | 0.020        | <b>0.485</b> | 0.064        | 0.603**  |
| Chl                         | 0.017         | -0.115       | 0.033         | 0.020                    | -0.007                    | -0.048                    | 0.059                      | 0.000        | 0.001        | -0.003        | -0.009       | 0.122        | <b>0.255</b> | 0.326**  |

*R SQUARE = 0.5292 RESIDUAL EFFECT = 0.686*

1

**Table 6. Genotypic path coefficient for 14 character in selected rice genotypes**

| Traits                           | DTFI         | DFF           | DHF           | PH@at<br>the time<br>of 15<br>DAS | PH<br>@at<br>the<br>time of<br>30<br>DAS | PH @at<br>the time<br>of 45<br>DAS | PH @at<br>the time<br>of<br>maturity | NOT           | PL           | SPP          | EBT    | GPP    | Chl    | GYP      |
|----------------------------------|--------------|---------------|---------------|-----------------------------------|--|------------------------------------|--------------------------------------|---------------|--------------|--------------|--------|--------|--------|----------|
| DTFI                             | <b>1.034</b> | -0.598        | -0.576        | -0.051                            | 0.017                                    | -0.032                             | 0.060                                | 0.057         | 0.018        | 0.044        | 0.152  | -0.016 | -0.016 | 0.100    |
| DFF                              | 1.004        | <b>-0.619</b> | -0.546        | -0.063                            | 0.009                                    | -0.039                             | 0.058                                | 0.112         | 0.060        | 0.046        | 0.151  | -0.004 | -0.014 | 0.154    |
| DHF                              | 0.905        | -0.525        | <b>-0.648</b> | -0.059                            | 0.076                                    | -0.035                             | 0.049                                | 0.092         | -0.004       | 0.044        | 0.236  | 0.004  | -0.016 | 0.122    |
| PH@at<br>the time of<br>15 DAS   | 0.308        | -0.237        | -0.233        | <b>-0.167</b>                     | 0.048                                    | -0.095                             | 0.064                                | 0.139         | -0.001       | -0.059       | 0.268  | -0.030 | -0.023 | -0.002   |
| PH@at<br>the time of<br>30 DAS   | 0.176        | -0.056        | -0.492        | -0.080                            | <b>0.100</b>                             | -0.101                             | 0.043                                | 0.332         | -0.125       | -0.211       | 0.350  | 0.007  | -0.018 | -0.070   |
| PH@at<br>the time of<br>45 DAT   | 0.188        | -0.145        | -0.143        | -0.094                            | 0.059                                    | <b>-0.170</b>                      | 0.142                                | 0.053         | 0.096        | -0.134       | 0.076  | -0.189 | -0.009 | -0.251*  |
| PH@at<br>the time of<br>Maturity | 0.330        | -0.191        | -0.170        | -0.057                            | 0.023                                    | -0.129                             | <b>0.187</b>                         | -0.038        | 0.011        | -0.174       | 0.011  | -0.170 | -0.009 | -0.375** |
| NOT                              | -0.192       | 0.243         | 0.195         | 0.076                             | -0.109                                   | 0.029                              | 0.023                                | <b>-0.306</b> | -0.084       | 0.350        | -0.230 | -0.212 | -0.013 | -0.246*  |
| PL                               | 0.064        | -0.125        | 0.009         | 0.001                             | -0.042                                   | -0.055                             | 0.007                                | 0.087         | <b>0.295</b> | 0.073        | -0.058 | -0.063 | 0.006  | 0.199    |
| SPP                              | 0.132        | -0.083        | -0.079        | 0.028                             | -0.059                                   | 0.063                              | -0.091                               | -0.299        | 0.060        | <b>0.358</b> | 0.080  | 0.005  | 0.002  | 0.117    |

|     |        |        |        |        |        |        |        |       |        |       |              |              |              |         |
|-----|--------|--------|--------|--------|--------|--------|--------|-------|--------|-------|--------------|--------------|--------------|---------|
| EBT | 1.173  | -0.644 | -1.143 | -0.332 | 0.261  | -0.095 | 0.015  | 0.522 | -0.127 | 0.212 | <b>0.135</b> | 0.417        | 0.008        | 0.356** |
| GPP | -0.045 | 0.006  | -0.067 | 0.013  | 0.002  | 0.087  | -0.086 | 0.174 | -0.050 | 0.005 | 0.151        | <b>0.372</b> | 0.013        | 0.637** |
| Chl | -0.364 | 0.193  | 0.220  | 0.081  | -0.039 | 0.034  | -0.034 | 0.086 | 0.041  | 0.015 | 0.024        | 0.104        | <b>0.046</b> | 0.417** |

*R SQUARE = 0.3750 RESIDUAL EFFECT = 0.7906*

The estimated residual effect was 0.790 indicating that about 80% of the variability in grain yield was contributed by the characters studied in path analysis. This residual effect towards yield in the present study might be due to many reasons, such as other characters, which are not included in the investigation, environmental factor and sampling errors. Within the scope of path analysis carried out in the present study, it is therefore, suggested that number of spikelets per panicle (SPP) and number of tillers (NOT), the main components of grain yield should be given high priority in the selection programme.

## References:

- Bagheri, N.; Babaeian-Jelodar, N.; Pasha, A. (2011); Path coefficient analysis for yield and yield components in diverse rice (*Oryza sativa* L.) genotypes. *Bih Biol.* **5**: 32-35.
- Bhadru, D.; Reddy, D. L. and Ramesha, M. S. (2011); Correlation and path coefficient analysis of yield and yield contributing traits in rice hybrids and their parental lines; *Electronic Journal of Plant Breeding*; **2**(1): 112-116.
- Borbor, T.K.; Hazarika, G.N.; Medhi, A.K. (2005); Correlation and path analysis for panicle characters in rice. *Crop Res.*; **30**(2):215-222.
- Chandra, B. S.; Reddy, T. D.; Ansari, N. A. and Kumar, S. S. (2009) Correlation and path coefficient analysis for yield and yield components in rice (*Oryza sativa* L.). *Agric. Sci. Digest* **29** (1): 45-47.
- Dewey, D.R.; Lu, K.H.A.(1959) Correlation and path coefficient analysis of components of crested wheat grass seed production. *Agron. J.*; **51**:515-518.
- Eidi kohnaki, M.; Kiani, G. and Nematzadeh. G. (2013) Relationship between Morphological Traits in Rice Restorer Lines at F3 Generation using Multivariate Analysis; *Int J Adv Biol Biom Res.* **1**(6):572-577.
- Khush, G.S. and Virk, P. S. (2000) Rice breeding achievements and future strategies. *Crop Improvement*; **27**(2):115-144.
- Kiani, G. and Nematzadeh. G. (2012) Correlation and Path Coefficient Studies in F Populations of Rice. *Not Sci Biol.* **4**(2):124-127.
- Mahto, R.N.; Yadava, M.S.; Mohan, K.S. (2003) Genetic variation, character association and path analysis in rainfed upland rice. *Indian J Dryland Agric. Res. and Devel.*; **18**(2):196-198.
- Qamar Zia, U.; Cheema, A.A.; Ashraf, M.; Rashid, M.; Tahir, G.R. (2005); Association analysis of some yield influencing traits in aromatic and non-aromatic rice. *Pak. J Bot.*; **37**(3):613-627.
- Ramakrishnan, S.; Hari Ananda Kumar C.R.; Sarvanan, S.; Malini, N. (2006); Association analysis of some yield traits in Rice (*Oryza sativa* L.). *Journal of Applied Sciences Research*; **2**(7):402-404.
- Seyoum, M.; Alamerew, S. and Bantte, K. (2012); Genetic Variability, Heritability, Correlation Coefficient and Path Analysis for Yield and Yield Related Traits in Upland Rice (*Oryza sativa* L.). *Journal of Plant Sciences* **7**(1): 13-22.
- Singh, R. K and Chaudhary, B. D. (1995), Biometrical methods in quantitative genetic analysis. Kalyani Publishers New Delhi ., pp. 215-218.
- Singh, R.; Yadav, V.; Mishra, D.N. and Yadav, A., (2018); Correlation and Path Analysis Studies in Rice (*Oryza sativa* L.); *Journal of Pharmacognosy and Phytochemistry*; SP1: 2084-2090.

**Comment [S3]:** Follow the journal style.