

Influence of microbial priming on germination and seedling growth traits of compact cotton CO17

ABSTRACT

Cotton, known as “the King of fibers”, is the predominant fiber in the Indian textile industry. Plant growth-promoting rhizobacteria (PGPR) represent a potential sustainable alternative for the enhancement and protection of crops. The germination and seedling growth of cotton can be optimized by inoculating with PGPR. An experiment was conducted to evaluate the effect of different PGPR strains on seed germination and seedling establishment characters on cotton. The highest germination percentage, maximum vigour index and leaf area was obtained with the *PPFM* TNAU1 strain inoculation. The maximum shoot and root length were observed with seeds treated with *Azospirillum* strain sp7 with an increase of 24.4 and 42.8 % over the control. Underground fresh and dry matter was ~~recorded~~ higher in ~~the~~ seedlings treated with *Azospirillum* sp7 strain compared to control, while the *PPFM* TNAU1 strain treatment increased the aerial fresh and dry matter content because of its larger leaf area. Seeds inoculated with individual strain of *Azospirillum* sp7 and *PPFM* TNAU1 outperformed the combined inoculation of PGPR strains. The increase in germination traits and seedling characters by PGPR strains indicates the positive influence on improving cotton seedling establishment traits associated to for higher better yield.

Keywords: Cotton, plant growth promoting rhizobacteria, microbial priming, germination and seedling emergence traits.

1. INTRODUCTION

Cotton is one of the predominant commercial ~~and a crops with of~~ global significance, playing ~~an important momentous~~ role in foreign exchange and industrial economy. Current production of cotton fiber is not sufficient to meet the gradually increasing demand of world population which arises due to many limiting factors, *i.e.*, ~~frequent droughts, soil degradation of soils,~~ salinity, and alkalinity. To date, many farming practices were employed to increase cotton production, as a consequence, toxic impacts were implied on water and soil resources. ~~In addition On top of that,~~ a major part of the chemical fertilizers applied to the crops remain in the soil as insoluble inorganic compounds ~~which and~~ promotes soil toxicity ~~to the soil~~. Lately, efforts have been focused on minimizing the use ~~age~~ of chemical fertilizers in order to optimize cost of production ~~cost~~ and protect the environment against pollution without compromising the seed cotton yield. The use of biological stimulators in Indian agriculture has many economic and ecological advantages. The search for alternative solutions has ~~stimulated prompted~~ researchers to take a second look at the range of microorganisms which provide benefits to agricultural production by stimulating plant growth and producing inge a higher yield [1,2].

Plant growth promoting rhizobacteria (PGPR) are ~~the~~ bacteria living in the rhizosphere ~~region~~ which interacts with plant metabolism and improving advances their growth ~~directly or indirectly~~. ~~Nowadays,~~ PGPR inoculants are considered as a part of integrated nutrient management system to improve plant growth and development. The use of beneficial bacteria such as *Azotobacter*, *Azospirillum*, *Acetobacter*, *Pseudomonas*, *Methylophs*, *Bacillus*, ~~pP~~ *Phosphobacteria* etc. colonizes plant root and promotes growth ~~through by~~ nitrogen fixation, phosphorus and potassium solubilization. Seed priming by the microbial inoculants favours increased germination ~~by the activating on of~~ germination related enzymes, increasing metabolism that helps in the rapid growth of radicle and plumules. These PGPRs helps in improve better seed germination and colonizes plant roots ~~to~~ improvinge seedling vigour, and modifies root morphology to facilitate better acquisition of nutrients increasing which ultimately enhances the crop yield and quality ~~of crop~~.

Formatted: Strikethrough

Formatted: Strikethrough

Higher production and productivity of the crop is achieved through the use of good quality seeds and following proper management practices in any cultivar, variety or hybrid. Good quality seeds implies vigour, uniformity and structural soundness, in addition to besides its genetic and physical purity. Seed priming, identified as an effective seed invigoration method has become a common seed treatment to increase the rate and uniformity of germination emergence and crop establishments. Therefore, the present study was conducted formulated with thean objective to evaluate the different PGPR strains on cotton seed germination, seedling emergence and growth traits.

2. MATERIAL AND METHODS

2.1. Experimental site and soil characteristics

The field experiment was conducted at Eastern block farm, Tamil Nadu Agricultural University, Coimbatore, located in the Western Agro-climatic zone of Tamil Nadu (11° 02' N latitude, 76° 93' E longitude, and at an altitude of 428.5 m asletres above mean sea level). The experiment was planted laid out following in thea randomized block design with four replications. Experimental soil was sandy clay loam with pH 8.49 and organic matter content 0.47%. Newly released compact cotton seeds (CO 17 cultivarvariety) seeds were obtained collected from the Cotton Department of Cotton, Tamil Nadu Agricultural University, Coimbatore. Planting Sewing was done in rows by dibbing at row to row 90 cm apart and plant to plant 15 cm spacing by placing the seeds at each hill. Irrigation was given as per requiredment of crop. Standard plant protection schedule was followed to protect the crop from diseases and pests as per recommended package of practices of Tamil Nadu Agricultural University, Coimbatore.

2.2. Seed treatment details

Four PGPR strains included i.e., Azospirillum sp7, Phosphobacteria PS1, Potash releasing bacteria KRB9 and Pink pigmented facultative methylotrophs TNAU 1 were obtained collected from the Department of Agricultural Microbiology, Tamil Nadu Agricultural University, Coimbatore. The treatments include T1 - Control, T2 - Azospirillum sp7 (50 ml/acre of seeds), T3 - Pink Pigmented Facultative Methylotrophs (PPFM) TNAU 1 (50 ml/acre of seeds), T4 - Azospirillum sp7 + Phosphobacteria PS1 + Potash releasing bacteria KRB9 (50 ml/acre of seeds), T5 - PPFM TNAU1+ Phosphobacteria PS1 + Potash releasing bacteria KRB9 (50 ml/acre of seeds), T6 - PPFM TNAU1+ Phosphobacteria PS1 + Azospirillum sp7 + Potash releasing bacteria KRB9 (50 ml/acre of seeds). Azospirillum, PPFM, Phosphobacteria and Potash releasing bacteria are the commercial formulations of biofertilizers which contains Azospirillum brasilense sp7 strain, PPFM TNAU1 strain, Bacillus megaterium PS1 strain and Bacillus mucilaginosus KRB9 strain, respectively. Cotton seeds were bioprimered using 2% CMC (Carboxymethyl Cellulose) solution and then shade dried before plantingprior to sowing.

Formatted: Font: Italic

Formatted: Font: Not Italic

2.3. Data collection

The efficiency of different PGPR strains as bio-inoculants as an individual and interactive effect on seed germination efficiency, seedling emergence and growth traits in cotton were recorded.

2.3.1. Germination percentage - GP (%)

Final germination percentage was calculated by dividing the number of seeds germinated by the total number of seeds and expressed in % [3].

2.3.2. Mean germination time - MGT (day)

Mean germination time (MGT) was given by [4].

$$MGT = \sum \left(\frac{n_i \cdot t_i}{n_i} \right)$$

where n_i is the number of germinated seeds on germination days, t_i is the number of days during the germination period (between 0 and 10 days)

2.3.3. Germination rate index - GRI (Germination %¹/day¹)

The germination rate index was computed by the formula given by [5]

Formatted: Superscript

$$GRI = \frac{G1}{1} + \frac{G2}{2} + \frac{GX}{X}$$

where G1=Germination percentage at the first day after sowing, G2= Germination percentage at the second day after sowing.

2.3.4. Coefficient of velocity of germination - CVG

The coefficient of velocity of germination was calculated by the formula given by [6]

$$CVG = \frac{\sum ni}{\sum ni \cdot ti} * 100$$

where ni is the number of germinated seeds on day ti; ti = the number of days during the germination period (between 0 and 10 days)

2.3.5. Germination index - GI

The germination index (GI) was calculated based on the formula of [7]

$$GI = (10 * n1) + (9 * n2) + \dots + (1 * n10)$$

where n1 = number of seedlings emerging on first day after planting; n2 = number of seedlings emerging on second day; n10 = number of seedlings emerging on tenth day.

2.3.7. Seedling vigour index - SVI

The vigour index value was ~~obtained~~ computed by multiplying germination of seeds in percentage and total seedling length in centimeter and expressed in whole number as described by [8].

2.3.8. Root length and shoot length (cm)

Root and shoot length were measured at 10 ~~days~~ old seedlings and expressed in centimeters (cm)

2.3.9. Leaf Area (cm²)

Leaf samples collected from each replication were cleaned ~~well~~ and inserted into a leaf area meter (LICOR, Model LI 3000) and leaf area measured was expressed as cm² per plant.

2.3.10. Aerial and underground weight (g seedling⁻¹)

Fresh aerial and underground weight of 10 ~~days~~ old seedlings were taken and expressed in g seedlings⁻¹. ~~The s~~Seedlings used for growth measurement were placed in a paper cover and dried in shade for 24 ~~hrs~~ and then ~~they were placed kept~~ in a hot air oven at 65 °C for 48 ~~hrs~~. The dried seedlings were weighed to estimate the aerial and underground dry matter production and the mean values were expressed in mg seedlings⁻¹.

2.4. Statistical analysis

Data were analysed ~~by~~ using the software SPSS Statistics (version 16.0) and XLSTAT version 2019.2.1 (XLSTAT, 2019) and ~~a~~ comparison of means were done at 5 % significance level with Duncan's multiple range test. The principal component analysis (PCA) [9] was performed with seed germination and growth traits as influenced by different PGPR inoculation in cotton.

3. RESULTS AND DISCUSSION

3.1. Effect of PGPR strains on ~~G~~germination efficiency

Inoculation with dDifferent PGPR strains ~~inoculation~~ significantly influenced the germination efficiency of cotton seeds. Experimental results ~~in~~ from Table 1, revealed that seeds treated with PGPR strains showed significant improvement in germination percentage. Cotton seeds inoculated with strain of *PPFM* TNAU 1 (T3) showed maximum germination percentage followed by seeds inoculated with *Azospirillum* sp7 strain (T2) ~~with e.~~ 96.8 and 95.3 %, respectively. Likewise, cotton seeds treated with different PGPR strains positively influenced the mean germination time (~~day~~) (Fig. 1). Mean germination time determines how faster the seed ~~population~~ germinates in shorter period

the less amount of time [10]. The mean germination time was found to be lower in the strains treated with PPFM TNAU1 (T3) (5.6 days) followed by *Azospirillum* sp7 (T2) strain (5.9 days) treated seeds whereas seeds co-inoculated with both *Azospirillum* sp7 + Phosphobacteria PS1 + Potash bacteria KRB9 (T4) strains resulted in recorded higher mean germination value of 6.3 days. The seeds treated with individual strains of PPFM TNAU1 and *Azospirillum* sp7 strain had recorded about 16.2 and 14.4% increase, respectively, over the control. Results obtained coincide The present study is in corroboration with those obtained e findings of by Prathibha and Siddalingeshwara, [11]n and, Sirohi et al. [12] where that seed inoculation of *Bacillus* sp. and *Pseudomonas* significantly improved germination percentage and the rate of germination in sorghum and wheat seeds. Similar findings were reported by Nehra et al. [13], El-Sheekh et al. [14], Gowtham et al. [15] and Roman-Ponce et al. [16]




Fig.1. Effect of PGPR strains on mean germination time (MGT)—(days) on cotton seeds.

Inoculation of PGPR strains showed exhibited significant improvement in germination index (GI), germination rate index (GRI) and coefficient of velocity of germination (CVG) (Fig. 2. and Fig. 3.). Cotton seeds inoculated with PPFM TNAU 1 (T3) strain showed maximum germination index (354.3) followed by the treatment T2, *Azospirillum* sp7 strain (337.3). GI accentuates both germination percentage and speed. Higher GI values indicates a higher germination percentage and rate at which germination occurs [7]. Germination rate index was recorded higher in the treatment T3, PPFM TNAU1 strain followed by the treatment T2 *Azospirillum* sp7 strain treated seedplants i.e., 17 %/day¹ and 16.2 %/day¹, respectively. The GRI measures the percentage of seeds that germinate on each day of the germination period, with higher GRI values indicateing faster germination [5]. Similarly, the coefficient of velocity of germination also recorded higher in the (T3) PPFM TNAU 1 strain (17.8) inoculated seedplants. The CVG indicates the speed at which seeds germinate. It increases as the number of germinated seeds increases whileand the time needed for germination declines [6]. The inoculation of PPFM TNAU1 and *Azospirillum* sp7 strain had recorded about 27.1 and 21.%, 23.5 and 18.3%, 10.7 and 4.8% increase ment in germination index, germination rate index and coefficient of velocity of germination over the uninoculated control, respectively. The increased germination efficiency is due to the synthesis of gibbereline, the hormone, gibberellin which stimulates the activity of alpha-amylase, the germination-specific enzyme and, alpha-amylase, protease and nuclease which is mainly involved in starch breakdown and assimilation [17], and mitochondrial enzyme activities [18]. The present study coincides is in corroboration with results of Xiao et al. [19], and; Hossain et al. [20] in which PGPR inoculation significantly improved germination percentage and germination rate index over the with reference to the control in rice. Similar findings were reported by Makhaye et al. [21], and Hamidi et al. [22], where that seed inoculation of PGPRs significantly improved all the germination traits such as germination percentage (GP), germination index (GI), germination rate index (GRI), mean germination time (MGT) and coefficient of velocity of germination (CVG) in maize. However, the increased germination traits were also observed in combined inoculants of PGPR strains. The results of the present study are similar to the findings of Anitha- [23] and Meena et al. [24].

Formatted: Font: Italic

Formatted: Font: Italic


Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Superscript

Formatted: Superscript



T1 – Control (Uninoculated treatment)
T2 – *Azospirillum* sp7
T3 – PPFM TNAU1
T4 – *Azospirillum* sp7 + *Phosphobacteria* PS1 + KRB9
T5 – PPFM TNAU1 + *Phosphobacteria* PS1 + KRB9
T6 – PPFM TNAU1 + *Phosphobacteria* PS1 + *Azospirillum* sp7 + KRB9


Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Fig. 2. Effect of PGPR strains on germination rate index GRI (Germination %/day¹) and coefficient of velocity of germination of cotton seeds.

Formatted: Superscript



T1 – Control (Uninoculated treatment)
T2 – *Azospirillum* sp7
T3 – PPFM TNAU1
T4 – *Azospirillum* sp7 + *Phosphobacteria* PS1 + KRB9
T5 – PPFM TNAU1 + *Phosphobacteria* PS1 + KRB9
T6 – PPFM TNAU1 + *Phosphobacteria* PS1 + *Azospirillum* sp7 + KRB9

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Fig. 3. Effect of PGPR strains on germination index (GI) of cotton seeds.

3.2. Effect of PGPR on seedling emergence traits

Cotton seeds treated with different PGPR strains exhibited significant improvement in the seedling emergence traits such as shoot and root length (Fig. 4), and seedling vigour index (Table 1). The experimental results obtained indicated that shoot and root length of PGPR inoculated seedlings were significantly higher compared to the uninoculated treated control (Fig. 4). The treatment of *Azospirillum* sp7 strain (T2) inoculated seeds resulted in a maximum shoot length and root length (12.0 and 6.0 cm, respectively) followed by the treatment 3 (PPFM TNAU1) (T3) with 11.9 and 5.8 cm, respectively). The combined inoculation of PGPR strains (T4, T5 and T6) resulted in higher shoot length and root length compared to the non-inoculated control. The percent increment in shoot and root length was 24.4 and 42.8% in the *Azospirillum* sp7 (T2) strain inoculated seeds plants over the uninoculated control. PGPR inoculation stimulates the production of the phytohormone IAA (Indole-3 Acetic Acid (IAA) and acquiring more nutrients such as nitrogen and phosphorus, which in turn increases root length and shoot length [25]. In addition of the increasing in shoot and root length, IAA is mainly due to the primary synthesis of plant hormone, IAA which is responsible for cell elongation in *Azospirillum* treated plants. The results present findings are similar to those obtained by in accordance with Efthimiadou et al. [26], Taha et al. [27] and El-Gamal et al. [28]. Similar findings were reported by Dhale et al. [29], Zamioudis et al. [30], and Pindi et al. [31], where plant growth promoting bacterial (PGPB) inoculation stimulates the production of plant growth hormones that favours root growth and alters root morphology. The plants inoculated with the treatment T3, PPFM TNAU1 strain documented resulted in higher seedling vigour index (1714.4) as it has higher germination percentage, followed by the treatment T2, *Azospirillum* sp7 strain (1709.6) which as it showed higher values in shoot and root length compared to other treatments and control. The findings Results obtained by by Noumavo et al. [32] are similar in accordance with the results obtained in the present study. This enhanced seedling vigour index could

Formatted: Font: Italic

be associated with greater production and metabolism of hormones, auxin and cytokinin which primarily promotes the cell elongation and cell division induced by PGPR inoculation [33,34].




Fig. 4. Effect of PGPR strains on shoot and root length (cm) on cotton seed.

3.3. Effect of PGPR strains on growth traits

Experimental results obtained showed-revealed that cotton seeds plants-treated with PGPR strains had showed-substantial-improved-ment-in leaf area, aerial fresh and dry weight and underground fresh and dry weight (Table-1). The treatment T3, PPFM TNAU1 and the treatment T2, Azospirillum sp7 strains inoculated seedsplants had-exhibited a higher foliage size compared to the combined inoculation of strains and control plants. PPFM TNAU1 strain (T3) inoculated seedsplants recorded maximum leaf area (17.4 cm²) followed by the treatment T2, Azospirillum sp7 strain inoculated seedsplants (16.1 cm²). The rate of cell division is higher in the PPFM TNAU1 inoculated seedsplants due to by-synthesis of cytokinin, the phytohormone, cytokinin-which that contributes to the wider leaf surface area. The inoculation of PPFM TNAU1 and Azospirillum sp7 strain resulted in an increase-ment of 57.3 and 45.6%, respectively, over the control. Our results are similar to those obtained by in-corroboration-with Wang et al. [35], and Namwongsa et al. [36].

Maximum aerial fresh and dry weight were recorded in the treatment T3, PPFM TNAU1 strain (3.25 g seedling⁻¹ and 72.8 mg seedling⁻¹), respectively. Likewise, the maximum underground fresh and dry weight recorded was to-be-higher in the treatment T2, Azospirillum sp7 strain inoculated plants (0.40 g seedling⁻¹ and 6.85 mg seedling⁻¹). This increase of aerial fresh and dry weight in the treatment T3, PPFM TNAU1 strain, is due to the higher leaf biomass. Azospirillum sp7 (T2) strain inoculated treatment is found to have maximum underground fresh and dry weight due to increased root length by the synthesis of phytohormones that acts on cell division and cell elongation. The maximum leaf area obtained in the treatment T3, PPFM TNAU1 (T3) inoculated seedlings, contributed to the higher total dry matter production (79.6 mg seedling⁻¹) compared to the (T2) Azospirillum sp7 (77.3 mg seedling⁻¹) strain treated seedlings. The present findings are similar to those obtained by in-line-with Wang et al. [35] where that-increased transcriptional levels of auxin biosynthesis genes enhanced the seedling growth thus improved the-biomass. Our results of increased seedling fresh and dry weight were similar in-harmony-with to those reported by-Etesami and Alikhani. [37], Turan et al. [38], Asari et al. [39], and Egamberdieva et al. [25]. This considerable improvement in seedling growth parameters suggests that selective inoculation of PGPRs could be considered as an effective alternative biofertilizer for promoting cotton seed germination, biomass, and yield.

Table 1. Effect of PGPR strains on germination percentage, seedling emergence and seedling growth traits

Treatments	Germination Percentage %	Seedling vigour index	Leaf Area (cm ²)	Fresh weight (g plant ⁻¹)		Dry weight (mg plant ⁻¹)	
				Aerial weight	Underground weight	Aerial weight	Underground weight
T1 – Control	83.3	1149.5	11.0	2.33	0.22	50.3	5.78
T2 – Azospirillum	95.3	1709.6	16.1	3.19	0.40	70.5	6.85
T3 – PPFM	96.8	1714.4	17.4	3.25	0.38	72.8	6.83
T4 – Azospirillum + PSB + KRB	88.5	1459.4	11.7	2.75	0.30	55.5	6.60

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Space After: 0 pt, Line spacing: single

Formatted: Left, Space After: 0 pt, Line spacing: single

Formatted Table

T5 – PPFM + PSB + KRB	88.8	1453.5	11.6	2.65	0.35	55.0	6.63
T6 – PPFM + PSB+ Azospirillum + KRB	91.3	1453.6	11.3	2.61	0.32	54.0	6.60
Mean	90.7	1490.0	13.2	2.8	0.30	59.7	6.50
SEd	1.06**	34.74**	0.63*	8.28**	5.44	1.74**	0.13**
CD (P=0.05)	2.26	74.04	1.33	0.18	0.12	3.72	0.29


Values are mean of replicate. ~~Vs~~ values followed by the same letter in each column are not significantly different from each other as determined by DMRT (P = 0.05).

Formatted: Left, Space After: 0 pt, Line spacing: single

3.4. Principal component analysis (PCA) of seed germination and growth traits in cotton influenced by PGPR strains

The PCA analysis was done using the data of seed germination and growth-related variables such as germination percentage (GP), mean germination time (MGT), germination index (GI), germination rate index (GRI), seedling vigour index (SVI), leaf area (LA), aerial and underground fresh (AFW and UFW) and dry weight (ADW and UDW) obtained from six treatments with four replications (Fig. 5.). The PCA ~~showed revealed~~ that the variables were correlated with the principal component value of 96.60 % (PC1 – 87.57 % and PC2 - 9.03 %). The variables ~~that were~~ obtained from seed treatment of *Azospirillum* sp7 (T2) and PPFM TNAU1 (T3) strains were located in the right side of the scoring plot, showing positive correlation between the components. ~~However, On the contrary,~~ variables of the control treatment, ~~control~~ (T1) were located in the bottom left quarter of the scoring plot in PCs (Fig. 5A). ~~In loading plot, The~~ plant growth variables were ~~influenced~~ positively ~~influenced~~ by PGPR strain inoculation, ~~positioned~~ at the right side of the plot (Fig. 5B). PCA results confirmed that PGPR strain inoculation positively influenced the germination traits and growth characters, while mean germination time was least correlated with the inoculation of different PGPR strains. Results from PCA recorded that the treatment *Azospirillum* sp7 (T2) strain, ~~which is~~ located in the right side of first quadrant is the best treatment followed by the treatment PPFM TNAU1 (T3), ~~which is~~ located in the right end of the second quadrant, which detailed that PGPR inoculation improved germination efficiency and seedling growth traits.

A



B




Fig. 5. Principal component analysis (PCA) of cotton as influenced by different PGPR strains (A) Scoring plot of treatments and (B) Loading plot of variables. Variables are GP, Germination percentage; MGT, mean germination time; GI, Germination index; GRI, Germination rate index; SVI, Seedling vigour index; SL, Shoot length; RL, Root length; LA, Leaf area; AFW, Aerial fresh

weight; UFW, Underground fresh weight; ADW, Aerial dry weight and UDW, Underground dry weight.

4. CONCLUSION

Single and combined inoculation of PGPR strain in cotton seeds considerably improved the germination efficiency and plant growth during the early growth stagephase. However, seeds inoculated with individual strain of *Azospirillum* sp7 and *PPFM* TNAU1 outperforms the combined inoculation of PGPR strains. These results indicate conclude that the PPGR strain inoculation has a positive stimulant effect on germination efficiency and plant growth promotion in the cotton crop. It is concluded Thus, the that the investigated beneficial microbes can be employed to increase the physiological potential of cotton to obtain achieve higher yield.

REFERENCES

1. Lehr P. Biopesticides: The Global Market. Report code CHM029B, BCC Research, Wellesley, Massachusetts; 2010.
2. Jangral J, Hamit L. Impact of fertilizers on the environment sustainability development and agriculture. GE-International Journal of Management Research. 2014;2(2):160–66.
3. Ranal MA, Santana DGD. How and why to measure the germination process?. Brazilian Journal of Botany. 2006;29:1-11.
4. Mauromicale G, Licandro P. Salinity and temperature effects on germination, emergence and seedling growth of globe artichoke. Agronomie. 2002;22(5):443-450.
5. Esechie HA. Interaction of Salinity and Temperature on the Germination of Sorghum. Journal of Agronomy and Crop Science. 1994;72:194-199.
6. Jones K, Sanders D. The influence of soaking pepper seed in water or potassium salt solutions on germination at three temperatures. Journal of Seed Technology. 1987;11:97–102.
7. Bench AR, Fenner M, Edwards P. Changes in germinability, ABA content and ABA embryonic sensitivity in developing seeds of *Sorghum bicolor* (L.) Moench induced by water stress during grain filling. New Phytologist. 1991;118:339–347.
8. Abdul-Baki AA, Anderson JD. Vigor determination in soybean seed by multiple criteria 1. Crop science. 1973;13(6), 630-633.
9. Wold S, Esbensen K, Geladi P. Principal component analysis. Chemometrics and intelligent laboratory systems. 1987;2(1-3):37-52.
10. Orchard T. Estimating the parameters of plant seedling emergence. Seed Science and Technology. 1977;5:61–69.
11. Prathibha KS, Siddalingeshwara KG. Effect of plant growth promoting *Bacillus subtilis* and *Pseudomonas fluorescence* as Rhizobacteria on seed quality of sorghum. International Journal of Microbiology and Applied Science. 2013;2:11-18.
12. Sirohi G, Upadhyay A, Srivastava PS, Srivastava S. PGPR mediated Zinc biofertilization of soil and its impact on growth and productivity of wheat. Journal of Soil Science and Plant Nutrition. 2015;15:202-216.
13. Nehra V, Saharan BS, Choudhary M. Evaluation of *Brevibacillus brevis* as a potential plant growth promoting rhizobacteria for cotton (*Gossypium hirsutum*) crop. Springerplus. 2016;5:948-959.
14. El-Sheekh MM, Ismail MM, Hamouda MM. Influence of some brown seaweed extracts on germination and cytological responses of *Trigonella foenum-graecum* L. BioTechnology: An Indian Journal. 2016;12:1-12
15. Gowtham HG, Murali M, Singh SB, Lakshmeesha TR, Narasimha Murthy K, Amruthesh KN. Plant growth promoting rhizobacteria- *Bacillus amyloliquefaciens* improves plant growth and induces resistance in chilli against anthracnose disease. Biological Control. 2018;126:209–217.
16. Roman-Ponce B, Reza-Vazquez DM, Gutierrez-Paredes S, De Haro-Cruz MDJ, Maldonado Hernandez J, Bahena-Osorio Y, Estrada-De Los Santos P, Wang ET, Vasquez Murrieta MS. Plant growth promoting traits in rhizobacteria of heavy metal-

- resistant plants and their effects on *Brassica nigra* seed germination. *Pedosphere*. 2017;27:511-526.
17. Gholami A, Shahsavani S, Nezarat S. The Effect of Plant Growth Promoting Rhizobacteria (PGPR) on Germination, Seedling Growth and Yield of Maize. *World Academy of Science, Engineering and Technology*. 2009;49:19-24.
 18. Fatima Z, Saleemi M, Zia M, Sultan T, Aslam M, Rehman R, FayyazChaudhary M. Antifungal Activity of Plant Growth-Promoting Rhizobacteria Isolates against *Rhizoctonia solani* in Wheat. *African Journal of Biotechnology*. 2009;8(2):219-225
 19. Xiao AW, Li Z, Li WC, Ye ZH. The effect of plant growth promoting rhizobacteria (PGPR) on arsenic accumulation and the growth of rice plants (*Oryza sativa* L.). *Chemosphere*. 2020;242.
 20. Hossain MM, Das KC, Yesmin S, Shahriar S. Effect of plant growth promoting Rhizobacteria (PGPR) in seed germination and root-shoot development of chickpea (*Cicer arietinum* L.) under different salinity conditions. *Research in Agriculture Livestock and Fisheries*. 2016;3:105-113.
 21. Makhaye G, Amoo SO, Gerrano AS, Aremu AO, Tesfay S. Effect of biostimulants on germination of okra (*Abelmoschus esculentus* L.) genotypes. *Combined Congress. University of the Free State, Bloemfontein, South Africa*; 2020.
 22. Hamidi A, Asgharzadeh A, Ahmadi A, Akbari Vala S, Choukan R. Effect of Plant Growth Promoting Bacteria (PGPB) and Mycorrhizae Fungi on three Maize (*Zea mays* L.) hybrids some seed germination and seedling vigour trait. *Journal of Agricultural Science and Sustainable Production*. 2021;31(3):149-167.
 23. Anitha KG. Enhancing seed germination of mono and dicotyledons through IAA production of *PPFM*. *Trends in Soil Science. Journal of Plant Nutrition*. 2010;1:14-18.
 24. Meena KK, Kumar M, Kalyuzhnaya MG, Yandigeri MS, Singh DP, Saxena AK, Arora DK. Epiphytic pink-pigmented methylotrophic bacteria enhance germination and seedling growth of wheat (*Triticum aestivum*) by producing phytohormone. *Antonie van Leeuwenhoek*. 2012;101(4):777-786.
 25. Egamberdieva D, Wirth S, Jabborova D, Rasanen L, Liao H. Coordination between *Bradyrhizobium* and *Pseudomonas* alleviates salt stress in soybean through altering root system architecture. *Journal of Plant Interactions*. 2017;12:100–107.
 26. Efthimiadou A, Katsenios N, Chanioti S, Giannoglou M, Djordjevic N, Katsaros N. Effect of foliar and soil application of plant growth promoting bacteria on growth, physiology, yield and seed quality of maize under Mediterranean conditions. *Scientific Reports*. 2020;10:21060.
 27. Taha RS, Alharby HF, Bamagoos AA, Medani RA, Rady MM. Elevating tolerance of drought stress in *Ocimum basilicum* using pollen grains extract; a natural biostimulant by regulation of plant performance and antioxidant defense system. *South African Journal of Botany*. 2020;128:42-53.
 28. El-Gamal AD, Ismail MA, Amin MA, Sayed AM. Comparative studies between seaweeds and commercial algae in alleviation of harmful effects of drought stress of faba bean (*Vicia faba* L.). *Plants*. 2020.
 29. Dhale DA, Chatte SN, Jadhav VT. Effect of Bioinoculents on Growth, Yield and Fibre Quality of Cotton under Irrigation. *Research Journal of Agriculture and Biological Sciences*. 2010;6(4): 542-547.
 30. Zamioudis C, Mastranesti P, Dhonukshe P, Blilou I, Pieterse CMJ. Unraveling root developmental programs initiated by beneficial *Pseudomonas spp.* bacteria. *Plant Physiology*. 2013;162:304–318.
 31. Pindi PK, Sultana T, Vootla PK. Plant growth regulation of Bt-cotton through *Bacillus* species. *Biotechnology*. 2014;4:305–315
 32. Noumavo PA, Kochoni E, Didagbe YO, Adjanohoun A, Allagbe M, Sikirou R, Baba-Moussa L. Effect of different plant growth promoting rhizobacteria on maize seed germination and seedling development. *American Journal of Plant Sciences*. 2013;4(5):1013.

33. Bharathi R, Vivekananthan R, Harish S, Ramanathan A, Samiyappan R. Rhizobacteria-Based Bio-Formulations for the Management of Fruit Rot Infection in Chillies. *Crop Protection*. 2004;23(6):835- 843.
34. Anzala FJ. Controle de la Vitesse de Germination chez le Maïs (*Zea mays*): Etude de la Voie de Biosynthèse des Acides Aminés issus de l'Aspartate et Recherche de QTLs. Thèse de Doctorat, Université de Angers, Angers; 2006.
35. Wang J, Zhang Y, Li Y, Wang X, Nan W, Hu Y. Endophytic microbes *Bacillus* sp. LZR216-regulated root development is dependent on polar auxin transport in *Arabidopsis* seedlings. *Plant Cell Reports*. 2015;34:1075–1087.
36. Namwongsa J, Jogloy S, Vorasoot N, Boonlue S, Riddech N, Mongkolthanaruk W. Endophytic bacteria improve root traits, biomass and yield of *Helianthus tuberosus* L. under normal and deficit water conditions. *Journal of Microbiology and Biotechnology*. 2019;29:1777–1789.
37. Etesami H, Alikhani HA. Co-inoculation with endophytic and rhizosphere bacteria allows reduced application rates of N-fertilizer for rice plant. *Rhizosphere*. 2016;2:5–12.
38. Turan M, Ekinci M, Yildirim E, Gunes A, Karagoz K, Kotan R. Plant growth-promoting rhizobacteria improved growth, nutrient, and hormone content of cabbage (*Brassica oleracea*) seedlings. *Turkish Journal of Agriculture and Forestry*. 2014;38:327–333.
39. Asari S, Tarkowska D, Rolcik J, Novak O, Palmero DV, Bejai S. Analysis of plant growth-promoting properties of *Bacillus amyloliquefaciens* UCMB5113 using *Arabidopsis thaliana* as host plant. *Planta*. 2016; 245:15–30.

Formatted: Spanish (Mexico)