

Short Research Article

Growth behaviour of *Bt* (*Bacillus thuringiensis*) cotton as influenced by mepiquat chloride under varying nitrogen levels

ABSTRACT

Cotton is indeterminate and long duration crop. For synchronising in growth and to prevent heavy shedding of bolls different growth regulators used in cotton field. The present field experiment was conducted at National Seed Production Area of CCS Haryana Agricultural University, Hisar, during *kharif* season 2018. The experiment comprised of three levels of nitrogen (100%, 125% and 150% RDF) and three spray of mepiquat chloride (control, single spray at 60 DAS and two spray at 60 and 75 DAS) was conducted in factorial randomized complete block design replicated thrice. Based on the research investigation, it was found that different phenological stages were delayed by 3 to 4 days with application of 150% RDF, while twice spray of mepiquat chloride resulted in 6 to 7 days earlier completion of these stages. Ginning out turn was statistically at par with different levels of nitrogen, while 100% RDF has shown maximum span length with lowest micronaire value. With single mepiquat chloride spray at 60 DAS ginning percentages decreases by 6.80% over control, it because of reducing growth stages period. While span length reduces with two sprays of mepiquat chloride at 60 and 75 DAS by 0.62% over the control and single spray of mepiquat chloride at 60 DAS. However finest fibre ($4.23 \times 10^{-6} \text{ g/inch}^{-1}$ micronaire value) obtained with single spray of mepiquat chloride. The result of present study reveals that, 100% RDF in *Bt* cotton along with single spray of mepiquat chloride at 60 DAS gives finest fibre with more span length by completing all phenological stages earlier than others.

Keywords: *Bt* cotton, mepiquat chloride, ginning percentages, micronaire value, span length.

1. INTRODUCTION

Cotton is one of the important cash crop in world. Cotton crop attains excessive vegetative growth mainly due to high soil fertility; coincidence of vegetative growth period with rainy season coupled with high humidity. Thick crop canopy prevents penetration of light besides shading of bolls and utilization of plant energy on the formation of vegetative stature which results in shedding of flower buds, flowers and immature bolls. Excessive vegetative growth often occurs at the expense of reproductive growth and a large fraction of squares and small bolls on the lower sympods either shed or open poorly resulting in low yield.

Cotton is used as raw material in textile industry and this industry contributes about 14% to the industrial production and 4% of the GDP [1]. Cotton is used as major raw material in this sector. Around 35 million people are directly dependent on this sector for their employment. The hand loom sector consumes around 12.50 % of raw cotton and power loom sector around 62.70 %. Mills and Hosiery sector consume nearly 3.40 % and 21.40 % of total raw cotton respectively. It is also the source of edible oil, cotton seed oil cake, linters, and huge biomass as dried cotton stalk used as fuel. It has the important character of high lint production and long staple length. Lint is the most important

Comment [U1]: Sentences without references (citations).

economical product, provides a source of quality fiber for textile industry and cotton seeds which is the primary byproduct of lint production, are an important source of oil for human consumption and also used as livestock feed. Cotton plants must have a harmonic balance between vegetative and reproductive growth for adequate photosynthates supply for healthy boll development leading to better productivity. The use of mepiquat chloride increases the N uptake resulting into higher seed cotton yield [2]. Quality parameters viz., ginning percent, lint index, fiber length, bundle strength and fiber fineness were not affected significantly by N management practices [3]. A field experiment on effect of different nitrogen levels (50, 100, 150 kg N/ha) on cotton shows that application of 150 kg N/ha gave significantly higher ginning out turn [4]. Under extreme cases, rotting of lower bolls also occurs. The loss of reproductive structures alters the physiological growth and development of the plant by redirecting assimilates which normally are incorporated into these abscised organs to other plant parts. Most source sink research has focused on leaf boll relationships (harmonic balance) with little study of vegetative storage reserves.

Source-sink balance can be altered by using plant growth regulator such as mepiquat chloride (MC) [5]. The best way to prevent excessive vegetative growth is to manage for early and high fruit set. If early fruit are not set, the crop may compensate by setting fruit at higher nodes and at outer fruiting positions leading to delayed crop maturity. Growth retardants like mepiquat chloride (MC) are known to reduce inter nodal length, thereby, reducing plant height and stimulating the translocation of photosynthesis towards reproductive sinks (developing cotton bolls), all of which result in higher yields. Plant growth regulators (PGRs) may enhance yield by increasing the retention of photosynthates into developing bolls. PGRs have been widely used in developed nations for increasing cotton production by adjusting plant growth and to improve lint yield and fiber quality. Mepiquat chloride (MC) is used in cotton production across the globe to control plant growth and maximize yield and quality. Application of MC at squaring stage or at both squaring and flowering stages significantly improved cotton quality parameters like fiber length (1.7%) and fiber strength (2.8%) without significant loss of yields [6]. Fiber strength increased by 1.5 to 2.8 g/tex with MC [7]. Application of MC increased fiber strength by 3.8% [8]. MC did not significantly affect fiber qualities [9,10] but micronaire increased with MC application [11]. Cycocel and Alar did not affect cotton fiber quality [12,13].

The main objective of this study was to find out the effect of nitrogen and mepiquat chloride on duration and fiber quality parameter of *Bt* cotton under semi-arid condition.

2. MATERIAL AND METHODS AND MATERIALS

2.1 Experimental Site

The study was conducted at National Seed Production Area of Choudhary Charan Singh Haryana Agricultural University, Hisar, during *kharif* season 2018. Hisar is situated in the sub-tropics at longitude (75°46'E, latitude 29°10'N) and altitude of 215.2 m above mean sea level in Haryana India state of India. Hisar has semi-arid climate with very hot summers (temperature rises up to 45°C or more) and extremely cool winter (temperature falls up to 1-2°C or less than this).

The initial soil characteristics of experimental sites like pH, soil texture, electrical conductivity, organic carbon, available NPK are presented in table 1.

Formatted: Highlight

Comment [U2]: This is not clear.

Comment [U3]: Please provide validated datas, that is to say, in agree with scientific literature.

Formatted: Highlight

Formatted: Highlight

Table 1. Initial physico-chemical analysis of experimental field soil (2018)

Formatted: Space After: 0 pt

Soil property	values	Method of determination
Sand (%)	73.8	International pipette method [14]
Silt (%)	15.9	
Clay (%)	10.3	
pH	7.9	Glass electrode pH meter [15]
Electrical conductivity (ds/m)	0.23	Solubridge conductivity meter 1:2 soil-water suspension [16]
Organic carbon (%)	0.44	Walkley and Black rapid titration method [15]
Available nitrogen (Kg/ha)	160	Alkaline permanganate method [17]
Available phosphorus (Kg/ha)	16	Olsen method [18]
Available potassium (Kg/ha)	280	Flame photometer method [16]

2.2 Experimental Details

The experiment was carried out during the 2018 kharif season by using *Bt* cotton genotype RCH 650 at CCS Haryana Agricultural University, Hisar. The experiment comprised of three levels of nitrogen (100%, 125% and 150% RDF) and three spray of mepiquat chloride (control, single spray @ 20_g a.i./ha at 60 DAS and two spray @ 20_g a.i./ha at 60 and 75 DAS) was conducted in Factorial Randomized Complete Block Design replicated thrice. Recommended dose of fertilizer 100% and 125%, 150% was applied as one third of N and full dose of P, K was applied at the time of seed bed preparation. Remaining dose of N was applied after 1st irrigation and at flowering stage. Recommended dose of fertilizer for RCH 650 are 175 kg N, 60 kg P₂O₅ and 60 kg K₂O ^{1per}hectare.

Formatted: Not Superscript/ Subscript

Formatted: Superscript

2.3 Phenological Stages

Phenological stages like days to squaring, 50% flowering, 50% boll opening and maturity was the total number of days noted after the sowing in which 50% of plants initiate squaring, flowering, boll opening and shows maturity.

2.4 Quality Parameters

2.4.1 Ginning out turn (GOT), %

500 g sample of seed cotton was taken from each plot and then ginned to get lint and cotton seed.

Formatted: Highlight

It was calculated by using following formula.

Comment [U4]: Please reviewed this space between lines.

Formatted: Highlight

$$\text{GOT (\%)} = \frac{\text{Weight of lint}}{\text{Weight of seed cotton}} \times 100$$

2.4.2 Micronaire value

It measures the fiber weight in 10^{-6} g/inch length of fiber. Fineness denotes the size of cross-sectional diameter of the fibre. A sample of 100_g lint was taken and measure micronaire value by using Precitronic Digital Mic Tester at CICR, Sirsa.

2.5 Statistical Analysis

Data used in the study are the mean values of the replicated observations. For the statistical analysis of all the research field data, online computer programme OPSTAT (<http://hau.ernet.in/sheoranop/>) was used.

3. RESULTS AND DISCUSSION

3.1 Phenological Stages

There was significant difference on phenological ~~stages~~growth—observed with different levels of nitrogen (Table 2). Higher dose of nitrogen *i.e* 150% RDF were significantly delayed the duration of each phenological stages compared to 100% RDF. It might be due to more vegetative growth which delays the maturity *i.e.*, it has taken more number of days to reach boll development stage at higher levels of nitrogen application. Similar results were reported by Dong *et al.* [19] and Munir *et al.* [20].

Earliness in phenological stages was significantly enhanced by MC application (Table 2). Increased earliness may be related to MC effect on biomass partitioning (inhibiting growth of branches and stems, expanding leaves, and extending stem internodes and petioles), which led to the development of a more compact canopy structure, this provides an improved microclimate, especially better light conditions, that results in earlier maturity. Similar findings were recorded by Oosterhuis *et al.*, [21].

Table 2. Effect of different nitrogen levels and mepiquat chloride dose on phenological stages of Bt cotton hybrid

Treatments	Days to squaring	Days to 50% flowering	Days to 50% boll opening	Days to maturity
Nitrogen levels				
N ₁ (100% RDF)	50	81	108	174
N ₂ (125% RDF)	52	80	109	177
N ₃ (150% RDF)	53	80	111	179
CD	1.01	0.91	1.01	1.01
Mepiquat chloride dose				
G ₁ (Control)	51	84	112	182
G ₂ (MC@ 20g a.i./ha at 60 DAS)	52	80	109	176
G ₃ (MC@ 20g a.i./ha at 60 and 75 DAS)	51	77	106	171
CD	1.01	0.90	1.01	1.01

Formatted: Highlight

Comment [U5]: Please review space between line and paragraph.

Formatted: Justified

Comment [U6]: Please it is necessary the complete name in all legends, for example Bt (*Bacillus thuringiensis*) cotton hybrid.

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)

Formatted: English (U.S.)




Fig. 1. Effect of different nitrogen levels and mepiquat chloride dose on phenological stages of *Bt* cotton hybrid

Formatted: Justified, Indent: Left: 0 cm, First line: 0 cm

3.2 Quality Parameters

3.2.1 Ginning out turn (%)

Ginning out turn was not affected with different levels of nitrogen. Ginning out turn was **minimally or not affected with increasing application rate of N from 100% to 150% RDF** (Table 3) [22]. But mepiquat chloride significantly effects the ginning out turn, it was decreases with mepiquat chloride application. Significantly higher ginning out turn was recorded in control compared to other two treatments of mepiquat chloride. While ginning out turn was at par with twice and single application of mepiquat chloride. It might be due to high seed cotton yield with mepiquat chloride [11].

Formatted: Highlight

Comment [U7]: This is not clear.

3.2.2 Span Length (mm)

Span length significantly differ with different levels of nitrogen and mepiquat chloride (Table 3). 100% RDF shows significantly higher span length followed by 150% and 125% RDF. Span length was significantly higher in control and single application of mepiquat chloride as compared to twice spray. Span length was reduced with higher levels of nitrogen but it too small to affect the quality of fiber. Span length not affected with mepiquat chloride, but twice of mepiquat chloride application decreases span length [11].

3.2.3 Micronaire value ($10^{-6} \text{g/inch}^{-1}$)

Micronaire value was significantly vary with different levels of nitrogen and mepiquat chloride. Significantly higher value of micronaire recorded with 125% RDF which was close to 150% RDF followed by 100% RDF (Table 3). ~~TheThis showed that~~ excess application of N can reduce fiber quality. Nitrogen rate had no effect on fiber uniformity [22]. Excess application of N than the required

for optimum crop performance can reduce fiber quality. With mepiquat chloride, significantly higher value of micronaire was recorded in control followed by twice spray of mepiquat chloride and single spray. Twice application of mepiquat chloride gave almost closer micronaire value to control. Very low effect of mepiquat chloride on micronaire value was observed.

Comment [U8]: It is relevant you place the ideal percentage (%)

Table 3. Effect of different nitrogen levels and mepiquat chloride dose on quality parameters of *Bt* cotton hybrid

Formatted: Justified

Treatments	Ginning out turn (%)	Span length (mm)	Micronaire value (10 ⁻⁶ g/inch)
Nitrogen levels			
N ₁ (100% RDF)	34	28.84	4.3
N ₂ (125% RDF)	34.78	28.55	4.44
N ₃ (150% RDF)	34.21	28.64	4.37
CD	NS	0.08	0.05
Mepiquat chloride dose			
G ₁ (Control)	36.16	28.74	4.47
G ₂ (MC@ 20g a.i./ha at 60 DAS)	33.7	28.74	4.23
G ₃ (MC@ 20g a.i./ha at 60 and 75 DAS)	33.13	28.56	4.41
CD	1.87	0.08	0.05




Fig. 2. Effect of different nitrogen levels and mepiquat chloride dose on quality parameters of *Bt* cotton hybrid

Formatted: Justified, Indent: Left: 0 cm, First line: 0 cm

4. CONCLUSION

Phenological stages of *Bt* cotton like days to squaring, boll formation and maturity extended by 2 to 5 days with increasing nitrogen dose from 175 kg per ha⁻¹ (100% RDF) to 218.75 kg per ha⁻¹ (125%

RDF), 262.5 kg per ha⁻¹ (150% RDF). However, flowering occurs 4 and 7 days earlier with single and two spray of mepiquat chloride respectively than no spray. Also, the boll opening and maturity was 3 to 11 days earlier with spray of mepiquat chloride than no spray. Finest quality fibre was recorded with 125% RDF which was 0.69% and 0.70% more fine than 150% RDF and 100% RDF respectively. Maximum lint yield was gained (36.16% ginning percent) by without mepiquat chloride spray, while finest fibre was recorded with single spray of mepiquat chloride at 60 DAS with 4.26 x 10⁻⁶ g/inch⁻¹ micronaire value. The results reveal that, for getting fine quality fibre single spray of mepiquat chloride at 60 DAS with 100% RDF is recommended in *Bt* cotton growing areas in semi-arid region.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

Formatted: Indent: Left: 0 cm, First line: 0 cm

REFERENCES

1. Anonymous. The Indian express, Tuesday, May 01, 2018.
2. Shekar K, Venkataramana M, Kumari SR. Response of hybrid cotton to chloromepiquat chloride and detopping under high density planting. Journal of Cotton Research Development. 2015;29(2): 84-86.
3. Das A, Prasad M, Guatam RC, Shivay YS. Productivity of cotton as influenced by organic and inorganic sources of nitrogen. Indian Journal of Agricultural Sciences. 2006;76(6): 354-357.
4. Kumbhar AM, Buriro UA, Junejo FC, Jamro GH. Impact of Different Nitrogen Levels on Cotton (*Gossypium hirsutum*L.) Growth, Yield and N-Uptake Planted In Legume Rotation. Pakistan Journal. 2008;40(2): 767-778.
5. Gwathmey CO, Clement JD. Alteration of cotton source sink relations with plant population density and mepiquat chloride. Field Crop Research. 2010;116(2): 101-107.
6. Ren X, Zhang L, Dua M, Evers JB, Werf W, Tiana X, Li Z et al. Managing mepiquat chloride and plant density for optimal yield and quality of cotton. Field Crop Research. 2013;149(2): 1-10.
7. Livingston SD, Anderson DJ, Wilde LB, Hickey JA. Use of foliar applications of Pix, PRG IV, and PCHA in lowrate multiple applications for cotton improvement under irrigated and dryland conditions. In Proc. Beltwide Cotton Conf., Nashville. 1992;1055-1056.
8. Boman RK, Westerman RL. Nitrogen and mepiquat chloride effects on the production of nonrank, irrigated, short-season cotton. Journal of Production Agriculture. 1994;7: 70-75.
9. Mert M, Caliskan ME. The effect of mepiquat chloride (pix) on yield, yield components and fiber characteristics of cotton. Turkish Journal of Field Crops. 1998;3(2): 68-72.
10. Karthikeyan PK, Jayakumar R. Nitrogen and chlormequat chloride on cotton cultivar. Kluwer Academic Publishers, Dordrecht, Netherlands. 2001;806-807.

Comment [U9]: inadequate reference.

Formatted: English (U.S.)

Comment [U10]: Please confirm this.

Comment [U11]: Please review this reference. It is a book reference? View i.e Rang HP, Dale MM, Ritter JM, Moore PK. Pharmacology. 5th ed. Edinburgh: Churchill Livingstone; 2003.

Comment [U12]: Please review this reference.

11. Mekki BB. Effect of mepiquat chloride on growth, yield and fiber properties of some Egyptian cotton cultivars. Arab University Journal of Agriculture Sciences.1999;7: 455-466.
12. Sawan ZM, Mahmoud HM, Momtaz O. Influence of nitrogen fertilization and foliar application of plant growth retardants and zinc on quantitative and qualitative properties of Egyptian cotton (*Gossypium barbadense* L. Var. Giza 75). Journal of Agricultural and Food Chemistry,1997;45:3331-3336
13. Sawan ZM. Plant growth retardants, plant nutrients, and cotton production. Communications in Soil Science and Plant Analysis. 2013;44(?): 1353–1398.
14. Piper CS. Soil and plant analysis. Hons Publishers, Bombay. 1966.
15. Jackson ML. Soil chemical analysis. Prentice Hall India Pvt.Limited, New Delhi. 1973;498.
16. Richards LA. Diagnosis and improvement of saline and alkali soils. USDA Hand Book No. 60, Washington, D.C. 1954.
17. Subbaiah BV, Asija GL. A rapid procedure of estimation of available nitrogen in soils. Current Science. 1956;65(7): 477-480.
18. Olsen SR, Cole CV, Watanable FS, Dean LA. Estimation of available phosphorus in soil by extraction with sodium bicarbonate, U.S.D.A. Circular. 1954;939.
19. Dong H, Li W, Eneji AE, Zhang D. Nitrogen rate and plant density effects on yield and late season leaf senescence of cotton raised on a saline field. Field Crops Research. 2012;126: 137-144.
20. Munir MK, Tahir M, Saleem MF, Yaseen M. Growth, yield and earliness response of cotton to row spacing and nitrogen management. The Journal of Animal and Plant Science. 2015;25(3): 729-738.
21. Oosterhuis DM, Coker DL, Gomez SK. Characterization of the fruiting growth curve used in crop monitoring. Spec. Rep. 198. Arkansas Exp. Stn., Fayetteville, AR. 2000.
22. Hussain SZ, Faid S, Anwar M, Gill MI, Baugh MD. Effect of plant density and nitrogen on the yield of seed cotton-variety CIM-443. Sarhad Journal of Agriculture. 2000;16: 143-147.

Comment [U13]: Please review this references.

Comment [U14]: Incomplete reference.

Comment [U15]: Please review this reference