

Original Research Article

Diversity of powdery mildew mycoparasite *Ampelomyces quisqualis* under natural ecosystem and its molecular characterization

Abstract

The powdery mildew ~~diseases~~ are the most common ~~diseases and which widely~~ widely affect ~~crops~~ in many countries. The mildew infection ~~appeared~~ on leaves, petiole, buds, inflorescence and other tender tissues of crops ~~resulting causing~~ 65% of the crop loss. Even though the fungicidal spray to control the disease, the residual and environmental effects are causing long term ecological imbalance in cropping system. ~~Recently, an The~~ alternative and eco-friendly ~~approaches were made in the present by exploring the~~ mycoparasite associated with powdery mildew pathogens ~~was used against~~ *Erysiphe* sp, *Leveillulla* sp, *Sphaerotheca* sp and *Oidium* sp. ~~This study aimed to mitigated the use of the expensive and harmful fungicides to save human health and to reduce the financial costs of controlling the fungal infections of crops in the field by using a commercial biofungicide naturally associated with different species of powdery midew fungi.~~ Our results exhibited that ~~The the~~ isolation of ~~the~~ mycoparasite *Ampelomyces quisqualis* from the mildew pathogens ~~revealed that the Ampelomyces quisqualis~~ was closely associated with most of the genera of powdery mildew pathogens. The natural mycoparasitization efficiency of *A. quisqualis* was observed and higher (81%) efficiency was recorded in *Erysiphe cichoracearum* followed by 76% in black gram powdery mildew caused by *Erysiphe polygoni*. A total of 20 mycoparasitic *Ampelomyces* isolates were enumerated from ~~mycelia-pycnidia~~ of 6 different powdery mildew species that naturally infected their host plants. The pycnidial morphological variations of *A. quisqualis* and the largest size ~~of~~ pycnidia was found in the *Erysiphe chichoracearum* (73.54µm length × 42.15µm width) shows maximum efficiency in natural parasitization. The molecular characterization of *A. quisqualis* isolates based on using rDNA ITS region was carried out and sequenced. The phylogenetic analysis was performed using the Maximum likelihood technique was shown the distinct relatedness with five *Ampelomyces* isolates made in the present study were clustered. This is the first and detailed study on diversity of *Ampelomyces* and quantification of natural mycoparasitism of different genera of powdery mildew of *A. quisqualis*.

Key words: bhendi, mycoparasitism, *Ampelomyces quisqualis*, biocontrol, powdery mildew

Introduction

The obligate biotrophic powdery mildew fungi are causing infection to more than 10,000 host plant species, including important vegetables and horticultural crops (Takamatsu 2013). To protect the crops from powdery mildew pathogens, fungicides are indiscriminately applied frequently with higher dosage which causes fungicide resistance. In addition to that the fungicides are causing harmful effect on biodiversity, natural ecosystem and ~~possess-causes~~ the residual fungicides problem in food (Fernandaz *et al.*, 2021). The physical and biological approaches have been proposed to support and replace chemical management of powdery mildews. The mycoparasites (fungi that parasitize other fungi) are naturally abundant in the majority of the powdery mildew infection in terrestrial environmental conditions especially in biotrophic interaction (Boddy and Watkinson 1995; Kiss *et al.*, 2004). Numerous mycoparasites have been investigated extensively and economically used as bio-control agents (Viterbo and Horwitz 2010). *Ampelomyces quisqualis* is an unique mycoparasite of Erysiphales fungi (Angeli *et al.*, 2012), and it is classified as an endoparasitic nature because of its conidia penetrate into *Pyllactinia xanthii* and generate pycnidia inside powdery mildew structures (Romero *et al.*, 2003). The mycoparasite inhibits the growth of *P. xanthii* haustoria, hence limiting the pathogen's nutrition absorption.

Formatted: Font: Italic

The cross-inoculation assay of *A. quisqualis* studied by Liang *et al.*, (2007) for mycohost specificity. The studies have shown that a strain of *A. quisqualis* isolated from a species of powdery mildew fungi can infect different powdery mildew species, suggesting that there is no mycohost-specificity. Variations in mycohost phenotypesphenology and differentiation in that specific of *Ampelomyces* sp. from bhendi host caused in the infections in different powdery mildew pathogens viz., *Leveillula taurica* and *Erysiphe polygoni* in bell pepper and black gram respectively (Kiss 2012), (Kiss 2012) [please this phrase must be rerewitten.]

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

~~In order to~~ The explore the mycoparasite viz., usage of *Ampelomyces* spp to as manage controller of the different strain of grapevine powdery mildew ~~with different strain werewas~~ successfully investigated by Kiss, (2012). Liyanage *et al.*, (2018) showed that the diversity of *Ampelomyces* strains

belong to genetically-distinguished groups, based on sequence of internal transcribed (rDNA-ITS) regions with nuclear ribosomal DNA. Legler *et al.*, (2016) ~~constitute different species of *Ampelomyces* and its morphological characterized and molecular variations were~~ formed a different distinct genetical genetic variations but the taxonomy of species level has not been carried out yet. (This phrase is not clear).

The main goals of this work were 1) to isolate the *Ampelomyces* mycoparasitic strains in the powdery mildew pathogen, 2) to investigate the morphological variation for identification and its mycoparasitic efficiency in natural conditions, 3) to study the genetic diversity of *Ampelomyces* isolates using ITS rDNA. (please rewrite this sentence)

Formatted: Highlight

Materials and methods

Collection of samples and processing

Powdery mildew-infected plant samples were collected from various parts-regions of Tamil Nadu, India. A total of 526 samples were collected from horticultural and agricultural cropping ecosystem in terrestrial-field environmental conditions and samples were stored in refrigerator. The presence-isolation of *Ampelomyces* pycnidia was ensured-carried out under stereo microscope with the magnification of 20X for isolation of mycoparasite. The live samples were stored in the growth chamber using the procedure proposed by Braun (1987) for one week with the modified temperature of $23\pm 1^{\circ}\text{C}$ and 83% RH, 12:12 dark to light ratio according to Braun (1987).

Enumeration of mycoparasite from powdery mildew infection

The presence of *Ampelomyces* pycnidia in the mycelium of the Erysiphales species was examined through microscopic studies. Pycnidia were observed by placing a spore suspension in a slide and covering it with a cover slip. The advanced microscopic equipments such as stereomicroscope, light microscope, phase contrast and Scanning electron microscope were used to document the host parasitic relationship. The qualitative and quantitative morphological structures of *Ampelomyces* pycnidia and pycnidiospores was measured according to (Angeli, 2014).

Isolation of *Ampelomyces* sp. using pycnidia picking method

The powdery mildew infected leaves parasitized by *Ampelomyces* spp. were used for isolation in potato dextrose agar medium. The pycnidia were examined using a stereo microscope were picked using a sterilized insulin needle and placed on potato dextrose agar (PDA; Himedia, Mumbai). Streptomycin

sulphate 0.3% was added to the culturing medium to avoid ~~the any~~ contamination. Plates were incubated at a temperature of 20±2°C and monitored the growth and development. A total of twenty isolates were obtained from different cropping system under natural ecosystem.

Morphological examination of *Ampelomyces* sp.

The ~~sub-cultured~~ twenty ~~ten days old culture~~ isolates of *Ampelomyces* spp were ~~subcultured and ten days old culture were~~ used for morphological studies using phase contrast microscope. The radial ~~development growth five replicates each of mycelia~~ was assessed, as well as the height, texture, and colour of both ~~Petri colony surfaces plate side for five replicated Petri plates per isolate~~. The morphological parameters such as pycnidia, pycnidiospores, and the presence of petiolate in each isolate were measured at 100X magnifications. To study the characteristics of mycelium scanning electron microscope, a ten days old culture was chopped, scraped with a needle and mounted on ~~aluminium~~aluminum stubs using double-sided adhesive tape, coated with gold palladium.

Extraction of genomic DNA

The twenty isolates of *Ampelomyces* sp. were cultured in 100ml Erlenmeyer flasks containing 20ml PDA broth, after 10 days incubation, mycelium was collected. Total fungal DNA was extracted from 100 mg of mycelium by cetyl trimethyl ammonium bromide CTAB method (Möller *et al.*, 1992). The purified DNA was dissolved in 50µl TE buffer (Tris 10mM + EDTA 1mM pH 8.0). Integrity of genomic DNA (gDNA) was checked in 1.5 per cent agarose gel (HiMedia, Mumbai). The quality and quantity of DNA was assessed by using NanoDrop1000 spectrophotometer (Thermo Fisher Scientific NanoDrop 2000c, USA). The concentration of DNA was adjusted to 50 ng/µl and stored at 4°C for further use (Sambrook and Russell, 2001).

PCR amplification

Ampelomyces sp. cultures were identified molecularly using the conserved ribosomal internal transcribed spacer (ITS) region. Using the universal primer pairs ITS1 (5'-TCCGTAGGTGAACCTGCGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3'), we amplified the ITS regions between the small nuclear 18S rDNA and the large nuclear 28S rDNA, including 5.8S rDNA (White *et al.*, 1990; Hirata and Takamasu, 1996). All PCR reactions were carried out using a Mastercycler® Nexus X2 PCR cycler (MA, USA) using the following parameters: 1) initial denaturation at 95°C for 10 min, followed by 35 cycles at 94°C for 30 s, 60°C for 45 sec and 72°C for 1 min and a

final extension at 72°C for 8 min, the electrophoresis was carried out on 1.0 percent agarose gels. The Bio-Rad Gel Doc EZ Imaging System was used to view the PCR products (Biotium, Hayward, CA).

Sequencing of ITS region and analysis

The PCR products were eluted and sequenced further at Barcode Scientific in Bangalore, India. Partial nucleotide sequences of rDNA ITS region of isolates of *Ampelomyces* sp were downloaded from NCBI database (www.ncbi.info). The program the Basic Local Alignment Search Tool-Nucleotide or BLASTn server, was used to edit and align the ITS sequences and the similarity between strains of 18S rRNA gene sequences was calculated using ClustalW (Thompson *et al.*, 1997; Hall, 1999). The aligned sequences were deposited in the GenBank database.

Phylogenetic analysis

The [phylogenetic](#) tree was built with 1000 bootstrap replications in Mega X (Kumar *et al.*, 2018) using the Maximum Likelihood approach based on the Tamura 3-parameter model (Tamura, 1992). The *Pythium insodium* reference sequence for 18S rRNA were obtained from GenBank data and used as an out group in phylogenetic tree analyses.

RESULTS

The powdery mildew diseased samples were collected from different locations of Tamilnadu, India during 2018–2020 and its severity along with presence of mycoparasitic infections were recorded. The mycoparasitic infections were noticed in different genera of powdery mildew pathogens viz., *Erysiphe*, *Uncinula*, *leveillula* and *Sphaerotheca*. The maximum mycoparasitization (81%) of *Ampelomyces* sp. pycnidia was recorded with genera of *Erysiphe* powdery mildews (Table 1).

Morphological characterization and identification of *Ampelomyces* sp

The morphology of *Ampelomyces* sp. pycnidia were studied and shown with different shape such as ovoid, ellipsoid, or globose. The size of the pycnidia and pycnidiospores were ranged in length from 72.09 to 119.57 μm (major axis) and width from 28.45 to 47.12 μm (minor axis) (Fig 1). The variations were in pycnidial shape depends on the genera of powdery mildew pathogens.

Pycnidiospores varied in length from 9.54 to 6.15 μm and in width from 5.71 to 3.21 μm . Pycnidial range in colour from light brown to dark brown, while pycnidiospores are olive green in colour. Out of 20 isolates two isolates were examined in depth for their morphological characters (Fig 2). The morphological parameters of pycnidial shape and size were listed in the Table 1.




Fig 1)Bhendi powdery mildew *Erysiphe chichoracearum* microstructures; **(a)** conidiophore of bhendi powdery mildew; **(b)**chain of conidia. (Notes must be mentioned about the used electron microscope).




Fig 2) Pycnidium on the surface of a bhendi powdery mildew; **(a)** pycnidiospores of *Ampelomyces* spp.; **(b)** conidia with germ_tube





Fig 3) *Ampelomyces* spp. microstructures **(a)** pycnidia produced in the conidiophores of bhendi powdery mildew arrow indicates conidia of bhendi powdery mildew; **(b)** dehiscent pycnidium by apical rupture.

Fig 4. Microscopic image of *Ampelomyces* pycnidia; (a) pycnidia and pycnidiospores produced in *Ampelomyces* isolate; (b) Scanning electron microscopic image of pycnidia of *Ampelomyces* isolate AQTNAU-DST01



The pycnidia are unicellular, hyaline, and elongated to pyriform in shape, measuring around $46.36 \times 10.82 \mu\text{m}$ of pale brown angular textured (**Fig 4**). The hyphal lengths [were](#) developed 48 hours after injection. Following inoculation of a single mature pycnidium in the middle of PDA medium, fungal colonies expand slowly and concentrically.

The Internal Transcribed Spacer (ITS) regions (ITS1 and ITS4) and 5.8S gene area of 18S rDNA were initially amplified with the primers ITS1 and ITS 4 to validate the initial identification and identify the clear taxonomic position. All twenty isolates were amplified with 560 bp([the unit must be written without abbreviation for the first time only](#)). The amplification were identical with prior identity and the amplified 18S-rDNA (ITS 1 and ITS 4) region was purified individually and sequenced by sangar dideoxy sequencing([the year and the reference must be mentioned please](#)) (Fig 5).




Fig 5) Molecular identification of *Ampelomyces* species of ITS region,1–16: M) DNA marker, 1) AQTNAU-DST1, 2) AQTNAU-DST2, 3) AQTNAU-DST3, 4) AQTNAU-DST4, 5) AQTNAU-DST5, 6) AQTNAU-DST6, 7) AQTNAU-DST7, 8) AQTNAU-DST8, 9) AQTNAU-DST9, 10) AQTNAU-DST10,11)AQTNAU-DST11, 12)AQTNAU-DST12, 13) AQTNAU-DST13,14) AQTNAU-DST14,15) AQTNAU-DST15, 16)AQTNAU-DST16, 17) AQTNAU-DST17, 18) AQTNAU-DST18, 19) AQTNAU-DST19,20) AQTNAU-DST20

Molecular identification and phylogenetic analysis

The sequence of ITS regions were shown 97% sequence homology with GenBank sequences with BLASTn analysis. The sequences were submitted in NCBI GenBank, and OM424627, OM424628, OK236008, OM190525 and OM190526 were assigned as accession number for AQTNAU-DST01, AQTNAU-DST02, AQTNAU-DST03, AQTNAU-DST04 and AQTNAU-DST05 isolates, respectively. The phylogenetic analysis shown that the isolated *Ampelomyces* were grouped with other reported *Ampelomyces* on Gen bank. The Maximum likelihood of 18s rDNA sequences of Species of *Ampelomyces* constructed with bootstrap values more than 500 (from 1000 ~~replaces~~replicates) are indicated at the nodes as percentage that ~~varies~~varied with each cluster has been shown in Fig 6. Phylogenetic analysis based on 18s rDNA sequences of different species of *Ampelomyces* isolates were ~~analysed~~analyzed and the results revealed that 6 different clusters were formed in phylogenetic tree. Cluster 1 was the biggest clade, with isolates, including two *Ampelomyces* isolates (AQTNAU-DST03, AQTNAU-DST04) isolated in this study. Further three isolates *Ampelomyces* (AQTNAU-DST02, AQTNAU-DST01 and AQTNAU-DST05) were clubbed together to form a cluster 2 with other different *Ampelomyces* isolates. There were multiple subgroups within this cluster. In the dendrogram, *Pythium insodium* has created a separate cluster (out_group) (Fig. 6). Using the ITS region, we discovered that the 10 isolates obtained from various areas in Tamilnadu had sequence homology with isolates from other regions, including India, China, and Korea.




Fig 6. Phylogenetic tree generated from 18s rDNA sequence of *Ampelomyces* sp. using Maximum likelihood analysis in MEGA10.

DISCUSSION

The powdery mildew is an economically important disease that affects a wide range of crops and caused significant loss in yield. The high occurrence of the infection was noticed throughout the season especially under micro climatic condition. Cortesi *et al.*, (1995) reported that the source of inoculum was high in spring season in bhendi crop and inoculum was survived in the weed hosts.

In this research, we isolated 20 *Ampelomyces* isolates from five different ~~pathogens~~ pathogenic isolates of ~~of~~ powdery mildew in Tamilnadu. ~~at~~ All the isolates are showing the potential powdery mildews infection (Kiss, 2003) The morphological characteristics of all the mycoparasitic activity against *A. quisqualis* isolates were identical and similar with earlier reports (Emmons, 1930; Chona and Munjal, 1956; Rudakov, 1979; Szejnberg *et al.*, 1989; Rankovic, 1997; Liang *et al.*, 2007; Angeli *et al.*, 2012; Jamali, 2015; Gautam and Avasthi, 2016). The investigations clearly indicated that the variations of pycnidia and its spores ~~shown~~ showed different species of powdery mildew pathogens in various plant species. It is supported by Puzanova (1991) and Pintye *et al.* (2012) who reported that there was no specific host specialization with *Ampelomyces* isolates and they mentioned that cross mycoparasitic activity with wide range powdery mildew pathogens. { you must foscused on this very important point}

In all the examined strains, the colour and shape of the pycnidia varied from light to dark (brown or grey) and from ovoid to ellipsoid. Similar results were reported by Kim *et al.*, (2009) who reported that the ~~colour~~ color of pycnidium ranged from light brown to dark brown. At present, the morphometric studies of the isolates of *A. quisqualis* revealed that the size of pycnidia was variable in the range 72.09 to 119.57 µm; pycnidiospore ranged length from 9.54 to 6.15 µm and in width from 5.71 to 3.21 µm. This was in accordance with the findings of Liang *et al.* (2007) who reported that the pycnidia were pyriform to globose measuring about 36-123 × 22-45 µm and it contained unicellular guttulated d conidia which measured about 4.2-7.5 µm in length.

The mycelia of powdery mildew-infecting *Ampelomyces* spp. were hyaline and the pycnidia were light brown to dark brown in ~~colour~~ color, with olive green pycnidiospores. However, Lee *et al.*, (2007) and Angeli *et al.*, (2011) found ~~colour~~ color differences in the mycelium and pycnidia of several mycohosts belonging to the genus *Erysiphe*, ranging from olive green to light and dark brown. Pycnidia and pycnidiospores in *Ampelomyces* isolates from powdery mildews varied in shape depending ~~upon~~ on the fungal structure in which they were formed. They were pear _shaped, spindle-shaped or nearly

spherical when they were formed inside *E. necator* conidiophores, hyphae or chasmothecia (Angeli *et al.*, 2009).

In the present investigation, ITS-PCR analysis were attempted using ITS 1 and 4 universal primers pertaining to 18s rDNA Intergenic Transcribed Spacer region. The isolates of *Ampelomyces quisqualis* collected from different powdery mildew hosts were identified as *Ampelomyces quisqualis* based on the amplified as a fragment of 560 bp and it has been confirmed that all the isolates were *Ampelomyces* sp. The sequencing of the isolates showed 94 – 98% homology with the respective *Ampelomyces* sp. ~~Our results were highly in agreement by those of~~ Similarly, the result of Kiss (1997), Angeli *et al.* (2012 and Jamali (2015) ~~those~~ who have reported that the size of the PCR product of *A. quisqualis* varied from 500-600 bp. The 18s rDNA gene sequencing analysis has ~~aids an important role~~ in the identification process of *Ampelomyces quisqualis* ~~and identified up to~~ species level through 18s rDNA sequencing (Liyange *et al.*, 2018).

The Neighbour joining method of phylogenetic tree revealed that the similarity in clustering pattern with other species *Ampelomyces quisqualis* reported earlier. The phylogenetic tree generated by our analysis contains six distinct Cluster. These results was confirmed further to conclude as *Ampelomyces quisqualis* and previous reports of Park *et al.*, (2010); Angeli *et al.*, 2011 and Nemeth *et al.*, (2021) were shown the molecular characterization and diversification of *Ampelomyces* isolates formed four distinct clades. Likewise, ITS rDNA region of *Ampelomyces* sp in our current investigation revealed the diversity of six different phylogenetic groups comprising of an outgroup of *Pythium insodium*. *Ampelomyces* isolate AQTNAU-DST03, AQTNAU-DST04 which formed cluster 2. The cluster 3 was comprising of AQTNAU-DST 02, AQTNAU-DST01 and AQTNAU-DST05.

Mycoparasitism has been shown to be an important mechanism of biological control (Brozova, 2004). The *Ampelomyces* sp parasites on powdery mildew host resulted in reduction of mycelial growth. It infected and produced pycnidia within the powdery mildew hyphae and conidiophores. This has been ~~witnessed endorsed by with~~ other studies of powdery mildew including grapes and various crops (Kiss, 2003). *A. quisqualis* with significantly reduced the powdery mildew symptom expression caused by *E. chichoracearum* in bhendi crop. Similarly, Shishkoff and Mcgrath (2002) showed that the parasitism reduces growth and may eventually kill the mildew colonies. (This paragraph is very important please check the language to be more clear)

These results shows that the *Ampelomyces* sp. mycoparasites and further disease progressions and suppress the sporulation rate of their mycohosts of their infected plants and enhances crop vigour

Formatted: Highlight

Formatted: Highlight

and growth after reduction of powdery mildew infection. A number of experiments demonstrated that *Ampelomyces* sp. are being used as a biocontrol agent for the management of powdery mildews (Falk *et al.*, 1995; Szentivanyi *et al.*, 2003; Angeli *et al.*, 2012). [\(Same notice as mentioned above\)](#)

Conclusion

Conclusively, the function of *A. quisqualis* in the control of powdery mildew infections is essential in integrated disease management. As AQ10 Biofungicide, this mycoparasite is presently one of the most advanced in terms of commercial development of a fungal bio control product for powdery mildew diseases. The ecological interaction between plants, powdery mildews and *Ampelomyces* might be explored further to better understand the role of fungal antagonists in plant parasite population dynamics. Continuous use of fungicides creates resurgence among the pathogens and also affects beneficial microflora. Under these circumstances exploitation of effective bio control agents for disease management are required. Future studies regarding the improvement of the efficacy of bio control agents and their delivery methods can further improve the yield.

Formatted: Highlight

[Please](#) [rewrite](#) [the](#) [conclusion](#) [linguistic](#) [style](#)

Table 1. *A.quisqualis* isolated from the mycoparasitized samples collected from different regions of Tamil Nadu

S. No	Isolates	Plant host	<i>A. quisqualis</i> parasitic on powdery mildew pathogens	Pycnidium			<i>A. quisqualis</i>			Mycoparasitism (%)
				Color	Length	width	Shape	Length	Shape	
1.	AQ-TNAU-DST-01	<i>Abelmoschus esculentus</i> L.	<i>Erysiphe cichoracearum</i> DC.	Brown	73.54	42.15	Round	9.54	Cylindrical and curved at both the edges	81
2.	AQ-TNAU-DST-02	<i>Vigna unguiculata</i> L.	<i>Erysiphe polygoni</i> DC.	Brown	56.18	41.22	Pyriform	8.27	Cylindrical shaped	76
3.	AQ-TNAU-DST-03	<i>Sesamum indicum</i> L.	<i>Oidium erysiphoides</i>	Dark brown	48.77	40.21	Round	8.42	Oval shaped	52
4.	AQ-TNAU-DST-04	<i>Abelmoschus esculentus</i> L.	<i>Erysiphe cichoracearum</i> DC.	Dark brown	52.08	38.96	Oval	7.27	Cylindrical and curved at both the edges	78
5.	AQ-TNAU-DST-05	<i>Capsicum annuum</i> L.	<i>Leveillula taurica</i> (Lev.)	Dark brown	49.25	36.27	Oval and tapering at both the edges	9.22	Oval shaped	43
6.	AQ-TNAU-DST-06	<i>Abelmoschus esculentus</i> L.	<i>Erysiphe cichoracearum</i> DC.	Brownish black	41.25	36.23	Oval	8.14	Cylindrical and curved at both the edges	72
7.	AQ-TNAU-DST-07	<i>Vitis vinifera</i> L.	<i>E. necator</i> Schwein.	Dark brown	42.61	29.05	Round	9.01	Cylindrical and curved at both the edges	61
8.	AQ-TNAU-DST-08	<i>Tagetes erecta</i> L.	<i>Leveillula taurica</i> (Lev.)	Dark brown	32.62	28.25	Cylindrical	8.74	Oval shaped	31
9.	AQ-TNAU-DST-09	<i>Capsicum annuum</i> L.	<i>Leveillula taurica</i> (Lev.)	Brownish black	42.15	34.31	Round	8.52	Cylindrical and curved at both the edges	36
10.	AQ-TNAU-DST-10	<i>Abelmoschus esculentus</i> L.	<i>Erysiphe cichoracearum</i> DC.	Brown	40.78	33.13	Oval	7.61	Oval shaped	73
11.	AQ-TNAU-DST-11	<i>Vitis vinifera</i> L.	<i>E. necator</i> Schwein.	Brown	43.01	28.47	Oval and tapering at both the edges	9.11	Oval shaped	69
12.	AQ-TNAU-DST-12	<i>Rosa sp.</i>	<i>Sphaerotheca pannosa</i>	Dark brown	31.77	29.15	Oval	8.55	Cylindrical and curved at both the edges	28
13.	AQ-TNAU-DST-13	<i>Morus alba</i>	<i>Phyllactinia corylea</i>	Dark brown	51.13	37.21	Round	8.31	Cylindrical and curved at both the edges	64
14.	AQ-TNAU-DST-14	<i>Vigna unguiculata</i> L.	<i>Erysiphe polygoni</i> DC.	Dark brown	48.05	37.14	Cylindrical	9.34	Oval shaped	66
15.	AQ-TNAU-DST-15	<i>Sesamum indicum</i> L.	<i>Oidium erysiphoides</i>	Brownish black	42.12	35.12	Round	8.16	Oval shaped	53
16.	AQ-TNAU-DST-16	<i>Abelmoschus esculentus</i> L.	<i>Erysiphe cichoracearum</i>	Whitish	41.02	45.1	Round	9.01	Oval shaped	70
17.	AQ-TNAU-DST-17	<i>Vitis vinifera</i> L.	<i>E. necator</i> Schwein.	Black	32.62	41.22	Cylindrical	8.74	Oval shaped	63
18.	AQ-TNAU-DST-18	<i>Tagetes erecta</i> L.	<i>Leveillula taurica</i> (Lev.)	Dark brown	42.15	40.21	Round	8.52	Oval shaped	52
19.	AQ-TNAU-DST-19	<i>Vigna unguiculata</i> L.	<i>Erysiphe polygoni</i> DC	Black	41.78	39.96	Oval	7.61	Oval shaped	66

20	AQ-TNAU-DST-20	<i>Cucumis sativus</i>	<i>Erysiphe cichoracearum</i>	Grey	45.01	37.27	Round	9.11	Oval shaped	58
----	----------------	------------------------	-------------------------------	------	-------	-------	-------	------	-------------	----

UNDER PEER REVIEW

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

Reference

1. Angeli D, Maurhofer M, Gessler C, Pertot I. Existence of different physiological forms within genetically diverse strains of *Ampelomyces quisqualis*. *Phytoparasitica*. 2011; 40:37–51.
2. Angeli D, Pellegrini E, Pertot I. Occurrence of *Erysiphe necator* chasmothecia and their natural parasitism by *Ampelomyces quisqualis*. *Phytopathology*. 2009;99(6):704-10.
3. Angeli D, Puopolo G, Maurhofer M, Gessler C, Pertot I. Is the mycoparasitic activity of *Ampelomyces quisqualis* biocontrol strains related to phylogeny and hydrolytic enzyme production?. *Biological control*. 2012;63(3):348-58.
4. Boddy L, Watkinson SC. Wood decomposition, higher fungi, and their role in nutrient redistribution. *Canadian journal of botany*. 1995;73(S1):1377-83.
5. Braun U. A monograph of the Erysiphales (powdery mildews). *Beihefte zur Nova Hedwigia*. 1987(89).
6. Brozova J. Mycoparasitic fungi *Trichoderma* spp. in plant protection—Review. *Plant Protection Science*. 2004;40(2):63.
7. Chona BL, Munjal RL. *Cicinnobolus cesatii* de Bary: Hyperparasite of powdery mildews in India. *Indian Phytopathol*. 1956;9:98-105.
8. Cortesi P, Rigling D, Heiniger U. Comparison of vegetative compatibility types in Italian and Swiss subpopulations of *Cryphonectria parasitica*. *European Journal of Forest Pathology*. 1998;28(3):167-76.
9. Emmons CW. *Cicinnobolus cesatii*, a study in host-parasite relationships. *Bulletin of the Torrey Botanical Club*. 1930:421-41.
10. Falk SP, Gadoury DM, Cortesi P, Pearson RC, Seem RC. Parasitism of cleistothecia of *Uncinula necator* by the mycoparasite *Ampelomyces quisqualis*. *Phytopathology*. 1995; 85:794–800.

11. Fernandes GW, Arantes-Garcia L, Barbosa M, Barbosa NP, Batista EK, Beiroz W, Resende FM, Abrahao A, Almada ED, Alves E, Alves NJ. Biodiversity and ecosystem services in the Campo Rupestre: A road map for the sustainability of the hottest Brazilian biodiversity hotspot. *Perspectives in Ecology and Conservation*. 2020;18(4):213-22.
12. Gautam AK, Avasthi S. *Ampelomyces quisqualis* Ces.—a mycoparasite of *Euphorbia hirta* powdery mildew in Himachal Pradesh, India. *Journal of Phytopathology and Pest Management*. 2016;20: 64-70.
13. Hall T. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. In *Nucleic Acids Symp. Ser.* 1999; 41:95-98).
14. Hirata T, Takamatsu S. Nucleotide sequence diversity of rDNA internal transcribed spacers extracted from conidia and cleistothecia of several powdery mildew fungi. *Mycoscience*. 1996;37(3):283-8.
15. Jamali S. Molecular phylogeny of endophytic isolates of *Ampelomyces* from Iran based on rDNA ITS sequences. *Molecular biology reports*. 2015;42(1):149-57.
16. Kim JY, Lee WH, Kim HM. Physical Characteristics and Antagonistic Effect of *Ampelomyces*. *Research in Plant Disease*. 2009;15(3):209-16.
17. Kiss L, Russell JC, Szentiványi O, Xu X, Jeffries P. Biology and biocontrol potential of *Ampelomyces* mycoparasites, natural antagonists of powdery mildew fungi. *Biocontrol Science and Technology*. 2004;14(7):635-51.
18. Kiss L. A review of fungal antagonists of powdery mildews and their potential as biocontrol agents. *Pest Management Science: formerly Pesticide Science*. 2003;59(4):475-83.
19. Kiss L. Genetic diversity in *Ampelomyces* isolates, hyperparasites of powdery mildew fungi, inferred from RFLP analysis of the rDNA ITS region. *Mycological Research*. 1997; 101(9):1073-80.
20. Kiss L. Limits of nuclear ribosomal DNA internal transcribed spacer (ITS) sequences as species barcodes for Fungi. *Proceedings of the National Academy of Sciences*. 2012;109 (27):E1811.
21. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Molecular biology and evolution*. 2018;35(6):1547.
22. Lee SY, Hong SK, Kim YK, Kim HG. Selection and identification of a hyperparasite, *Ampelomyces quisqualis* 94013 for biocontrol of cucumber powdery mildew. *The Korean Journal of Mycology*. 2007;35(2):121-7.

23. Legler SE, Pintye A, Caffi T, Gulyás S, Bohár G, Rossi V, Kiss L. Sporulation rate in culture and mycoparasitic activity, but not mycohost specificity, are the key factors for selecting *Ampelomyces* strains for biocontrol of grapevine powdery mildew (*Erysiphe necator*). *European Journal of Plant Pathology*. 2016;144(4):723-36.
24. Liang C, Yang J, Kovács GM, Szentiványi O, Li B, Xu X, Kiss L. Genetic diversity of *Ampelomyces* mycoparasites isolated from different powdery mildew species in China inferred from analyses of rDNA ITS sequences. *Fungal Diversity*. 2007; 24:225-40.
25. Liyanage KK, Khan S, Brooks S, Mortimer PE, Karunarathna SC, Xu J, Hyde KD. Morpho-molecular characterization of two *Ampelomyces* spp.(Pleosporales) strains mycoparasites of powdery mildew of *Hevea brasiliensis*. *Frontiers in microbiology*. 2018;9:12.
26. Nemeth MZ, Mizuno Y, Kobayashi H, Seress D, Shishido N, Kimura Y, Takamatsu S, Suzuki T, Takikawa Y, Kakutani K, Matsuda Y. *Ampelomyces* strains isolated from diverse powdery mildew hosts in Japan: their phylogeny and mycoparasitic activity, including timing and quantifying mycoparasitism of *Pseudoidium neolycopersici* on tomato. *PloS one*. 2021;16(5):e0251444.
27. Park MJ, Choi YJ, Hong SB, Shin HD. Genetic variability and mycohost association of *Ampelomyces quisqualis* isolates inferred from phylogenetic analyses of ITS rDNA and actin gene sequences. *Fungal Biology*. 2010;114(2-3):235-47.
28. Pintye A, Bereczky Z, Kovacs GM, Nagy LG, Xu X, Legler SE, Vaczy Z, Vaczy KZ, Caffi T, Rossi V, Kiss L. No indication of strict host associations in a widespread mycoparasite: grapevine powdery mildew (*Erysiphe necator*) is attacked by phylogenetically distant *Ampelomyces* strains in the field. *Phytopathology*. 2012;102(7):707-16.
29. Puzanova LA. Distribution of hyperparasites from the genus *Ampelomyces* ces-ex-schlecht on the powdery mildew fungi in northern caucasus and their importance in disease-control. *Mikologiya i Fitopatologiya*. 1991;25(5):438-42.
30. Rankovic B. Hyperparasites of the genus shape *Ampelomyces* on powdery mildew fungi in Serbia. *Mycopathologia*. 1997;139(3):157-64.
31. Romero D, Rivera ME, Cazorla FM, De Vicente A, Perez-Garcia A. Effect of mycoparasitic fungi on the development of *Sphaerotheca fusca* in melon leaves. *Mycological research*. 2003;107(1):64-71.

32. Rudakov OL. Fungi of the genus *Ampelomyces* Ces. ex Schlecht. Mikologiya i Fitopatologiya. 1979;13:104-10.
33. Sambrook J, Russell DW. Fragmentation of DNA by sonication. Cold spring harbor protocols. 2006;(4):4538.
34. Shishkoff N, McGrath MT. AQ10 biofungicide combined with chemical fungicides or AddQ spray adjuvant for control of cucurbit powdery mildew in detached leaf culture. Plant Disease. 2002;86(8):915-8.
35. Szentivanyi O, Kiss L. Overwintering of *Ampelomyces* mycoparasites on apple trees and other plants infected with powdery mildews. Plant pathology. 2003;52(6):737-46.
36. Szejnberg A, Galper S, Mazar S, Lisker N. *Ampelomyces quisqualis* for biological and integrated control of powdery mildews in Israel. Journal of Phytopathology. 1989;124(4):285-95.
37. Takamatsu S. Origin and evolution of the powdery mildews (Ascomycota, Erysiphales). Mycoscience. 2013;54(1):75-86.
38. Tamura K. Estimation of the number of nucleotide substitutions when there are strong transition-transversion and G+ C-content biases. Mol Biol Evol. 1992;9(4):678-87.
39. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. Nucleic acids research. 1997;25(24):4876-82.
40. Viterbo A, Horwitz BA. Mycoparasitism. Cellular and molecular biology of filamentous fungi. 2010:676-93.
41. White TJ, Bruns T, Lee SJ, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. PCR protocols: a guide to methods and applications. 1990;18(1):315-22.