

Impact of different arbuscular mycorrhizal fungi on nutrient status and their uptake by *Melia azedarach* seedling

Comment [H1]:

Comment [H2R1]: Capitalize each word please

ABSTRACT

Melia azedarach belongs to the family *Meliaceae*, is a deciduous tree with a short bole and a spreading crown and one of the most important trees in arid and semi-arid area. The arbuscular mycorrhizal fungi (AMF) are soil microorganisms composing the essential components of the sustainable soil-plant system. These AM fungi form extensive extraradical mycelia which increases accessible soil volume for the plant to absorb phosphorus and water. The effect of three different species of *Glomus* (*G. mosseae*, *G. intraradices* and *G. fasciculatum*) of AM fungi inoculated with *Melia azedarach* drupes was observed in the nursery of the Forestry Department, CCS Haryana Agricultural University, Hisar during 2019. The results revealed that application of three *Glomus* species in combination produced synergistic increase in soil macro and micro nutrient contents after harvest i.e. N = 119.86 kg ha⁻¹, P = 16.91 kg ha⁻¹, K = 260.86 kg ha⁻¹, Fe = 5.95 ppm, Zn = 0.66 ppm, Cu = 0.42 ppm and Mn = 1.63 ppm. Similarly, there was significantly higher macro and micronutrient (N = 1.39 %, P = 0.27 %, K = 0.63 %, Fe = 90.43 ppm, Zn = 32.24 ppm, Cu = 9.96 ppm and Mn = 31.28 ppm) concentrations in leaves after 180 days in the treatments when soil was inoculated with all three *Glomus* species. The seedlings uptake of macro and micronutrient was also found highest when three *Glomus* species were inoculated artificially and simultaneously in soil. It can be summarized that inoculation of the AM fungi improved soil fertility and seedlings growth of *Melia azedarach*.

Comment [H3]: You need to mention it in full with the abbreviation in bracket before subsequent use of abbreviation please

Comment [H4]: Remove this statement

Keywords: AMF, *G. mosseae*, *G. intraradices*, *G. fasciculatum*, Macro and Micro nutrient

INTRODUCTION

India is the seventh-largest country globally, with a total area of 7, 67,419 square km under the tree. The country's total forest area is 24.62 per cent of its geographical area comprising 21.71 per cent of the forest cover and 2.91 per cent tree cover and most growing states are Madhya Pradesh, Arunachal Pradesh, Maharashtra, Orissa, Chhatisgarh and Haryana. In Haryana, forests cover share only 3.63 per cent of the total geographical area (ISFR, 2021). *Melia azedarach* (family *Meliaceae*) is a moderate deciduous tree with a short bole and a spreading crown, bi-pinnate or tri-pinnate leaves and a dark grey bark with shallow longitudinal furrows (Troup, 1921). This plant has a reasonably coarse root system with very few root hairs, which are common characteristics of plants that are responsive to arbuscular mycorrhizal symbiosis. The arbuscular mycorrhizal fungi (AMF) are soil microorganisms composing the essential components of the sustainable soil-plant system. The AM fungi forms extensive extraradical mycelia which increases accessible soil volume for the plant to absorb phosphorus and water (Dierks *et al.*, 2021). These fungi provide their host with various benefits including higher phosphorus absorption (Gossus and Mohammad, 2009), defending root against diseases (Bakhtiar *et al.*, 2010), plant growth-promoting hormone (Herrera-Medina *et al.*, 2007) and boosting plant growth and productivity (Duponnoisa *et al.*, 2005). The mycorrhizal fungi are naturally occurring species that form symbiotic relationships with almost all plants inside the roots. This also provides a perfect ecological niche that is necessary for fungal growth and development including the completion of the sexual cycle (Sandeep *et al.*, 2015). The arbuscules are branched hyphae, within the root cells and this is a connection for nutrient exchange between fungi and the host plant (Afzal *et al.*, 2011). Fungus improves nutrition, growth, drought resistance and provides nutrients and water to the host plant (Luyindula and Haque, 1992). Besides the direct effects of AM fungi on the cycling of nutrients and plant performance, there are several indirect effects of AM F fungi for improving soil fertility via enmeshment of soil particles for better soil aggregation (Rillig *et al.*, 2015) and nutrient maintenance (Cavagnaro *et al.*, 2015). Therefore, farmers' field management through a greater abundance of AM fungi could have improved nutrient accessibility and availability (Sato *et al.*, 2015). However, sustaining AM fungi especially

mycelia, needs a supply of plentiful carbohydrates from host plants (Raven *et al.*, 2018). In Haryana, agroforestry is distributed throughout arid and semi-arid regions (Handa *et al.*, 2020), which could act as perennial AM fungi hosts and thus satisfy AM fungi continuous carbohydrate demand (Bainard *et al.*, 2011). Some trees are exhaustive for nutrients but not all so, it depends on tree species for their effect on soil nutrient status, and differences in their relative availability which may further affect the AM fungi symbiosis relationship functioning (Johnson, 2010) and AM fungi growth strategies (Chagnon *et al.*, 2013). Furthermore, it is vital to contemplate the overall effect of tree species with AM fungi inoculation on nutrients trade-off ultimately affecting soil fertility status, tree growth, and biomass production, including potential trade-offs regarding competition for light, water, and nutrients (Blaser *et al.*, 2018; Wartenberg *et al.*, 2017). Globally, it remains an open question whether trees grown with AM fungi can rejuvenate and preserve soil fertility and, thus, boost crop performance. However, few studies have been made to isolate and select effective AM fungi for inoculation in *Melia azedarach* in nursery conditions. Keeping in view of all these factors the present study was planned to find out the effect of different AM fungi (*Glomus* sp.) on the growth and nutrient uptake of *Melia azedarach* seedling.

MATERIAL AND METHODS

The study was carried out in the nursery of Forestry Department, CCS Haryana Agricultural University, Hisar (20° 10' N lat., 75° 46' E long., alt. 215 m msl), situated in the semi-arid region of North-Western India. The soil types were Typic Ustochrept with sandy loam texture. The climate of Hisar (Haryana) is semi-arid with hot and dry desiccating winds accompanied by frequent dust storms with high velocity in summer months, severe cold during in winter months and humid warm during monsoon rainy season. The mean monthly maximum and minimum temperature sometimes exceed 48° C on hot summer days. The relative humidity varies from 5 to 100 per cent, while temperature below freezing point accompanied by frost in winter is usually experienced in this region. The characteristics of mother plant i.e. plant height; DBH and clear bole height were 9.1 m, 40 cm and 4.2 m, respectively. An average of 100 drupes were used for the experiment and traits such as fruit length (15 mm), fruit breadth (10 mm), fruit weight (0.32 g), fruit thickness (1.7 mm) and test weight (24 g) were recorded. The soil and rootlets from the root horizon of *Glomus mosseae*, *Glomus fasciculatum* and *Glomus intraradices* inoculated wheat plants were used to inoculate soil before sowing of drupes of *Melia azedarach* in seven treatments. The experimental soil was collected from Balsamand Research Farm, CCS HAU, Hisar. The soil was drenched with formaldehyde, and immediately covered with an air-tight polythene sheet for 5 days. The drupes of *Melia azedarach* were collected from plus tree at Hisar and were sown at about 2-3 cm deep in polybags of one kg soil capacity inoculated with different *Glomus* species as per detail given below. This experiment was carried out with completely randomized design and each treatment having three replications.

There were seven treatments of soil inoculation of three different *Glomus* sp. inoculated singly as well as in combination before sowing and control (uninoculated AM fungi) as shown below:-

- T₁ - Treated soil + *G. mosseae* (450-500 sporocarps kg⁻¹ soil)
- T₂ - Treated soil + *G. intraradices* (450-500 sporocarps kg⁻¹ soil)
- T₃ - Treated soil + *G. fasciculatum* (450-500 sporocarps kg⁻¹ soil)

Comment [H5]: MATERIALS

Comment [H6]: Write in full then you use the abbreviation in subsequent statements

Comment [H7]: Any source from whom you adapted this procedure?

Comment [H8]: Any source?

- T₄** - Treated soil + *G. mosseae* + *G. intraradices* (225-250 sporocarps of each AM fungi kg⁻¹ soil)
T₅ - Treated soil + *G. mosseae* + *G. Fasciculatum* (225-250 sporocarps of each AM fungi kg⁻¹ soil)
T₆ - Treated soil + *G. intraradices* + *G. Fasciculatum* (225-250 sporocarps of each AM fungi kg⁻¹ soil)
T₇ - Treated soil + *G. mosseae* + *G. intraradices* + *G. fasciculatum* (150-165 sporocarps of each AM fungi kg⁻¹ soil)
T₈ - Control (un-inoculated)

The soil samples were collected before and after harvesting of *M. azedarach* seedlings. These samples were air-dried and passed through a 2 mm sieve before determining macro and micronutrients (Antil *et al.*, 2002). The alkaline permanganate method described by Subbiah and Asija (1956) was used to determine available nitrogen in soil. The available phosphorus was determined spectrophotometrically by Olsen's method (Olsen, 1954) and blue colour intensity was measured at 660 nm wavelength using a red filter. Neutral normal NH₄OAC solution was used for estimation of available Potassium on flame photometer (Hanway and Heidal, 1952). DTPA-TEA buffer (0.005 M DTPA + 0.001 M CaCl₂ + 0.1 M TEA, pH 7.3) as proposed by Lindsay and Norvell (1978) was used for the determination of available micronutrients (Fe, Cu, Zn and Mn) concentration using Atomic absorption spectrophotometer. The plant leaf samples were ground and digested with the di-acid mixture (nitric acid and perchloric acid). The colorimetric method proposed by Lindner (1944) was used to estimate total nitrogen in the plant. Blue filter at 440 nm wavelength was used to read the intensity of orange colour on spectrophotometer. In the plant sample, total phosphorus was determined by Vanado-molybdophosphoric yellow colour method proposed by Koenig and Johnson (1942). The potassium was determined by a Flame photometer using the standard curve (Jackson, 1973). The micronutrient concentration in aliquot was determined spectrophotometrically at most sensitive wavelengths (Fe-248.7 nm, Zn-213.7 nm, Cu-324.6 nm and Mn-279.5 nm), Elwell and Gridley, 1967.

$$\text{Macronutrient uptake (g plant}^{-1}\text{)} = \frac{\text{Macronutrient content in plant (\%)} \times \text{total dry weight of corresponding plant (g)}}{100}$$

$$\text{Micronutrient uptake (mg plant}^{-1}\text{)} = \frac{\text{Micronutrient content in plant (ppm)} \times \text{total dry weight of corresponding plant (g)}}{10^3}$$

Hyphae contribution (HC) was calculated using the following formula (Kothari *et al.*, 1991):

$$\text{HC} = \frac{(\text{X uptake of the mycorrhizal plants} - \text{X uptake of the non-mycorrhizal plants})}{\text{X uptake of the mycorrhizal plants} \times 100}$$

Where, X= respective nutrient (P/Zn)

RESULTS AND DISCUSSION

The data presented in Fig. 1 showed that available N, P and K contents were found maximum 119.86, 16.91 and 260.86 kg ha⁻¹, respectively in treatment T₇ followed by treatment (T₅) when *G. mosseae* and *G. fasciculatum* were applied in soil together i.e. 118.99, 15.81 and 258.87 kg ha⁻¹, respectively after harvest of seedlings. However, the nutrient content of N and K were found non-significant, when drupes were inoculated individually with *G. mosseae* (T₁), *G. intraradices* (T₂) and *G. fasciculatum* (T₃). The phosphorus was also non-significant

Comment [H9]: You may wish to separate the RESULTS FROM DISCUSSION

Comment [H10]: You need to enrich the discussion aspect. Its too scanty please . Give it a detailed scientific interpretation referring to similar or contrary findings

Fig 2: Soil DTPA extractable Fe, Zn, Cu and Mn content before sowing and after harvest of *Melia azedarach* seedlings

Table 1: Macro and micronutrient content in *Melia azedarach* seedlings after 180 days of sowing

Comment [H11]: Capitalize each word

Treatments		Nutrients content						
		N (%)	P (%)	K (%)	Fe (ppm)	Cu (ppm)	Zn (ppm)	Mn (ppm)
T ₁	<i>Glomus mosseae</i>	1.31	0.17	0.55	85.56	7.15	21.87	27.61
T ₂	<i>G. intraradices</i>	1.30	0.14	0.54	85.23	6.45	18.67	27.45
T ₃	<i>G. fasciculatum</i>	1.31	0.17	0.54	85.43	7.11	21.71	27.52
T ₄	<i>G. mosseae</i> + <i>G. intraradices</i>	1.34	0.20	0.58	86.81	7.82	24.97	28.83
T ₅	<i>G. mosseae</i> + <i>G. fasciculatum</i>	1.35	0.23	0.59	87.02	8.77	27.91	28.99
T ₆	<i>G. intraradices</i> + <i>G. fasciculatum</i>	1.34	0.20	0.58	86.27	7.78	24.91	28.76
T ₇	<i>G. mosseae</i> + <i>G. intraradices</i> + <i>G. fasciculatum</i>	1.39	0.27	0.63	90.43	9.96	32.24	31.28
T ₈	Control (un-inoculated)	1.27	0.10	0.51	77.79	5.58	15.45	25.12
CD at 5%		0.02	0.02	0.02	2.13	0.46	2.25	1.45

A significantly higher N, P and K uptake was recorded in six months old seedlings in T₇ (0.370, 0.11 and 0.19 (g plant⁻¹), respectively and minimum in control with 0.19, 0.01 and 0.08 (g plant⁻¹), respectively (Table 2). Caglar and Akgun (2006) reported that uptake of nitrogen in the leaf of *Pistacia* seedlings was enhanced upon inoculation with *Glomus* species. Among all the treatments Fe, Cu, Zn and Mn uptake were recorded maximum in interactive treatments of all species (T₇) (2.41, 0.27, 0.86 and 0.83 mg plant⁻¹, respectively) whereas, minimum was recorded in control (1.22, 0.09, 0.24 and 0.39 mg plant⁻¹, respectively). These result are in corroboration with the findings of many researchers viz., Jamaluddin and Shukla (2012), Muthukumar *et al.* (2012), Mohan and Sandeep (2015) and Ortiz *et al.* (2017) who reported an increase in nutrient uptake in different plant species with the application of different AM fungi species. The maximum Hyphal contribution (%) for P and Zn was found in T₇ followed by T₅ while under control there was no HC recorded (Table 3).

Table 2: Macro and micronutrient uptake in *Melia azedarach* after 180 days of sowing

Treatments		Nutrients uptake						
		g/plant			mg/plant			
		N	P	K	Fe	Cu	Zn	Mn
T ₁	<i>G. mosseae</i>	0.26	0.04	0.10	1.70	0.14	0.43	0.55
T ₂	<i>G. intraradices</i>	0.23	0.02	0.09	1.52	0.12	0.33	0.49

T ₃	<i>G. fasciculatum</i>	0.25	0.04	0.10	1.64	0.14	0.42	0.54
T ₄	<i>G. mosseae</i> + <i>G. intraradices</i>	0.30	0.06	0.13	1.91	0.17	0.55	0.60
T ₅	<i>G. mosseae</i> + <i>G. fasciculatum</i>	0.33	0.08	0.16	2.01	0.22	0.69	0.67
T ₆	<i>G. intraradices</i> + <i>G. fasciculatum</i>	0.29	0.06	0.13	1.88	0.17	0.54	0.63
T ₇	<i>G. mosseae</i> + <i>G. intraradices</i> + <i>G. fasciculatum</i>	0.37	0.11	0.19	2.41	0.27	0.86	0.83
T ₈	Control (un-inoculated)	0.19	0.01	0.08	1.22	0.09	0.24	0.39
CD at 5%		0.03	0.01	0.02	0.21	0.01	0.05	0.07

Table 3: Hyphal contribution (%) of AM fungi for Phosphorus and Zinc

Treatments		180 DAS	
		HC (%) for P	HC (%) for Zn
T ₁	<i>G. mosseae</i>	75.0	44.18
T ₂	<i>G. intraradices</i>	50.0	27.27
T ₃	<i>G. fasciculatum</i>	70.0	42.85
T ₄	<i>G. mosseae</i> + <i>G. intraradices</i>	83.3	56.36
T ₅	<i>G. mosseae</i> + <i>G. fasciculatum</i>	87.5	65.21
T ₆	<i>G. intraradices</i> + <i>G. fasciculatum</i>	83.3	55.55
T ₇	<i>G. mosseae</i> + <i>G. intraradices</i> + <i>G. fasciculatum</i>	90.9	72.09
T ₈	Control (un-inoculated)	00.0	0

CONCLUSION

The drupes of *M. azedarach* treated concomitantly with *G. mosseae*, *G. intraradices*, *G. fasciculatum* enhanced the soil nutrients especially P and Zn. The nutrients content in *M. azedarach* seedlings such as N (1.39 %), P (0.27 %), K (0.63 %), Fe (90.43 ppm), Cu (9.96 ppm), Zn (32.24 ppm) and Mn (31.28 ppm) were also found significantly higher in drupes sown in soil concomitantly treated with *G. mosseae*, *G. intraradices* and *G. fasciculatum*. The macro and micronutrient uptake was also found significantly higher in drupes sown in soil treated with all three *Glomus* species as cited above. Thus, it is concluded that the concomitant application of all three *Glomus* species have the potential to improve soil fertility, nutrient content and biomass of *M. azedarach*.

REFERENCES

- Afzal, M., Yousaf, S., Reichenauer, T.G., Kuffner, M. and Sessitsch, A. (2011). Soil type affects plant colonization, activity and catabolic gene expression of inoculated bacterial strains during phytoremediation of diesel. *Journal of Hazardous Materials* **186**(2): 1568-1575.
- Antil R.S., Singh, A. and Dahiya, S.S. (2002). Practical manual for soil and plant analysis. Department of Soil Science, CCS Haryana Agricultural University, Hisar-125004.
- Bainard, L.D., Klironomos, J.N. and Gordon, A.M. (2011). Arbuscular mycorrhizal fungi in tree-based intercropping systems: a review of their abundance and diversity. *Pedobiologia* **54**(2): 57-61.

- Bakhtiar, Y., Yahya, S., Sumaryono, W., Sinaga, M.S., Budi, S.W. and Tajuddin, T. (2010). Isolation and identification of Mycorrhizosphere bacteria and their antagonistic effects towards *Ganoderma boninense* in vitro. *Microbiology Indonesia* **4**(2): 9.
- Basumatary, N., Parkash, V., Tamuli, A.K., Saikia, A.J. and Teron, R. (2014). Arbuscular mycorrhizal inoculation affects growth and rhizospheric nutrient availability in *Hevea brasiliensis* (Willd. ex A. Juss.) Mull. Arg. clones. *International Journal of Current Biotechnology* **2**(7): 14-21.
- Blaser, W.J., Oppong, J., Hart, S.P., Landolt, J., Yeboah, E. and Six, J. (2018). Climate-smart sustainable agriculture in low-to-intermediate shade agroforests. *Nature Sustainability* **1**(5): 234-239.
- Caglar, S. and Akgun, A. (2006). Effects of vesicular- arbuscular mycorrhizal (VAM) fungi on the seedling growth of three *Pistacia* species. *Journal of Environmental Biology* **27**(3): 485-489.
- Cavagnaro, T.R., Bender, S.F., Asghari, H.R. and Van der Heijden, M.G. (2015). The role of arbuscular mycorrhizas in reducing soil nutrient loss. *Trends in Plant Science* **20**(5): 283-290.
- Chagnon, P.L., Bradley, R.L., Maherali, H. and Klironomos, J.N. (2013). A trait-based framework to understand life history of mycorrhizal fungi. *Trends in Plant Science* **18**(9): 484-491.
- Dhar, P.P. and Mridha, M.A.U. (2012). Arbuscular mycorrhizal associations in different forest tree species of Hazarikhil forest of Chittagong, Bangladesh. *Journal of Forestry Research* **23**(1): 115-122.
- Dierks, J., Blaser-Hart, W.J., Gamper, H.A., Nyoka, I.B., Barrios, E. and Six, J. (2021). Trees enhance abundance of arbuscular mycorrhizal fungi, soil structure, and nutrient retention in low-input maize cropping systems. *Agriculture, Ecosystems and Environment* **318**: 107487.
- Duponnoisa, R., Colombeta, A., Hienb, V. and Thioulouse, J. (2005). The mycorrhizal fungus *Glomus intraradices* and rock phosphate amendment influence plant growth and microbial activity in the rhizosphere of *Acacia holosericea*. *Soil Biochemistry* **37**(8): 1460-1468.
- Elwell, W.T. and Gridley, J.A.F. (1967). Atomic absorption spectrophotometer ypergamon press Ltd. London, W-1.
- Filho, J.A.V., Mendonça Freitas, M.S., Martins, M.A., dos Santos, P.C. and Cordeiro de Carvalho, A.J. (2017). Arbuscular mycorrhizal fungi and phosphate fertilization on star fruit tree seedlings. *Revista Brasileira de Ciencias Agrárias* **12**(1): 14-19.
- Goussous, S.J. and Mohammad, M.J. (2009). Comparative effect of two arbuscular mycorrhizae and N and P fertilizers on growth and nutrient uptake of onions. *International Journal of Agriculture and Biology* **11**: 463-467.
- Handa, A.K., Sirohi, C., Chavan, S.B., Dhillon, R.S., Ahlawat, K S. and Rizvi, R.H. (2020). Agroforestry in Haryana: Status and way forward. *Indian Journal of Agroforestry* **22**(1): 1-10.
- Hanway, J.J. and Heidel, H. (1952). Soil analysis methods as used in Iowa state college soil testing laboratory. *Iowa Agriculture* **57**: 1-31.
- Herrera-Medina, M. J., Steinkellner, S., Vierheilig, H., Bote, J.A.O. and Garrido, J.M.G. (2007). Absciscic acid determines arbuscule development and functionality in the tomato arbuscular mycorrhiza. *New phytologist* **175**(3): 554-564.
- India State of Forest Report (2021). Forest Survey of India, Ministry of Environment and Forests, released by Ministry of Environment Forest and Climate change.
- Jackson, M.L. (1973). Soil chemical analysis. *Prentice Hall of India*, pp. 498.
- Jamaluddin and Shukla, R. (2012). Effect of root exudates on colonization of AM fungi in *Jatropha curcas*. *Indian Forester* **138**: 113-115.
- Johnson, N.C. (2010). Resource stoichiometry elucidates the structure and function of arbuscular mycorrhizas across scales. *New Phytologist* **185**(3): 631-647.
- Koenig, R. and Johnson, C. (1942). Colorimetric determination of phosphorus in biological materials. *Industrial and Engineering Chemistry Analytical Edition* **14**(2): 155-156.
- Kothari, S.K., Marschner, H. and Römheld, V. (1991). Contribution of the VA mycorrhizal hyphae in acquisition of phosphorus and zinc by maize grown in a calcareous soil. *Plant and Soil* **131**(2): 177-185.
- Lindner, R.C. (1944). Rapid analytical methods for some of the more common inorganic constituents of plant tissues. *Plant Physiology* **19**(1): 76.
- Lindsay, W.L. and Norvell, W.A. (1978). Development of a DTPA soil test for zinc, iron, manganese, and copper. *Soil Science Society of America Journal* **42**(3): 421-428.
- Luyindula, N. and Haque, I. (1992). Effect of rhizobium inoculation and phosphorus on growth and nitrogen fixation in tree grown on vertisols. In K. Mulongoy, M. Gueye and D.S.F. Spencer. *Biological nitrogenfixation and sustainability of tropical Agriculture*, New York, pp 109 -113.
- Mohan, V. and Sandeep, C. (2015). Biochemical analysis and growth enhancement studies of important medicinal plant, *Rauvolfia serpentina* inoculated with Arbuscular Mycorrhiza fungi in nursery. *International Journal of Current Microbiology and Applied Sciences* **4**(6): 811- 820.

Comment [H12]: This is missing on the body of the work. Please check

- Muthukumar, T., Kathikeyan, A. and Udaiyan, K. (2012). Evaluation of the growth and nutrient uptake of *Casuarina equisetifolia* on alfisol soil inoculated with different arbuscular mycorrhizal fungi. In: Jayaraj, R.S.C., Warriar, R.R., Nicodemus, A. and Krishnakumar, N. (Eds.). *Advances in Casuarina Research in India: Proceedings of the 2nd National Seminar on Casuarina*. Institute of Forest Genetics and Tree Breeding, Coimbatore, India, pp. 49-58.
- Olsen, S.R. (1954). Estimation of available phosphorus in soils by extraction with sodium bicarbonate. *United States Department of Agriculture; Washington* 2: 25-43.
- Ortiz, Y.R., Bello, C.H.Á., Alarcón, A. and Cerrato, R.F. (2017). Effectiveness of native arbuscular mycorrhiza on the growth of four tree forest species from the Santa Marta Mountain, Veracruz (Mexico). *Forest Systems* 26(1): 3.
- Raven, J.A., Lambers, H., Smith, S.E. and Westoby, M. (2018). Costs of acquiring phosphorus by vascular land plants: patterns and implications for plant coexistence. *New Phytologist* 217(4): 1420-1427.
- Rillig, M.C., Aguilar-Trigueros, C.A., Bergmann, J., Verbruggen, E., Veresoglou, S.D. and Lehmann, A. (2015). Plant root and mycorrhizal fungal traits for understanding soil aggregation. *New Phytologist* 205(4): 1385-1388.
- Sandeep, C., Mohan, V. and Viswanath, S. (2015). Significance of ectomycorrhizae in forest ecosystems of India. *International Journal of Plant, Animal and Environment Sciences* 5: 23-31.
- Sato, T., Ezawa, T., Cheng, W. and Tawarayama, K. (2015). Release of acid phosphatase from extraradical hyphae of arbuscular mycorrhizal fungus *Rhizophagus sclarus*. *Soil Science and Plant Nutrition* 61(2): 269-274.
- Subbiah, B.V. and Asija, G.L. (1956). A rapid procedure for assessment of available nitrogen in soils. *Current Science* 25: 259-260.
- Troup, R.S. (1921). *The silviculture of Indian Trees*. Clarendon Press, Oxford, pp. 1195.
- Wartenberg, A.C., Blaser, W.J., Gättinger, A., Roshetko, J.M., Van Noordwijk, M. and Six, J. (2017). Does shade tree diversity increase soil fertility in cocoa plantations?. *Agriculture, Ecosystems and Environment* 248: 190-199.