

Original Research Article

Assessment of Physico-Chemical Properties on Soil of Vizianagaram District, Andhra Pradesh, India.

Abstract

The study was conducted at Sam Higginbottom University of Agriculture Technology and Sciences' Department of Soil Science and Agricultural Chemistry Lab. A total of 27 samples were collected from 3 blocks (Pusapatirega, Denkada and Bhogapuram), 9 villages, and analysed for physico-chemical properties using standard laboratory at [varying varied](#) depths (0-15, 15-30, and 30-45 cm). The soil colour ranged from yellowish red to yellowish brown (dry) and reddish brown to [yellowish-yellowish](#)-brown (wet). It had a red sandy loam texture. Bulk density $1.402 \text{ (Mg m}^{-3}\text{)}$, Particle density $2.603 \text{ (Mg m}^{-3}\text{)}$, Pore space 43.44(%), Water Holding Capacity 40.23(%), Specific gravity 2.42, pH 7.63 neutral, Electrical Conductivity $0.25 \text{ (dS m}^{-1}\text{)}$ non-saline, and Soil Organic Carbon 0.37(%). Available Nitrogen $269.88 \text{ (kg ha}^{-1}\text{)}$ is low, Available Phosphorous and Available Potassium 17.34 and 191.09 (kg ha^{-1}) are medium. Exchangeable Calcium $4.63 \text{ (cmol kg}^{-1}\text{)}$ and Exchangeable Magnesium $1.91 \text{ (cmol kg}^{-1}\text{)}$ medium to high. Farmers need to maintain a soil health card, adopt proper management techniques, and provide adequate nutrients to the soil in order to overcome pollution.

Keywords: Soil pH, EC, Available Nitrogen, Available Phosphorous, Available Potassium, Exchangeable Calcium and Magnesium, Soil Health.

INTRODUCTION

Soil is the cradle for all crops and plants, it is the most crucial part of every farming sector. Every 250-1200 years, non-renewable resources appear. It could take another 3000-12,000 years for agriculture to become productive. This natural resource is scarce and cannot be replenished within a human lifetime (Desavathu, *et al.* 2018). The thin layer of material that covers the earth's surface is referred to as soil. It is made up of Organic matter, minerals, gases, water, and living beings that all work together to sustain life. Water storage, nutrient supply, and purification are all important functions of soil (Gilluly James, 1975). Pollutants affect minerals, Organic matter and the microbial ecology of the soil. Industrial effluent

discharge, especially if it is not treated, can have a major impact on soil physico-chemical and biological features that are linked to soil fertility (Pushpanjali, *et al.* 2017). Because the physico-chemical qualities of soil affected food productivity and environmental quality, it's critical to have a fundamental understanding of these properties (Tale, *et al.* 2015).

Physical attributes include soil texture, Bulk density, Particle density, Pore space-, Water Holding Capacity, Soil structure and Soil colour. Chemical qualities include pH, EC, OC, Nitrogen, Phosphorus, Potassium, Exchangeable Calcium and Magnesium. Soil health is inextricably tied to soil microbial diversity and activity, which are important components of soil health. Healthy soils are essential for long-term development, not only for increased agricultural production as the world's population expands, but also for the long-term survival of essential ecosystem services (Sahu, *et al.* 2021).

MATERIALS AND METHODS

The present study, entitled – “ Assessment of Physico-chemical properties of Soil in Vizianagaram district, Andhra Pradesh, India” was carried out during 2021-22 and comprised of a lab experiment which was carried out in [the](#) Department of Soil Science and Agricultural Chemistry, Naini, Agricultural institute, Sam Higginbottom University of ~~Agrielture~~[Agriculture](#) Technology and Sciences, Prayagraj (U.P.), India.

Study area

The district is located in Andhra Pradesh's Northern Coastal plains and covers geographical area 6,539 Sq. kms. ~~between~~[Between](#) 17°-15¹ and 19°-15¹ North latitude and 83°-0⁰ to 83°-45¹ East longitude. To the north, the state of Orissa, to the west and south, the district of Srikakulam and to the south, the Bay of Bengal. The district's agriculture is rainfed, with an average annual rainfall of 1131mm. The main crops grown in the district are Paddy, Ragi, Bajra, Sugarcane, Pulses, Mesta, Cotton and Groundnut.

Soil Sampling

Total [of](#) 27 samples are collected from 3 blocks (Pusapatirega, Denkada, Bhogapuram) and 9 villages at different depths of 0-15cm, 15-30 cm, ~~and~~ [30-45cm](#) ~~in the~~[of](#) Vizianagaram district. At first foreign were removed from [the](#) sampling place and 'V'-shaped pit was dug out with the help of spade / khurpi / crowbar. After that, the soils were properly mixed before being divided into four equal pieces. After eliminating two opposite quarters, the two remaining soil quarters were mixed up. Before being delivered to the

laboratory, the soil was packed in a poly bag and labelled with the sample number, sample location, area, farmer information, crop data and sampling date. The soil samples were dried in the shade and manually crushed with a mallet before being sieved in the laboratory with a 2mm sieve [mesh](#). The acquired material's physical and chemical characteristics were investigated.

ANALYSIS OF PHYSICO-CHEMICAL PRARAMETERS

The hydrometer method was used to analyse the texture of soil particles less than 2mm (Bouyoucos, 1927) and the Soil Colour was tested by using [the](#) Munsell colour chart (Munsell, A. H. 1921). By using [a](#) graduated 100ml measuring cylinder, the bulk density, Particle density, Pore space, and Water Holding Capacity were calculated (Muthuvel, *et al.* 1992). The Specific gravity of soil was determined using the relative density bottle or pycnometer method (Black, 1965). A digital pH metre and the 1:2.5 soil-water suspension method were used to determine the EC (Wilcox, 1950). Wet oxidation was used to measure Organic Carbon (Walkley and Black, 1947). Using the alkaline Potassium permanganate method, the Available Nitrogen was measured (Subbiah and Asija, 1956). Using the colorimetric approach, the Available Phosphorus was determined (Olsen, *et al.* 1954). The flame Photometer method was used to determine the amount of Available Potassium (Toth and Prince, 1949). A digital EC metre and the 1:2 Soil-Water suspension method were used to determine the pH. The neutral Ammonium Acetate extraction method or the EDTA method was used to determine Exchangeable Calcium and Magnesium (Cheng and Bray, 1951).

STASTICAL ANALYSIS

The analysis of variance (ANOVA) approach was used to statistically analyse the data obtained throughout the study (Fisher, 1960). The experiment used ANOVA with two components and no replication. The analysis used a completely randomised experiment design (CRD). It is utilised when the experimental units are homogeneous since it just requires two basic experimental design principles: replication and randomization. Researchers used the F'9 variance ratio test to determine which treatment effects were significant and which were not.

RESULTS AND DISCUSSION

Physical Properties

Yellowish red to Pale brown, yellowish red to yellowish brown and yellowish red to yellowish brown were among the soil colours (dry condition). The soil was a mix of reddish brown and yellowish red, yellowish red and yellowish brown, yellowish red and yellowish brown (wet condition). The texture was red sandy loam, with sand content ranging from 62.56 to 83.78%, silt from 19.16 to 42.15%, and clay from 5.42 to 16.04% and the similar findings were reported in (Raju, 2015). The Bulk density ranged from 1.33 to 1.50 (Mg m^{-3}), with a mean of 1.402 (Mg m^{-3}). In (Karanam, 2020), non-significant changes related to depth and site were discovered. As soil depth increases, Bulk density increases. The Particle density ranged from 2.54 to 2.69 (Mg m^{-3}), with 2.603 (Mg m^{-3}) being the average. Non-significant differences related to depth and significant differences due to site were found in (Karam, 2020). Particle density is influenced by the mineral content of soil particles (Yuvarani, *et al.* 2019). Pore space (%) ranged from 26.39 to 50.78 %, with 43.44% as the mean. Significant differences were found related to depth and site, and similar findings were reported in (Srilakshmi, *et al.* 2019). Water Holding Capacity (%) ranged from 32.73 to 48.41%, with 40.23% as the mean. Significant differences were found related to depth and site, and similar findings were reported in (Venkateswarlu, *et al.* 2020). Because of soil compaction and pore space contraction, WHC values decrease as depth increases. The Specific gravity ranged from 2.16 to 2.61, with 2.42 as the mean. A significant difference was found due to depth and site and similar findings were reported in (Karnam, 2020).

Chemical Properties

The pH of the soils ranged from 6.65 to 8.63, indicating neutral to slightly alkaline conditions, with a mean of 7.63. The change in depth isn't considerable, but the difference in location. Because the top horizons absorb the most leaching from rainfall and dissolved carbonic acids, as well as a high volume of Exchangeable Sodium ions, the pH value rises with depth. Similar research was conducted by (Okolo, *et al.* 2016). Non-saline soils were identified by Electrical Conductivity ranging from 0.17 to 0.36 (dS m^{-1}), with a mean of 0.25 (dS m^{-1}). The Soil Organic Carbon (%) ranged from 0.25 to 0.52 %, suggesting productive soils, with 0.37% being the mean. Because surface soil includes undecomposed and partially degraded Organic Matter, whereas subsoil contains decomposed Organic Matter that has undergone chemical and biological changes, Organic Carbon declines with depth (Singh, *et al.* 2012).

Available Nitrogen ranged from 215.00 to 322.00 (kg ha⁻¹), with a mean of 269.88 (kg ha⁻¹). Because Available Nitrogen is positively connected with Organic Matter concentration, which reduces with depth, and may be owing to greater pH at deep, Available Nitrogen decreases with depth. Similar research was conducted by (Ho, *et al.* 2019). Available Phosphorous ranged from 12.64 to 24.82 (kg ha⁻¹) with increasing depth, the amount of Available Phosphorus decreases. The presence of more accessible Phosphorous in surface soil may be due to a favorable soil pH and Organic Matter concentration. Similar research was conducted by (Wani, *et al.* 2017). The Available Potassium ranged from 136.62 to 235.65(kg ha⁻¹) with a mean of 191.09 (kg ha⁻¹). The amount of Available Potassium reduces as you go deeper into the earth. The release of Available Potassium from Organic residues and the application of Potassium fertilizers may be responsible for the high quantity of accessible Potassium on [the](#) surface soil. A similar result was found by (Singh, *et al.* 2020). The ~~of~~-Exchangeable Calcium ranged from 3.23 to 5.66 (cmol kg⁻¹) with a mean of 4.63 (cmol kg⁻¹). The content of Exchangeable Magnesium ranged from 1.24 to 2.69 (cmol kg⁻¹) with a mean of 1.91 (cmol kg⁻¹) Pushpanjali. (2017), Prasadini (1995), Raju (2015) and Ramana, et al. (2015) all found significant differences in depth and site.

Comment [K1]: Please reference to table

Soil Health Card for the farmers of Vizianagaram district-

A Soil health card is being produced for the ~~villages'~~[villages's](#) farmers. This would aware them, about the fertility status of their farm soil, help them to make practical decisions in adopting the farming techniques, [and](#) guide them to incorporate suitable doses of fertilizers (nutrients) based on the soil test there by improving soil productivity and crop yield leading towards increasing farm income.

CONCLUSION

The present investigation was done on the soils of [the](#) -Vizianagaram district, and it revealed that the soils were red sandy loam soils with a considerable amount of Bulk density and good physical condition. It has a neutral pH and all soil samples are non-saline, making it good for all crops. Available Nitrogen is low because soil Organic Carbon is low. Available Phosphorous and Available Potassium are both available in moderate amounts. Exchangeable Calcium and Magnesium levels are medium, while macronutrients ~~s~~ levels are low to medium. At some locations, [the](#) secondary nutritional deficiency was discovered. Nutrient deficits can be alleviated by using organic and in-organic fertilisers. It demonstrates that the soils are suitable for Paddy, Ragi, Bajra, Sugarcane, Pulses, Mesta, Cotton and Groundnut agriculture.

Farmers must keep a Soil Health Card in accordance with central, ICAR, and state government recommendations for crop production, and are advised to use appropriate management strategies and supply proper nourishment to soil health.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES:

- Black, C. A. Methods of soil analysis. American Society of Agronomy, Madison, Wisconsin, USA, 1965;2.
- Bouyoucos, G.J., (1927) The hydrometer as a new method for the mechanical analysis of soils. *Soil Science* 23, 343-353
- Cheng, K.L. and Bray, R. H. (1951). Determination of Calcium and Magnesium in soil and plant material. *Soil*; 72:449-458.
- Desavathu, R. N., Nadipena, A. R., Peddada, J, R (2018) Assessment of soil fertility status in Paderu mandal, Visakhapatnam district of Andhra Pradesh through. Geospatial techniques; *The Egyptian Journal of Remote Sensing and Space Sciences*, 21: 73-81.
- Fisher, R. A. (1960). Statistical methods and scientific induction. *Journal of the royal statistical society series*. 17:69-78.
- Gilluly, J., Waters, A. C., Woodford, A. O. (1975). *Principles of geology (4th ed.)*. San Francisco, California: W.H. Freeman. ISBN 978-0-7167- 0269-6.
- Ho, S. Y., Wasli, M. E. B., Perumal, M. (2019). Evaluation of Physicochemical Properties of Sandy- Textured Soils under smallholder Agricultural Land Use Practices in Sarawak, East Malaysia. *Applied and Environmental Soil Science*.
- Karanam, H.K. (2020) Integrated Methodology of Geoengineering Investigation for Determination of Subsurface Lithology and Suggest Founding Levels of the Civil Structures at Road Over Bridge Locations - A Case Study in Vizianagaram District, Andhra Pradesh. *International Journal of Management, Technology and Engineering*.8(12).

- Munsell, A. H.(1921) Munsell's description of his colour system, from a lecture to the American Psychological Association. *American journal of psychology* 23, 236- 244.
- Muthuval, P., Udaysoorian, C., R. Natesan, R., Ramaswami, P.R. (1992) Introduction to Soil Analysis, Tamil Nadu Agricultural University, Coimbatore-641002.
- Okolo, V. N., Olowolafe, E. A., Akawu, I., Okoduwa, S. I. R. (2016). Effects of Industrial Effluents on Soil Resource in Challawa Industrial Area, Kano, Nigeria. *Journal of Global Ecology and Environment*, 5(1):1-10.
- Olsen, S.R., Cole, C.V., Watanhe, F.S. and Dean, L.A. (1954) Estimation of Available Phosphorus in soils by extraction with sodium bicarbonate *U.S. Dept. Agr. Circ.*, 939.
- Prasadini, R.P., Rao, M.S. (1995) Soils of Srikakulam Properties and Problems. Andhra Pradesh Agricultural University Hyderabad.
- Pushpanjali, Tejaswi, P.B., Sharma, K.L., Mandal, U.K., Raju, A.J.S., Venkanna, K., Masanne, R., Samuel, J. and Karthikeyan, K. (2017) Soil Quality and Heavy Metal Contamination of Soils in Mindi industrial Area, Visakhapatnam, Andhra Pradesh, India. *Indian Journal of Ecology* ,44(4):774-777.
- Raju, N. A. (2015) Land Capability and Suitability in Vizianagaram District of Andhra Pradesh Using Remote Sensing and GIS Techniques. *Journal Of Humanities and Social Science*. 20(7):56-64.
- Sahu, G. and Mohanty, S. (2021). Soil Health and Sustainability *Current research in soil science*, 9(4): 978- 93.
- Singh, B. and Sharma, K. N. (2012). Depth Wise Distribution of Soil Organic Carbon and Nutrients under some Tree Species after Seventeen Years of Plantation. *Journal of the Indian Society of Soil Sciences*, 60(3):198-203.
- Singh, D., Singh, A. K., Singh, A. K., Guptha, S. K. (2020). Charaterization of rice growing soil of Nagara block of Ballia District (U.P.), India. *International Journal of Current Microbiology and Applied Sciences*, 9(04):575-581.
- Srilakshmi, M., Thomas, T., Rao, P.S. (2019) Assessment of soil physico- chemical properties of nearby coastal area paddy soils of Prakasam district, Andhra Pradesh. *International Journal of Chemical Studies*. 7(3): 1605-1608.
- Subbiah, B. V. and Asija, E.C., (1956) A rapid procedure for estimation of available Nitrogen in soil *Current Science* 25(8), 259- 260.

- Tale, K. S. and Ingole, S. (2015). A Review on Role of Physico-Chemical Properties in Soil Quality. *Chem Sci Rev Lett*, 4(13), 57 – 66.
- Toth, S. J., and Prince, A. L. (1949). Estimation of cation exchange capacity and exchangeable Ca, K and Na content of soil by flame photometer technique. *Soil Sci.* 67:439-445.
- Venkata Ramana, C.H., Bhaskar, C.H., Prasada Rao, P.V.V., Byragi Reddy, T. (2015). Soil quality in four different areas of Visakhapatnam city, Andhra Pradesh, India. *International Journal of Current Microbiology and Applied Sciences*. 4(1):528–532.
- Venkateswarlu, V. And Venkatrayulu, C. (2019) Assessment of Heavy Metals in Water of Selected Places of Nellore Coast of Andhra Pradesh Through Atomic Absorption Spectroscopy. *ISSN: 2581-8740*.
- Walkley, A. And Black, I. A., (1934) Estimation of soil Organic Carbon by the chronic acid titration method. *Soil Science*. 47: 775-776.
- Wani, S. A., Najar, G. R., Padder, B. A., Akhter, F., Chand, S. (2017). Altitudinal and depth-wise variation of Soil Physico-Chemical Properties and available Nutrients of Pear Orchards in Jammu and Kashmir, India. *Chemical Science Review and Letters*, 6(23): 1638-1645.
- Wilcox, L. V. (1950) Electrical conductivity Am. *Water works, Assoc. J.* 42: 775-776.
- Yuvarani, R., Swarrop, N., Rao, P. S. and Thomas, T. (2019). Impact of tannery effluent on physical properties of soils of Dindigul district of Tamil Nadu state. *International Journal of Chemical Studies*, 7(3): 420-422.




Fig 1: Bulk density (Mg m⁻³), Particle density (Mg m⁻³), Porosity (%), WHC (%) and Specific gravity are all represented graphically

Comment [K2]: Need to be separate. The unit is not the same
And also show the adequate amount of each factors




Fig 2: Graphical representation of pH, EC (dS m⁻¹), Organic Carbon (%)

Comment [K3]: Need to be separate
And also show the adequate amount of each elements and compare to the result




Fig 3: Nitrogen (kg ha⁻¹), Phosphorus (kg ha⁻¹), Potassium (kg ha⁻¹), Exchangeable Calcium and Magnesium (cmol kg⁻¹) are represented graphically.

Comment [K4]: Need to be separate
Need to be separate
And also show the adequate amount of each elements and compare to the result

Table 1: Assessment of Db, Dp, Porosity, WHC and Specific Gravity of Vizianagaram district

Sample No.	Bulk Density(Mg m ⁻³)			Particle Density (Mg m ⁻³)			Porosity(%)			Water Holding Capacity (%)			Specific Gravity		
	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm
B ₁ V ₁	1.4	1.43	1.46	2.61	2.64	2.68	41.37	35.98	33.2	43.14	40.72	38.64	2.42	2.37	2.33
B ₁ V ₂	1.42	1.45	1.48	2.56	2.58	2.6	42.57	34.49	30.38	40.31	38.62	36.91	2.28	2.25	2.16
B ₁ V ₃	1.43	1.46	1.5	2.65	2.67	2.69	42.64	35.2	26.39	39.86	36.91	32.73	2.68	2.64	2.59
B ₂ V ₁	1.39	1.41	1.41	2.56	2.59	2.64	49.6	48.64	45.83	40.06	37.91	36.13	2.31	2.28	2.18
B ₂ V ₂	1.37	1.39	1.4	2.55	2.58	2.61	50.19	48.44	44.44	42.89	40.05	38.78	2.4	2.36	2.31
B ₂ V ₃	1.33	1.35	1.39	2.54	2.57	2.6	50.78	49.02	46.53	48.41	46.75	41.47	2.52	2.47	2.44
B ₃ V ₁	1.42	1.44	1.46	2.56	2.6	2.63	47.26	45.38	41.44	44.76	42.13	39.01	2.57	2.53	2.49
B ₃ V ₂	1.43	1.45	1.48	2.55	2.59	2.62	48.62	48.64	44.65	43.87	41.72	39.43	2.61	2.56	2.5
B ₃ V ₃	1.43	1.46	1.49	2.58	2.61	2.64	50	49.8	41.66	40.16	38.91	36.13	2.48	2.43	2.39
		S.Ed.			S.Ed.			S.Ed.			S.Ed.			S.Ed.	
	F-test	(+)	C.D.@5%	F-test	(+)	C.D.@5%	F-test	(+)	C.D.@5%	F-test	(+)	C.D.@5%	F-test	(+)	C.D.@5%
Due to depth	NS	-	-	NS	-	-	S	3.831	1.44	S	2.461	3.08	S	0.048	0.00026
Due to site	NS	-	-	S	0.032	1E-10	S	5.863	8.83	S	2.704	1.27	S	0.136	4.41

Comment [K5]: Show it in the text

Comment [K6]: Need to be explained the below of table for interpreted

Table 2: Assessment of pH, EC, Organic Carbon, Nitrogen, Phosphorus, Potassium, Calcium, Magnesium of Vizianagaram district

Sample No.	pH			EC (dS m ⁻¹)			Organic Carbon (%)			Nitrogen (kg ha ⁻¹)			Phosphorus (kg ha ⁻¹)			Potassium (kg ha ⁻¹)			Calcium (cmol kg ⁻¹)			Magnesium (cmol kg ⁻¹)		
	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm	0-15 cm	15-30 cm	30-45 cm
B ₁ V ₁	6.65	6.95	7.03	0.25	0.27	0.32	0.32	0.3	0.27	243	237	230	17.23	15.47	13.87	212.98	197.65	172.14	3.82	3.66	3.23	1.68	1.42	1.24
B ₁ V ₂	6.7	6.94	7.1	0.23	0.23	0.29	0.3	0.28	0.26	255	248	241	19.82	16.67	14.53	184.45	178.04	152.87	4.25	4.04	3.68	1.76	1.54	1.4
B ₁ V ₃	7.7	7.77	7.9	0.3	0.33	0.36	0.31	0.28	0.25	234	226	215	18.12	14.87	12.64	167.05	146.23	136.62	4.16	4.02	3.74	1.72	1.63	1.44
B ₂ V ₁	8.46	8.56	8.63	0.22	0.25	0.29	0.44	0.4	0.37	289	275	269	22.07	19.46	17.72	198.75	182.16	176.34	5.66	5.39	5.02	2.18	1.86	1.74
B ₂ V ₂	8.13	8.17	8.19	0.2	0.23	0.26	0.47	0.43	0.4	276	264	258	20.63	16.69	14.37	202.75	184.19	175.09	5.37	5.14	4.96	2.69	2.44	2.18
B ₂ V ₃	7.48	7.72	7.85	0.17	0.2	0.22	0.52	0.49	0.46	322	293	286	24.82	20.71	18.63	235.65	208.02	191.31	5.45	5.11	4.84	2.38	2.18	1.94
B ₃ V ₁	7.36	7.47	7.63	0.19	0.21	0.24	0.45	0.41	0.39	304	289	276	19.98	18.07	15.04	226.07	196.94	172.18	5.18	4.93	4.76	2.24	2.14	2.02
B ₃ V ₂	7.43	7.75	7.81	0.2	0.23	0.25	0.41	0.38	0.35	316	302	282	17.26	16.41	13.8	212.63	209.13	192	5.04	4.84	4.62	2.14	2.06	1.9
B ₃ V ₃	7.54	7.55	7.58	0.23	0.27	0.33	0.46	0.43	0.4	297	286	274	18.84	16.73	14.06	230.34	219.65	198.46	4.95	4.74	4.41	2.02	1.93	1.89
	F-test	S.Ed.	C.D.@5%	F-test	S.Ed.	C.D.@5%	F-test	S.Ed.	C.D.@5%	F-test	S.Ed.	C.D.@5%	F-test	S.Ed.	C.D.@5%	F-test	S.Ed.	C.D.@5%	F-test	S.Ed.	C.D.@5%	F-test	S.Ed.	C.D.@5%
		(+)			(+)			(+)			(+)			(+)			(+)			(+)			(+)	
Due to depth	NS	-	-	S	0.0318	2.31	S	0.076	3.428	S	11.421	1.07	S	11.421	1.07	S	16.87	3.04	S	0.257	1.41	S	0.17	0.013
Due to site	S	3.391	0.00002	S	0.0405	1.041	S	0.0294	1.762	S	27.71	3.16	S	27.719	3.16	S	20.62	2.74	S	0.63	1.89	S	0.324	0.001

Comment [K7]: Need to be explained the below of table for interpreted

Comment [K8]: Show it in the text