

## Impact of Integrated Nutrient Management on Physico-chemical properties of soil in Pea (*Pisum sativum* L.) var. GS 10

### Abstract

During the *Rabi* season of 2021-2022, the field experiment was conducted at the soil science research farm of the Sam Higginbottom University of Agriculture, Technology and Sciences in Prayagraj, Uttar Pradesh, India. Three different parameters viz., three levels of NPK, FYM at 0%, 50%, and 100% ha<sup>-1</sup>, and three levels of *Rhizobium* inoculation at 0%, 25%, and 50% ha<sup>-1</sup> were used in the study. The result obtained with treatment T<sub>9</sub>[I<sub>3</sub>@100% + F<sub>3</sub>@ 100% + R<sub>3</sub>@50%] had showed a Bulk density (1.17 Mg m<sup>-3</sup>) at 0-15cm and (1.18 Mg m<sup>-3</sup>) at 15-30 cm, Particle density (2.41 Mg m<sup>-3</sup>) at 0-15cm and (2.42 Mg m<sup>-3</sup>) at 15-30, Pore space (58.26%) at 0-15cm and (58.09%) at 15-30cm, Water holding capacity (58.60%) at 0-15cm and (58.13%) at 15-30, pH (7.75) at 0-15cm and (7.75) at 15-30cm, EC (0.47dSm<sup>-1</sup>) at 0-15cm and (0.48 dSm<sup>-1</sup>) at 15-30cm, soil Organic Carbon (0.58%) at 0-15cm and (0.49%) at 15-30cm. As regards soil available nutrients, Available nitrogen (280.86 kg ha<sup>-1</sup>) at 0-15cm and (286.40) at 15-30cm, Available phosphorus (16.56 kg ha<sup>-1</sup>) at 0-15cm and (17.26 kg ha<sup>-1</sup>), Available potassium (178.13 kg ha<sup>-1</sup>) at 0-15cm and (172.80 kg ha<sup>-1</sup>). Under control and full NPK fertilizer application, no significant differences in Pea yield and growth were detected. The use of FYM and *Rhizobium*, as well as its blend with complete NPK, significantly improves the growth and overall production of Pea.

**Keywords:** Physico-chemical Properties of soil, N<sub>2</sub>P<sub>2</sub>K, FYM, *Rhizobium*, Pea, Yield attributes

### Introduction

Pea (*Pisum sativum* L.) belongs to Fabaceae family and has a chromosome number 2n= 14. Peas are native to Central or Southeast Asia and are grown all around the world. In India, Garden pea is grown as a winter vegetable crop in hilly and plain areas. It is very rich source of protein, carbohydrates, vit. A & C, calcium, phosphorus whose nutritive value of fresh green pea per 100g contain Energy 339KJ, Dietary fiber 5.1g, Protein: 5.42 g, Carbohydrates: 14.45 g, Sugars: 5.67 g, Fat: 0.4 g, Vitamin C: 40 mg, Folic acid: 50.7 mg, Iron: 1.47 mg, Potassium: 217 mg, Magnesium: 33 mg, and Phosphorus: 108 mg. Temperature favorable for grown of pea is 15-25°C. It can be cultivated in various types of soil condition like loam, sandy loam to clay soil. Pea needs well drained, loose and friable

soil condition. Pea does not thrive on acidic condition and very sensitive to saline and alkaline soil. The pH ideal for it is 6.0-7.0.

Chemical fertilizers provide instant nutrient supply and to get good crop yield but it can have harmful effect for environment. To sustain soil fertility and productivity, it's critical to utilize a combination of inorganic, organic, and biofertilizers. Biofertilizers improve soil fertility by symbiotically fixing atmospheric nitrogen with plant roots, solubilizing insoluble soil phosphates, and producing necessary plant development chemicals. The organic source of manure like FYM, is a type of bulky organic manure that is widely employed. Since it is derived from plant and animal residues therefore, it supplies the entire nutrient to the plant in easily available form in slow mineralization. It has a significant positive impact on soil's physical, chemical, and biological qualities besides, it pulverising the soil and improves the structure of the soil. Thus, an integrated strategy to nutrient supply that includes chemical fertilisers, organic manure, and biofertilizer which not only minimises inorganic fertiliser consumption, but also improves soil health and is ecologically friendly. This study aimed to evaluate the effect of integrated application of biofertilizer, organic manure and inorganic fertilizers on pea in terms of physico-chemical properties.

## Materials and Method

The field investigation was carried out with garden pea variety GS-10 during Rabi season, 2021 in the research area Department of Soil science and Agricultural Chemistry, SHUATS, Prayagraj, located at 25°24'30" North latitude, 81°51'10" East longitude and 98 m above mean sea level. The experimental soil in the experimental area is classified as Inceptisol, which and the soil in the experimental plots is alluvial in character. The soil texture (% of sand, silt, and clay) of the departmental research farm, with soil samples taken at depths of 0-15cm and 15-30cm. The soil had a sandy loam texture, with 55% sand, 30% silt, and 15% clay. The soil color (dry and wet method) sample was taken on depth 0-15cm and the soil color- yellowish brown was found at dry condition and at wet condition the soil color- brown was found and on depth 15-30cm the soil color- light yellowish brown was found at dry condition and at wet condition the soil color- yellowish brown was found. FOR ANALYSIS of physic-chemical properties when the soil samples were drawn???? A The trial used a randomised block design (RBD) with three replications and nine treatments,

Formatted: Font: Italic

TREATMENT DETAILS NOT CLEAR

NINE TREATMENTs T1 to T9.

What is T1, T2 ..... T9.

Indicate clearly

using varied levels of FYM (0, 50, and 100 percent) and Rhizobium inoculation (0, 25 and 50 percent).

On what basis rhizobium was applied?Quantity

Rewrite treatments and methodology clearly

varied levels of FYM (0, 50, and 100 percent means what? Whether on RDF if so what quantity?

Basal doses of nitrogen, phosphorus and potassium are applied to the field where RDF was found 30:50:50 NPK kg ha<sup>-1</sup>. ~~The sources of NPK were Urea, SSP, MOP.~~ FYM were applied at their recommended dose 5 t ha<sup>-1</sup> of soil depth 0-15cm and 15-30 cm both were taken for analysis of soil physico-chemical properties.

Chart 1: Soil texture property

Particulars	Method employed	Results
Sand (%)	Bouyoucos Hydrometer (1927)	55%
Silt (%)		30%
Clay (%)		15%
Textural class	Munsell color chart	Sandy loam
Soil Colour		Yellowish brown
Bulk density (Mg m <sup>-3</sup> )	Graduated measuring cylinder method Muthuval <i>et al.</i> , (1992)	1.24%
Particle density (Mg m <sup>-3</sup> )		2.48
Pore Space (%)		51.56%

Chart 2: Study methods

Particulars	Method employed	Results
Soil pH (1:2)	Jackson (1958)	7.41
Soil EC (dSm <sup>-1</sup> )	Wilcox (1950)	0.34
Organic Carbon (%)	Walkley and Black's (1947)	0.51

Formatted: Font color: Red

Available Nitrogen (kg ha <sup>-1</sup> )	Subbiah and Asija (1956)	239.96
Available Phosphorus (kg ha <sup>-1</sup> )	Olsen <i>et al.</i> (1950)	14.06
Available Potassium (kg ha <sup>-1</sup> )	Toth and Prince (1949)	158.28

## Results and Discussion

Effects of various treatment combinations on soil physical properties

During the trail of field experiment, a perusal of data reveals the application of NPK, FYM and Rhizobium inoculation was observed that Bulk density ranged from 1.24 Mg m<sup>-3</sup> to 1.17 Mg m<sup>-3</sup> at 0-15cm soil depth and 1.26 Mg m<sup>-3</sup> to 1.18 Mg m<sup>-3</sup> at 15-30cm soil depth. In both soil depths, the lowest bulk density was found in T<sub>9</sub>. Soil Particle density ranged from 2.48 Mg m<sup>-3</sup> to 2.41 Mg m<sup>-3</sup> in 0-15cm and 2.50 Mg m<sup>-3</sup> to 2.42 Mg m<sup>-3</sup> in 15-30cm soil depth. FYM impact on particle density positively means lowest particle density observed in T<sub>9</sub>. Porosity ranged from 51.56% to 58.26% and Water holding capacity ranged from 52.73% to 58.60% in 0-15cm soil depth respectively. However, as soil depth increased, porosity and water holding capacity decreased, resulting in a range of 50.66 % to 58.09 % porosity and 51.46 % to 58.13 % water holding capacity at 15-30 cm. Similar results were also reported by Yadav *et. al.*, (2018), Bhambhu *et al.*, (2016). [What is T<sub>9</sub>](#).

[Compare with control and other treatments. The author discussed only T<sub>9</sub>.](#)

[The parameter. Pore space was discussed and data presented](#)

Effects of various treatment combinations on soil chemical properties

The application of NPK, FYM and Rhizobium inoculation significantly, affected the soil pH at 0-15cm and 15-30cm soil depth. A Minimum soil pH was recorded in under the treatment T<sub>1</sub> i.e., 100% NPK + % 100FYM + 50% Rhizobium. EC (dSm<sup>-1</sup>) was influenced significantly it is ranged from 0.34 dSm<sup>-1</sup> to 0.47 dSm<sup>-1</sup>. A Maximum EC was recorded in T<sub>9</sub> and Minimum in T<sub>1</sub>, recorded in both the soil depths. Soil % Organic carbon content was maximum found in T<sub>9</sub> with a value of 0.58% and 0.49% in 0-15cm and 15-30cm soil depth, respectively it was followed by T<sub>7</sub>. The Available Nitrogen content in soil ranged from 239.96 kg ha<sup>-1</sup> to 280.86 kg ha<sup>-1</sup> at 0-15cm soil depth and 234.26 kg ha<sup>-1</sup> to 286.40 kg ha<sup>-1</sup> at 15-30cm soil depth. While Maximum was recorded in T<sub>9</sub> and minimum in T<sub>1</sub>, recorded in both soil depths. Available Phosphorus at 0-15cm soil depth, phosphorus levels ranged from 14.06 kg ha<sup>-1</sup> to 16.56 kg ha<sup>-1</sup> while at 15-30cm soil depth it was 13.70 kg ha<sup>-1</sup> to 17.26 kg ha<sup>-1</sup> and T<sub>9</sub> had the most accessible phosphorus in both soil depths, owing to

increased soil organic carbon, which boosted the activity of phosphorus solubilizing microorganism in the soil. The maximum available potassium in 0-15cm and 15-30cm soil depth i.e., 158.2 kg ha<sup>-1</sup> and 178.1 kg ha<sup>-1</sup> and 156.7 kg ha<sup>-1</sup>, 172.8 kg ha<sup>-1</sup> respectively (which was at par with T<sub>8</sub> and T<sub>7</sub>) followed by T<sub>6</sub> (which was at par with T<sub>5</sub> and T<sub>4</sub>) followed by T<sub>3</sub> (which was at par with T<sub>2</sub> and T<sub>1</sub>) in both soil depth by Rhizobium inoculation, FYM and NPK application. Similar results were also reported by Sharma and Thakur *et. al.*, (2016).


UNDER PEER REVIEW

Table 1: Effect of NPK, FYM and Rhizobium on Physico-chemical properties of post-harvest soil of Pea

Soil Parameters →	Bulk density (Mg m <sup>-3</sup> )		Particle density (Mg m <sup>-3</sup> )		Pore space (%)		WHC (%)		pH (1:2)		EC (dSm <sup>-1</sup> )		OC (%)		Nitrogen (Kg ha <sup>-1</sup> )		Phosphorus (Kg ha <sup>-1</sup> )		Potassium (Kg ha <sup>-1</sup> )	
	0-15	15-30	0-15	15-30	0-15	15-30	0-15	15-30	0-15	15-30	0-15	15-30	0-15	15-30	0-15	15-30	0-15	15-30	0-15	15-30
Depths (cm) → Treatments ↓																				
T1	1.24	1.26	2.48	2.50	51.56	50.66	52.73	51.46	7.41	7.49	0.34	0.35	0.51	0.40	239.96	234.26	14.06	13.70	158.2	156.7
T2	1.23	1.25	2.47	2.49	52.65	51.60	53.40	52.96	7.53	7.48	0.35	0.38	0.53	0.43	242.90	237.86	14.66	14.20	164.3	163.2
T3	1.22	1.23	2.47	2.48	53.55	52.66	54.50	53.36	7.57	7.52	0.35	0.38	0.53	0.44	247.56	240.70	15.26	14.63	167.2	164.6
T4	1.22	1.23	2.45	2.46	54.58	53.33	55.43	54.43	7.52	7.52	0.40	0.41	0.54	0.44	250.50	245.36	15.20	15.00	167.6	166.4
T5	1.22	1.22	2.45	2.46	55.54	54.47	55.50	55.40	7.69	7.68	0.40	0.42	0.55	0.45	254.96	262.30	15.43	15.53	168.6	166.6
T6	1.21	1.20	2.45	2.45	56.32	55.61	56.66	55.90	7.70	7.70	0.41	0.42	0.56	0.47	259.16	264.30	15.80	15.86	171.9	167.2
T7	1.20	1.20	2.44	2.44	56.81	56.72	57.03	56.13	7.71	7.71	0.44	0.44	0.56	0.47	261.23	272.76	16.03	16.26	174.9	167.5
T8	1.18	1.19	2.43	2.44	57.14	57.44	57.73	56.70	7.72	7.74	0.44	0.45	0.57	0.48	276.06	281.10	16.36	16.83	175.9	169.5
T9	1.17	1.18	2.41	2.42	58.26	58.09	58.60	58.13	7.75	7.75	0.47	0.48	0.58	0.49	280.86	286.40	16.56	17.26	178.1	172.8
F-Test	NS	NS	NS	NS	S	S	S	S	NS	NS	S	S	S	S	S	S	S	S	S	S
S. Em. (±)	-	-	-	-	0.33	0.25	0.17	0.18	-	-	0.006	0.006	0.007	0.008	2.01	1.28	0.19	0.21	1.18	0.84
C.D.(P=0.05)	-	-	-	-	0.99	0.78	0.52	0.54	-	-	0.019	0.019	0.023	0.025	6.06	3.38	0.57	0.63	2.52	1.80

[GIVE LEGEND FOR T1 ..... T9](#)

UNDER PEER REVIEW



## Conclusion

The effect of different levels in the experiment was concluded based on the trail. Treatment  $T_9$  ( $I_3@100\% + @100\% F_3 + @50\% R_3$ ) was shown to be the best in terms of improving soil physico-chemical parameters like of soil such as Bulk density, Particle density, % pore space, Water holding capacity, pH, EC, soil % Organic carbon, soil Available Nitrogen, Available Phosphorus, Available Potassium. Nevertheless, Fertilizer requirements in pea are critical for early development and overall yield generation. But, Crop productivity can be improved by combining biofertilizer, organic and inorganic fertilizers also enhance nutrient absorption, which accelerates cell division, cell elongation and hence plant metabolic activity.

## References

1. Bouyoucos, G. J. (1927) The hydrometer as the new method for the mechanical analysis of the soils, *Journal of Soil Science*, 2: 343 – 353.
2. Das, D., David, A. A., Swaroop, N., Hasan, A., Thomas, T. (2020). Response of different levels of inorganic fertilizer, organic manure and bio-fertilizer on physico-chemical properties of soil in pea (*Pisum sativum* L.) var. Kashi Ageti. *International Journal of current microbiological and applied sciences*, 9(10): 468-474.
3. Fischer, R. A. (1950). Techniques of analysis of variance, *Hand book of Agricultural statistics*. Anchal Prakashan Mandir. 332-334.
4. Jackson, M. L. (1958) *Soil Chemical Analysis*, Prentice – Hall India, New Delhi.
5. Kaur, H., Gosal, S. K. and Walia, S. S. (2017) Integrated application of biofertilizers with different fertilizers affects soil health in pea crop. *Chemical Science Review and Letters*, 6(23): 1646-1651.
6. Kimi, Z. S., David, A. A., Thomas, T., Swaroop, N. and Hassan, A., (2021). Response of Integrated nutrient management on soil health, Yield attributes and yield of pea (*Pisum sativum* L.). *The pharma innovation*, 10(10): 1815-1818.
7. Muthuvel, P.C., Udaysooriyan, R., Natesa, P.P. and Ramaswami (1992) *Introduction to Soil Analysis*, Tamil Nadu Agriculture University, Coimbatore- 641002. Muche, M., Kokeb, A. and Molla, E. Assessing the Physiochemical Properties of Soil Under Different Land Use Type.



8. Olsen, S.R., Cole, C.V., Watanabe, F.S. and Dean, L. A., (1954) Estimation of available phosphorus in soil by extraction with sodium bicarbonate ( $\text{NaHCO}_3$ ), *U.S.D.A Circular*. 939: 1-19.
9. Pandey, V., Dahiya, O. S., Mor, V. S., Yadav, R., Jitendra, Peerzada, O. H., and Brar, A. (2017). Impact of Integrated Nutrient Management on seed yield and its attributes in Field Pea (*Pisum sativum* L.). *Chemical science review and letters*, 6(23): 1428-1431.
10. Sharma, N. and Thakur, K. S. (2016). Effect of Integrated Nutrient Management on soil properties and nutrient content in pea (*Pisum sativum* L.). *The Bioscan* 11(1): 455-458.
11. Sharma, N. and Thakur, K. S. (2016). Effect of Integrated Nutrient Management on soil properties and nutrient content in pea (*Pisum sativum* L.). *The Bioscan* 11(1): 455-458.
12. Subbiah, B.V. and Asija, C.L. (1956) A rapid procedure for the estimation of available nitrogen in soils, *Current Science*, 25: 259-260.
13. Varsha, U., Hemlata, V. and Devidas, N. (2015). Influence of organic, chemical and biofertilizer on growth and yield of pea. *Agricultural Science Digest*, 35(3): 237-240.
14. Walkley, A. and Black, I. A., (1934) Estimation of soil organic carbon by the chronic acid titration method. *Soil Science*. 47: 775-776.
15. Wilcox, L.V. (1950) Electrical Conductivity. *American Water works Association Journal*. 42-776.