

## **Original Research Article**

### **Outcome of Tuberculosis Case Surveillance at Kano Central Correctional Center, North-west Nigeria: A Need for Routine Active Case Findings for TB in Nigerian Correctional Centers.**

#### **ABSTRACT**

**Background:** Tuberculosis is endemic in the Nigerian general population with higher incidence in correctional centers. The high incidence of TB is generally due to the overcrowded nature of the Nigerian correctional centres that made it high-risk environments for the transmission of the disease therefore, the routine active case finding (ACF) for TB in the centres is critical. To demonstrate the need for ACF in the Nigerian prisons, the KNCV Tuberculosis Foundation Nigeria carried out TB case surveillance at the largest correctional centre in the most populous state in Nigeria.

**Objective:** To conduct mass TB screening of inmates of Kano central correctional centre in Kano state Nigeria, identify TB cases, and place them on treatment.

**Methods:** A digital X-ray with artificial intelligence (AI) was used for mass TB screening of 1,967 consenting inmates at the Kano central correctional centre in Kano state Nigeria, from 21st September to 2nd October 2020. Participants with CAD4TB score  $\geq 60$  had a GeneXpert assessment of their sputa for TB diagnosis. Where sputum production was not possible or GeneXpert result was negative, expert clinical assessment of the presumptive radiogram was carried out by experienced radiologist. Data from the project were extracted and analysed for this report.

**Results:** Overall, 1,967 inmates were screened for TB and 92 (4.7%) presumptive were identified - males (4.8%, 91/92), females (1.9%, 1/92). Out of the 92 presumptive, 21 males were diagnosed as TB cases giving a TB prevalence of 1.1% among the inmates and 22.8%

**Comment [h1]:** The summary has more than 300 words such as the magazine rules, the type of study carried out is missing, no statistical tests applied in the student are indicated and I consider the objective of the same

among presumptive. One of the TB cases had multi-drug resistant TB. The number needed to screen (NNS) was 94. All TB cases were enrolled on treatment.

**Conclusion:** The prevalence of TB at the Kano central correctional centre during the mass TB screening project was high. The National Tuberculosis Control Programme of Nigeria should accelerate the implementation of the planned paradigm shift from passive to active case-finding for TB in Nigerian correctional centers.

**Key-words:** Mass Screening for TB, Wellness on Wheels Truck, Chest X-ray, CAD4TB, GeneXpert.

## INTRODUCTION

Monitoring of tuberculosis disease in a geographic area or population is an important public health program,<sup>1</sup> because the incidence of TB in prisons (correctional centers) is generally higher than in the general population.<sup>2</sup> TB case surveillance should be more critical in the overcrowded correctional centres in settings such as Nigeria with a huge burden of TB infection. Nigeria is the sixth country in the world with the largest burden of TB, and ranks second among all countries as regards the global missed TB burden of 2.9 million cases.<sup>3</sup>

Tuberculosis is transmitted through the inhalation of airborne droplets containing nuclei of tubercle bacilli generated from individuals with the pulmonary disease.<sup>4</sup> Therefore, poor ventilation and reduced immune status of exposed individuals are known predispositions to TB transmission. Unfortunately, these two predispositions are prevalent in the Nigerian correctional centres majorly due to overcrowding, and inadequate nutrition. Likewise, some independent risk factors for TB in West Africa such as household crowding, male sex, and smoking are prevalent in our correctional centers.<sup>5</sup> Furthermore, a longer duration of imprisonment is also associated with TB infection among inmates.<sup>6</sup> Since the risk of re-activation of latent TB is higher among the immunocompromised, the poor nutrition and its attendant low host immunity among prison inmates can also predispose to the re-activation of a latent TB infection.<sup>7</sup>

These prison-related factors and inadequate prison TB services explain the higher risk of developing active TB by inmates when compared with the general population.<sup>8,9</sup> A recent systematic review from Iran shows TB prevalence among inmates ranging from 25/100,000

to 517/100,000.<sup>10</sup> Another study from Romania which used a one-stop mobile X-ray with computer-aided TB detection software followed by Xpert assessment for presumptive found a TB prevalence of 97/100 000 of prisoners screened.<sup>11</sup>

In Nigeria, inmates of correctional centres constitute a socially excluded population characterized by overcrowding, poor nutrition, and inadequate health services, which make them a vulnerable group that qualifies for active case finding (ACF) for TB. For instance, a symptom-based TB ACF at the Aba prison, Abia State Nigeria found a TB prevalence of 21.2% among inmates with cough of at least two weeks duration, which constituted an alarming 1100% increase in TB detection rate in that centre.<sup>12</sup> As in the Aba study, a few other studies among Nigerian prison inmates relied on sputum microscopy and showed a varying TB prevalence of 1.2% to 2.4%.<sup>13,14</sup>

Due to the lack of evidence on TB ACF in most sub-Saharan African (SSA) countries, a recent systematic review has called for increased ACF interventions in SSA prisons, particularly countries included in the three high-burden country lists for TB, TB/HIV, and MDR-TB.<sup>15</sup> Nigeria inclusion in the three high-burden lists justified the ACF intervention in Nigeria Prisons. The KNCV Tuberculosis Foundation Nigeria (KNCV Nigeria) as part of its active case finding interventions for TB amongst most at risk populations in Nigeria carried out mass screening for TB at the largest correctional centre in the most populous State in Nigeria using a Wellness on Wheels (WoW) truck which is a mobile TB diagnostic unit housing a digital X-ray with computer-aided TB detection software (CAD4TB) and two GeneXpert instruments. The project findings would drive health policy shift in Nigeria Correctional centres aimed at improving TB prevention and services in centres.

UNDER PEER REVIEW

## MATERIAL AND METHODS

**Comment [h2]:** The sample must be defined

### Study design:

This was the report of the active TB case finding intervention conducted by KNCV Tuberculosis Foundation Nigeria at the Kano Central correctional centre, Kano state, Nigeria. The two weeks intervention lasted from 21st September to 2nd October 2020.

### Study setting:

Kano central correctional centre is the largest of 10 prisons in Kano State Nigeria. It is located in Kurmawa Quarters of Kano City and is managed by the Nigerian Correctional Services. It has 27 cells with a design capacity of 600 inmates;<sup>16</sup> however, at the time of the intervention, there were 2,000 inmates, and each cell accommodated 50 to 140 inmates. It has a medical clinic staffed by two medical practitioners, three nurses, and other support staff. The clinic has a functional DOTS service for TB diagnosis and treatment of TB. Passive TB case finding is the routine model in the Nigerian prison health system. Within the year preceding the intervention, the correctional centre reported seven TB cases.

### Study population:

All consenting inmates of the correctional centre were eligible to be screened irrespective of the presence or absence of respiratory and constitutional TB symptoms.

### Methods:

The project was conducted using the Wellness on Wheels (WoW) truck which is a mobile TB diagnostic unit housing a digital X-ray with computer-aided TB detection software (CAD4TB) and two GeneXpert instruments. The WHO 3b screening/diagnostic algorithm was used for the project.<sup>17</sup> The X-ray and GeneXpert instruments were operated by a radiographer and laboratory scientists trained by KNCV Nigeria for the project. The KNCV Nigeria team paid an advocacy visit to the management of the correctional center and got their buy-in and

support for the TB surveillance project. Following a group education of inmates on the high burden of TB in correctional centres and the need for the project as a means of TB prevention, individual counselling was conducted by the DOTS officer. As shown in the case surveillance flow diagram (Figure 1), all consenting inmates had X-ray screening for TB. Participants with a CAD4TB score of  $\geq 60$  were defined as presumptive for TB and encouraged to produce quality spot sample of sputum. The spot sputum specimens of the presumptive were processed and assayed the same day, with Xpert instruments. Presumptive that could not produce sputum had their chest radiograms sent electronically to two independent Radiologists for clinical review – where there was a discordant report; a third radiologist review served as a tiebreaker. All confirmed TB cases (bacteriological or clinical) were commenced on appropriate TB treatment. The prison authority was also informed of all TB-positive cases for case isolation.

#### **Outcome measures:**

The primary outcome measure was the prevalence of TB cases among inmates of Kano Correctional centre, Kano state, Nigeria. The secondary measures were the proportion of TB cases among the presumptive identified after X-ray, and the number needed to screen (NNS) to diagnose one person with active TB.

#### **Data management:**

Relevant data of participants were retrieved for this report including age, sex, X-ray CAD4TB scores, Xpert result, clinical X-ray report. The data were extracted into an Excel datasheet and de-identified. Data analysis was conducted with IBM SPSS version 21. Proportions and means were compared where necessary with Fisher Exact test and Student t-test respectively. Results were presented as appropriate using median (and interquartile range),

frequencies, percentages, tables, and charts. A p-value of  $< 0.05$  was considered statistically significant.

UNDER PEER REVIEW

## RESULTS

**Comment [h3]:** The tables have the title above and also the source from which the data in them was extracted was missing

There were 2,000 inmates in the correctional centre at the time of the TB-case surveillance program [males: 1,948 (97.4%), females: 52 (2.6%)], which gave an occupancy level of 333.3%. As shown in Table 1, out of all the inmates, 1,967 (98.4%) were screened for TB during the project (males: 97.4%, females: 2.6%). The 33 (1.7%) male inmates that opted out of the project were on death row. No inmate was below 16 years of age. The mean age of all participants was  $32.0 \pm 11.5$  (range: 16 – 89) years. In all, the average age of male participants [ $31.9 \pm 11.4$  (range: 16-89) years] was lower than females [ $36.4 \pm 14.9$  (range: 18-80) years],  $p = 0.003$ . The modal age group for all participants, presumptive, and confirmed TB cases were 25 – 34 years (38.0%), 35 – 44 years (29.3%), and 25 – 34 years (38.1%), respectively.

As shown in Table 2, the proportion of presumptive identified among all participants was 4.7% (92/1964). The prevalence of presumptive among male participants was 4.8% while that of female participants was (1.9%). The observed difference was not significant, OR = 2.6 (95%CI: 0.35 – 18.83),  $p = 0.513$ .

As shown in Figure 1 and Table 2, all identified presumptive TB were evaluated for TB. Twenty-one TB cases were confirmed during the project which gave a TB prevalence of 1.1% among all participants, and 22.8% among the presumptive. All confirmed TB cases were males while the only female presumptive was negative for TB following Xpert assessment. One of the cases had multidrug-resistant TB (MDR TB) and further evaluation showed resistance to a second-line injectable drug i.e., pre-extensively drug-resistant TB (Pre-XDR TB). All confirmed TB cases were separated from other inmates in different cells and placed on appropriate treatment.

Table 2 also shows that the number needed to screen (NNS) with CAD4TB chest X-ray to diagnose one person with active TB in the Kano central correctional centre was 94. On the other hand, the number needed to test (NNT) to diagnose one TB case for all participants was 5.

## DISCUSSION

In line with the strategic plan of the National Tuberculosis, Leprosy, and Buruli Ulcer Control Programme (NTBLCP) to move from passive to active case-finding in key affected populations including the prisons, this TB-case surveillance project targeted the largest correctional centre in Kano State, the most populous state in Nigeria. The main finding of the project was a high TB prevalence of 1.1% among all participants and 23% among presumptives in the prison.

The TB prevalence for all participants in this project translates to a case rate of 1,100/100,000 population which is far above the 2019 actual and targeted national TB case notification rates of 60/100,000 and 235/100,000 population respectively.<sup>20</sup> Also, the TB prevalence among presumptive in this study was similar to the 21% reported from Aba prison in Abia state Nigeria<sup>12</sup> but, higher than 9% from Ethiopia.<sup>6</sup> However, these related studies used a symptom-based ACF model which is associated with more number of missed cases when compared to the CAD4TB chest X-ray that was used in this project.<sup>21</sup> Furthermore, Xpert testing used for this study is superior to sputum microscopy for the diagnosis of TB.<sup>21</sup>

A prison is considered overcrowded when the occupancy level is above 100%.<sup>23</sup> The Kano central correctional centre occupancy level of 333% during the project was more than twice

**Comment [h4]:** I believe that in the discussion so many results should not be mentioned and if comparing with similar studies carried out in other countries

the national average of 147%<sup>24</sup> which suggests that it might be more crowded than other correctional centres in Nigeria. This finding supports an earlier report of the Service Compact with all Nigerians (SERVICOM) compliance evaluation of the prison which identified congested cells and poor-quality foods to inmates as part of the weaknesses of the correctional center.<sup>16</sup> Because, these two attributes are known predispositions to TB spread in the community, their impacts are likely to be higher in a prison environment which may explain the high TB rate identified by the project. In fact, it has been established that in countries such as Nigeria, with a high TB prevalence in the general population, TB prevalence in the prisons is as much as 100 times higher than in the outside communities.<sup>23</sup> The increased TB prevalence in the prisons when compared to the outside communities is demonstrated in this project and other related studies in Nigerian prisons.<sup>12-14</sup> It is therefore important that while efforts are being made to implement the strategies outlined by the United Nations' office on crime and drugs to decongest the prisons,<sup>23</sup> routine TB ACF approaches including entry and routine mass screening,<sup>15</sup> are implemented in Nigerian Prisons.

It is noteworthy that all inmates that opted out of the TB screening were on death row. This attitude of the death row inmates to TB screening may not be out of place because such inmates are characterized by an increased incidence of psychological disorders including depression.<sup>25</sup> Major depressive disorder, for instance, is often associated with loss of interest and diminished capacity to think which might make it difficult for an affected individual to comprehend health information and make an informed choice.<sup>26</sup> Unfortunately, one or more of the death row inmates could be harbouring active TB thereby still exposing the post-screening TB-free inmates to the risk of tuberculosis infection.

Because of the grave public health concern of missing TB case(s) among non-consenting inmates after a mass screening, a mandatory TB screening may therefore be an appropriate approach for inmates in Nigerian correctional centres.

As was the case with Kano central prison, the proportion of female prisoners in Nigerian prisons is low with an average of 1.9%. Despite this usual lower proportion of female prisoners, this project demonstrated that females were as likely to be TB presumptive as the male inmates (Table 2). Therefore, the ACF for TB in correctional centres must target all inmates irrespective of sex. The project found that 94 inmates needed to be tested to identify an active TB case (Table 2) so, the small number of female inmates in the Kano central prison ( $n = 52$ ) may explain the finding that all TB cases identified in the project were males. A case of MDR-TB was identified during the project which is not surprising for a country listed among the top MDR-TB high-burden countries. However, because of the difficulty in the treatment of the MDR-TB cases, prevention of its infection and spread within the correctional centre through routine ACF, including routine entrance TB screening, should be a top priority of the National TB programme.

The strength of this project lies in the fact that it is the first mass TB screening in Nigerian correctional centres using digital X-ray with CAD4TB artificial intelligence, and Xpert testing for presumptive. However, the opt-out of some inmates from the mass screening limited the project because of the possibility of undiagnosed TB cases among the cohort.

## CONCLUSION

The prevalence of TB at the overcrowded Kano central prison during the TB-case surveillance project was high. Though all TB cases were males, inmates' sex had no relationship with the odds of being presumptive. The NTBLCP should accelerate the implementation of its planned shift from passive to active case-finding in Nigerian correctional centers. To minimize missed TB cases, the routine ACF for TB approach should include entry screening and intermittent mass screening using chest x-ray.

#### **CONSENT AND ETHICAL CONSIDERATIONS**

Provider initiated counselling and testing with opt-out approach was used for the project.<sup>18,19</sup> Each participant received information on the objectives, procedure, and benefits of the project after which s/he proceeded with TB screening using the X-ray if s/he did not specifically decline the intervention. Participants were assured of their confidentiality however, there was a need to isolate those found to have TB to protect other inmates. They were informed that their screening results and de-identified bio-data would be used for disseminating project results.

#### **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## REFERENCES

1. Centers for Disease Control and Prevention (CDC). Self-Study Modules on Tuberculosis, 6-9 Module 9: Tuberculosis Outbreak Detection and Response. CDC; 2014 [cited 2021 1<sup>st</sup> October]. Available from:  
<https://www.cdc.gov/tb/education/ssmodules/pdfs/module9.pdf>
2. Velen K, Charalambous S. Tuberculosis in prisons: an unintended sentence? The Lancet Public Health. 2021; 6(5):e263-4
3. World Health Organization (WHO). Global Tuberculosis Report 2020. Geneva: WHO; 2020 [cited 2021 1<sup>st</sup> October]. Available from:  
<https://www.who.int/publications/i/item/9789240013131>
4. American Thoracic Society, CDC, Infectious Disease Society of America. Diagnostic standards and classification of tuberculosis in adults and children. Am J Respir Crit Care Med 2000; 161: 1376–95
5. Lienhardt C, Fielding K, Sillah JS, Bah B, Gustafson P, Warndorff D, et al. Investigation of the risk factors for tuberculosis: a case-control study in three countries in West Africa. Int J Epidemiol. 2005; 34(4): 914-23.
6. Addis Z, Adem E, Alemu A, Birhan W, Mathewos B, Tachebele B, et al. Prevalence of smear positive pulmonary tuberculosis in Gondar prisoners, North West Ethiopia. Asian Pacific journal of tropical medicine. 2015; 8(2): 127-31.

**Comment [h5]:** The bibliographies are limited according to the order of appearance, there are some (25) that are not limited, more than 50% are outdated, are more than 5 years old and not all of them are limited according to Vancouver international standards.

7. CDC. Guidelines for Preventing the Transmission of Mycobacterium tuberculosis in Health-Care Settings. MMWR 2005; 54(No. RR -17):1-141
8. Dara M, Acosta CD, Melchers NV, Al-Darraji HA, Chorgoliani D, Reyes H, et al. Tuberculosis control in prisons: current situation and research gaps. Int J Infect Dis. 2015; 32:111-7
9. Dadu A, Ciobanu A, Hovhannesyan A, Alikhanova N, Korotych O, Gurbanova E, et al. Tuberculosis Notification Trends and Treatment Outcomes in Penitentiary and Civilian Health Care Sectors in the WHO European Region. Int J Environ Res Public Health. 2021; 18(18): 9566
10. Taherpour S, Bazzaz MM, Naderi H, Samarghandian S, Amirabadizadeh A, Farkhondeh T, et al. A systematic and meta-analysis study on the prevalence of tuberculosis and relative risk factors for prisoners in Iran. Infect Disord Drug Targets. 2021 Sep 12. Epub ahead of print.
11. Mahler B, de Vries G, van Hest R, Gainaru D, Menezes D, Popescu G, et al. Use of targeted mobile X-ray screening and computer-aided detection software to identify tuberculosis among high-risk groups in Romania: descriptive results of the E-DETECT TB active case-finding project. BMJ Open. 2021; 11(8): e045289
12. Ekundayo EO, Onuka O, Mustapha G, Geoffrey M. Active case finding of pulmonary tuberculosis among prison inmates in aba Federal prison, Abia state, Nigeria. Advances in Infectious Diseases. 2015; 5(01): 57
13. Adesokan HK, Cadmus EO, Adeyemi WB, Lawal O, Ogunlade CO, Osman E, et al. Prevalence of previously undetected tuberculosis and underlying risk factors for transmission in a prison setting in Ibadan, south-western Nigeria. Afr J Med Med Sci. 2014; 43(Suppl 1): 45-50

14. Lawal MA, Omili M, Bello TO, Onuha L, Haruna A. Tuberculosis in a Nigerian medium security prison. *Benin Journal of Postgraduate Medicine*. 2009;11(1).
15. Kuupiel D, Vezi P, Bawontuo V, Osei E, Mashamba-Thompson TP. Tuberculosis active case-finding interventions and approaches for prisoners in sub-Saharan Africa: a systematic scoping review. *BMC Infect Dis*. 2020; 20(1): 570
16. Federal Ministry of Interior, Nigeria. Report of SERVICOM Compliance Evaluation of Nigerian Prisons Service, Kano Central Prison. 2011. [cited 2021 2<sup>nd</sup> October]. Available from: <https://servicom.gov.ng/wp-content/uploads/2017/08/Nigerian-Prisons-Service-Kano-Central-Prison.pdf>
17. WHO. Systematic screening for active tuberculosis: an operational guide. WHO/HTM/TB/2015.16. Geneva: WHO, 2015 [cited 2021 29th September]. Available from: [http://apps.who.int/iris/bitstream/handle/10665/181164/9789241549172\\_eng.pdf](http://apps.who.int/iris/bitstream/handle/10665/181164/9789241549172_eng.pdf)
18. WHO. Guidance on provider-initiated HIV testing and counselling in health facilities. Geneva: WHO; 2007. [cited 2021 2nd October]. Available from: [https://www.who.int/hiv/pub/guidelines/9789241595568\\_en.pdf](https://www.who.int/hiv/pub/guidelines/9789241595568_en.pdf)
19. Dim CC, Nwagha UI, Ezegwui HU, Dim NR. The need to incorporate routine cervical cancer counselling and screening in the management of women at the outpatient clinics in Nigeria. *J Obstet Gynaecol* 2009; 29: 754-6
20. Nigeria Federal Ministry of Health (FMoH). National Tuberculosis and Leprosy Control Programme: Annual report 2019 Abuja: FMoH, 2020.
21. Nadiah Z, et al. Early chest X-ray in persons with presumptive tuberculosis increases Xpert® MTB/RIF diagnostic yield and efficiency. *Public health action*. 2020 Mar 21;10(1):17-20.

22. Odume B, Nwokoye N, Spruijt I, Slyzkyi A, Dim C, Chukwuogo O, et al. Diagnostic Accuracy of TB-LAMP for Diagnosis of Pulmonary Tuberculosis among Adult Presumptive TB in Nigeria. *Greener Journal of Medical Sciences*, 2021; 11(2): 122-129
23. United Nations Office on Drugs and Crime (UNOC). Handbook on Strategies to Reduce Overcrowding in Prisons. 2013. [cited 2021 23 October]. Available from: [https://www.unodc.org/documents/justice-and-prison-reform/Overcrowding\\_in\\_prisons\\_Ebook.pdf](https://www.unodc.org/documents/justice-and-prison-reform/Overcrowding_in_prisons_Ebook.pdf)
24. World Prison Brief. World Prison Brief Data: Nigeria. [cited 2021 21 October]. Available from: <https://www.prisonstudies.org/country/nigeria>
25. Cunningham MD, Vigen MP. Death row inmate characteristics, adjustment, and confinement: a critical review of the literature. *Behav Sci Law*. 2002;20(1-2):191-210.
26. Halverson JL, Bhalla RN, Moraille-Bhalla P, Andrew LB, Leonard RC. Depression. *E-Medicine*. [cited 2021 26 October]. Available from: <https://emedicine.medscape.com/article/286759-overview#a1>

## TABLES AND FIGURES




Figure 1: Kano Central Prison's TB-Case Surveillance Flow diagram

Category	Inmates screened (n = 1967) Freq. (%)	Presumptive TB (n = 92) Freq. (%)	TB patient (n = 21) Freq. (%)
Gender			
Male	1915 (97.4)	91 (98.9)	21 (100.0)
Female	52 (2.6)	1 (1.1)	0 (0.0)
Age (years)			
0-14	0 (0.0)	0 (0.0)	0 (0.0)
15-24	504 (25.6)	8 (8.7)	2 (9.5)
25-34	747 (38.0)	24 (26.1)	8 (38.1)
35-44	400 (20.3)	27 (29.3)	6 (28.6)
45-54	190 (9.7)	20 (21.7)	4 (19.0)
55-64	85 (4.3)	9 (9.8)	1 (4.8)
65+	41 (2.1)	4 (4.3)	0 (0.0)

Table 1: Demographic characteristics of inmates screened, presumptive identified and TB patients detected

Table 2: The number needed to screen (NNS) and number needed to test (NNT) to diagnose one person with active TB in Kano Central Correctional Center

	Total n (%)	Male n (%)	Female n (%)	p-value
Persons screened	1967 (100.0)	1915 (47.2)	52 (2.6)	-
Presumptive identified	92 (4.7)	91 (4.8)	1 (1.9)	0.513*
Presumptive evaluated for TB	92 (100.0)	91 (100.0)	1 (100.0)	0.513
TB cases diagnosed	21 (22.6)	21 (22.8)	0 (0.0)	1.0
TB patients put on treatment	21 (100.0)	21 (100.0)	-	-
NNS	94	92	-	-
NNT	5	5	-	-

NNS = (Persons screened / TB patients diagnosed)

NNT = (Presumptive patients evaluated for TB / number of TB patients diagnosed)

\* OR = 2.6 (95%CI: 0.35 – 18.83)