

Prevalence of Intestinal Parasites and Their Transmission Risk Factors among Inmates of Naivasha and Kisii Prisons in Kenya.

Comment [p1]: Centralize

Runing Title : Intestinal parasitic infections in selected Kenya prisons and their risk factors.

Comment [p2]: Not necessary

ABSTRACT

Parasitic infections in endemic countries shorten lives as well as impose a lifelong burden on the country's potential for advancement. Very few studies have been done to establish the prevalence of intestinal parasites among neglected and institutionalized groups of the population such as prison inmates; therefore, there was a need to establish the prevalence of IPs among inmates of Kisii and Naivasha prisons in Kenya. A cross-sectional study of 400 sampled inmates provided stool samples for formal ether concentration techniques for identification of helminths eggs and cysts of protozoa while direct wet mount using saline and iodine for microscopic examination of loose stool for identification of trophozoites of protozoa. Data on the predisposing factors was collected using structured questionnaire and focus group discussion. Chi Square test was used to determine the association between age groups and sex while odds ratio was used to determine the association of personal hygiene and infection by intestinal parasites. The prevalence rate of intestinal parasites for the two prisons was 84(20.84%). Intestinal protozoa infections were higher 50(12.4%) than helminthes 28(10.4%). Intestinal protozoa identified were *Entamoeba histolytica* 30(7.4%) and *Giardia lamblia* 20(4.9%) while the intestinal helminthes identified were *Ascaris lumbricoides* 22(5.5%), *Strongyloides stercoralis* 2(0.5%) and hookworms 4(1.0%). Chi square test showed that significantly more females were infected than males ($\chi^2 = 12.53$, $P=0.016$, $df = 2$) as well as inmates of 18-27 years in both prisons ($\chi^2 = 9.36$, $P=0.007$, $df = 28$). Correlation coefficient indicated that the infections increased with lack of footwear (OR 2.68, CI 0.94-3.41), failures to hand wash before meals (OR 2.88, CI 0.93-3.36) and after toilet use (OR 2.19, CI 1.02-4.44). The age of the respondent also had a significant contribution to the high prevalence of IPs. Provision of health education and personal hygiene and sanitation, more pit latrines with enough water points for hand washing as well as sufficient footwear is recommended to reduce the IPs in the prisons. This can be facilitated by the officers-in-charge in collaboration with the Ministry of health.

Keywords: inmates; intestinal parasitic infections; prevalent; sanitation; prison.

Comment [p3]: It will be more appropriate to begin by defining parasitic infections before stating the effects of their burden.

Comment [p4]: Replace with and

Comment [p5]: Tautology; You could make it Impose lifelong burden on the potential for advancement

Comment [p6]: Use limited instead

Comment [p7]: Carried out

Comment [p8]: Which justified the need

Comment [p9]: There is need to mention in full being the first time and then make the abbreviations to be in the bracket.

Comment [p10]: This is too complex, rephrase by splitting the statements,

Comment [p11]: were

Comment [p12]: There is need to state the prevalence for each; for the Kisii and Naivasha prison separately

Comment [p13]: Where? This seems ambiguous; there is need to state weather the comparsion is with respect to the two prisons or not; it is difficult to follow.

Comment [p14]: Has this been backed up with a reference?

Comment [p15]: Replace with comma

Comment [p16]: The use of

Comment [p17]: are

Comment [p18]: Prevalence?

1. INTRODUCTION

Significant level of morbidity and mortality across the globe especially in developing countries is caused by intestinal parasitic infections and more so in persons with additional disorders [1]. There

Comment [p19]: There is need to better up this first statement, so as to engage the readers first with what the intestinal parasitic infections are.

are two types of intestinal parasites infesting humans; helminthes and protozoa. The main species of soil transmitted helminthes are *Ascaris lumbricoides*, *Trichuris trichura* and hookworms (*Ancylostoma duodenale* and *Necator americanus*). Transmission of *Ascaris lumbricoides* and *Trichuris trichura* occurs through ingestion of eggs passed out in stool by an infected person which becomes infective in soil where sanitation is poor, whereas Hookworm larvae cause infection by penetrating the skin. Approximately 1.7billion people are infected with soil transmitted helminthes worldwide [2]. There is also a striking relationship between the prevalence of hookworm and low socio-economic status [3]. Pathogenic intestinal protozoa frequently isolated from diarrheal patients include *Giardia lamblia* and *Entamoeba histolytica*. Some of the epidemiological factors that promote the prevalence of intestinal parasites include poor personal hygiene, poor sanitation and waterborne epidemics due to water treatment failure, climatic condition and socio-cultural practices [4].

Comment [p20]: Reference?

Comment [p21]: Reference?

Comment [p22]: consider

Comment [p23]: Reference?

Comment [p24]: Reference

Comment [p25]: transmission

Prisoners harbor diseases that are determined by their home environment before imprisonment and the prison environment. Prisoners are vulnerable to diseases through poor health care delivery, overcrowding, high risk behavior such as homosexuality and reduced immunity compared to the general population [5, 6]. Prisoners in Kenya just like those in California are faced with problems such as severe drug and substance abuse, HIV/AIDS, STDs, TB, skin and helminthes infections among others [6]. These prison facilities are few and insufficient due to high growth rate of Kenya population and as such, the problem of congestion and overcrowding, degrading beddings and clothing, poor sanitation is experienced [8]. Most of these factors predispose the inmates to parasitic infections and because most of these infections are asymptomatic, they are neglected to a point of chronic features appearing [7]. Those imprisoned represent a medically underserved population and are always at a higher risk of medical disorders [8]. Many studies have been carried out in children and mental institutions but very little work has been carried out in these neglected prison institutions and therefore, results from this study will help to develop policies to improve the health aspects of inmates among the other ongoing reform programmes. Kisii and Naivasha Prisons were used for this study as the prisons have both male and female sections making the study all inclusive.

Comment [p26]: get infected with

Comment [p27]: either due to their home environment prior to imprisonment

Comment [p28]: or due to the prison environment

Comment [p29]: due to

Comment [p30]:

Comment [p31]: Replace with furthermore

Comment [p32]: ,

Comment [p33]: are

Comment [p34]: ,

Comment [p35]: Split the statements atleast into two

1. MATERIALS AND METHODS

1.1 Study Area

Comment [p36]: Same numbering as in the introduction?

This study was carried out in two prisons in Kenya which have both male and female sections. 1) Naivasha prison is situated 2km north of the central business district of Naivasha town which is within Nakuru County in Kenya. It has an attitude of 2084m above sea level. on average, it receives an annual rainfall of 677mm and average temperature of 17°C [9]. 2) Kisii prison is found in Kisii County located in the western Kenya highlands at an altitude of 1660 M above sea level and an annual rainfall of 1500 mm, distributed throughout the year. It has annual temperatures range from 10 to 30°C and 88% relative humidity [10].

Comment [p37]: Capitalize

Comment [p38]: Situated

Comment [p39]: with

1.2 Study Design

A cross-sectional study of inmates from Kisii and Naivasha prisons was done to establish the prevalence of IPIs among the inmates. Four hundred inmates were sampled, stratified into three based on their jail terms; SJT, MJT and LJT. Inmates that attended the dispensary for treatment were picked by simple random sampling. Participation was voluntary after signing an informed consent.

Comment [p40]: carried out

Each of the sampled inmates recruited at the prison health facility was given dry, clean, leak proof container for 3gms stool collection and requested to bring early morning stool samples the following day. Diarrheic samples were examined immediately at Kisii and Naivasha prison dispensaries by wet mount technique [12]. The remaining stool samples of approximately 2gm were mixed thoroughly with 10 % formalin to preserve the trophozoites and emulsify formed stool before shipment and laboratory examination within Kisii level 5 hospital and Naivasha maximum prison's health facilities. Formalin-Ethyl Acetate Sedimentation concentration technique was used [13].

Comment [p41]: containers?

Comment [p42]: Split the statements

Comment [p43]: To do what particularly?

The questionnaires were filled by inmates the morning of returning the stool specimen which provided data on demographic and socio-environmental factors that potentially predispose the inmates to parasitic infections. Chi Square test and ANOVA were used to determine the association between age and sex while adjusted odds ratio was used to determine the association of personal hygiene and infection by intestinal parasites. The differences were considered significant when the p -value was ≤ 0.05 .

Comment [p44]: between

2. RESULTS

Four hundred inmates filled the questionnaires and provided stool specimen for the study of which 185(46.25%) were from Kisii prison while 215(53.75%) were from Naivasha prison. The male inmates

sampled were more 311(77.75%) than females 89(22.25%) as the population of male inmates was higher than females in these prisons.

Comment [p45]: There should be a title for this.... Is it the sex of inmates with respect to prisons?

2.1 Prevalence of IPIs by Demographic Characteristics of Inmates

Comment [p46]: 2 .Seems the above results could be included here (1)

The infection rates of intestinal parasitic infections with respect to sex was significantly higher among females than males ($\chi^2 = 12.53$, $P=0.016$, $df = 2$). According to table 1, age was a statistically significant factor to the infection rate ($\chi^2 = 9.36$, $P=0.007$, $df = 28$) where infection rate of intestinal parasites was higher among inmates of ages 18-27 years (41.67%) but hookworm infections were much higher among inmates of above 58 years 11.11%.

Comment [p47]: remove

Comment [p48]: Comparison about the parasites should be done separately, not merged with comparisons about the overall parasite prevalence.

Table 1: Age of inmates and type of parasite infection in the two prisons

Age	Age of respondents					
	18 – 27	28 – 37	38 – 47	48 - 57	Above 58	ANOVA
Intestinal parasites	N=84 (%)	N= 161 (%)	N=103 (%)	N= 34 (%)	N= 18 (%)	P-value
<i>E. histolytica</i>	17(20.24) ^a	5(3.11)	5(4.85)	2(5.88)	1(5.56)	0.014
<i>G.lambliia</i>	6(7.14)	6(3.73)	5(4.85)	3(8.82)	0(0.00)	0.057
Total protozoa	23(27.38)	11(6.83)	10(9.71)	5(14.7)	1(5.56)	0.061
<i>A. lumbricoides</i>	12(14.28)	6(3.73)	3(2.91)	0(0.00)	1(5.56)	0.068
<i>S. stercoraries</i>	0(0.00)	1(0.62)	1(0.97)	0(0.00)	0(0.00)	0.159
Hookworms	0(0.00)	1(0.62)	1(0.97)	0(0.00)	2(11.11)	0.101
Total helminthes	12(14.28)	8(4.97)	5(4.85)	0(0.00)	3(16.67)	0.395
Total infection	35(41.67)^b	19(11.8)	15(14.6)	5(14.7)	4(22.22)	0.028

Comment [p49]: ≥

Superscript a and b show results where infection rate was significantly higher among inmates aged 18-27years as compared to other age groups ($\chi^2 = 9.36$, $P=0.007$, $df = 28$).

Protozoa infections were proportionately higher among female inmates (13.48%) than the male counterparts (12.1%) and so were helminthic infections (7.86%) and (6.68%) as shown in table 2. The infection rates of intestinal parasitic infections with respect to sex was significantly higher among females than males ($\chi^2 = 12.53$, $P=0.016$, $df = 2$).

Table 2: Type of parasite infection relative to sex of the study subjects

Sex	Female	Male	ANOVA
Intestinal parasites	n=89 (%)	n= 314 (%)	P VALUE
<i>E. histolytica</i>	9(10.11)	21(6.68)	0.067
<i>G.lamblia</i>	3(3.37)	17(5.41)	0.088
Total protozoa	12(13.48)^a	38(12.1)	0.025
<i>A. lumbricoides</i>	5(5.62)	17(5.41)	0.113
<i>S.stercoralis</i>	1(1.12)	1(0.318)	0.143
Hookworms	1(1.12)	3(0.95)	0.207
Total helminthes	7(7.86)^b	21(6.68)	0.014
Total infection	19 (21.34)^c	59 (18.79)	0.041

Superscript a, b and c shows results where protozoa and helminth infections were significantly higher among female inmates than their male counterparts ($\chi^2 = 12.53$, $P=0.016$, $df = 6$).

Comment [p50]: -

2.2 Risk factors associated with protozoa and helminthes infections among inmates

Table 3 shows some of the personal hygiene practices that were found to influence transmission of intestinal parasites among the inmates as recorded from questionnaires and focus group discussions. Almost 52% of the subjects had no shoes out of which 22% had at least one intestinal parasitic infection. The odds of being infected with an intestinal parasite for a person not wearing shoes were 2.68 (i.e 3 to 1 for a person wearing shoes). Lack of hand washing before meals had an odds ratio of 2.88 showing that 3 inmates not washing hands were at risk of infection with IPIs while hand washing after visiting the toilet had an odds ratio of 2.19 which indicate that twice the number of inmates who do not practice hand washing after toilet use were predisposed to infections compared to those who washed hands.

Comment [p51]: Could

Comment [p52]: Subjects/ inmates.... It would be better to use one word

Comment [p53]: Too complex... Split or rephrase

Table 3: Risk factors of transmission of intestinal parasites infections.

Environmental factor n= 403	Number of inmates (%)	Infected inmates (%)	Odds ratio (95% ci)
Wearing shoes	194 (48.5)	38 (19.6)	OR 2.68 CI (0.94- 3.41)
Not wearing shoes	206 (51.5)^a	46 (22.3)	
Washing hands before meals	288 (72.0)	58 (20.1)	OR 2.88 CI (0.93-3.36)
Not washing hands before meals	112 (28.0)	26 (23.21)^b	
Using soap while hand washing before meals	33 (8.2)	6 (18.2)	OR 1.47 CI (0.22-2.19)
Not using soap while hand washing before meals	255 (63.3)	52 (20.4)	
Washing hands after visiting the toilet	152 (37.7)	27 (17.8)	OR 2.19, CI (1.02-4.44)
Not washing hands after visiting the toilet	247 (61.8)^c	57 (23.1)	
Using soap while washing hands after visiting the toilet	33 (8.25)	4 (21.1)	OR 1.24, CI (0.12-2.19)
Not using soap while washing hands after visiting the toilet	116 (29.0)	23 (19.8)	

Superscript a, b and c shows lack of shoe wear and failure to wash hands as some of the factors that significantly increased the risk of transmission of intestinal parasites infections.

3. DISCUSSION

The results in this study indicate that intestinal parasitic infections are prevalent among inmates of Kenya prisons where 20% and 21.57% of the inmates in Kisii and Naivasha prisons respectively were infected with one or more species of intestinal parasites. This rate was lower than that reported by Rop [14] where 24.7% of inmates in Kisii prison in Kenya were parasitized by IPIs. *E. histolytica* cysts are infectious when passed in stool or shortly afterward unlike helminth eggs which has to undergo development to infective forms in the soil or where the egg has to hatch into larvae and therefore protozoa can indefinitely propagate the infections unless treated. This could explain the high prevalence of *E. histolytica* among the inmates [15].

Comment [p54]: ,

Comment [p55]: were

Comment [p56]: prevalence

Comment [p57]: This could be the reason why *E. histolytica* was the most prevalent among the inmates

Intestinal parasitic infections were significantly higher in females 21.34% than in males 18.97%. This could be attributed to the fact that females were engaged more in garden activities. From focus group discussions, it was noted that females had the habit of eating unwashed fruits and vegetables/salads which could be contaminated with intestinal parasites [25]. From these garden activities, helminthes eggs present in soils contaminated by stool can adhere to hands and finger nails, be ingested and thus causing infections [22]. This study is contrary to Acuna-soto [16] who noted that ,asymptomatic *E.histolytica* infection is distributed equally among both sexes.

In the current study, as age increased (from 18 years) the burden of intestinal parasite infection reduced. Inmates aged 18-27 years (27.38%) had higher prevalence of intestinal protozoa compared to those of ages> 58years (5.56%). These age groups of the inmates are the most active when it comes to duties assigned to them such as carpentry, farming and others. They are also mobile inmates to be taken to duties outside the prison walls thus high exposure to the parasites, increasing their risk of infection [17]. This is because the older inmates are confined and thus their movements are restricted. Hookworm infections were notably higher among inmates of age 58years and above (11.11%). Hookworm frequency and intensity remained high in elderly people probably due to a drop in immunity among elderly people [18, 19].

High prevalence of various infections observed in this study may be attributed to the direct effect of the prison environment. This study shows that, most of the inmates that walked barefoot (51.9%) were 3 times more likely to get infected with intestinal parasites unlike those wearing shoes (OR=2.68, CI 0.94-3.41). Wearing footwear protects the feet against penetration of hookworms and *S. stercoralis* larvae through the skin especially when working on farms and playing sports such as football barefoot [20]. Three times more inmates are likely to get infected with IPIs due to poor hygiene practices such as failure to wash hands before meals (OR 2.88, CI 0.93-3.36). Failure to wash hands and eating raw, unwashed fruits and vegetables increases chances of infection by IPIs especially among immigrants and institutionalized people [21]. Overcrowding and high risk behaviors such as homosexuality predispose inmates to infections by IPIs [22]. Naivasha and Kisii prisons whose cells are recommended to house approximately 25 male inmates were housing 40-45 inmates. Kisii prison is one of the congested prisons in Kenya with poor living conditions for the prisoners. The dilapidated

Comment [p58]: ,

Comment [p59]: ,

Comment [p60]: This reference should be used as a suport to the report, since this is your report, not that of the author quoated.

Comment [p61]: Consider removing and use comma instead

Comment [p62]: and

Comment [p63]: reported

Comment [p64]: was?

Comment [p65]: ≥

Comment [p66]: Taken should be used

Comment [p67]: Replace with contrary

Comment [p68]: ≥58years

Comment [p69]: Hookworm infection or hookworm?

Comment [p70]: Was higher

Comment [p71]: possibly

Comment [p72]: ,

Comment [p73]: It was reported that,.....

Comment [p74]: Furthermore,

state of the prison toilets, buildings, infrastructure and environment may have contributed greatly to the declined health of the inmate as reported previously [15].

4. CONCLUSIONS

Intestinal parasitic infections are common in prisons with a prevalence of 20.8%. The most common protozoa were *Entamoeba histolytica* (7.5%) while the most frequently occurring helminth was *Ascaris lumbricoides* (5.5%). The least occurring helminth was *Strongyloides stercoralis* with a prevalence of 0.5% in both prisons. Poor personal hygiene practices including failure to wash hands before meals and after toilet use and not wearing shoes were some of the factors found to predispose inmates to intestinal parasitic infections in the study sites.

Comment [p75]: Be specific; which prisons? Those under your study?

Comment [p76]: *S. stercoralis*: since is not first time mentioning

Comment [p77]: As well as not

5. RECOMMENDATIONS

Due to the high prevalence of IPIs recorded, constant medical checkups should be provided to all inmates as well as treatment with anti-helminthes to all positive inmates as well as new inmates. Public awareness on the transmission factors of intestinal parasites especially relating to environmental sanitation and personal hygiene in prison should be done to help reduce transmission of parasites. The conditions of the facilities in prisons such as toilets and sewerage disposal should be improved as well as provision of constant and clean water supply.

Comment [p78]: ambiguous

Comment [p79]: sewage?

ETHICAL CONSIDERATION AND CLEARANCE

Ethical clearance was sought from the ethical review committee of Kenyatta University while the research permit was acquired from the National Commission for Research Technology and Innovation (NACOSTI) alongside the Ministry of Health in Naivasha and Kisii and the concerned officers-in-charge of the prison facilities.

References

- [1] Curval LG, Franca AO, Fernandes HJ, Mendes RP, Carvalho L, et al. Prevalence of intestinal parasites among inmates of Midwest Brazil. *Plos one*. 2017; 12(19).
- [2] Mara D, Lane J, Scott B, Trouba D. Sanitation and health. *Plos Med* 2010 7(11): e1000363
- [3] Okolie N. Intestinal Parasites Distribution among Inmates of Owerri Prison. *International Journal on Parasitic diseases* 2008; 4(1).
- [4] Mamman AS, Gyar SD, Reuben CR. Malaria among Inmates of Jos prison, Plateau state, Nigeria. *International journal of microbiology and immunology research* 2014; 3 (3)1-4.

Comment [p80]: italics

Comment [p81]: of?

- [5] Ishaleku D, Mamman SA. Co-infection of Malaria and Helminthes Infection among Prison Inmates. *Journal of microbiology research and reviews*. 2014; 2 (1) 1-5.
- [6] Escola -verge L, Arando M, Vall M, Rovira R, et al. Outbreak of intestinal amoebiasis among men who have sex with men, Barcelona (spain). *Euro surveillance*. 2017; 22 (30).
- [7] KHRC. Improving Prison in Kenya. Nairobi: Foundation Press, 2002.
- [8] Omboto JO. The Challenges Facing Rehabilitation of Prisoners in Kenya and the Mitigation Strategies. *International journal of research in social sciences* 2013; 2 (2)39-40.
- [9] Krienitz L, Dadheech PK, Fastner and Kotut. The rise of potentially toxin producing cyanobacteria in Lake Naivasha, Great African Riftvalley, Kenya. *Science Direct*. 2013; 27: 42-51.
- [10] Nyarango RM, Peninah AA, Ephantus WK and Nyanchongi BO. The risk of pathogenic intestinal parasite infections in Kisii Municipality, Kenya. *Biomedical central Public Health* 2008; (8)237-243.
- [11] Mugenda OM and Mugenda AG. Research Methods: Quantitative and qualitative Approaches. Nairobi. African Center for Technology Studies, 2003.
- [12] Arora DR and Arora B. Medical Parasitology (2nd edition). New Delhi, India: CBS Publishers and Distributors, 2005; 76-77.
- [13] CDC. Laboratory identification of parasitic diseases of public health concern. 2013. www.cdc.gov/dpdx/diagnostic_procedures/stool/morphcomp.html.
- [14] Rop DC, Nyanchongi BO and Orucho VO. Risk factors associated with intestinal parasitic infections among inmates of Kisii prison, Kisii County, Kenya. *Bio Medical Central* 2016 (9); 384-385.
- [15] Nnaemeka AM, Iyioku UU, Yaro SA and Mohammed AH. Microbial Status of Prison inmates in Abakalki Prison, Ebonyi State, South - East Nigeria. *Global Journal of Medicine Researches and Studies* 2015; (1) 007-011.
- [16] Acuna-Soto R, Maguire JH and Wirth DF. Gender distribution in asymptomatic and invasive amoebiasis." *Am journal Gastroenerol* 2005; (5) 1277-83.
- [17] Colman S, Mangoro ZM and Isa L. Incidence of Intestinal and Urinary parasites among prison inmates. *Academia Journal of Microbiology Research* 2013; 1(1) 011-015.
- [18] Angal L, Mahumud R, Samin S, Yap N, Ngui R, Amir A, Ithoi I, Kamarulzaman A and Lim AY. Determining intestinal parasitic infections (IPIs) in inmates from Kajang prison, Selangor, Malaysia for improved prison management. *BMC infectious Diseases*. 2015; (15) 467- 498.
- [19] Belyhun Y, Medhin G, Amberbir A, Erko B, Hanlon et.al. Prevalence and risk factors for soil transmitted helminth infection in mothers and their infants in Butajira, Ethiopia. *BMC Public health* 2010; (10) 21.
- [20] Odieri MR and Opisa S. Geographical distribution of Schistosoma and STH among school children in informal settlement in Kisumu city, Western Kenya. *Parasitology*. 2011; 138(12) 1569-1577.
- [21] Nagata N, Shimbo T, Akiyama J, Nakashima R, Nishimura S and Yada T. Risk facors for intestinal invasive amoebiasis in Japan, 2003-2009. *Emerging infectious diseases*. 2012; 18 (5) 717-724.
- [22] Mamo H. Intestinal parasitic infections among prison inmates and tobacco farm workers in Shewa Robit, North-Central, Ethiopia. *Public Library of Science one*. 2014; 9(6): e99559.

Comment [p82]: Complete?

Comment [p83]:italize

- [23] Wanyama FW, Wanjala AM and Okoth OE. Predisposing factors contributing to the prevalence of intestinal parasitic infections (IPIs) among the HIV/AIDS patients in Bungoma County, Kenya.” *International journal of science research* 2014; 3: 627-631.
- [24] Pullan RL, Smith JL, Jasasaria R and Brooker SJ. Global numbers of infection and disease burden of soil transmitted helminth infections in 2010. *Parasites and Vectors* 2014; 7(1) 37

UNDER PEER REVIEW