

Review Article

Prevalence of Antibiotic Resistant Gram-negative Bacteria in the Street Vended Foods of Bangladesh: An Overview.

ABSTRACT

Background: Street vended foods contaminated with antibiotic resistant bacteria poses a great public health concern in developing countries like Bangladesh as foodborne diseases caused by such bacteria are difficult, sometimes impossible to treat. The aim of this review is to investigate the present scenario of the antibiotic resistance status of five selected Gram-negative bacteria isolated from a variety of street vended foods of Bangladesh.

Methodology: A methodical literature search was performed to identify relevant studies on Google Scholar, ScienceDirect and ResearchGate using suitable keywords arranged in different manners to produce a meaningful search string. Weighted mean resistance was calculated to evaluate the resistance status from 20 studies selected through the PRISMA procedure.

Results: *Escherichia coli*, *Klebsiella* spp., *Salmonella* spp., *Pseudomonas* spp., and *Vibrio* spp. showed relatively low resistance towards Imipenem and Gentamycin, whereas the antibiotics of penicillins and macrolids class listed in this review were mostly ineffective. Relatively high resistance was found against Tetracycline, Vancomycin, and Rifampicin.

Comment [DS1]: Correct macrolides

Conclusion: A considerable number of bacterial isolates present in the SVF are resistant to most of the commonly used antibiotics. Therefore, there is an urgent need for public awareness along with a robust national action plan to combat antibiotic resistance, before the situation worsens.

Keywords: Street vended foods, Bangladesh, Gram-negative bacteria, Contamination, Antibiotic resistance.

1. INTRODUCTION

The World Health Organization (WHO) has defined Street Vended Foods (SVF) as, “Foods and beverages prepared and/or sold by vendors in streets and other public places for immediate consumption or consumption at a later time without further processing or preparation”. Due to inexpensiveness, unique flavor, variation and other conveniences the demand of street vended foods is increasing among the urban people [1]. There are about 200,000 street food vendors in Dhaka city alone and the number is growing due to rapid urbanization [2,3]. Most vendors set up their stalls near bus terminals, busy roads, in front of schools, market places and railway stations in hopes of high potential customers. Vending carts are also placed by the side of municipal drains and sewage which attract insects and flies [2]. All these places have limited access to basic sanitary facilities such as running water, garbage disposal and clean toilets which puts the cleanliness of food preparing places and personal hygiene of vendors in question [4]. Inappropriate food handling, contaminated raw materials, cross contaminants from utensils and equipment are important sources of bacterial contamination in street foods. Contaminated foods act as a vehicle for transmitting foodborne diseases like diarrhea, cholera, typhoid fever and food poisoning [5]. In Bangladesh, about 30 million people suffer from food borne illnesses each year and approximately 2.2 million people including many children die of diarrheal diseases [6]. To battle against infections such as these, antibiotics are a blessing to human civilization that have saved millions of lives [7]. However, widespread availability and uncontrolled application of antibiotics in humans, food-producing animals, veterinary practices, and agriculture are causing a gradual uprising of antibiotic resistant bacteria [8,9]. The foodborne diseases become fatal when the pathogens causing them are also antibiotic resistant which elevate the duration of hospitalization, cost of treatment and the risk of mortality [10]. Antibiotic resistance is a property of bacteria that confers the capacity to inactivate or exclude antibiotics, or a mechanism that

Comment [DS2]: Correct for

Comment [DS3]: Delete the

Comment [DS4]: Correct markets

Comment [DS5]: Correct elevate

blocks the inhibitory or killing effects of antibiotics [11]. Resistance mechanisms may develop over months or years. Once accomplished, a single mechanism can allow a bacterium to become multi-drug resistant [12]. The 2017 WHO report shows a serious lack of new antibiotics under development to keep pace with the threat posed by the resistant bacteria. As a result, the world is gradually running out of antibiotics and approaching towards a time when people will dread common infections [13]. Developing countries like Bangladesh are more vulnerable to this issue for their underprivileged healthcare infrastructure [14]. Therefore, this review was conducted to reveal a comprehensive scenario of antibiotic resistance status of *Escherichia coli*, *Klebsiella* spp., *Salmonella* spp., *Pseudomonas* spp. and *Vibrio* spp. isolated from a wide range of street vended foods including drinking water, fruit juices, ready-to-eat (RTE) fruits and vegetables, fried items and a variety of other traditional items available in Bangladesh.


Comment [DS6]: Delete

2. METHODOLOGY

A methodical literature search was performed to identify studies associated with antibiotic resistance of Gram-negative bacteria found in various street vended foods of Bangladesh. Multitudinous searches were conducted on Google Scholar, ScienceDirect and ResearchGate to identify relevant studies using keywords such as microbial quality, street food, antibiotic resistant bacteria, Bangladesh, RTE foods etc. arranged in different manners to produce a meaningful search string. Studies were also added through manual searching from the INFS Library of the University of Dhaka. Figure 1 shows the PRISMA procedure through which 20 studies were selected to be used in this review article [15]. These studies were thoroughly revised by the authors and the extracted data was cross-checked multiple times.

The criteria of data extraction were publication year, study location, sources, antibiotic susceptibility testing method, antibiotic susceptibility testing standard, type and isolate number

of bacteria identified and resistance status of the bacteria identified, as depicted in table 1. The percentages of resistance were calculated by following the procedure used by Sabuj et al. 2018 for those studies where percentages were not mentioned but sufficient data was available [16]. The resistance status of each bacterial genre to different antibiotics is presented as the weighted mean resistance (percentage), combining all of the studies. The weighted mean resistance is calculated so that the resistance values corresponding to a larger number of isolates can contribute more towards the final mean. Data collection and analysis were conducted on Microsoft Word 2016 and Microsoft Excel 2016.



3. RESULT

Antibiotic resistance pattern of *Escherichia coli*, *Klebsiella* spp., *Pseudomonas* spp., *Vibrio* spp. and *Salmonella* spp. analyzed in this review article are extracted from the studies published between 2011-2019. Weighted mean was calculated only if antibiotic susceptibility test (AST) data of a significant number of isolates (ten or more) of the bacteria to a specific antibiotic were found after combining all the studies (table 2). The resistance status can be classified into: i) high resistance (>50% resistant isolates), ii) intermediate resistance (20-50% resistant isolates) and iii) low resistance (<20 % resistant isolates) [17].

All five bacteria showed high resistance to Amoxicillin. All bacteria except *Vibrio* spp. showed high resistance to Vancomycin and Erythromycin and low resistance to Gentamycin. Most of the bacteria showed high resistance to Ampicillin, Tetracycline, Azithromycin and Rifampicin. All the isolates of *Escherichia coli* and *Pseudomonas* spp. were resistant to Penicillin. *Escherichia coli*, *Klebsiella* and *Pseudomonas* spp. were completely susceptible towards Ceftriaxone, Chloramphenicol and Neomycin respectively. The bacteria were found to have a varying degree of resistance (high, low & intermediate) to quinolones (Nalidixic acid, Ciprofloxacin). *Escherichia coli* and *Pseudomonas* spp. showed relatively low resistance towards Ceftriaxone, a third generation Cephalosporin, compared to Cephalexin, a first generation Cephalosporin. However, *Salmonella* spp. showed relatively high resistance to Ceftriaxone. Low or intermediate resistance was observed towards the aminoglycosides class (Neomycin, Kanamycin, Streptomycin, and Gentamycin), carbapenems class (Imipenem), Chloramphenicol and Doxycycline.

Comment [DS8]: Correct Ciprofloxacin

Table 1: Characteristics of the Studies Included in the Review.

Class	Frequency of papers (total n=20)	Percentage of papers (%)	Reference
Publication Year			
2011-2013	3	15	[18–20]
2014-2016	3	15	[21–23]
2017-2019	14	70	[9,16,32–35,24–31]
Antibiotic Susceptibility Testing Methods			
Disk Diffusion Method	20	100	[9,16,26–35,18–25]
Antibiotic Susceptibility Testing Standards			
Clinical Laboratory Standards Institute (CLSI)/National Committee for Clinical and Laboratory Standards (NCCLS)	10	50	[9,16,20,26,28,29,31,32,34,35]
Not mentioned	10	50	[18,19,21–25,27,30,33]
Food Type			
Juices e.g. sugarcane, lemon, papaya, woodapple, mango, orange etc.	4	20	[21,29,34,35]
Ready to Eat Vegetables & Fruits e.g. cucumber, carrot, hog plum etc.	2	10	[18,31]
Drinking Water	2	10	[28,33]
Traditional and Fried Street Food e.g. chatpati, jhalmuri, fuchka, shingara, samosa, chittoi pitha etc.	7	35	[16,22–25,27,30]
Mixed Food Groups	5	25	[9,19,20,26,32]
Bacteria			
<i>Escherichia coli</i>	16	80	[16,18,27,30–33,35,19–26]
<i>Vibrio</i> spp.	5	25	[21,27,29,33,34]
<i>Salmonella</i> spp.	11	55	[9,16,33,18,19,21,22,26,28,30,32]
<i>Klebsiella</i> spp.	12	60	[18,19,33,34,21–23,25–27,29,30]
<i>Pseudomonas</i> spp.	5	25	[19,21,23,27,28]
Location			
Dhaka	11	55	[18,19,30,20–23,25,27–29]
Chittagong	5	25	[9,24,32–34]
Mymensingh	2	10	[16,31]
Gazipur	1	5	[35]
Dinajpur	1	5	[26]

Table 2: Antibiotic Resistance Status (percentage) of Five Selected Gram-negative Bacteria.

Class and Name of Antibiotics	Others	C (30µg)	25.00 (32)	0 (12)	-	16.67 (24)	30.55 (36)
		RD (5µg)	50 (12)	76.92 (13)	-	-	75.00 (20)
	Glycopeptides	VA (30µg)	95.62 (146)	96.19 (21)	26 (10)	79.17 (12)	73.45 (29)
	Carbapenems	IMP (30µg)	1.01 (134)	20 (10)	16.36 (11)	-	-
	Tetracyclines	DOX (30µg)	1.49 (67)	-	-	-	40 (25)
		TE (30µg)	54.85 (225)	-	-	83.74 (123)	60 (15)
	Cephalosporins	CEF (30µg)	0 (21)	-	-	49.05 (116)	41.5 (10)
		CL (30µg)	69.74 (95)	-	-	7.69 (13)	66.67 (15)
	Macrolides	AZM (15µg)	-	-	-	91.43 (105)	60 (15)
		ERY (10µg)	81.79 (56)	56.67 (18)	50 (10)	93.58 (109)	61.71 (44)
	Quinolones	NA (30µg)	22.08 (161)	18.46 (13)	-	71 (25)	60.43 (44)
		CIP (5µg)	9.76 (146)	20 (23)	57.15 (14)	30.40 (136)	27.33 (30)
	Aminoglycosides	CN (10µg)	7.08 (146)	3.48 (23)	40 (13)	18.16 (38)	3.67 (49)
		NEO (30µg)	27.08 (48)	-	-	36 (25)	0 (25)
		STR (10µg)	28.15 (60)	30 (20)	-	14.55 (22)	37.24 (29)
		KAN (30µg)	7.43 (142)	14.12 (17)	-	-	33.86 (44)
	Penicillins	P (10µg)	100 (131)	-	-	-	100 (20)
		AMX (10µg)	81.49 (121)	58.75 (16)	52.31 (13)	89.06 (123)	83.30 (44)
		AMP (10µg)	93.14 ^a (78) ^b	41.42 (14)	-	97.69 (108)	76.40 (25)

Name of Bacteria	<i>Escherichia coli</i>	<i>Klebsiella</i> spp.	<i>Vibrio</i> spp.	<i>Salmonella</i> spp.	<i>Pseudomonas</i> spp.
------------------	-------------------------	------------------------	--------------------	------------------------	-------------------------

Ampicillin = AMP; Amoxicillin = AMX; Penicillin = P; Kanamycin = KAN; Streptomycin = STR; Neomycin = NEO; Gentamicin = CN; Ciprofloxacin = CIP;

Nalidixic Acid = NA; Erythromycin = ERY; Azithromycin = AZM; Cephalexin = CL; Ceftriaxone = CEF; Tetracycline = TE; Doxycycline = DOX;

Imipenem = IMP; Vancomycin = VA; Rifampicin = RD; Chloramphenicol = C.

^a = Weighted mean resistance (%)

^b = values in parenthesis represent the number of isolates

Hyphen (-) represents number of isolates less than 10 (ten), hence not included.

4. DISCUSSION

Our study revealed high resistance of the selected bacteria to a considerable number of antibiotics especially against Ampicillin, Amoxicillin and Penicillin whereas relatively lower resistance was detected towards Ciprofloxacin, Gentamicin and Imipenem. Several studies carried out in different parts of the world showed a result which is in compliance with our present findings. A study at Tumkur, India found a significant number of Ampicillin resistant isolates of *Escherichia coli*, *Salmonella* spp. and *Vibrio* spp. from street vended foods [36]. Corresponding result was also observed in Delhi where *Escherichia coli*, *Salmonella* spp., *Vibrio* spp. isolated from different fruit juices exhibited a relatively higher resistance to Ampicillin [37]. Apart from the Indian subcontinent, the outcome of this review is upheld by different studies

Comment [DS9]: Correct that complies

Comment [DS10]: Add the

Comment [DS11]: Correct were

conducted in Africa as well. A study held in Akure Metropolis, Nigeria revealed the high resistance capability of *Escherichia coli*, *Salmonella* spp., *Pseudomonas* spp. and *Vibrio* spp. against Amoxicillin, all of these bacteria were also highly resistant to Tetracycline except for the latter one [38]. Another study conducted in Ethiopia revealed quite similar result where isolates of *Escherichia coli* found in street foods were comparatively less resistant to Ceftriaxone, Ciprofloxacin, Gentamycin, Kanamycin compared to Ampicillin [39]. Tadesse et al. (2019) reported *Salmonella* spp. isolated from street foods of Eastern Ethiopia were detected to be predominant over Ampicillin, Amoxicillin and Tetracycline [40].

Comment [DS12]: Add a

In developing countries like Bangladesh, people can obtain most of the antibiotics over the counter at a cheaper price without any medical prescription [7]. Overuse and misuse of these antibiotics is linked to the emergence of resistant bacteria [41]. For example, Amoxicillin is frequently used among low income people as it is relatively cheaper which might be a reason why bacteria are highly resistant to it [42]. Bacteria also get exposed to antibiotics from the environment as most of the antibiotics used in human and food producing animals ultimately make their way to the environment [9]. Moreover, most of the pharmaceutical companies do not treat their wastes properly before disposing in the ponds, rivers and other water systems [43]. Repeated low dose exposure to antibiotics triggers bacteria to start developing resistance mechanisms and hence its presence in the environment plays a crucial role in the emergence of resistant bacteria [9,43].

Comment [DS13]: Correct are

Comment [DS14]: Delete

Comment [DS15]: Add of them in

Resistant bacteria enter the food chain through animal based foods [44]. *Escherichia coli*, *Salmonella* spp. isolated from some commonly used raw animal food like meat, chicken egg and milk were found resistant to most of the antibiotics [8,45,46]. Therefore, it can be a reason for the presence of antibiotic resistant bacteria in street foods. Also, cross contamination can occur

through water sources as street vendors frequently use municipal water for preparing fruit juices and for washing purposes [37]. *Escherichia coli* isolated from tap water collected from Dhaka, Jamalpur, Tangail, Netrokona, Kishoreganj showed resistance against Tetracycline, Erythromycin, Amoxicillin, Streptomycin and some other antibiotics [47,48]. Unhygienic condition of the vendors and the food preparing places are also accountable for cross contamination. Hassan et al. reported *Escherichia coli* and *Klebsiella* spp. isolated from hand-rinsed water of street vendors were found resistant against Amoxicillin, Ciprofloxacin, Gentamycin and Azithromycin [42].

Comment [DS16]: Add the

Given the fact that most of the studies included in this review were concentrated around Dhaka, there is a likelihood of selection bias and this review might not portray the complete scenario of Bangladesh. Moreover, due to the insufficiency of the studies available, authors were compelled to include even the studies that performed AST on a single isolate. However, weighted mean was calculated only if the isolate number was 10 or more, combining all the studies. While extracting the percentage values of resistance from a graph, some data had to be excluded for not being sufficiently clear. That being said, the authors confronted this issue only once. The authors combined data acquired from all the selected studies even though half the studies did not mention their testing standard. However, since the other half followed the CLSI guidelines and all the studies employed disc diffusion method, the extent of variation should be negligible.

Comment [DS17]: Add the

Comment [DS18]: Add the

5. CONCLUSION

More studies should be carried out in different parts of Bangladesh as data was only available for five districts out of the sixty-four districts. Nevertheless, despite the data gaps and other limitations, it is apparent that the prevalence of antibiotic resistant bacteria is considerably high in the street-vended foods of Bangladesh. Therefore, improved regulatory frameworks need to

be established and the surveillance system should be strengthened to control the misuse of antibiotics. Alongside, public awareness programs should be held. The street food vending needs to be taken under licensing system and only the qualified ones, having been trained on basic food safety and hygiene, should be provided with a license. If adequate actions to combat antibiotic resistance is not considered immediately, it will have a devastating impact on the public health in the coming decades.

REFERENCES

1. World Health Organization. Essential Safety Requirements for Street-Vended Foods. World Heal Organ. 1996;96.7:36.
2. Hassan MZ, Islam MS, Salauddin M, Zafor AHA, Scott ML, Alam S. Detection of enteric bacteria in the popular street food chotpoti in Dhaka, Bangladesh. *Asian J Med Biol Res.* 2016;2(4):596–602.
3. Fatema K, Rahman S, Ahmed S, Datta S. Microbiological Quality Assessment of Handmade Juice in Street of The Dhaka City. *Hendun Res Access.* 2016;1(1):1–7.
4. Feglo P, Sakyi K. Bacterial contamination of street vending food in Kumasi, Ghana. *J Med Biomed Sci.* 2012;1(1):1–8.
5. Rane S. Street Vended Food in Developing World: Hazard Analyses. *Indian J Microbiol.* 2011;51(1):100–6.
6. Hossain MB, Mahbub NDB, Chowdhury MMK, Rahaman MM. Molecular Characterization of Enterobacter and Escherichia coli Pathotypes Prevalent in the Popular Street Foods of Dhaka City and their Multidrug Resistance. *Bangladesh J Microbiol.* 2017;34(2):67–72.
7. Zaman S Bin, Hussain MA, Nye R, Mehta V, Mamun KT, Hossain N. A Review on Antibiotic Resistance: Alarm Bells are Ringing. *Cureus.* 2017 Jun 28;9(6):1–9.
8. Ahmed MM, Rahman MM, Mahbub KR, Wahiduzzaman M. Characterization of Antibiotic Resistant Salmonella spp Isolated from Chicken Eggs of Dhaka City. *J Sci Res.* 2011 Dec 20;3(1):191.
9. Hassan MM, Begum S, Faruq A Al, Alam M, Mahmud T, Islam A. Multidrug Resistant Salmonella Isolated from Street Foods in Chittagong, Bangladesh. *Microbiol Res J Int.* 2018;26(6):1–8.
10. Khan SA, Imtiaz MA, Sayeed MA, Shaikat AH, Hassan MM. Antimicrobial resistance pattern in domestic animal - Wildlife - environmental niche via the food chain to humans

with a Bangladesh perspective; A systematic review. *BMC Vet Res.* 2020;16(1):1–13.

11. Harrison PF, Lederberg J, Institute of Medicine (U.S.). Forum on Emerging Infections., Institute of Medicine (U.S.). Division of Health Sciences Policy. Antimicrobial resistance : issues and options : workshop report. Washington, D.C.: National Academy Press; 1998. 115 p.
12. Zhang R, Eggleston K, Rotimi V, Zeckhauser RJ. Antibiotic resistance as a global threat: Evidence from China, Kuwait and the United States. *Global Health.* 2006;2(6):1–14.
13. WHO. ANTIBACTERIAL AGENTS IN CLINICAL DEVELOPMENT. *World Heal Organ.* 2017;31–5.
14. Byarugaba DK. Antimicrobial resistance in developing countries and responsible risk factors. *Int J Antimicrob Agents.* 2004;24(2):105–10.
15. Moher D, Liberati A, Tetzlaff J, Altman DG, Altman D, Antes G, Atkins D, Barbour V, Barrowman N, Berlin JA, Clark J, Clarke M, Cook D, D'Amico R, Deeks JJ, Devereaux PJ, Dickersin K, Egger M, Ernst E, Gøtzsche PC, Grimshaw J, Guyatt G, Higgins J, Ioannidis JPA, Kleijnen J, Lang T, Magrini N, McNamee D, Moja L, Mulrow C, Napoli M, Oxman A, Pham B, Rennie D, Sampson M, Schulz KF, Shekelle PG, Tovey D, Tugwell P. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med.* 2009;6(7).
16. Sabuj AAM, Haque ZF, Barua N, Islam MA, Saha S. Assessment of Bacteriological Quality of Street Vended Fast Foods and Their Antimicrobial Resistance. *Int J Curr Microbiol Appl Sci.* 2018 Nov 20;7(11):3049–59.
17. Jahan M, Rahman M, Rahman M, Sikder T, Uson-Lopez RA, Selim ASM, Saito T, Kurasaki M. Microbiological safety of street-vended foods in Bangladesh. Vol. 13, *Journal of Consumer Protection and Food Safety.* Birkhauser Verlag AG; 2018. p. 257–69.
18. Ali M, Khan M, Saha ML. Antibiotic resistant patterns of bacterial isolates from ready-to-eat (RTE) street vended fresh vegetables and fruits in Dhaka City. *Bangladesh J Sci Res.* 2011;24(2):127–34.
19. Tabashsum Z, Khalil I, Nazimuddin MD, Mollah AKM, Inatsu Y, Bari ML. Prevalence of Foodborne Pathogens and Spoilage Microorganisms and Their Drug Resistant Status in Different Street Foods of Dhaka city. *Agric Food Anal Bacteriol* [Internet]. 2013;3(4):281–92. Available from: www.afabjournal.com
20. Sarker N. Use of Multiplex PCR Assay for Detection of Diarrheagenic Escherichia Coli in Street Vended Food Items. *Am J Life Sci.* 2013;1(6):267.
21. Lucky NA, Nur IT, Ahmed T. Microbiological quality assessment for drug resistant pathogenic microorganisms from the fresh vended fruit juices. *Stamford J Microbiol.* 2016;6(1):7–10.
22. Khan FI, Saha ML. Bacteria Laden street food (Chatpati) and their multiple antibiotic resistance index. *Bangladesh J Bot.* 2015;44(4):599–604.

23. Tabassum A, Saha ML, Islam MN. Prevalence of multi-drug resistant bacteria in selected street food and water samples. *Bangladesh J Bot* [Internet]. 2015;44(4):621–7. Available from: <http://bioinformatics.psb.ugent.be/cgi-bin/rRNA/>
24. Akter, H., Shaikat, A. H., Imtiaz, M. A., Islam, A., Khan, S. A. and Hassan MM. Prevalence of multidrug resistance pattern of *Escherichia coli* isolated from street food.pdf. Vol. 7, *Bangladesh Journal of Veterinary and Animal Sciences*. 2019. p. 69–74.
25. Lal Saha M, Dilara Akter M, Khan T, Ansari A, Nurul Islam M. BACTERIAL LOAD AND MULTI-DRUG RESISTANCE PATTERNS OF SOME READY-TO-EAT STREET FOODS OF DHAKA CITY [Internet]. Vol. 27, *Dhaka University Journal of Biological Sciences*. 2018. Available from: <http://blast.ncbi.nlm.nih.gov/>
26. Khalif MA, Hossain MK, Rumi NA, Rahman MS, Hosen MA. Identification and antibiogram study of bacteria isolated from different street food. *Asian J Med Biol Res*. 2018;4(3):279–87.
27. Moumita Dey, Mansura Mokbul, Ila Ismail SRA. Identification of Antibiotic Resistant Gram-Negative Bacteria in a Popular Street-Food Item (Chatpati) in Dhaka University Campus, Bangladesh. *Front Environ Microbiol*. 2018;4(2):75.
28. Ahmed S, Mahbub KR, Ahmed MM, Rahman M, Hoque MM. Microbiological Quality of Street Vended Drinking Water in Dhaka City and Screening for Antibiotics Resistance of Isolated *Salmonella spp* and *Pseudomonas spp*. *J Sci Res*. 2014 Apr 24;6(2):359–71.
29. Afroz K, Shammi T, Kabir MS. Study of Microbiological Quality of Fresh Juices Vended in Dhaka City. *Bangladesh J Microbiol*. 2019;36(1):49–53.
30. Happy AH, Alam MG, Mahmud S, Imran MAS, Rony MH, Azim MAA, Islam MM, Sarker MKD, Akter P, Mondol GC, Hossain T, Rahman MM, Islam MM, Roy A, Das S, Ahmed MR, Uddin ME. Isolation, Identification and Characterization of Gram Negative Bacteria from Popular Street Food (Chotpoti) at Savar Area, Dhaka, Bangladesh. *OALib*. 2018;05(11):1–11.
31. Siddiqui A, Nasrin S, Moonmoon M, Islam M, Khatun M. Bacterial assessment of street-vended hog plum (*Spondias mombin*) and its public health importance. *Bangladesh Vet*. 2015;32(1):19–26.
32. Paul P, Ahad A, Meher MM, Asgar A, Mostafa AKM. Antibiotic Susceptibility Profile of Microbes Isolated from Street Food Vended in Chittagong , Bangladesh. *Int Invent Sci J*. 2018;02(06):208–12.
33. Maesha AN, Marzan LW, Akter Y. Identification and Antibiogram Studies of Pathogenic Bacteria Isolated From Drinking Water (Unpackaged) In Chittagong City, Bangladesh. *J Bangladesh Acad Sci*. 2018;42(2):137–53.
34. Sultana F, Marzan L, Mina S. Microbiological quality assessment of locally vended and commercially packed fruit juices in Chittagong city of Bangladesh. *J Bio-Science*. 2019 Dec 26;27:43–58.
35. Jahan M, Rahaman Sumon SMM, Md. Selim AS, Rahman MM. Occurrence of antibiotic resistant *Escherichia coli* and *Staphylococcus aureus* in streetvended beverages accessible

in an industrial zone of Bangladesh. *J Food Saf Hyg* [Internet]. 2019;5(4):220–9. Available from: <http://jfsht.tums.ac.ir>

36. Kumar M DS, Krishnamurthy DV, Nagaraj DE. Microbial profile of street food from different locations at Tumkur, India. *Trop J Pathol Microbiol* [Internet]. 2017;3(2):84–9. Available from: www.pathologyreview.in
37. Sharma N, Singh K, Toor D, Pai SS, Chakraborty R, Khan KM. Antibiotic resistance in microbes from street fruit drinks and hygiene behavior of the vendors in Delhi, India. *Int J Environ Res Public Health*. 2020 Jul 1;17(13):1–12.
38. Ogidi OC, Oyetayo VO, Akinyele BJ. Microbial Quality and Antibiotic Sensitivity Pattern of Isolated Microorganisms from Street Foods Sold in Akure Metropolis, Nigeria. Vol. 9, *Jordan Journal of Biological Sciences*. 2016.
39. Eromo T, Tassew H, Daka D, Kibru G. Bacteriological Quality of Street Foods and Antimicrobial Resistance of Isolates in Hawassa, Ethiopia. *Ethiop J Health Sci*. 2016 Nov 1;26(6):533–42.
40. Tadesse G, Mitiku H, Teklemariam Z, Marami D. Salmonella and Shigella Among Asymptomatic Street Food Vendors in the Dire Dawa city, Eastern Ethiopia: Prevalence, Antimicrobial Susceptibility Pattern, and Associated Factors. *Environ Health Insights*. 2019;13:1–8.
41. Komolafe O. Antibiotic resistance in bacteria - an emerging public health problem. Vol. 15, *Malawi Medical Journal*. 2003. p. 63–7.
42. Hassan MM, Chakrabarty RP, Siddique MA, Rahaman MM. Prevalence of antibiotic resistant enteric bacteria in the hands of street food vendors in Dhaka city. *Bangladesh J Microbiol*. 2017 Dec 31;34(1):33–8.
43. Nasreen M, Sarker A, Malek MA, Ansaruzzaman M, Rahman M. Prevalence and Resistance Pattern of *Pseudomonas aeruginosa* Isolated from Surface Water. *Adv Microbiol*. 2015;05(01):74–81.
44. Abdullahel M, Islam B, Hasan AKMM, Kabir E, Shekhar HU. Pattern of Drug Resistance of Pathogenic Microbes in the Street Foods of Dhaka City, Bangladesh. *Bioresearch Commun*. 2015;1(1):48–52.
45. M. A. Rahman, A. K. M. A. Rahman, M. A. Islam MMA. ANTIMICROBIAL RESISTANCE OF *ESCHERICHIA COLI* ISOLATED FROM MILK, BEEF AND CHICKEN MEAT IN BANGLADESH. *Bangl J Vet Med*. 2017;15(2):141–6.
46. Islam M, Sabrin MS, Kabir MH Bin, Karim SJI, Sikder T. Prevalence of multidrug resistant (MDR) food-borne pathogens in raw chicken meat in Dhaka city, Bangladesh: an increasing food safety concern. *Asian-Australasian Journal of Prevalence of multidrug resistant (MDR) food-borne pathogens in raw chicken m*. *Asian-Australasian J Biosci Biotechnol* [Internet]. 2018;3(1):17–27. Available from: www.ebupress.com/journal/aajbbArticle
47. Davison HC, Woolhouse MEJ, Low JC. What is antibiotic resistance and how can we measure it? *Trends Microbiol*. 2000;8(12):554–9.

48. Champa H, Kabir SL. Microbial analysis of tap water collected from selected upazillas of Jamalpur, Tangail, Kishoreganj and Netrokona districts of Bangladesh. Asian J Med Biol Res. 2018 Sep 30;4(2):193–200.

UNDER PEER REVIEW